

Gastric Microbiota and Helicobacter Pylori in Indonesian Population

by Muhammad Miftahussurur

Submission date: 19-May-2020 01:14PM (UTC+0800)

Submission ID: 1327544665

File name: Gastric_Microbiota.pdf (1.29M)

Word count: 6104

Character count: 34447



Gastric microbiota and *Helicobacter pylori* in Indonesian population

Muhammad Miftahussurur^{1,2} | Langgeng Agung Waskito^{2,3} | Hashem B El-Serag⁴ | Nadim J. Ajami⁴ | Iswan Abbas Nusi¹ | Ari Fahrial Syam⁵ | Takashi Matsumoto³ | Yudith Annisa Ayu Rezkitha⁶ | Dalla Doohan^{2,3} | Kartika Afrida Fauzia^{2,3} | Ummi Maimunah¹ | Titong Sugihartono¹ | Tomohisa Uchida⁷ | Yoshio Yamaoka^{1,3,4,8}

¹Division of Gastroentero-Hepatology, Department of Internal Medicine, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

²Institute of Tropical Diseases, Universitas Airlangga, Surabaya, Indonesia

³Department of Environmental and Preventive Medicine, Oita University, Faculty of Medicine, Yufu, Japan

⁴Gastroenterology and Hepatology Section, Department of Medicine, Baylor College of Medicine, Houston, TX, USA

⁵Division of Gastroenterology, Department of Internal Medicine, Faculty of Medicine, University of Indonesia, Jakarta, Indonesia

⁶Faculty of Medicine, University of Muhammadiyah Surabaya, Surabaya, Indonesia

⁷Department of Molecular Pathology, Faculty of Medicine, Oita University, Oita, Japan

⁸Global Oita Medical Advanced Research Center for Health, Yufu, Japan

Correspondence

Yoshio Yamaoka, Department of Environmental and Preventive Medicine, Institute of Tropical Diseases, Faculty of Medicine, 1-1 Idaigaoka, Hasama-machi, Yufu-City, Oita 879-5593, Japan.
Email: yyamaoka@oita-u.ac.jp

Muhammad Miftahussurur, Gastroenterology Division, Department of Internal Medicine, Faculty of Medicine-Dr. Soetomo Teaching Hospital, Universitas Airlangga, Surabaya, Jalan Mayjend Prof. Dr. Moestopo No. 6-8 Surabaya 60131, Indonesia.
Email: muhammad-m@fk.unair.ac.id

Funding information

This study was funded by grants from the National Institutes of Health (DK62813) and Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT) of Japan (15H02657 and 221S0002, 16H05191, 16H06279, 18KK0266 and 18K03473) (YY). It was also supported by the Japanese Society for the Promotion of Science Institutional Program for Young Researcher Overseas Visits (YY) and Strategic Funds for the Promotion of Science and Technology from Japan Science and Technology Agency (JST) (YY). It was also supported by Riset Mandat Grant from Universitas Airlangga (340/

Abstract

Background: The profile of gastric mucosal microbiota has not yet been described in the Indonesian population where the prevalence of *Helicobacter pylori* is low.

Methods: This is a cross-sectional study analyzing 16S rRNA of 137 gastric biopsy specimens. We analyzed the association between gastric microbiota, *H. pylori* infection, and gastric mucosal damage.

Result: Among 137 analyzed samples, 27 were *H. pylori*-positive and 110 were *H. pylori*-negative based on culture, histology, and 16S rRNA gene analysis. Significantly lower α -diversity parameters, including Pielou's index, was observed in *H. pylori*-infected individuals compared with noninfected individuals (all $P < .001$). Among *H. pylori*-negative individuals, the permutational analysis of variance of Bray-Curtis dissimilarity distances showed a significant association with different ethnicities, suggesting some ethnic groups had specific microbiota profiles based on the presence of different operational taxonomic units. The linear discriminant analysis effect size (LEfSe) of the *H. pylori*-negative group showed significant associations between the presence of *Micrococcus luteus* and *Sphingomonas yabuuchiae* with Timor and Papuan ethnicities, respectively. The presence of *Bulleidia* sp and *Atopobium* sp was associated with the Javanese ethnicity. We observed lower α -diversity scores in individuals with gastric mucosal damage and profiles with high abundances of *Paludibacter* sp and *Dialister* sp based on LEfSe analysis.

Muhammad Miftahussurur and Langgeng Agung Waskito contributed equally.

UN3.14/LT/2019). All authors have read and approved the final version of the manuscript.

Conclusion: Our findings suggest the presence of *H. pylori* is more correlated with a distinct microbiome profile than ethnic precedence.

KEYWORDS

Helicobacter pylori, Indonesia, microbiome

1 | INTRODUCTION

The gastric mucosal environment was thought to be sterile due to the presence of gastric ²⁴ until the discovery of *Helicobacter pylori*,¹ a causal agent for gastritis, peptic ulcer diseases, and gastric cancer.² The use of 16S rRNA next-generation sequencing revealed trillions of gastric mucosal microbiota, most of which cannot be cultivated.³ Further, there are possibly different microbial communities between stomachs that are healthy and those with diseases (eg, gastric cancer).^{4,5} The presence of *H. pylori* greatly influences microbial dysbiosis^{6,7} and the risk of gastric cancer.⁶

The composition of the gastric microbiota is influenced by age,⁸ diet,⁹ and ethnicity.¹⁰ Ethnicity has been proposed to influence the gastric microbiome through diet patterns and possibly genetic factors.⁴ A study conducted in China reported a significant difference in the richness and diversity of the gut microbiome among three ethnic groups (Yao, Zhuang, and Han),¹¹ and similar findings were reported in Europe and the United State, indicating that ethnicity contributed to the intergroup dissimilarity of gut microbiota composition.^{12,13} However, most of these studies identified gut microbiota using stools samples.¹³ Few studies have evaluated the gastric microbiome using gastric mucosa samples, which provide a more accurate representation of the resident gastric microbiome compared with the transient one.^{14,15}

Indonesia is a country with several hundred ethnicities and a low prevalence of *H. pylori* infection.¹⁶ The Javanese ethnicity is the largest in Indonesia, contributing ~40% of the population, most of whom live on Java Island. Several other ethnic groups live on other islands, such as the Kupang, Timorese, and Papuan ethnicities in eastern areas of Indonesia and the Nias and Batak tribes in the western tip of Indonesia. These ethnic groups have distinctive cultures and diets that could influence the composition of gut microbiota. The lower prevalence of *H. pylori* allows us to examine the role of other organisms in the pathogenesis of gastric disorders. Therefore, we investigated the composition of gastric mucosal microbiota in an Indonesian population and analyzed the associations between microbiome composition, ethnicity, and gastric disease status.

2 | METHODS

2.1 | Patients and samples

From January 2014 to August 2016, we collected 784 gastric mucosal specimens from 784 individuals in 11 cities across Indonesia³³ including sites located on the five largest islands (Figure S1). We

excluded patients with a history of *H. pylori* eradication therapy, partial or total gastrectomy, or a contraindication to upper endoscopy with biopsies. Gastric mucosal specimens for microbiome analysis were obtained from the lesser curvature of the antrum, ~3 cm from the pyloric ring. They were preserved in transport medium and homogenized in 500 μ L of phosphate-buffered saline for *H. pylori* culture. The remaining homogenized specimens were stored⁵ into -80°C and used for DNA extraction. We⁵ took two additional biopsy specimens from the antrum and corpus for histologic examination. All specimens were obtained using Radial Jaw 4 forceps (Boston Scientific).

We excluded 97 patients who did not have a complete histology result. The remaining 687 samples comprised patients who were Javanese (203 patients), Batakese (113 patients), Bugis (87 patients), Chinese Indonesian (100 patients), Balinese (60 patients), Dayak (47 patients), Papuan (47 patients), and Timor (30 patients). We randomly selected 192 samples with the selection algorithm weighted for each ethnic group and the location regardless of *H. pylori* status. These samples included patients who were Javanese (48 samples); Batak (30 samples); Bugis (35 samples); Chinese Indonesian (17 samples); Balinese (19 samples); Dayak (14 samples); Papuan (13 samples); and Timor (16 samples).

2.2 | DNA extraction and PCR amplification of 16S rRNA sequences

³⁰ DNA was extracted using the DNeasy Blood & Tissue Kit (QIAGEN) and was concentrated using a DNA Clean & Concentrator (Zymo Research). Extracted DNA was stored at -20°C .

We followed the manufacturer's protocol for preparing the ²⁰ rRNA gene DNA library (Illumina Inc, San Diego, CA). The V3-V4 region of the bacterial 16S rRNA gene was amplified using the universal primers 341F (5'-TCGTCGGCAGCGTCAGATGTATAAGAGACAGCCTACGGGNGGCWGCAG-3') and 805R (5'-GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGGACTACHVGGGTATCTAATCC3-3').¹⁷ PCR amplification was performed using KAPA³ HiFi HotStart Ready Mix (KAPA Biosystems Inc) with denaturing at 95°C for 3 minutes; 35 cycles of amplification at 95°C for 30 seconds, 55°C for 30 seconds, and ¹¹ 72°C for 30 seconds; and final elongation at 72°C for 5 minutes. The Nextera XT Index kit (Illumina, ²⁹ was added, and samples underwent an additional eight cycles of PCR. Amplicons were purified using Agencourt AMPure XP magnetic beads (Beckman Coulter). We validated the DNA library using a bioanalyzer MCE-202 MultiNA system (Shimadzu) and QuantiFluor dsDNA system (Promega Corporation). Finally,

the pooled 5 pM DNA library was denatured with 0.2 N NaOH and mixed with PhiX Control v3 (Illumina Inc) to 15% of the final concentration as described in Illumina's procedure. Paired-end sequencing was performed using the next-generation sequencer MiSeq platform (Illumina Inc) with the MiSeq Reagent Kit version 3 (2 × 300 bp Paired-End Reads; Illumina Inc).

2.3 | Sequence data analysis

We removed index and adapter sequences and low-quality reads using Trimmomatic.¹⁸ The read-retention parameter was having a Phred quality score ≥ 20 . The high-quality reads underwent close reference operational taxonomic unit (OTU) picking using vsearch.¹⁹ The reference database was Greengenes version 13.5.²⁰ Using a 99% identity parameters, we generated an OTU table and the representative sequences of each OTUs. These sequences were aligned using MAFFT,^{21,22} and a phylogenetic tree was constructed using FastTree.²³ We imported the OTU table, representative sequences, and phylogenetic tree as artifacts to the QIIME 2 analysis platform (<https://qiime2.org>). The α -diversity analysis, including species richness, Pielou's evenness index,²⁴ Faith's phylogenetic diversity index,²⁵ and Shannon's diversity index,²⁶ was analyzed using QIIME2. The β -diversity analysis using Bray-Curtis dissimilarity matrix²⁷ was also performed using QIIME 2. The Bray-Curtis dissimilarity matrix was used to draw principal coordinate analysis figures in R. The permutational analysis of variance (PERMANOVA) with Bonferroni correction was calculated using the Adonis function from the vegan package in R.²⁸

We performed linear discriminant analysis effect size (LEfSe) to identify OTUs that are associated with that are likely to explain differences between groups, specifically those based on ethnicity and disease status.²⁹ In addition, we also performed analysis of variance (ANOVA) to identify differences in relative abundance of specific OTU between ethnic groups and diseases. The LEfSe and ANOVA analysis were performed using the Calypso web-based analysis platform version 8.84,³⁰ available at (<http://cgenome.net:8080/calypso-8.84/faces/uploadFiles.xhtml>).

2.4 | Determination of *H. pylori* infection

We determined the *H. pylori* infection status based on *H. pylori* culture and histologic examination.

2.4.1 | Culture

Antral biopsy specimens were homogenized, inoculated on an *H. pylori* -selective agar plate,⁴⁵ and incubated up to 10 days in microaerophilic condition (10% O₂, 5% CO₂, and 85% N₂) at 37°C.²⁸ Subsequently, colonies were sub-cultured in antibiotic-free Mueller-Hinton II agar medium (Becton Dickinson) supplemented with 10%

Horse Blood and incubate the same microaerophilic conditions. *H. pylori* stock was stored in Brucella Broth (Difco) with 10% dimethyl sulfoxide and 10% horse serum at -80°C.

2.4.2 | Histologic analysis

All biopsy material for histologic testing was fixed in 10% buffered formalin and embedded in paraffin. Serial sections were stained with hematoxylin and eosin and May-Giemsa stain. The degree of neutrophil activity, inflammation, atrophy, intestinal metaplasia, and *H. pylori* bacterial density was classified into four grades based on the updated Sydney system (0, normal; 1, mild; 2, moderate; and 3, marked).³¹ Samples with bacterial loads of grade 1 or higher were considered positive for *H. pylori*. We defined gastric mucosal damage, that is, an observation of both neutrophil or monocyte infiltration, as gastritis. Precancerous lesions were defined as an observation of either atrophic gastritis or intestinal metaplasia. To increase the accuracy of *H. pylori* detection, we performed immunohistochemistry of all samples, as described previously.³² The experienced pathologist (TU) who evaluated the specimens in this study also performed studies for Myanmar, Vietnam, Bhutan, the Dominican Republic, and Indonesia.³³⁻³⁸

3 | RESULTS

Among the gastric mucosal samples from 192 patients, 14 samples produced suboptimal 16S rRNA gene libraries for sequencing. Therefore, 178 samples were sequenced for microbiome profiling targeted at the V3-V4 regions of the 16S rRNA gene. We found 16.3% (29/178) of specimens were *H. pylori* -positive based on histology and confirmed by immunohistochemistry or culture.

We performed OTU selection and identified 68 classes, 135 orders, and 253 families of bacteria. We excluded samples that had conflicting *H. pylori* status results between histology/culture and 16S rRNA. Among excluded samples, 39 were designated *H. pylori*-negative based on histology/culture but had relative abundances of *H. pylori* greater than 2% (range 2.02%-83.12%) and 2 were *H. pylori*-positive based on histology/culture but had low relative abundance of *H. pylori* (0.67% and 0.46%).

A total of 137 samples were included in the final analysis; 27 were *H. pylori*-positive and 110 were *H. pylori*-negative based on histologic examination, culture, and 16S rRNA gene sequencing. The mean age of this group was 48.34 ± 14.8 years, and 79 samples were from male patients and 58 (42.3%) from female patients. *H. pylori*-positive patients were significantly older than *H. pylori*-negative ones (52.2 vs 46.5 years, respectively; $P = .032$). There were 22 (16.7%) samples from patients with gastritis; 20 (90.9%) were *H. pylori*-negative, and 2 (9.1%) were *H. pylori*-positive. Further, 56 (40.9%) samples were from patients with atrophic gastritis or intestinal metaplasia; 31 (55.4%) were *H. pylori*-negative, and 25 (44.6%) were *H. pylori*-positive.

3.1 | Gastric microbiota diversity and *H. pylori* positivity

We observed significantly lower species richness, Pielou's evenness, Shannon's diversity index, and Faith's phylogenetic diversity in *H. pylori*-positive samples ($n = 22$) compared with *H. pylori*-negative samples ($n = 110$) (all $P < .001$, Figure 1A). The *H. pylori*-positive group had a lower number of observed OTUs, suggesting lower diversity in *H. pylori*-positive samples compared with *H. pylori*-negative samples. The *H. pylori*-positive group also had a lower Pielou's index score for evenness, suggesting dominance by one or more OTUs. We observed higher OTU similarity in the *H. pylori*-positive than the *H. pylori*-negative group as indicated by Faith's phylogenetic diversity. Shannon's diversity index also indicated that the *H. pylori*-positive group was less diverse than the *H. pylori*-negative one. There was a significant negative correlation between Shannon's diversity index and age ($r = -0.31$, $P < .001$, Figure 1B).

Similarly, the *H. pylori*-negative group showed a significant negative correlation between Shannon's diversity index and age ($r = -0.25$, $P = .0089$; Figure S2).

Next, we evaluated differences in β -diversity between *H. pylori*-negative and *H. pylori*-positive groups using the Bray-Curtis dissimilarity matrix. *H. pylori*-positive and *H. pylori*-negative groups exhibited different clustering positions (Figure 2). PERMANOVA analysis of the Bray-Curtis dissimilarity matrix supports our observation of distinct clustering of the *H. pylori*-positive compared with the *H. pylori*-negative groups, highlighting their differences in microbial composition ($R^2 = 0.248$, $P < .001$). The relative abundance of Family *Helicobacteraceae* (74.6%) predominated in *H. pylori*-positive samples (Figure 3), whereas the Families *Streptococcaceae* (25.9%), *Prevotellaceae* (13.5%), and *Veillonellaceae* (10.8%) predominated in the *H. pylori*-negative group. On the species level, *H. pylori* had 76.2% relative abundance in the *H. pylori*-positive group. In the *H. pylori*-negative group, *Prevotella melaninogenica* had 23.5% relative





FIGURE 1 Association of α -diversity with the *Helicobacter pylori* infection and age. A. *H. pylori* infection significantly reduced α -diversity parameters and Pielou's phylogenetic diversity. B. Shannon's diversity index was significantly reduced as age increased

abundance, *Rothia mucilaginosa* had 19.5%, *Veillonella dispar* had 18.6%, and *Haemophilus parainfluenzae* had 7.2%.

3.2 | Microbiome diversity and ethnic groups


There were significant differences in species richness, evenness, Faith's phylogenetic diversity, and Shannon's diversity indexes (all



18
17 **FIGURE 2** The β -diversity of 137 analyzed samples between *Helicobacter pylori*-positive and *H. pylori*-negative subjects. *H. pylori*-positive and *H. pylori*-negative samples were clustered separately. PERMANOVA analysis showed a significant difference in β -diversity between *H. pylori*-positive and *H. pylori*-negative groups ($P < .001$)

$P < .001$) among the different ethnic groups (Figure 4A). Pairwise analysis showed the Timor ($n = 9$), Balinese ($n = 7$), and Papuan ($n = 9$) ethnic groups had significantly lower richness, evenness, Faith's phylogenetic diversity index, and Shannon's diversity index than Javanese ($n = 30$), Dayak ($n = 12$), Chinese ($n = 11$), Bugis ($n = 33$), and Batak ($n = 26$; Figure S3). Interestingly, in the *H. pylori*-negative group, there were no significant differences in richness, evenness, Faith's phylogenetic diversity index, or Shannon's diversity index ($P = .087$, $P = .065$, $P = .222$, and $P = .078$, respectively) among ethnic groups (Figure 4B), but the β -diversity was significantly associated with ethnic group based on the PERMANOVA analysis of Bray-Curtis dissimilarity matrix ($P = .023$, $R^2 = 0.086$). This result suggests a predominant OTU in some ethnic groups. We performed alpha and phylogenetic diversity analyses comparing *H. pylori*-positive and *H. pylori*-negative samples in each ethnic group and observed significantly lower species richness, Pielou's evenness, Shannon's diversity index, and Faith's phylogenetic diversity index in the positive group compared with the negative group (all $P < .001$; Figure S3).

To find candidate OTUs related to each ethnic group, we performed LEfSe analysis of only *H. pylori*-negative samples. We observed that *Streptococcus* sp and *Micrococcus luteus* were associated with the Timor ethnicity (linear discriminant analysis [LDA] score = 4.78 and 3.56, respectively). *Sphingomonas yabuuchiae* was associated with the Papuan ethnicity (LDA score = 3.82). *Bulleidia* sp and *Atopobium* sp were associated with the Javanese ethnicity (LDA score = 3.89 and 3.87, respectively). *Neisseria subflava*, *Leptotrichia* sp, *Prevotella nanciensis*



2
FIGURE 3 Relative abundance of gastric microbiota between *Helicobacter pylori*-positive and *H. pylori*-negative subjects. At the family level, we observed that Helicobacteraceae accounted for 74.6% of the relative abundance in the *H. pylori*-positive group. At the species level, *H. pylori* was the most dominant OTU, accounting for 76.2% of the relative abundance in the *H. pylori*-positive group




FIGURE 4 The α -diversity among ethnic group in Indonesia. The α -diversity among all Indonesian samples was significantly different, suggesting that ethnic group influences gastric microbial composition in Indonesia. When we analyzed only the *Helicobacter pylori*-negative samples, there were no observed differences among ethnic groups, suggesting that *H. pylori* infection is a stronger influence on microbial composition than ethnic group

(LDA score = 4.16, 3.91 and 3.77, respectively), and *Actinobacillus parahaemolyticus* (LDA score = 3.90) were associated with the Dayak ethnicity. *Schlegelella* sp was associated with the Batakese ethnicity (LDA score = 3.83). *Capnocytophaga* sp, *Prevotella intermedia*, and *Mycoplasma* sp were associated with the Balinese ethnicity (LDA score = 3.94, 3.94 and 3.48, respectively). ANOVA analysis showed significant differences in the relative abundances of *Bulleidia* sp, *Prevotella nanciensis*, and *Prevotella intermedia* among the ethnic groups ($P = .022$, $P = .048$ and $P < .001$, respectively; Figure S4).

3.3 | The microbiome diversity and gastroduodenal diseases

Analysis of all included samples showed significant lower Pielou's phylogenetic diversity index and α -diversity parameters, including Shannon's diversity index, OTU evenness, and species richness, among individuals with gastric mucosal damages (either gastritis ($n = 22$) or precancerous lesions ($n = 56$) group) than those with normal mucosa ($n = 54$) ($P < .001$). When only analyzing the *H. pylori*-negative group, there was no significant difference in species richness between the gastritis ($n = 20$) group and the normal group ($n = 59$; $P = .16$). However, the gastritis group had significantly lower OTU evenness and Shannon's diversity index ($P = .02$ and $P = .046$, respectively).

The LefSe analysis showed *H. pylori* and *Helicobacter* sp OTUs are likely to be factors in gastric mucosal damage (LDA score = 4.83 and 3.69, respectively; Figure 5A). ANOVA analysis confirmed a higher abundance of both *H. pylori* and *Helicobacter* sp in the gastritis group versus the normal group ($P < .001$ and $P = .012$, respectively). In the *H. pylori*-negative group, *Lactobacillus* sp, *Paludibacter* sp, *Dialister* sp, and *Scardovia* sp are likely to be factors in gastric mucosal damage (LDA score = 3.86, 3.56, 3.52 and 3.52, respectively; Figure 5B). Furthermore, ANOVA analysis showed significantly higher relative abundances of *Lactobacillus* sp, *Paludibacter* sp, *Dialister* sp, and *Scardovia* sp in the gastritis group versus the normal group ($P = .03$, $P = .025$, $P = .03$, and $P = .019$, respectively).

There were significant differences in species richness, OTU evenness, and Shannon's diversity index observed between the precancerous (ie, atrophic gastritis, intestinal metaplasia), gastritis, and normal groups (all $P < .001$). The lowest diversity was observed in the precancerous group, followed by the gastritis and normal groups. Among *H. pylori*-positive samples, the species richness, OTU evenness, and Shannon's diversity index were significantly lower in the precancerous group compared with the gastritis group ($P = .02$, $P = .023$, and $P = .012$, respectively). In the *H. pylori*-negative group, the evenness was reduced in the gastritis and precancerous groups compared with the normal group ($P = .067$), whereas there were no significant differences





FIGURE 5 The LefSe analysis of gastric abnormality among Indonesian patients. A. Among all analyzed data, *Helicobacter pylori* and other *Helicobacter* sp showed the most associated to the gastric abnormalities, regardless the type of the abnormality. B. When classified into three groups, the *H. pylori* and other *Helicobacter* sp. showed the most associated to the gastric cancer precursor group (either atrophic gastritis or intestinal metaplasia)

in Shannon's diversity index and species richness. LefSe analysis showed *Corynebacterium* sp and *Rothia dentocariosa* are likely to be factors in the precancerous group (LDA score = 3.56 and 3.47, respectively), and *Paludibacter* sp is likely to be a factor in with the gastritis group (LDA score = 3.43; Figure 6). ANOVA analysis showed *Rothia dentocariosa* had significantly higher relative abundance in the precancerous group compared with the other groups ($P = .037$).

4 | DISCUSSION

This is the 5th study to analyze how gastric mucosal microbiota is related to *H. pylori* infection and ethnicity in Indonesia. Similar to previous studies,^{39,40} we showed that *H. pylori* infection status was an important factor in determining 16 gastric microbiota diversity in Indonesia, with lower α -diversity in the positive group compared with the negative group. In addition to *H. pylori* infection, increasing



18

FIGURE 6 The LEfSe analysis of gastric abnormalities among the *Helicobacter pylori*-negative group

age was also significantly associated with a lower diversity index, consistent with previous observations.^{41,42} As expected, β -diversity analysis showed family Helicobacteraceae had extremely high relative abundance in *H. pylori*-positive group, representing more than three-quarters of the total family output.³⁹ Also consistent with previous studies,^{39,43} we found that *Prevotella* and *Rothia* were the two highest abundance bacterial families within the *H. pylori*-negative group in Indonesia.

In this study, we found and excluded from the analysis 39 cases with discordance in *H. pylori* finding between histologic analysis and 16s rRNA analysis which was excluded for further analysis. This discordance might due to very low bacterial load in the stomach which could not be observed by histologic analysis. Additionally, there are several cases that *H. pylori* colonized in the patchy state.

We found that the Timor, Balinese, and Papuan ethnic groups had significantly less microbiota richness and evenness compared with the Javanese, Dayak, Chinese, Bugis, and Batakese ethnic groups. When these analyses were stratified by *H. pylori* status, there were no significant differences in gastric microbiota composition between the different ethnic groups among *H. pylori*-positive individuals, likely due to *H. pylori* dominating the microbiota composition. Additionally, we found no significant differences in α -diversity between the different ethnic groups among *H. pylori*-negative individuals. Therefore, these differences were more likely explained by the higher prevalence of *H. pylori* infection in these ethnic groups as the prevalence had been shown in previous studies.^{16,44} However, among *H. pylori*-negative individuals, we found a significant association between β -diversity and ethnic groups based on PERMANOVA analysis, suggesting the presence of a predominant OTU in each ethnic group. *Streptococcus* sp was the predominant species in *H. pylori*-negative subjects of Timor ethnicity, *Sphingomonas yabuuchiae* was predominant in those of Papuan ethnicity, and *Bulledia* sp was predominant in those of Javanese ethnicity.

The analysis of the microbiota factors that could explain the differences between healthy tissue and gastric mucosal damage showed that the *H. pylori* and *Helicobacter* sp OTUs had the largest effect in with respect to gastric mucosal damage. However, Indonesia has a low prevalence of *H. pylori* infection but a high prevalence of dyspepsia cases in outpatient clinics.^{16,45,46} Therefore, we focused on *H. pylori*-negative subjects when analyzing association between histologic gastric disease and gastric microbiota. While there was no

significant difference in species richness between subjects with and without gastric mucosal damage, we found significant difference in OTU evenness, which could be caused by the predominance of specific bacteria in the stomachs of subjects with abnormalities. This finding was also supported by the significant differences observed in β -diversity analysis. We found that *Lactobacillus* sp, *Paludibacter* sp, and *Dialister* sp were significantly higher in subjects with gastric abnormalities. Additionally, LEfSe analysis showed that *Paludibacter* sp had the largest effect size in with respect to the gastritis group. *Paludibacter* is a genus of a Gram-negative bacteria that are nonmotile; strictly anaerobic; and lack oxidase, catalase, and nitrate reducing activity.⁴⁷ One previous study reported that *Paludibacter* sp was significantly reduced in subjects receiving proton-pump inhibitor treatment.⁴⁸ Therefore, there could be a possible role for these bacteria in the development of gastric disease. *Dialister* sp found in gastric mucosa was hypothesized to be a members of oral pathogenic taxa that could have a role in the development of gastric cancer.⁴⁹ *Dialister* sp was previously reported to form co-occurring interactions that strengthen with gastric disease progression.⁴⁹

The study was limited by the relatively small sample size for some of the ethnic groups. This small sample size resulted in small numbers in some groups. To mitigate this effect, we combined atrophic gastritis and intestinal metaplasia into the gastric cancer precursor group. Therefore, the associated microorganism needs to be interpreted with caution. The further study with bigger sample number is necessary for more clear association. In addition, we did not include transcriptome, proteomic, or metabolomic profiles; therefore, we cannot draw inferences regarding functional differences. The use of DNA analysis does not confirm whether microorganisms are alive or not. Finally, the findings reflect significant association that may not be interpreted as the causal. Future studies are needed to determine the correlations between gastric microbiota and the natural history of gastric disease and get a better understanding of the role of gastric microbiota.

5 | CONCLUSIONS

Our study confirmed that *H. pylori* infection overwhelms other gastric microbiota, dominating the gastric microbial composition. The effect of *H. pylori* infection on the gastric microbiomes of various ethnic groups is stronger than the effect of the ethnic group

designation itself. Among *H. pylori*-negative individuals, *Paludibacter* sp and *Dialister* sp were the species that could explain gastric mucosal damage in Indonesia.

ACKNOWLEDGEMENTS

None.

9

CONFLICT OF INTEREST

The authors declare there was no potential competing interest.

AUTHORS' CONTRIBUTIONS

YY, MM, and AFS conceived and designed the experiments. MM, TM, TU, and LAW performed the experiments. MM, YY, LAW, HBE, and NJA analyzed the data. AFS, HBE, NJA, DD, PBS, YAAR, and IAN contributed reagents/materials/analysis tools. MM, LAW, YY, and HBE wrote the manuscript.

15

ETHICS APPROVAL

Written informed consent was obtained from all patients. The study protocol was approved by the Ethics Committee of Dr Cipto Mangunkusumo Teaching Hospital (Jakarta, Indonesia), Dr Soetomo Teaching Hospital (Surabaya, Indonesia), Dr Rahidin Sudirohusodo Teaching Hospital (Makassar, Indonesia), and Oita University Faculty of Medicine (Yufu, Japan).

2

FORMED CONSENT

Not applicable.

DATA AVAILABILITY STATEMENT

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

ORCID

Muhammad Miftahussurur  <https://orcid.org/0000-0003-1415-6033>

Langgeng Agung Waskito  <https://orcid.org/0000-0002-9400-1973>

Yoshio Yamaoka  <https://orcid.org/0000-0002-1222-5819>

REFERENCES

- Marshall B, Warren JR. Unidentified curved bacilli in the stomach of patients with gastritis and peptic ulceration. *Lancet*. 1984;323(8390):1311-1315.
- Uemura N, Okamoto S, Yamamoto S, et al. *Helicobacter pylori* infection and the development of gastric cancer. *N Engl J Med*. 2001;345(11):784-789.
- Petrosino JF, Highlander S, Luna RA, Gibbs RA, Versalovic J. Metagenomic pyrosequencing and microbial identification. *Clin Chem*. 2009;55(5):856-866.
- Nardone G, Compare D. The human gastric microbiota: Is it time to rethink the pathogenesis of stomach diseases? *United European Gastroenterol J*. 2015;3(3):255-260.
- Abreu MT, Peek RM. Gastrointestinal malignancy and the microbiome. *Gastroenterology*. 2014;146(6):1534-1546.e1533.
- Noto JM, Peek RM Jr. The gastric microbiome, its interaction with *Helicobacter pylori*, and its potential role in the progression to stomach cancer. *PLoS Pathog*. 2017;13(10):e1006573-e1006573.
- Castano-Rodriguez N, Goh KL, Fock KM, Mitchell HM, Kaakoush NO. Dysbiosis of the microbiome in gastric carcinogenesis. *Sci Rep*. 2017;7(1):15957.
- The Human Microbiome Project C, Huttenhower C, Gevers D, et al. Structure, function and diversity of the healthy human microbiome. *Nature*. 2012;486:207.
- Wu GD, Chen J, Hoffmann C, et al. Linking long-term dietary patterns with gut microbial enterotypes. *Science*. 2011;334(6052):105-108.
- Chong CW, Ahmad AF, Lim YAL, et al. Effect of ethnicity and socioeconomic variation to the gut microbiota composition among pre-adolescent in Malaysia. *Sci Rep*. 2015;5:13338-13338.
- Liao M, Xie Y, Mao Y, et al. Comparative analyses of fecal microbiota in Chinese isolated Yao population, minority Zhuang and rural Han by 16sRNA sequencing. *Sci Rep*. 2018;8(1):1142-1142.
- Deschasaux M, Bouter KE, Prodan A, et al. Depicting the composition of gut microbiota in a population with varied ethnic origins but shared geography. *Nat Med*. 2018;24(10):1526-1531.
- Brooks AW, Priya S, Blekhan R, Bordenstein SR. Gut microbiota diversity across ethnicities in the United States. *PLoS Biol*. 2018;16(12):e2006842.
- Das A, Pereira V, Saxena S, et al. Gastric microbiome of Indian patients with *Helicobacter pylori* infection, and their interaction networks. *Sci Rep*. 2017;7(1):15438-15438.
- Khosravi Y, Gan HM, Jing P, et al. The gastric microbiome of four Malaysian gastroduodenal disease patients. *Arch Gene Genome Res* 1(1):1-9.
- Syam AF, Miftahussurur M, Makmun D, et al. Risk factors and prevalence of *Helicobacter pylori* in five largest islands of Indonesia: a preliminary study. *PLoS ONE*. 2015;10(11):e0140186.
- Klindworth A, Pruesse E, Schweer T, et al. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Res*. 2013;41(1):e1.
- Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*. 2014;30(15):2114-2120.
- Rognes T, Flouri T, Nichols B, Quince C, Mahe F. VSEARCH: a versatile open source tool for metagenomics. *PeerJ*. 2016;4:e2584.
- DeSantis TZ, Hugenholtz P, Larsen N, et al. Greengenes, a chimeric-checked 16S rRNA gene database and workbench compatible with ARB. *Appl Environ Microbiol*. 2006;72(7):5069-5072.
- Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol*. 2013;30(4):772-780.
- Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res*. 2002;30(14):3059-3066.
- Price MN, Dehal PS, Arkin AP. FastTree 2—approximately maximum-likelihood trees for large alignments. *PLoS ONE*. 2010;5(3):e9490.
- Pielou EC. The measurement of diversity in different types of biological collections. *J Theor Biol*. 1966;13:131-144.
- Faith DP. Conservation evaluation and phylogenetic diversity. *Biol Cons*. 1992;61(1):1-10.
- Shannon CE. A mathematical theory of communication. *Bell Syst Tech J*. 1948;27(3):379-423.
- Bray JR, Curtis JT. An ordination of the upland forest communities of southern Wisconsin. *Ecol Monogr*. 1957;27(4):325-349.
- Oksanen J, Blanchet FG, Kindt R, et al. Vegan: Community Ecology Package. R package version 2.0-2. 2012.
- Segata N, Izard J, Waldron L, et al. Metagenomic biomarker discovery and explanation. *Genome Biol*. 2011;12(6):R60.

30. Zakrzewski M, Proietti C, Ellis JJ, et al. Calypso: a user-friendly web-server for mining and visualizing microbiome-environment interactions. *Bioinformatics*. 2017;33(5):782-783.
31. Dixon M, Genta R, Yardley J, Correa P. Classification and grading of gastritis. The updated Sydney System. International Workshop on the Histopathology of Gastritis, Houston 1994. *Am J Surg Pathol*. 1996;20(10):1161-1181.
32. Uchida T, Kanada R, Tsukamoto Y, et al. Immunohistochemical diagnosis of the cagA-gene genotype of *Helicobacter pylori* with anti-East Asian CagA-specific antibody. *Cancer Sci*. 2007;98(4):521-528.
33. Vilaichone RK, Mahachai V, Shiota S, et al. Extremely high prevalence of *Helicobacter pylori* infection in Bhutan. *World J Gastroenterol*. 2013;19(18):2806-2810.
34. Shiota S, Murakami K, Fujioka T, Yamaoka Y. Population-based strategies for *Helicobacter pylori*-associated disease management: a Japanese perspective. *Expert Rev Gastroenterol Hepatol*. 2010;4(2):149-156.
35. Shiota S, Cruz M, Abreu JA, et al. Virulence genes of *Helicobacter pylori* in the Dominican Republic. *J Med Microbiol*. 2014;63(Pt 9):1189-1196.
36. Nguyen TL, Uchida T, Tsukamoto Y, et al. *Helicobacter pylori* infection and gastroduodenal diseases in Vietnam: a cross-sectional, hospital-based study. *BMC Gastroenterol*. 2010;10:114.
37. Nguyen LT, Uchida T, Tsukamoto Y, et al. Clinical relevance of cagPAI intactness in *Helicobacter pylori* isolates from Vietnam. *Eur J Clin Microbiol Infect Dis*. 2010;29(6):651-660.
38. Miftahussurur M, Shiota S, Suzuki R, et al. Identification of *Helicobacter pylori* infection in symptomatic patients in Surabaya, Indonesia, using five diagnostic tests. *Epidemiol Infect*. 2015;143(5):986-996.
39. Andersson AF, Lindberg M, Jakobsson H, Backhed F, Nyren P, Engstrand L. Comparative analysis of human gut microbiota by bar-coded pyrosequencing. *PLoS ONE*. 2008;3(7):e2836.
40. Li TH, Qin Y, Sham PC, Lau KS, Chu KM, Leung WK. Alterations in gastric microbiota after *H. Pylori* eradication and in different histological stages of gastric carcinogenesis. *Sci Rep*. 2017;7:44935.
41. Jeffery IB, Lynch DB, O'Toole PW. Composition and temporal stability of the gut microbiota in older persons. *ISME J*. 2016;10(1):170-182.
42. O'Toole PW, Jeffery IB. Gut microbiota and aging. *Science*. 2015;350(6265):1214-1215.
43. Li XX, Wong GL, To KF, et al. Bacterial microbiota profiling in gastritis without *Helicobacter pylori* infection or non-steroidal anti-inflammatory drug use. *PLoS ONE*. 2009;4(11):e7985.
44. Miftahussurur M, Syam AF, Nusi IA, et al. Surveillance of *Helicobacter pylori* antibiotic susceptibility in Indonesia: different resistance types among regions and with novel genetic mutations. *PLoS ONE*. 2016;11(12):e0166199.
45. Makmun D. Present status of endoscopy, therapeutic endoscopy and the endoscopy training system in Indonesia. *Dig Endosc*. 2014;26(S2):2-9.
46. Miftahussurur M, Waskito LA, Syam AF, et al. Analysis of risks of gastric cancer by gastric mucosa among Indonesian ethnic groups. *PLoS ONE*. 2019;14(5):e0216670.
47. Ueki A, Akasaka H, Suzuki D, Ueki K *Paludibacter propionicigenes* gen. nov., sp. nov., a novel strictly anaerobic, Gram-negative, propionate-producing bacterium isolated from plant residue in irrigated rice-field soil in Japan. *Int J Syst Evol Microbiol*. 2006;56(Pt 1):39-44.
48. Parsons BN, Ijaz UZ, D'Amore R, et al. Comparison of the human gastric microbiota in hypochlorhydric states arising as a result of *Helicobacter pylori*-induced atrophic gastritis, autoimmune atrophic gastritis and proton pump inhibitor use. *PLoS Pathogens*. 2017;13(11):e1006653.
49. Coker OO, Dai Z, Nie Y, et al. Mucosal microbiome dysbiosis in gastric carcinogenesis. *Gut*. 2018;67(6):1024-1032.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Miftahussurur M, Waskito LA, El-Serag HB, et al. Gastric microbiota and *Helicobacter pylori* in Indonesian population. *Helicobacter*. 2020;00:e12695. <https://doi.org/10.1111/hel.12695>

Gastric Microbiota and Helicobacter Pylori in Indonesian Population

ORIGINALITY REPORT

18%

SIMILARITY INDEX

10%

INTERNET SOURCES

16%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Osamu Matsunari, Muhammad Miftahussurur, Seiji Shiota, Rumiko Suzuki et al. "Rare Helicobacter pylori Virulence Genotypes in Bhutan", Scientific Reports, 2016
Publication 2%
- 2 www.eurosurveillance.org
Internet Source 1%
- 3 repository.essex.ac.uk
Internet Source 1%
- 4 Seyedeh Azra Shamsdin, Abdolvahab Alborzi, Manoochehr Rasouli, Abbas Ghaderi et al. " The importance of TH22 and TC22 cells in the pathogenesis of associated gastric diseases ", Helicobacter, 2017
Publication 1%
- 5 H. Isomoto, Y. Mizuta, K. Inoue, T.. "A Close Relationship between Helicobacter pylori Infection and Gastric Xanthoma", Scandinavian Journal of Gastroenterology, 2009 1%

- | | | |
|----|--|-----|
| 6 | www.ajtmh.org
Internet Source | 1% |
| 7 | M. Barr. "Non-steroidal anti-inflammatory drugs and Helicobacter pylori", Alimentary Pharmacology and Therapeutics, 10/2000
Publication | 1% |
| 8 | T. Ohkusa. "Improvement in serum pepsinogens and gastrin in long-term monitoring after eradication of Helicobacter pylori: comparison with H. pylori-negative patients", Alimentary Pharmacology and Therapeutics, 7/2004
Publication | 1% |
| 9 | tel.archives-ouvertes.fr
Internet Source | <1% |
| 10 | Shamsdin, Seyedeh Azra, Abdolvahab Alborzi, Manoochehr Rasouli, Mohamad Kazem Hosseini, Kamran Bagheri Lankrani, and Mehdi Kalani. "Alterations in Th17 and the Respective Cytokine Levels in Helicobacter pylori -Induced Stomach Diseases", Helicobacter, 2015.
Publication | <1% |
| 11 | Keng-Po Lai, Yan-Tung Chung, Rong Li, Hin-Ting Wan, Chris Kong-Chu Wong. "Bisphenol A alters gut microbiome: Comparative metagenomics analysis", Environmental | <1% |

Pollution, 2016

Publication

12

trepo.tuni.fi

Internet Source

<1%

13

Vilaichone, Ratha-korn. "Antibiotics resistance rate of Helicobacter pylori in Bhutan", World Journal of Gastroenterology, 2013.

Publication

<1%

14

www.eurekaselect.com

Internet Source

<1%

15

journals.sagepub.com

Internet Source

<1%

16

gastrores.org

Internet Source

<1%

17

Bibi Fm Werdmuller, Toon Bmm Van Der Putten, Ton G Balk, Cornelius Bhw Lamers, Ruud Jlf Loffeld. " Clinical presentation of - positive and -negative functional dyspepsia ", Journal of Gastroenterology and Hepatology, 2001

Publication

<1%

18

es.scribd.com

Internet Source

<1%

19

Shuai Wang, Rene Martins, Megan C. Sullivan, Elliot S. Friedman et al. "Diet-induced remission

<1%

in chronic enteropathy is associated with altered microbial community structure and synthesis of secondary bile acids", *Microbiome*, 2019

Publication

20

Alistair R. Legione, Jemima Amery-Gale, Michael Lynch, Leesa Haynes, James R. Gilkerson, Fiona M. Sansom, Joanne M. Devlin. "Variation in the microbiome of the urogenital tract of Chlamydia-free female koalas (*Phascolarctos cinereus*) with and without 'wet bottom'", *PLOS ONE*, 2018

Publication

21

Submitted to Universitas Airlangga

Student Paper

22

jneuroengrehab.biomedcentral.com

Internet Source

23

Atsuki Fukushima, Zhang Jian, Waka Ishida, Kazuyo Fukata, Akemi Ozaki, Hisayuki Ueno. "Interleukin-4–Mediated Infiltration of Eosinophils into the Conjunctiva and Its Suppression by Interferon- γ ", *Current Eye Research*, 2009

Publication

24

Shamshul Ansari, Yoshio Yamaoka. "Helicobacter pylori Virulence Factors Exploiting Gastric Colonization and its Pathogenicity", *Toxins*, 2019

<1%

<1%

<1%

<1%

<1%

25

Muhammad Miftahussurur, Manu Tandan, Dadang Makmun, Zaheer Nabi. "Peroral Endoscopic Myotomy in a Patient with Achalasia Cardia with Prior Heller's Myotomy", Case Reports in Gastroenterology, 2020

Publication

<1%

26

Erin R. Shanahan, Ayesha Shah, Natasha Koloski, Marjorie M. Walker, Nicholas J. Talley, Mark Morrison, Gerald J. Holtmann. "Influence of cigarette smoking on the human duodenal mucosa-associated microbiota", Microbiome, 2018

Publication

<1%

27

Jinhee Choi, Sun-hee Yoon, Ja-eun Kim, Kwang-ho Rhee, Hee-sang Youn, Myung-hee Chung. "Gene-specific oxidative DNA damage in *Helicobacter pylori*-infected human gastric mucosa", International Journal of Cancer, 2002

Publication

<1%

28

bmcevolbiol.biomedcentral.com

Internet Source

<1%

29

pure.sruc.ac.uk

Internet Source

<1%

30

bmcbiol.biomedcentral.com

Internet Source

<1%

31

María Elena Rodríguez-García, Francisco Javier Cotrina-Vinagre, María de los Ángeles Gómez-Cano, Ana Martínez de Aragón et al. " variant causes mega-corporis-callosum syndrome with cortical malformations but without cerebellar hypoplasia ", American Journal of Medical Genetics Part A, 2020

Publication

<1%

32

J M Parry. "Fluorescence in situ hybridisation analysis of chromosomal aberrations in gastric tissue: the potential involvement of Helicobacter pylori", British Journal of Cancer, 05/09/2005

Publication

<1%

33

www.thieme-connect.com

Internet Source

<1%

34

tessera.spandidos-publications.com

Internet Source

<1%

35

wecaremobility.co.uk

Internet Source

<1%

36

gut.bmj.com

Internet Source

<1%

37

T Koike. "Helicobacter pylori infection prevents erosive reflux oesophagitis by decreasing gastric acid secretion", Gut, 2001

Publication

<1%

38

www.termedia.pl

Internet Source

<1%

39

Siobhan F. Clarke, Eileen F. Murphy, Kanishka Nilaweera, Paul R. Ross, Fergus Shanahan, Paul W. O'Toole, Paul D. Cotter. "The gut microbiota and its relationship to diet and obesity", Gut Microbes, 2014

Publication

<1%

40

espace.library.uq.edu.au

Internet Source

<1%

41

Peters. "The influence of Helicobacter pylori on oesophageal acid exposure in GERD during acid suppressive therapy", Alimentary Pharmacology and Therapeutics, 7/1999

Publication

<1%

42

Sicheng Wen, Christian P. Felley, Hanifa Bouzourene, Mark Reimers, Pierre Michetti, Qiang Pan-Hammarström. " Inflammatory Gene Profiles in Gastric Mucosa during Infection in Humans ", The Journal of Immunology, 2004

Publication

<1%

43

Carlotta Catozzi, Armand Sanchez Bonastre, Olga Francino, Cristina Lecchi et al. "The microbiota of water buffalo milk during mastitis", PLOS ONE, 2017

Publication

<1%

- 44 L. T. Nguyen, T. Uchida, A. Kuroda, Y. Tsukamoto et al. "Evaluation of the Anti-East Asian CagA-Specific Antibody for CagA Phenotyping", *Clinical and Vaccine Immunology*, 2009
Publication <1%
-
- 45 malariajournal.biomedcentral.com
Internet Source <1%
-
- 46 www.wpclt.com
Internet Source <1%
-
- 47 Katja Spiess, Suzan Fares, Alexander H. Sparre-Ulrich, Ellen Hilgenberg et al. "Identification and Functional Comparison of Seven-Transmembrane G-Protein-Coupled BILF1 Receptors in Recently Discovered Nonhuman Primate Lymphocryptoviruses", *Journal of Virology*, 2015
Publication <1%
-
- 48 Cuicui Sun, Yanan Xie, Xiaoli Hu, Jieni Fu, Jie Zhou, Ligeng Wu. "Relationship between Clinical Symptoms and the Microbiota in Advanced Caries", *Journal of Endodontics*, 2020
Publication <1%
-

Exclude bibliography On

Gastric Microbiota and Helicobacter Pylori in Indonesian Population

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10
