

Volume - 10

No. 11

November-2017

ISSN: 0972-8988

EISSN: 2231-0916



NAAS Rating : 5.71

Indexed in ESCI-Thomson Reuters, PubMed,  
PubMed Central, DOAJ, Scopus, CABI, CAS etc.

---

---

# Veterinary World

Open access and peer reviewed journal

---

---



**Editorial office**

Veterinary World,  
Star, Gulshan Park,  
NH-8A, Chandrapur Road,  
Wankaner - 363621,  
Dist. Morbi, Gujarat, India  
Website: [www.veterinaryworld.org](http://www.veterinaryworld.org)  
E-mail: [editorveterinaryworld@gmail.com](mailto:editorveterinaryworld@gmail.com)

# Veterinary World

**Editorial Office:** Veterinary World, Star, Gulshan Park, NH-8A, Chandrapur Road, Wankaner - 363621, Dist. Morbi, Gujarat, India

ISSN: 0972-8988, EISSN: 2231-0916, [www.veterinaryworld.org](http://www.veterinaryworld.org)

## Editor-in-Chief

Anjum V. Sherasiya - Ex-Veterinary Officer, Department of Animal Husbandry, Gujarat State, India

## Associate Editors

Shambhunath Choudhary - Department of Biomedical & Diagnostic Sciences, College of Veterinary Medicine, The University of Tennessee, 2407 River Drive, Room A 201, Knoxville, TN 37996, U.S.A.

Suresh H. Basagoudanavar - FMD Vaccine Research Laboratory, IVRI, Bangalore, Bangalore- 560024, Karnataka, India

## Editorial board

R. G. Jani - Ex-Coordinator Wildlife Health, Western Region Centre, Indo-US Project, Department of Veterinary Medicine, Veterinary College, Anand - 388001, Gujarat, India

G. N. Gongal - Technical Officer, WHO South-East Asia Regional Office, New Delhi -110002, India

Ranganath Mamidi - Dr. Julian E Stelzer's Lab, Department of Physiology & Biophysics, Medical School, Case Western Reserve University, Cleveland, OH - 44106, U.S.A.

Md. Tanvir Rahman - Department of Microbiology and Hygiene, Faculty of Veterinary Science, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh

Deepmala Agarwal - Cancer Prevention Laboratory, Pennington Biomedical Research Center, Baton Rouge, LA, U.S.A.

Foud Kasim Mohammad - Professor, Department of Pharmacology & Toxicology, Vice President for Administrative & Financial Affairs, University of Mosul, P.O. Box 11136, Mosul, Iraq

Abdel-Baset Nasr Sayed Ahmed - Professor and Head, Department of Animal Nutrition and Clinical Nutrition, Faculty of Veterinary Medicine, Assiut University, Assiut, Egypt

Nicole Borel - Department of Pathology, Vetsuisse Faculty, University of Zurich, CH-8057 Zürich, Switzerland

B. A. Lubisi - Virology, MED Programme, ARC - Onderstepoort Veterinary Institute, No. 100 Old Soutpan Road, Onderstepoort, Tshwane, 0110, South Africa

Kumar Venkitanarayan - Associate Professor, Graduate Programs Chair, Honors and Pre-Vet Programs Advisor, Department of Animal Science, University of Connecticut, Storrs, CT 06269, U.S.A.

Kemin Xu - Department of Veterinary Medicine, University of Maryland, College Park College Park, MD, 20742, U.S.A.

Vassilis Papatziros - Faculty of Veterinary Medicine, Department of Medicine (Porcine Medicine), University of Thessaly, Thessaly, Greece

Mathias Devreese - Laboratory of Pharmacology and Toxicology, Faculty of Veterinary Medicine, Ghent University, Belgium

Sumeet Sharma - Edmonton North Animal Hospital, Edmonton, Alberta, T5X 3Y7, Canada

K. P. Singh - School of Medicine and Dentistry, University of Rochester, Department of Environmental Medicine, Room: 4- 6820, 601 Elmwood Avenue, Box-EHSC, Rochester, New York-14620, U.S.A.

Raj Mohan Raja Muthiah - Massachusetts General Hospital, Harvard Medical School, Boston, Massachusetts, U.S.A. Ashok K. Chockalingam - Division of Applied Regulatory Science, U.S. Food and Drug Administration, 10903, New Hampshire Avenue, Silver Spring, Maryland 20993, U.S.A.

Ashutosh Wadhwa - Poxvirus and Rabies Branch, Division of High-Consequence Pathogens and Pathology, National Center for Emerging and Zoonotic Infectious Diseases, Centers for Disease Control and Prevention, 1600 Clifton Road, NE, Mail Stop G33, Atlanta, GA 30333, U.S.A.

Luiz Otavio Pereira Carvalho - Laboratory of Immunomodulation and Protozoology, Oswaldo Cruz Institute, Ministry of Health (Brazil), Pavilhão "108" - Sala: 09, Av. Brasil, 4365 - Manguinhos, Rio de Janeiro - RJ, CEP: 21040-360, Brazil.

Editorial office contact person: Anjum V. Sherasiya, email: [editorveterinaryworld@gmail.com](mailto:editorveterinaryworld@gmail.com)

**Ratings of Veterinary World:** NAAS - National Academy of Agricultural Sciences - 5.71, Scimago Journal Rank -0.284, Citescore - 0.57, SNIP - Source Normalized Impact per Paper - 0.570

## Indexing and abstracting

Academic Journals Database, AGORA, AGRICOLA, AGRIS, CABI, CAS, DOAJ, EBSCO, ESCI - Thomson Reuters, Gale, Google Scholar, HINARI, Index Scholar, Indian Animal Science Abstracts, Indian Science Abstracts, JournalSeek, Open J-gate, ProQuest, PubMed, PubMed Central, SCOPUS, TEEAL

## Publisher: Veterinary World

Veterinary World is an open access journal, each issue available free of cost at [www.veterinaryworld.org](http://www.veterinaryworld.org).

We accept online submission only. For more information regarding submission and publication charges, please visit [www.veterinaryworld.org](http://www.veterinaryworld.org)

**Rs.7000 for Indian/USD 175 for Abroad per copy**

Printed and Published by Dr. Anjum V. Sherasiya on behalf of Veterinary World. Printed and Published at Star, Gulshan Park, N.H. 8A, Chandrapur Road, Wankaner-363621, Dist. Morbi, Gujarat, India.  
Editor-in-Chief: Dr. Anjum V. Sherasiya

---

# Veterinary World

---

ISSN: 0972-8988, EISSN: 2231-0916, [www.veterinaryworld.org](http://www.veterinaryworld.org)

---

**Volume-10**

**No.11**

**November-2017**

---

The articles in Veterinary World are open access articles licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

---

Research (Published online: 01-11-2017)

1. Genotyping of virulent *Escherichia coli* obtained from poultry and poultry farm workers using enterobacterial repetitive intergenic consensus-polymerase chain reaction

M. Soma Sekhar, N. Mohammad Sharif, T. Srinivasa Rao and M. Metta

Veterinary World, 10(11): 1292-1296

Research (Published online: 02-11-2017)

2. Preliminary study of Malaysian fruit bats species diversity in Lenggong Livestock Breeding Center, Perak: Potential risk of spill over infection

Muhammed Mikail, T. A. Tengku Rinalfi Putra, Arshad Siti Suri, Mohd Noor Mohd Hezme and M. T. Marina

Veterinary World, 10(11): 1297-1300

Research (Published online: 03-11-2017)

3. Nutritional enrichment of broiler breast meat through dietary supplementation of Indian ginseng *Withania somnifera* and synbiotic substances under semi-arid climatic conditions

Sonal Thakur, Tribhuwan Sharma, Radhe Shyam Arya, Basant Bais and Vijay Kumar Agrawal

Veterinary World, 10(11): 1301-1306

Research (Published online: 08-11-2017)

4. Immunocytochemical study of canine lymphomas and its correlation with exposure to tobacco smoke

K. C. Pinello, M. Santos, L. Leite-Martins, J. Niza-Ribeiro and A. J. de Matos

Veterinary World, 10(11): 1307-1313

Research (Published online: 09-11-2017)

5. *Dermatophilus congolensis* infection in sheep and goats in Delta region of Tamil Nadu

M. Ananda Chitra, K. Jayalakshmi, P. Ponnusamy, R. Manickam and B. S. M. Ronald

Veterinary World, 10(11): 1314-1318

Research (Published online: 10-11-2017)

6. High prevalence of small *Babesia* species in canines of Kerala, South India

Kollannur Jose Jain, Bindu Lakshmanan, Karunakaran Syamala, Jose E Praveena and Thazhathuveetil Aravindakshan

Veterinary World, 10(11): 1319-1323

Research (Published online: 11-11-2017)

7. The phylogenetics of *Leucocytozoon caulleryi* infecting broiler chickens in endemic areas in Indonesia

Endang Suprihati and Wiwik Misaco Yuniarti

Veterinary World, 10(11): 1324-1328

Research (Published online: 12-11-2017)

8. Production of aldosterone in cardiac tissues of healthy dogs and with dilated cardiomyopathy

Alejandro Reynoso Palomar, Georgina Mena Aguilar, Marisol Cruz Garcia, Cesar Pastelin Rojas and Abel Villa Mancera

Veterinary World, 10(11): 1329-1332

Research (Published online: 13-11-2017)

9. Immunoglobulin subclass in experimental murine *Toxocara cati* infection

Kusnoto, Setiawan Koesdarto, Sri Mumpuni and R. Heru Prasetyo

Veterinary World, 10(11): 1333-1336

Research (Published online: 14-11-2017)

10. Effects of hyperbaric therapy on liver morphofunctional of rabbits (*Oryctolagus cuniculus*) after hind limb ischemia-reperfusion injury

Bambang Sektiari Lukiswanto, Wiwik Misaco Yuniarti and Y. Yosis Motulo

Veterinary World, 10(11): 1337-1342

---

Research (Published online: 15-11-2017)

11. Determination of serum adenosine deaminase and xanthine oxidase activity in Kangal dogs with maternal cannibalism  
N. Ercan, M. Kockaya, S. Kapancik and D. Bakir  
Veterinary World, 10(11): 1343-1346

Research (Published online: 16-11-2017)

12. Reducing zoonotic and internal parasite burdens in pigs using a pig confinement system  
Kadek Karang Agustina, Ida Bagus Ngurah Swacita, Ida Bagus Made Oka, I Made Dwinata, Rebecca Justin Traub, Colin Cargill and I Made Damriyasa  
Veterinary World, 10(11): 1347-1352

Research (Published online: 20-11-2017)

13. The clinical impact of antimicrobial resistance genomics in competition with she-camels recurrent mastitis metabolomics due to heterogeneous *Bacillus licheniformis* field isolates  
Nesreen Allam Tantawy Allam, Doaa Sedky and Enshrah Khalil Mira  
Veterinary World, 10(11): 1353-1360

Review (Published online: 21-11-2017)

14. Advances in genome editing for improved animal breeding: A review  
Shakil Ahmad Bhat, Abrar Ahad Malik, Syed Mudasir Ahmad, Riaz Ahmad Shah, Nazir Ahmad Ganai, Syed Shanaz Shafi and Nadeem Shabir  
Veterinary World, 10(11): 1361-1366

Review (Published online: 24-11-2017)

15. Metabolic and immunological changes in transition dairy cows: A review  
Pratik Ramesh Wankhade, A. Manimaran, A. Kumaresan, S. Jeyakumar, K. P. Ramesha, V. Sejian, D. Rajendran and Minu Rachel Varghese  
Veterinary World, 10(11): 1367-1377

Research (Published online: 25-11-2017)

16. Clinical, pathological, and molecular investigation of *Mycoplasma pulmonis*-induced murine respiratory mycoplasmosis in a rat (*Rattus norvegicus*) colony  
Saurabh Chawla, Sarita Jena, Balaji Venkatsan, Kuna Mahara and Nilanjan Sahu  
Veterinary World, 10(11): 1378-1382

Research (Published online: 26-11-2017)

17. Application of loop-mediated isothermal amplification assay in the detection of herpesvirus of turkey (FC 126 strain) from chicken samples in Nigeria  
A. J. Adedeji, P. A. Abdu, P. D. Luka, A. A. Owoade and T. M. Joannis  
Veterinary World, 10(11): 1383-1388

Research (Published online: 28-11-2017)

18. Molecular analysis of genome segment-3 of bluetongue virus serotype 12 isolates from Haryana  
Anita Dalal, Sushila Maan, Nitish Bansal, Vinay Kumar, Aman Kumar, Narender Singh Maan and Naresh Kumar Kakker  
Veterinary World, 10(11): 1389-1393

Research (Published online: 29-11-2017)

19. Plastination of macroparasites: An eco-friendly method of long-term preservation  
Niranjan Kumar, Bhupamani Das, Jayesh B. Solanki, Mehul M. Jadav and Ramasamy Menaka  
Veterinary World, 10(11): 1394-1400

Research (Published online: 30-11-2017)

20. Pathology and polymerase chain reaction detection of ovine progressive pneumonia (maedi) cases in slaughtered sheep in India  
Rahul Singh, Pawan Kumar, Rajendra Singh, Kuldeep Dhama, Swati Kumari, Jay Prakash Yadav, Gayatri Kashyap, Karam Pal Singh, Vidya Singh and Monalisa Sahoo  
Veterinary World, 10(11): 1401-1406

\*\*\*\*\*

## Effects of hyperbaric therapy on liver morphofunctional of rabbits (*Oryctolagus cuniculus*) after hind limb ischemia-reperfusion injury

Bambang Sektiari Lukiswanto<sup>1</sup>, Wiwik Misaco Yuniarti<sup>1</sup> and Y. Yosis Motulo<sup>2</sup>

1. Department of Veterinary Clinic, Faculty of Veterinary Medicine, Universitas Airlangga, Jl. Mulyorejo, Kampus C, Surabaya 60115, Indonesia; 2. Department of Surgery, Division of Thoracic, Cardiac, and Vascular Surgery, Dr. Soetomo General Hospital, Surabaya 60115, Indonesia.

**Corresponding author:** Bambang Sektiari Lukiswanto, e-mail: [bamsekti@yahoo.com](mailto:bamsekti@yahoo.com)

**Co-authors:** WMY: [wiwikmisaco@yahoo.com](mailto:wiwikmisaco@yahoo.com), YYM: [yosis\\_motulo@yahoo.com](mailto:yosis_motulo@yahoo.com)

**Received:** 11-06-2017, **Accepted:** 18-10-2017, **Published online:** 14-11-2017

**doi:** 10.14202/vetworld.2017.1337-1342 **How to cite this article:** Lukiswanto BS, Yuniarti WM, Motulo YY (2017) Effects of hyperbaric therapy on liver morphofunctional of rabbits (*Oryctolagus cuniculus*) after hind limb ischemia-reperfusion injury, *Veterinary World*, 10(11): 1337-1342.

### Abstract

**Aim:** The objective of this research was to study and to prove the effectiveness of hyperbaric oxygen therapy (HBOT) starting time on liver morphofunctional changes after ischemia-reperfusion in the hind limb of rabbits.

**Materials and Methods:** This research used a complete randomized design with 4 groups and 6 repetitions on each. After 6 h artery femoral is ligation, reperfusion was performed for 100 min (G1), HBOT for 90 min after 10 min reperfusion (G2), 10 min reperfusion (G3), and HBOT 90 min after 60 min reperfusion (G4). Then, all of the rabbits were sacrificed. The liver and blood were taken for histopathological changes examination as well as for measuring the level of aspartate aminotransferase (AST) and alanine aminotransferase (ALT). The statistical test using Kruskal–Wallis and Mann–Whitney showed that the score of degeneration, necrosis, and portal inflammation in groups without HBOT (G1 and G3) were not significantly different, as well as in group with HBOT (G2 and G4) ( $p > 0.05$ ). However, the scores of histopathological changes in G1 and G3 were significantly different from those in G2 and G4 ( $p < 0.05$ ). The levels of AST and ALT in the groups without hyperbaric therapy (G1 and G3) were not significantly different from those in the groups treated with hyperbaric therapy (G2 and G4) ( $p > 0.05$ ).

**Results:** Hind limb ischemia injury reperfusion can trigger damage for liver morphology, but not lead to liver dysfunction. Reperfusion can trigger increased activity of neutrophils, while neutrophil infiltration in the organ will lead to dysfunction. HBOT can inhibit the activity of neutrophils and the dysfunction of organs caused by ischemic reperfusion.

**Conclusion:** HBOT for 90 min, both 10 and 60 min after the reperfusion, can protect hepatocytes from damage.

**Keywords:** hyperbaric oxygen therapy, ischemia, liver, morphofunctional, reperfusion.

### Introduction

Ischemic condition can occur in a wide variety of disease status. Nevertheless, an ischemic condition often occurs in a variety of surgical procedures such as in a surgical process of vascular or organ transplantation. In such circumstances, ischemic injury can occur when blood flow to a tissue is reduced or stopped in a long time, resulting changes in the metabolism of muscles, from aerobic state into anaerobic one. Consequently, adenosine triphosphate (ATP) depletion and calcium (Ca) extracellular transfer into muscle cells occur, leading to abnormalities in metabolism and thrombosis, as well as tissue inflammation [1].

Blood flow improvement (reperfusion) is the main effort to overcome tissue ischemia (ischemia-reperfusion [IR]). However, reperfusion has negative side effects because it can cause damage and

impaired organ function when it is not given appropriately. Inappropriate reperfusion then can trigger activation of proinflammatory mediators as well as formation of reactive oxygen species (ROS), resulting injury to the tissue [2]. In severe ischemic conditions, excessive inflammatory response after reperfusion can lead to a systemic inflammatory response syndrome and multiple organ dysfunction syndrome (MODS) [3]. MODS generally starts from the lungs and then spreads to the liver, gastrointestinal tract, and kidneys. Bone marrow and liver dysfunctions are in the final stages of MODS. Meanwhile, the dysfunction of the nervous system can occur at the beginning or end of MODS [4].

Many researchers actually have already been conducted on the effects of various drugs in reducing the negative impacts of reperfusion. Those previous researchers focused on the impacts of reperfusion used hyperbaric oxygen therapy (HBOT), a method of therapy using 100% oxygen with a high atmospheric pressure in a special room. HBOT has been proved to be very helpful to reduce the negative impacts of reperfusion in various tissues by reducing the activity of mediators of inflammation and oxidative stress induced by the reperfusion process [5].

Copyright: Lukiswanto, et al. Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Unfortunately, there have not been a lot of researchers aimed to know the appropriate time for the administration of HBOT in patients with various organ function disorders after reperfusion, especially its impact on liver organ. In patients with stroke, HBOT is known to be effective if given exactly 3 h after reperfusion since if given more than 3 h, it will not give any effect [6]. This research aimed to investigate effects of time-based HBOT administration on liver morphofunctional state of rabbits (*Oryctolagus cuniculus*) as animal models after hind limb reperfusion.

## Materials and Methods

### Ethical approval

This research was conducted under the ethical use of laboratory animals and also was approved by the Ethical Committee of Faculty of Veterinary Medicine, Universitas Airlangga.

### Research sites

This research was conducted at the Faculty of Veterinary Medicine, Universitas Airlangga, and also at the Department of Underwater and Hyperbaric Medicine, Hospital Dr. Ramelan, Surabaya. Preparations for histopathological examination and blood tests were made in the Department of Pathology and Veterinary Hospital, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya.

### Experimental animals

Experimental animals used in this research were 24 male New Zealand rabbits aged 6 months old and weighed 2-2.5 kg. Most studies use male animal models because of their stable hormonal status during treatment, and these animals are not at risk of being affected by hormonal (estrogen/progesterone) fluctuation related to the reproductive cycle that may confound the results of the study. Reperfusion injury using the rabbit as animal model showed significantly less infarct in female hearts as compared to males [7].

### Research materials

Materials used in this research were mineral water and food for rabbits, 70% alcohol, 10% ketamine, 2% xylazine hydrochloride, atropine sulfate, sterile thump, 10% buffered formalin, materials for staining hematoxylin-eosin (HE), materials for aspartate aminotransferase (AST) and alanine aminotransferase (ALT) examinations, and materials for the euthanasia of rabbits at the end of the research.

### Research tools

Tools used in this research were rabbit hutches, minor surgery equipment, masks, gloves, 10 cc syringe, syringes 3 cc, silastic band, hyperbaric chamber, a place to store organs, light microscope, object, cover glass, as well as blood container tube without anticoagulants.

### Experimental animal preparation

Those rabbits were adapted for 1 week before the research. They were fed and given to drink *ad libitum*.

Next, femoral artery ligation was conducted on all the experimental animals. They were then divided into four groups, namely, G1, G2, G3, and G4.

Afterward, before anesthesia was conducted for surgery, all those rabbits were fasted for 12 h. Next, anesthesia was performed by weighing each rabbit and then injecting them with 0.02 mg/kg of atropine sulfate intramuscularly. After 10 min, anesthesia was carried out by injecting a combination of ketamine-xylazine with doses of 35 mg/kg and 5 mg/kg intramuscularly [7].

Although 30-50% of rabbits produce endogenous atropinase enzymes, because using atropine sulfate 0.1-0.2 mg/kg bw/sc, im. in some studies often helpful to decrease salivary production, bronchial secretion and preventing the occurrence of vagal which can cause bradycardia [8].

After all of those rabbits were anesthetized, an incision was made in the groin area of their sinister. Next, dissection was performed to separate the femoral artery from the nerve and the vein. Ischemia was induced by ligating the femoral artery with silastic band for 6 h, then followed with reperfusion by opening or removing the ligation to restore normal blood circulation in accordance with the treatment group.


Group 1 (G1): Femoral artery ligation for 6 h, followed with reperfusion by releasing ligation for 100 min (IR 100 min).

Group 2 (G2): Ligation for 6 h, followed with reperfusion by releasing ligation for 100 min. HBOT was given 10 min after the reperfusion for 90 min (IR 100 min with HBOT 90 min).

Group 3 (G3): Ligation for 6 h, followed with reperfusion by releasing ligation for 150 min (IR 150 min).

Group 4 (G4): Ligation for 6 h, followed with reperfusion by releasing ligation for 150 min. Hyperbaric therapy was given 60 min after the reperfusion for 90 min (IR 150 min with HBOT 90 min).

At the end of the research, all the rabbits were sacrificed. Their liver and blood were taken for examining histopathological changes in their liver using HE staining as well as for measuring the levels of AST and ALT. Histopathological changes were evaluated to assess the degree of liver damage as a result of the treatment given. Scoring method used in this research was a modification of Knodell method [9]. The type



**Figure-1:** Congestion on group 3.

of the lesions observed and scoring technique used are as following:

Type of lesions	Score	Note
Necrosis	0	No necrosis
	1	Necrosis on 1/3 of lobular
	3	Necrosis on 1/3-2/3 of lobular
	4	Necrosis on >2/3 of lobular
	5	Necrosis on 1/3-2/3 of lobular+bridging necrosis
	6	Necrosis on >2/3 of lobular+bridging necrosis
Degeneration	10	Multilobular necrosis
	0	No Degeneration
	1	Degeneration on 1/3 of lobular
	3	Degeneration on 1/3-2/3 of lobular
Portal inflammation	4	Degeneration on >2/3 of lobular
	0	No inflammation
	1	Light inflammation on 1/3 of lobular
Congestion	3	Mild inflammation on 1/3-2/3 of lobular
	4	Severe inflammation on >2/3 of lobular
	0	No hemorrhage
Congestion	1	Light (hemorrhage on 1/3 of lobular)
	2	Mild (hemorrhage on 1/3-2/3 of lobular)
	3	Severe (hemorrhage on >2/3 of lobular)

### Results

Scoring of histopathological changes in the livers was focused on necrotic lesions, degeneration, portal inflammation, and congestion. The results showed that within groups without hyperbaric therapy (G1 and G3), the scores of necrosis, degeneration, and portal inflammation were not significantly different ( $p>0.05$ ). Similarly, within groups with hyperbaric therapy, namely, G2 and G4, the scores of necrosis, degeneration, and portal inflammation were not significantly different ( $p>0.05$ ). However, the scores of histopathological changes in G1 and G3 were significantly different from those in G2 and G4 ( $p<0.05$ ).

Moreover, scoring results of congestion degree had a different pattern, in which there was no congestion in G1, G2, and G4. Unlike the other three groups, congestion only appeared in G3 ( $p<0.05$ ) (Figure-1). The degree of the total score (necrosis, degeneration, and portal inflammation) in the group without hyperbaric therapy was higher than in the group with hyperbaric therapy (Figures-2-4 and Table-1).

Differences in the scoring results of histopathological changes between the groups without hyperbaric therapy, furthermore, did not seem affected by the length of reperfusion duration. For instance, there was no difference in histopathological scoring results and AST/ALT levels between Group G2 and Group G4 although both had different reperfusion duration, namely, 10 and 60 min before the administration of hyperbaric therapy.

In addition, results of the analysis of AST and ALT levels in each treatment group also showed a different pattern of results derived from the scoring




Figure-2: Degeneration of hepatocyte on each group.




Figure-3: Necrosis of hepatocyte on each group.




Figure-4: Portal inflammation on each group.

results of histopathological changes in the livers. The results showed that the levels of AST and ALT in the groups without hyperbaric therapy (G1 and G3) were not significantly different from those in the groups treated with hyperbaric therapy (G2 and G4) ( $p>0.05$ ) (Table-2). The levels of AST/ALT in all treatment groups, nevertheless, were still within normal values of AST/ALT levels in rabbits.

**Table-1:** Scoring results of histopathological changes in the livers (mean±SD) of each treatment group.

Treatment group	Necrosis	Degeneration	Inflammation portal	Congestion
G1	1.2 <sup>b</sup> ±0.2	1.0 <sup>b</sup> ±0.0	1.1 <sup>b</sup> ±0.6	0.0 <sup>a</sup> ±0.0
G2	0.8 <sup>a</sup> ±0.2	0.3 <sup>a</sup> ±0.1	0.5 <sup>a</sup> ±0.1	0.0 <sup>a</sup> ±0.0
G3	1.3 <sup>b</sup> ±0.17	0.9 <sup>b</sup> ±0.5	1.1 <sup>b</sup> ±0.6	0.3 <sup>b</sup> ±0.1
G4	1.0 <sup>a</sup> ±0.00	0.5 <sup>a</sup> ±0.1	0.6 <sup>a</sup> ±0.0	0.0 <sup>a</sup> ±0.0

Means with different superscripts letters are significant at  $p < 0.05$ . SD=Standard deviation

**Table-2:** Levels of AST and ALT in each group.

Treatment group	AST (IU/L)	ALT (IU/L)
G1	36.76 <sup>a</sup> ±3.11	12.60 <sup>a</sup> ±0.67
G2	38.08 <sup>a</sup> ±3.11	13.07 <sup>a</sup> ±0.67
G3	42.32 <sup>a</sup> ±3.11	12.05 <sup>a</sup> ±0.67
G4	40.07 <sup>a</sup> ±3.11	12.30 <sup>a</sup> ±0.67

Means with different superscripts letters are significant at  $p < 0.05$ . AST=Aspartate aminotransferase, ALT=Alanine aminotransferase

## Discussion

Hepatic ischemia-reperfusion injury is a multifactorial pathophysiological process. Numerous studies have shown that this process includes excessive intracellular Ca, increased free oxygen radical, and increased of apoptosis or necrotic liver parenchymal cells [10,11]. The presence of oxidatives stress caused disturbances in the cell cycle and initiates apoptosis [12].

In this research, necrotic, degenerative, and inflammatory lesions on those livers were not accompanied with changes in the function of those livers by examining the levels of AST/ALT. This may be caused by the degree of hepatocyte damage classified into mild to moderate one. In addition, it can also be caused due to compensatory mechanism owned by those livers, where healthy hepatocytes would take over the function of hepatocytes suffering from the injury.

Similarly, a previous research also showed that limb ischemia suffered by mice for 2 h, then followed by reperfusion can cause injury to their lungs, but not in their liver and kidneys [13]. Unlike in the previous research, other researchers have also shown that limb ischemia suffered by mice for 3 h, then followed by reperfusion can cause injury to their lungs and kidneys, as well as the muscle of their lower limb [14,15].

Studies conducted directly with hepatic ischemia-reperfusion injury (HIRI) showed that this treatment cause hepatocytes apoptosis accompanied by elevate levels of AST and ALT, reflect severe hepatocyte damage [10,11]. These facts support an assumption that many negative effects of IR can trigger ischemia injury to vital organs near the ischemic area. Therefore, it can certainly be said that the improvement in the histopathological feature of G2 and G4 is not due to self-limited, but due to HBOT therapy given since IR was conducted 6 h after the induction of ischemia.

Vital organs, including liver, are very sensitive to ischemia that occurs on injury due to IR.

Ischemia-reperfusion is usually accompanied by oxidative stress and then continued with stimulation of inflammatory response, eventually leading to injury to the liver [16]. Oxidation and inflammatory are mediators for the occurrence of injury in various organs after IR, which will end with a dysfunction of the organs, including liver [17].

Various factors, such as oxygen free radicals, migration and activation, abnormalities in the micro-circulation, damage to endothelial cells sinusoid, coagulation cascade activation, and Kupffer cell activities to release inflammatory mediators and proteolytic enzymes, plays a role in causing injury to the liver due to IR [18,19]. Lesions on the liver caused by IR can occur in two phases, causing damage to hepatocytes. First, acute phase is characterized by the production of ROS by Kupffer cells, causing mild injury in the hepatocytes. In the next phase, there will be infiltration of leukocytes to the area of injury as a form of inflammatory cascade commencement. Neutrophil accumulation in the liver may cause injury to hepatocytes through oxidants and proteases released by neutrophils [20]. ROS after reperfusion in the initial phase is considered to have an important role in initiating and aggravating oxidative stress in the liver after reperfusion [21].

In this research, furthermore, HBOT could improve morphological function of liver, namely, in Group G2 and Group G4. On the other hand, according to Kihara [22], HBOT can trigger increased activity of neutrophils, while neutrophil infiltration in the organ will lead to dysfunction. In such conditions, HBOT can inhibit the activity of neutrophils, and the dysfunction of organs caused.

Da Silveira *et al.* [23] stated that the supply of oxygen after reperfusion can act as a protector of mitochondrial membrane. Oxygen is an agent that plays a role in the formation of ATP by oxidative phosphorylation. The energy generated is very necessary for the flow of both various ions out of the mitochondria as well as liquid into the mitochondria through the mitochondrial membrane. This process then will help to decrease the production and release of ROS, so the mitochondria can function normally. Therefore, an imbalance in this process can make oxidative damage become more extensive. This similar condition was also found in Group G1 and Group G3, experiencing ischemia in the same period without the administration of HBOT.

Biochemistry profile parameters, including AST/ALT, in rabbits, were influenced by various

factors such as breed, age, sex, feed, liver disease, environmental conditions, stress, pregnancy, and heart rhythm [24-27]. Nevertheless, in this research, there was no difference in AST/ALT level between the rabbits without treatment and the rabbits with hyperbaric-treatment. In other words, all of those rabbits had no liver function impairment. Research on pigs was subjected left hemihepatectomy suggests that portal vein arterialization promotes early and improved liver regeneration without impairing liver function [28].

### Conclusion

Hind limb ischemia injury reperfusion can trigger a damage for liver morphology, but not lead to liver dysfunction. HBOT for 90 min, 10-60 min after the reperfusion, can protect hepatocytes from damage.

### Authors' Contributions

BSL and YM carried out the main research works, WMY performed the statistical analysis and analyzed the main data in the experiments and BSL, WMY and YM approved the final manuscript. All authors read and approved the final manuscript.

### Acknowledgments

The authors are very thankful to Faculty of Veterinary Medicine, Universitas Airlangga; Department of Surgery, Division of Thoracic, Cardiac, and Vascular Surgery, Dr. Soetomo General Hospital, Surabaya; and the Department of Underwater and Hyperbaric Medicine, Dr. Ramelan Naval Hospital, Surabaya, for providing all the type of facilities to conduct this research.

### Competing Interests

The authors declare that they have no competing interests.

### References

- Fiolet, J.W.T. and Baartscheer, A. (2000) Update review cellular calcium homeostasis during ischemia: A thermodynamic approach. *Cardiovasc. Res.*, 45: 100-106.
- Neil-Granger, D. and Kvietys, P.R. (2015) Reperfusion injury and reactive oxygen species: The evolution of a concept. *Redox Biol.*, 6: 524-551.
- Jaffer, U., Wade, R.G. and Gourlay, T. (2010) Cytokines in the systemic inflammatory response syndrome: A review. *HSR Proc. Intensive Care Cardiovasc. Anesth.*, 2(3): 161-175.
- Gyurkovics, E., Aranyi, P., Stangl, R., Onody, P., Ferreira, G., Lotz, G., Kupcsulik, P. and Szijarto, A. (2011) Post conditioning of the lower limb-protection against the reperfusion syndrome. *J. Surg. Res.*, 169(1): 139-147.
- Bozok, S., Ilhan, G., Yilmaz, Y., Dokumcu, Z., Tumkaya, L., Karamustafa, H., Ozan, K., Karakisi, S., Ergene, S. and Sener, E. (2012) Protective effects of hyperbaric oxygen and iloprost on ischemia/reperfusion-induced lung injury in a rabbit model. *Eur. J. Med. Res.*, 17(1): 14.
- McCormick, J.G., Houle, T.T., Saltzman, H.A., Whaley, R.C. and Roy, R.C. (2011) Treatment of acute stroke with hyperbaric oxygen: Time window for efficacy. *Undersea Hyperb. Med.*, 38: 321-334.
- Mahmoodzadeh, S., Fliegner, D. and Dworatzek, D. (2012) Sex differences in animal models for cardiovascular diseases and the role of estrogen. *Handb. Exp. Pharmacol.*, 214: 23-48.
- Flecknell, P. (2009) *Laboratory Animal Anaesthesia*. 3<sup>rd</sup> ed. Academic Press, USA. p204-210.
- Brunet, E. (2000) Grading and staging the histopathological lesions of chronic hepatitis: The knodell histology activity index and beyond. *Hepatology*, 31: 241-246.
- Yang, J., Sun, H., Takacs, P., Zhang, Y., Liu, J., Chang, Y. and Candioti, K.A. (2013) The effect of octreotide on hepatic ischemia-reperfusion injury in a rabbit model. *Transplant. Proc.*, 45: 2433-2438.
- Weigand, K., Brost, S., Steinebrunner, N., Buchler, M., Schemmer, P. and Muller, M. (2012) Ischemia reperfusion injury in liver surgery and transplantation: Pathophysiology. *HPB Surg.*, 2012: 1-9.
- Vali, L., Taba, G., Szentmihalyi, K., Febel, H., Kurucs, T., Pallai, Z., Kupcsulik, P. and Blazovics, A. (2006) Reduced antioxidant level and increased oxidative damage in intact liver lobes during ischemia reperfusion. *World J. Gastroenterol.*, 21: 1086-1091.
- Mansour, Z., Charles, A.L., Kindo, M., Pottecher, J., Chamaraux-Tran, T.N., Lejay, A., Zoll, J., Mazzucotelli, J.P. and Geny, B. (2014) Remote effects of lower limb ischemia reperfusion: Impaired lung, unchanged liver, and stimulated kidney oxidative capacities. *Biomed. Res. Int.*, 4: 390-392.
- Kao, M.C., Yang, C.H., Chou, W.C., Sheu, J.R. and Huang, C.J. (2015) Cepharanthine mitigates lung injury in lower limb ischemia-reperfusion. *J. Surg. Res.*, 199: 647-656.
- Kao, M.C., Chung, C.Y., Chang, Y.Y., Lin, C.K., Sheu, J.R. and Huang, C.J. (2015) Salutary effects of cepharanthine against skeletal muscle and kidney injuries following limb ischemia/reperfusion. *Evid. Based Complement. Alternat. Med.*, 50: 40-46.
- Duehrkop, C., De noyelle, J., Shaw, S. and Rieben, R. (2014) Use of dextran sulfate in tourniquet-induced skeletal muscle reperfusion injury. *J. Surg. Res.*, 187: 150-161.
- Chang, Y.K., Huang, S.C., Kao, M.C. and Huang, C.J. (2016) Cepharanthine alleviates liver injury in a rodent model of limb ischemia-reperfusion. *Acta Anaesthesiologica Taiwanica*, 54: 11-15.
- Serracino-Inglott, F., Habib, N.A. and Mathie, R.T. (2001) Hepatic ischemia-reperfusion injury. *Am. J. Surg.*, 181: 160-166.
- Giakoustidis, D., Papageorgiou, G., Iliadis, S., Kontos, N., Kostopoulou, E., Papachrestou, A., Tsantilas, D., Spyridis, C., Takoudas, D., Botsoglou, N., Dimitriadou, A. and Giakoustidis, E. (2002) Intramuscular administration of very high dose of alpha-tocopherol protects liver from severe ischemia/reperfusion injury. *World J. Surg.*, 26: 872-877.
- Okaya, T. and Lentsch, A.B. (2003) Cytokine cascades and the hepatic inflammatory response to ischemia and reperfusion. *J. Invest. Surg.*, 16(3): 141-143.
- Furukawa, H., Hayashi, T. and Todo, S. (2000) A novel hydroxyl radical scavenger, nicaraven, protects the liver from warm ischemia and reperfusion injury. *Surgery*, 127: 661-669.
- Kihara, K., Ueno, S., Sakoda, M. and Aikou, T. (2005) Effects of hyperbaric oxygen exposure on experimental hepatic ischemia reperfusion injury: Relationship between its timing and neutrophil sequestration. *Liver Transplant.*, 11: 1574-1580.
- Silveira, M.R.G., Margarido, M.R., Vanni, J.C., Junior, R.N. and Silva, O. (2014) Effects of hyperbaric oxygen therapy on the liver after injury caused by the hepatic ischemia-reperfusion process. *Acta Cirúrgica Brasileira*, 29(1): 29-33.
- Chineke, C.A., Ologun, A.G. and Ikeobi, C.O.N. (2006) Haematological parameters in rabbit breeds and crosses in humid tropics. *Pak. J. Biol. Sci.*, 9: 2102-2106.

25. Melillo, A. (2007). Rabbit clinical pathology. *J. Exot. Pet Med.*, 16: 135-145.
26. Jeklova E., Leva L., Knotigova P. and Faldyna, M. (2009) Age-related changes in selected haematology parameteres in rabbits. *Res. Vet. Sci.*, 86: 525-528.
27. Abdel-Azeem, A.S., Abdel-Azim, A.M., Darwish, A.A. and Omar E.M. (2010) Haematological and biochemical observations in four pure breeds of rabbits and their crosses under Egyptian environmental conditions. *World Rabbit Sci.*, 18: 103-110.
28. Ott, R., Schuppan, D., Tannapfel, A., Wittekind, C., Erhardt, W., Henke, J., Kilic, N., Kockerling, F., Reck, T., Hohenberger, W. and Muller, W. (2003) Portal vein arterialization as a technical option in liver transplantation: Impact on function, regeneration, and morphology of the liver following hemihepatectomy in pigs. *Liver Int.*, 23: 54-62.

\*\*\*\*\*