

Brotowali Extract (*Tinospora Crispa*) for Oral Traumatic Ulcer in diabetes Mellitus Wistar Rat.

by Retno Indrawati.,drg.,msi

Submission date: 13-Jun-2018 02:09PM (UTC+0800)

Submission ID: 975271268

File name: pa_for_Oral_Traumatic_Ulcer_in_diabetes_Mellitus_Wistar_Rat..pdf (481.08K)

Word count: 3670

Character count: 19826

Brotowali Extract (*Tinospora Crispa*) for Oral Traumatic Ulcer in Diabetes Mellitus Wistar Rat

Retno Indrawati Roestamadji¹, Ira Arundina¹, Indeswati Diyatri¹,
Dewi Tamara Sambodo², Wahyuning Ratih Irmalia³

1. Lecturer of Departement of Oral Biology, Faculty of Dental Medicine, Universitas Airlangga, Surabaya, Indonesia.
2. Undergraduate Student, Faculty of Dental Medicine, Universitas Airlangga, Surabaya, Indonesia.
3. Graduate Student, Faculty of Dental Medicine, Universitas Airlangga, Surabaya, Indonesia.

Abstract

One of the diabetes complication is delayed wound healing in oral traumatic ulcer. *Tinospora crispa* contains flavonoid and terpenoid which can help to control blood glucose level and accelerate wound healing.

The aim of this study is to analyze brotowali (*Tinospora crispa*) extract potential in controlling blood glucose level and accelerating ulcer healing process. The extract was made with oven-dried method at 50°C and macerated with ethanol 80% (1:10 w/v). The rats were divided into 3 groups, control group I (K1) normal Wistar Rat with traumatic ulcer, control group II (K2) diabetes mellitus wistar rat with traumatic ulcer, and treatment group (P) diabetes mellitus wistar rat with traumatic ulcer treated with 250 mg/kg *tinospora crispa* extract once a day. Sample were euthanized on 3rd, 5th, and, 7th day after traumatic ulcer was made then a histopathology preparation was made to count fibroblast cell. Blood glucose level measurement was conducted on 3rd, 5th, 7th, and 14th after traumatic ulcer was made.

Brotowali extract can affect blood glucose levels but make no significant difference and affect the number of fibroblasts in traumatic ulcers healing diabetes mellitus Wistar rats with significant difference.

Experimental article (J Int Dent Med Res 2017; 10(3): pp. 991-996)

Keywords: Traumatic ulcer, Wound healing, *Tinosporacrispa*, Diabetes mellitus, Fibroblast cell.

Received date: 31 August 2017

Accept date: 13 October 2017

Introduction

Diabetes mellitus defined as a multi factorial disease or chronic metabolism disorder, characterized by elevated blood glucose level, carbohydrate, lipids and protein metabolism disturbance in consequence of insulin insufficiency. Insulin insufficiency may be caused by disturbance or insufficient insulin production by beta Langerhans cells of pancreas¹.

Indonesia is the fourth country that has the highest diabetes mellitus prevalence². According to report from Basic Health Research (Riskesdas) 2013, the prevalence of diabetes mellitus in Indonesia is 6,9% or equivalent to 12,1 million patients³. According to research by Diabcare, type 2 uncontrolled diabetes mellitus

patient reached 47,2% with plasma glucose level >130mg/dL⁴.

Diabetes Mellitus have various complications related to dentistry, one of them is traumatic ulcer^{5,6}. Traumatic ulcer is single mucosa ulcer that can be caused by mechanic, thermal, chemical, or direct physical trauma to the mucosa^{7,8}. Traumatic ulcers were more common in diabetes mellitus patient than in normal person. This may be caused by wound healing impairment⁹. In patients with diabetes mellitus, many aspects are disrupted during wound healing process, such as inflammatory response dysfunction, reduced granulation tissue formation, angiogenesis disorders, and increased cell apoptosis¹⁰.

Brotowali is widely known as traditional medicine as antidiabetic agent and antioxidant that accelerate wound healing process¹¹. Brotowali stem contains flavonoids which have antioxidant effect better than ascorbic acid¹². Flavonoid as antioxidant can bind free radical and decrease ROS level which is advantageous for wound healing process¹³. Low ROS level will

*Corresponding author:

Retno Indrawati Roestamadji,
Departement of Oral Biology Faculty of Dental Medicine
Universitas Airlangga. Jl. Mayjen Prof. Dr. Moestopo 47 Surabaya
60132, Indonesia.
E-mail: retnoindrawati@fkg.unair.ac.id; rathirmalia@yahoo.com

lead to decrease of excessive cell apoptosis¹⁴. Moreover, flavonoid ability to activate macrophage has been proven¹². Macrophage plays an important role in wound healing as a resource of various growth factors, such as *Transforming Growth Factor (TGF-β)*, *Platelet-Derived Growth Factor (PDGF)* and *Fibroblast Growth Factor (FGF)* which play role in fibroblast proliferation¹⁵. Brotowali stem also contains terpenoid that contains borapetosid, with ability to enhance insulin receptor fosforilation, and increase protein kinase B and GLUT2 expression in hepar. This may lead to increase of peripheral utilization of glucose and decrease of gluconeogenesis in hepar^{16,17}.

Hyperglycemic condition in diabetic patient stimulate excessive glucose in blood to enter sorbitol pathway that followed by elevated ROS formation. Due to antihyperglycemic activity of brotowali, sorbitol pathway stimulation does not occur, so that ROS formation can be reduced, and prevent fibroblas apoptosis caused by ROS¹⁸. This research is conducted to analyze the effect of brotowali extract on blood glucose level and the number of fibroblasts in traumatic ulcer healing process in Wistar rats with diabetes mellitus.

Materials and methods

This was an experimental laboratory with post test only control group design. This study was approved by Ethical Comittee of Faculty of Dental Medicine Universitas Airlangga (NO 59/KKEPK.FKGM/2016).

Brotowali Plant Extraction

Fresh specimen of brotowali stem washed and cut into small pieces and dried with oven in of 50°C temperature. Dry specimen was made into powder and macerated with ethanol 80% (1:10 w/v) for 24 hours. The extract was filtered, dregs and filtrate were subsequently separated. The dregs were macerated again with new ethanol, this procedure is repeated until TLC (Thin Layer Chromatography) test with H₂SO₄ is not showing pink color. Then the dregs and the filtrate from first to the last day is collected and evaporated with a rotary evaporator to remove the solvent. Extracts were stored at temperature of -20°C¹⁹.

Preparation of Experimental Animals

Male Wistar rats 150-200g weights were collected in the same cage with a room

temperature of 25±2°C and given food pellets and a standard drink distilled water ad libitum for 7 days prior the experiment¹⁹. Animals were divided into 3 groups, the control group 1 (K1), control 2 (K2), and treatment (P). Each group contains 8 experimental animals.

Diabetes Mellitus Induction with Alloxan

³⁰ For rats in groups K2 and P1, alloxan (120 mg/kg) dissolved in 0.05 M citrate buffer pH 3 was used to induce experimental diabetes. K1 group (control group) were injected with sterile saline solution 72 hours after injection, blood from lateral vein was drawn to determine the blood glucose levels by "on call plus" ACON Santiago USA. Animals stated diabetes when blood glucose levels ≥200 mg/dL after 72 hours post-induction of alloxan^{20,21}.

Wistar rats in all treatment groups were anesthetized using injection of ketamine (Biosynth, Switzerland) (60mg/kg) and xylazine (PT Merck Chemicals & Life Sciences. Jakarta) (60mg/kg) intraperitoneal²². The buccal mucosa is sterilized with a swab moistened with a 0.12% chlorhexidine digluconate. Buccal mucosa ulceration created by wounding the mucosa using a scalpel blade no. 15 with a cross-section diameter wound 8 mm²³. Brotowali extract administered per oral (250 mg/kg) once a day for the treatment group P1 after traumatic ulcer is formed. Extracts brotowali given until day 3, 5, 7 and 14.

Euthanasia Procedure at Day 3, 5, 7, and 14 Post Traumatic Ulcer Creation

Prior sacrificed, blood glucose levels of experimental animals in all groups are checked using a glucometer. Then euthanasia of experimental animals is performed with ketamine injection of 180mg/kg and xylazine 180mg/kg. Buccal mucosa tissue removal was done in all groups and put in formalin prior to the procedure of making preparat. The number of fibroblasts cells is determined by observation of HE preparations under light microscope (Olympus, Tokyo Jepang) with 400x magnification. Fibroblasts cell calculation using ocular graticulae tool mounted on the lens on a microscope with a visual field technique conducted on three field²⁴.

Results

Level of blood glucose on day 3 in control I group is significantly lower than in

control II group and treatment group ($p < \alpha = 0.05$). On day 5, blood glucose decreased in control 1 group, increased in control group 2 and decreased in treatment group ($p < \alpha = 0.05$). On the 7th day blood glucose levels decreased in control 1 group, increased in control 2 group and treatment groups remained the same ($p < \alpha = 0.05$). On the 14th day blood glucose level increased in control 1 group and decreased in control 2 group and treatment group (Figure 1).




Figure 1. Blood glucose level diagram at day 3, 5, 7 and 14.




Figure 2. Average number of fibroblasts on day 3, 5, and 7.

On day 3, there is no significant difference ($p > \alpha = 0.05$) between the number of fibroblast cells control in control 1 group than in treatment group. On day 5, the number of fibroblasts in control 1 group slightly lower while in treatment group is higher ($p > \alpha = 0.05$). On day 7, the number of fibroblasts increased in all groups ($p > \alpha = 0.05$) (Figure 2). The number of fibroblasts in treatment group continues to increase and has the highest average among each group (Figure 3).

Based on the result of fibroblast cell count, there are no significant difference between each group in day 3 (Figure 3). The number of Fibroblast is increased in day 5, and the number of fibroblast in treatment group is significantly higher than in control I and 2 group ($p > \alpha = 0.05$) (Figure 4). On day 7, the number of fibroblast cell increase in all group, with the highest number of fibroblasts are found in treatment group ($p > \alpha = 0.05$) (Figure 5).




Figure 3. Fibroblast cells (arrows) on day 3. (A) control 1 group, (B) control 2 group and (C) treatment groups on day 3 with HE staining and 400x magnification.




Figure 4. Fibroblast cells (arrows) on day 5. (A) control 1 group, (B) control 2 group and (C) treatment groups on day 5 with HE staining and 400x magnification.




Figure 5. Fibroblast cells (arrows) on day 7. (A) control 1 group, (B) control 2 group and (C) treatment groups on day 7 with HE staining and 400x magnification.

Discussion

The average blood glucose level in treatment group decreased in each observation days and lower than blood glucose in control 2 group but the difference was not significant. This means that the blood glucose in treatment groups had not back to normal. This result is contrary with the previous research that said there was significant decrease in blood glucose by 10% at day 14 after the administration brotowali extract at a dose of 250 mg/kg¹⁹. Other studies suggested that there were significant differences on blood glucose levels on day 7 and 14 after administration brotowali extract at a dose of 250 mg/kg²⁵.

Blood glucose levels were not significantly decreased may be caused by the duration of extract administration. Brotowali has ability to reduce blood glucose because contains terpenoids that consists of borapetosid A which can enhances insulin receptor phosphorylation and protein kinase B as well as increases the expression of GLUT2 in liver. This may lead to increase of peripheral utilization of glucose and decrease of gluconeogenesis in heparsoglucose levels in blood will decrease¹⁷. To obtain significant results, duration of 60 days is required²⁶. Previous study stated no significant difference in blood glucose after administration brotowali extract²⁷. These different results might be attributed to the stability of the material, so it is important to note the time between

manufacture of extracts and time of treatment²⁸. Monitoring the concentration of active ingredients is important because it also affects the quality, effectiveness and life of natural medicines during storage²⁹. In this study, the time between manufacture of extracts and treatment is a month. This may lead to the stability of the extract brotowali reduced.

In wound healing process, fibroblast cells maintain structural integrity of connective tissue by secreting extracellular matrix precursor such as collagen, reticulum, and elastin continuously³⁰. Catechins in brotowali can accelerate wound healing process, through the intermediary role of macrophages that stimulates growth factors to increase fibroblasts proliferation. Macrophages phagocytes bacteria that can interfere wound healing process in patients with diabetes mellitus and eliminate tissue debris, then release cytokines and growth factors such as PDGF, TGF- β and FGF. Those cytokine products stimulate cell migration and fibroblasts proliferation, as well as production and modulation of extracellular matrix²⁴.

Catechins also increase the activity of superoxide dismutase and catalase. Superoxide dismutase is an enzyme that catalyzes superoxide into oxygen and hydrogen peroxide. Catalase is an enzyme that catalyzes hydrogen peroxide to water (H₂O). Other content of brotowali extract is luteolin that binds superoxide radical in the body and with catechins make antioxidant effects that reduce ROS level in the body. This leads to decrease of fibroblasts apoptosis rate¹⁰.

Wound healing process can be observed histologically by counting the number of fibroblast cells to determine cell proliferation. From the research data, there is no significant difference in the number of fibroblasts in control I group and treatment group on day 3. This is because day 3 is an early phase of traumatic ulcer healing²³. Proliferation in fibroblast cell count began to increase on day 5 and day 7. This study used Wistar rats which have healing time faster than humans and synthesis of collagen by fibroblasts begin relatively early³¹. Synthesis of collagen by fibroblasts begin relatively early in day 3-5 and continue until a few weeks depending on the size of the wound³². From the data results, the number of fibroblasts in the control 1 and 2 group is less than the treatment group. The treatment group had an average number of fibroblasts at

most in all the groups. On the 7th day fibroblast cell number continues to increase and has the highest average when compared with the control group. Supposedly at day 7 the proliferation of fibroblasts was decreased for the formation of collagen fibers. This indicates proliferation of fibroblasts still high due to brotowali extract. Excessive fibroblasts proliferation may cause a build up of collagen so that the wound will heal with scars or fibrous tissue³².

The number of fibroblasts which remained high at day 7 may be caused by circadian rhythms. Wistar rats are nocturnal rodents which active at night and relatively inactive during day³¹. Diabetes induction, extract administration and euthanasia procedures which is not suitable with rats' circadian rhythm could affect the research results. Circadian rhythm affects the immune system. One of hormones associated with circadian rhythm is cortisol, that also regulate inflammation process³³. In this research, all procedures performed at 8-12 am. There was no cortisol activity as anti-inflammatory agent during that time. Lack of cortisol lead to lack of ability to prevent severe chronic inflammation. As a result, immune system goes out of control. Low cortisol level may increase the production of pro-inflammatory cytokines, which lead to over-activation of immune system and inflammation³³. Low cortisol level may lead to the absence of a regulatory function growth factors secretion that results in excessive secretion of growth factor. In this research, high level of fibroblasts proliferation is caused by over-activation of fibroblasts during proliferation phase of healing process in the morning¹⁰.

Conclusions

This research showed that administration of brotowali (*Tinospora crispera*) extract dose of 250 mg/kg for 14 days can reduce blood glucose wistar rats with diabetes mellitus and increase the number of fibroblasts on day 3, 5 and 7 of oral traumatic ulcer healing on diabetes mellitus Wistar rats.

Acknowledgements

Acknowledgements given to Dr. Bambang. A., dr., MS medical faculty of Universitas Airlangga on guidance during the process of this research.

Declaration of Interest

The authors report no conflict of interest and the article is not funded or supported by any research grant.

References

1. Wong N, Shaista M. Diabetes and cardiovascular disease: evaluation, prevention, and management. New Delhi: Jaypee Medical Publishers (P) Ltd. 2014: 1-13.
2. Global report on diabetes. Geneva.: WHO Library Cataloguing-in-Publication Data, World Health Organization. ISBN 978 92 4 156525 7. 2016. Vol. 978.
3. Laporan hasil riset kesehatan dasar nasional 2013. Jakarta.: Departemen Kesehatan RI; 2013.
4. Soewondo P, Soegondo S, Suastika K, Pranoto A, Soeatmadji DW, Tjokroprawiro A. The DiabCare Asia 2008 study – Outcomes on control and complications of type 2 diabetic patients in Indonesia. Med J Indones. 2010;19(4):235.
5. Al-Maskari AY, Al-Maskari MY, Al-Sudairy S. Oral manifestations and complications of diabetes mellitus: A review. Sultan Qaboos University Medical Journal. 2011;11: 179–86.
6. Kaomongkolgit R, Wongviriyi A, Daroonpan P, Chansamat R, Tantanapornkul W, Palasuk J. Denture stomatitis and its predisposing factors in denture wearers. J Int Dent Med Res. 2017;10(1):89–94.
7. Greenberg M. Burket's Oral Medicine. 12th ed. Glick M, editor. Connecticut: People's Medical Publishing House. 2015.86.
8. Arundina I, Soesilawati P, Damaiyanti DW, Maharani D. The effects of golden sea cucumber extract (*Stichopus hermanii*) on the number of lymphocytes during the healing process of traumatic ulcer on wistar rat's oral mucous. Dent J (Majalah Kedokt Gigi). 2015;100(56):100–3.
9. Saini R, Al-Maweri SA, Saini D, Ismail NM, Ismail AR. Oral mucosal lesions in non oral habit diabetic patients and association of diabetes mellitus with oral precancerous lesions. Diabetes Res Clin Pract. 2010;89(3):320–6.
10. Desta T, Li J, Chino T, Graves DT. Altered fibroblast proliferation and apoptosis in diabetic gingival wounds. J Dent Res. 2010;89(6):609–14.
11. Koay YC, Amir F. A review of the secondary metabolites and biological activities of *Tinospora crispera* (Menispermaceae). Tropical Journal of Pharmaceutical Research. 2013;12:641–9.
12. Abood WN, Fahmi I, Abdulla MA, Ismail S. Immunomodulatory effect of an isolated fraction from *Tinospora crispera* on intracellular expression of INF- γ , IL-6 and IL-8. BMC Complement Altern Med. 2014;14(1):205.
13. Mirjana P, Jasmin F, Vera R-N, Cena D, Lidija P, Aneta M, et al. The effects of nbf gingival gel in the treatment of recurrent oral ulcers: case report. J Int Dent Med Res. 2016;9(1):81–5.
14. Graves DT, Liu R, Alikhani M, Al-Mashat H, Trackman PC. Diabetes-enhanced inflammation and apoptosis–impact on periodontal pathology. J Dent Res. 2006;85(1):15–21.
15. Hom D, Patricia A, Arun K, Craig D. Essential tissue healing of head and neck. Connecticut: People's Medical Publishing House. 2009:388-95.
16. Lam SH, Ruan CT, Hsieh PH, Su MJ, Lee SS. Hypoglycemic diterpenoids from *Tinospora crispera*. J Nat Prod. 2012;75(2):153–9.
17. Ruan CT, Lam SH, Chi TC, Lee SS, Su MJ. Borapetoside C from *Tinospora crispera* improves insulin sensitivity in diabetic mice. Phytomedicine. 2012;19(8–9):719–24.
18. Chung SSM. Contribution of Polyol Pathway to Diabetes-Induced Oxidative Stress. J Am Soc Nephrol. 2003;14(90003):233S–236.
19. Talubmook C, Nopparat B. Bioactivities of Extract from *Tinospora crispera* stems, *Annona squamosa* Leaves, *Musa sapientum* Flowers, and *Piper Sarmentosum* Leaves in Diabetic Rats. Int J Adv Res Technol. 2013;2(6):1–6.

- 4
20. Asgary S, Rafeian-Kopaei M, Shamsi F, Najafi S, Sahebkar A. Biochemical and histopathological study of the anti-hyperglycemic and anti-hyperlipidemic effects of cornelian cherry (*Cornus mas L.*) in alloxan-induced diabetic rats. *J Complement Integr Med.* 2014;11(2):63–9.
21. Purwanto B, Paulus L. Model hewan coba untuk penelitian diabetes. Surabaya: PT Revka Petra, Media. 2014:9-11. 6
22. Brizenzo L, Ana M, Ana P, Fabricio B, Paulo G, Suzana C, et al. Delayed healing of oral mucosa in a diabetic rat model: Implication of TNF- α , IL-1 β and FGF-2. *Life Sci.* 2016;155:36–47.
23. Cavalcante G, Sousa R, Peres L, Sousa F, Lima M. Experimental model of traumatic ulcer in the cheek mucosa of rats. *Acta Cirúrgica Bras.* 2011;26(3):227–34.
24. Pradita A, Agung P, Catur A, Ali T. Periodontal dressing-containing green tea epigallocatechin gallate increases fibroblasts number in gingival artificial wound Model. *J Dent Indones.* 2013;20(3):68–72.
25. Arcueno R, Jinger L, Retumban, Joycelyn E, Jonathan J. Wound healing potential of *Tinospora Crispa* (Willd.) Miers [Menispermaceae] stem on diabetic mice. *J Med Plants Stud.* 2015;3(2):106–9.
26. Sharma R, Amin H, Galib, Prajapati PK. Antidiabetic claims of *Tinospora cordifolia* (Willd.) Miers: critical appraisal and role in therapy. *Asian Pac J Trop Biomed.* 2015;5(1):68–78.
27. Klangjareonchai T, Roongpisuthipong C. The effect of *tinospora crispa* on serum glucose and insulin levels in patients with type 2 diabetes mellitus. *J Biomed Biotechnol.* 2012;2012:808762.
28. Daburkar M, Vikram L, Arvind S, Pravin B, Shrikant T. An in vivo and in vitro investigation of the effect of Aloe vera gel ethanolic extract using animal model with diabetic foot ulcer. *J Pharm Bioallied. Sci* 2013;1:1–9.
29. Thakur L, Ghodasra U, Patel N, Dabhi M. Novel approaches for stability improvement in natural medicines. *Pharmacogn Rev.* 2011;5(9):48–54.
30. Velnar T, Bailey T, Smrkolj V. The wound healing process : an overview of the cellular and molecular mechanisms. *J Int Med Res.* 2009;37(5):1528–42.
31. Samuelson D. Textbook of veterinary histology. St. Louis: Saunders Elsevier. 2007. 71.
32. Deyhimi P, Khademi H, Birang R, Akhoondzadeh M, Author C. Histological evaluation of wound healing process after photodynamic therapy of rat oral mucosal ulcer. *J Dent Shiraz Univ Med Sci J Dent Shiraz Univ Med Sci.* 2016;17(171):43–8.
33. Evans JA, Davidson AJ. Health consequences of circadian disruption in humans and animal models. *Progress in Molecular Biology and Translational Science.* 2013;119:283–323.

Brotowali Extract (*Tinospora Crispa*) for Oral Traumatic Ulcer in diabetes Mellitus Wistar Rat.

ORIGINALITY REPORT

20%

SIMILARITY INDEX

15%

INTERNET SOURCES

14%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

1	"Oral Manifestations in Diabetes", Principles of Diabetes Mellitus, 2016. Publication	1%
2	posnf.sites.uff.br Internet Source	1%
3	eprints.lse.ac.uk Internet Source	1%
4	dept-biotech.mums.ac.ir Internet Source	1%
5	www.biomeddefine.com Internet Source	1%
6	Fawaz Alzoubi, Bobby Joseph, Lars Andersson. "Healing of soft tissue lacerations in diabetic-induced rats", Dental Traumatology, 2017 Publication	1%
7	www.ajol.info Internet Source	1%

Sami K. Saikaly, Amor Khachemoune. "Honey

8	and Wound Healing: An Update", American Journal of Clinical Dermatology, 2017	1%
Publication		
9	www.labome.com	1%
Internet Source		
10	Submitted to Misr International University	1%
Student Paper		
11	www.hoajonline.com	1%
Internet Source		
12	repositorio.uchile.cl	1%
Internet Source		
13	He, Jingsong, Ni Xie, Jianbo Yang, Hong Guan, Weicai Chen, Huisheng Wu, Zishan Yuan, Kun Wang, Guojin Li, Jie Sun, and Limin Yu. "siRNA-Mediated Suppression of Synuclein γ Inhibits MDA-MB-231 Cell Migration and Proliferation by Downregulating the Phosphorylation of AKT and ERK", Journal of Breast Cancer, 2014.	1%
Publication		
14	Lu-Lu Xu, Feng-Xia Guo, Sen-Sen Chi, Zi-Jian Wang, Yan-Yan Jiang, Bin Liu, Jia-Yu Zhang. "Rapid Screening and Identification of Diterpenoids in <i>Tinospora sinensis</i> Based on High-Performance Liquid Chromatography Coupled with Linear Ion Trap-Orbitrap Mass	1%

Spectrometry", *Molecules*, 2017

Publication

15

Zhang, Li-Ying. "Correlation between ¹³C-urea breath test value and severity of gastric mucosal lesions", *World Chinese Journal of Digestology*, 2013.

Publication

1%

16

Ye Zhang, Hong Deng, Zhouping Tang. "Efficacy of Cellular Therapy for Diabetic Foot Ulcer", *Cell Transplantation*, 2018

Publication

1%

17

www.irjpas.com

Internet Source

1%

18

clinicaltrials.gov

Internet Source

1%

19

wrap.warwick.ac.uk

Internet Source

<1%

20

Kibe, Toshiro, Takashi Koga, Kazuhide Nishihara, Takao Fuchigami, Takuya Yoshimura, Tetsushi Taguchi, and Norifumi Nakamura. "Examination of the early wound healing process under different wound dressing conditions", *Oral Surgery Oral Medicine Oral Pathology and Oral Radiology*, 2016.

Publication

<1%

21

www.ijoart.org

Internet Source

<1%

22

content.karger.com

Internet Source

<1%

23

www.jdentistry.ui.ac.id

Internet Source

<1%

24

www.omicsonline.org

Internet Source

<1%

25

Submitted to Westside High School

Student Paper

<1%

26

H. N. Santosh, Chaya M. David. "Role of ascorbic acid in diabetes mellitus: A comprehensive review", Journal of Medicine, Radiology, Pathology & Surgery, 2017

Publication

<1%

27

cdn.intechopen.com

Internet Source

<1%

28

Submitted to Grand Canyon University

Student Paper

<1%

29

ejournal.umm.ac.id

Internet Source

<1%

30

www.icsoffice.org

Internet Source

<1%

31

S Gupta, S Ghosh, S Gupta. "Interventions for

the management of oral lichen planus: a review of the conventional and novel therapies", Oral Diseases, 2017

Publication

<1%

32

www.ukessays.com

Internet Source

<1%

33

Kudo, Tetsuji, Takayuki Matsumoto, Ikuo Nakamichi, Shinichiro Yada, Motohiro Esaki, Yukihiro Jo, Yutaka Ohji, Takashi Yao, and Mitsuo Iida. "Recombinant human granulocyte colony-stimulating factor reduces colonic epithelial cell apoptosis and ameliorates murine dextran sulfate sodium-induced colitis", Scandinavian Journal of Gastroenterology, 2008.

Publication

<1%

34

bksikmikpikkfki.net

Internet Source

<1%

35

Sio-Hong Lam, Chi-Tun Ruan, Po-Hung Hsieh, Ming-Jai Su, Shoei-Sheng Lee. " Hypoglycemic Diterpenoids from ", Journal of Natural Products, 2012

Publication

<1%

Exclude bibliography Off

Brotowali Extract (Tinospora Crispa) for Oral Traumatic Ulcer in diabetes Mellitus Wistar Rat.

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
