

PAPER • OPEN ACCESS

Effects of hyperbaric oxygen therapy in enhancing expressions of e-NOS, TNF- α and VEGF in wound healing

To cite this article: Imam Susilo *et al* 2017 *J. Phys.: Conf. Ser.* **853** 012030

View the [article online](#) for updates and enhancements.

Related content

- [Diving Centre's Hyperbaric Complex Modernization for Medical Recovery and Treatment Throughout Hyperbaric Oxygen Therapy](#)
Mihai Diaconu, Simona Rus, Ctlin Fril *et al.*
- [Application of 670 nm InGaP laser and microcurrent favors the healing of second-degree burns in Wistar rats](#)
M Belli, C R Fernandes, L M G Neves *et al.*
- [Effects of microcurrent application and 670 nm InGaP low-level laser irradiation on experimental wound healing in healthy and diabetic Wistar rats](#)
L M G Neves, R L Matheus, G M T Santos *et al.*



IOP | ebooks™

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

Effects of hyperbaric oxygen therapy in enhancing expressions of e-NOS, TNF- α and VEGF in wound healing

Imam Susilo^{1*}, Anita Devi², Azham Purwandhono³ and Sunaryo Hadi Warsito⁴

¹Department of Pathology, Faculty of Medicine, Airlangga University, Surabaya, Indonesia

²Hyperbaric Division of Sanglah Hospital, Denpasar, Bali, Indonesia

³Department of Pathology, Faculty of Medicine, Jember University, Jember, Indonesia

⁴Department of Animal Husbandry, Faculty of Veterinary Medicine, Airlangga University, Surabaya, Indonesia

*Corresponding email address : imam_susilo@fk.unair.ac.id

Abstract. Wound healing is a physiological process that occurs progressively through overlapping phases. Tissue oxygenation is an important part of the complex regulation for wound healing. Hyperbaric Oxygen (HBO) therapy is a method of increasing oxygen delivery to tissues. The therapy improves tissue oxygenation and stimulates the formation of H₂O₂ as a secondary messenger for Tumour Necrosis Factor alpha (TNF α), e-NOS, VEGF and Nuclear Factor Kappa Beta phosphorylation (NF-Kb) which play an important role in the rapid transcription of a wide variety of genes in response to extracellular stimuli. This study aims to determine the effects of Hyperbaric Oxygen therapy in enhancing the expressions of e-NOS, TNF- α , VEGF and wound healing. This study is an animal study with a 'randomized control group of pre-test and post test design' on 28 Wistar rats. Randomly, the rats were divided into 4 groups with 7 rats in each group. The HBO treatment group 1 received 5 sessions of HBO 2.4 ATA in 3x30 minutes; the HBO treatment group 2 received 10 sessions of HBO 2.4 ATA in 3x30 minutes; and each of the control groups were without HBO. Each of the 28 male rats were given a full thickness excisional wound of 1x1cm. Examinations of e-NOS, TNF- α , VEGF expressions and wound healing were performed on day-0 (pre-HBO) and day-5 HBO or on day-0 (pre-HBO) and day-10 HBO. The results show that the Hyperbaric Oxygen therapy can improve e-NOS (p=0.02), TNF- α (p= 0.02), VEGF expression (p=0.02) and wound healing (p=0.002) significantly in the provision of HBO 2.4 ATA for 3x30 minutes in 5 sessions over 5 consecutive days. While the 10 sessions of HBO 2.4 ATA for 3x30 minutes over 10 consecutive days only increase e-NOS (p=0.02), TNF- α (p=0.04), VEGF expression significantly (p=0.03) but do not improve wound healing significantly (p=0.3) compared with no HBO. The study concludes that HBO can improve the expressions of e-NOS, TNF- α , VEGF and wound healing in the provision of HBO 2.4 ATA for 3x30 minutes in 5 sessions, while the 10 sessions of HBO 2.4 ATA for 3x30 minutes only increase e-NOS, TNF- α , VEGF expression but do not improve wound healing.



1. Introduction

Wound healing is a normal process through a highly complex mechanism. The process occurs progressively through overlapping phases; haemostasis, inflammatory, proliferation, contraction and tissue remodelling [1]. The healing process can be ended by fibrosis or scarring even in optimal conditions, resulting in multiple disabilities [2] Acute injuries can become chronic wounds. The problem of chronic wounds is now a challenge for modern society resulting economically and socially significant medical impact. A Chronic wound is one of the most frequent causes of morbidity in developing countries [3].

An estimated 1.5% incident of the population suffered injuries at any points in time. Laceration or wound often encountered daily (estimated at 20 million cases each year) as a result of cut. A post operation wound is the biggest cause of skin trauma. There are more than 110 million surgical incisions every year, some which can't recover completely due to various causes. The wound developed into chronic wounds .The large number of the population and the burden of costs require serious attention on early detection, prevention, diagnosis and treatment of chronic wounds [2].

The Hyperbaric Oxygen therapy, as an adjuvant therapy in wound healing, results an increase in RNS mainly in NO through iNOS and e-NOS, as well as an increase in ROS, especially H₂O₂. Excessive doses of RNS and ROS are a threat to the survival of the cells, while the therapeutic doses (2.4 ATA) of HBO provide positive benefits for wound healing [4]. In clinical use, the dose of HBO is restricted from 2-2.5 ATA to a maximum of 3 ATA for 60-120 minutes. The HBO therapy varies between 5-30 times [5]. The provision of 100% oxygen at a pressure of 2 ATA improves tissue oxygenation of 30-40 mmHg in normal circumstances up to 250-300 mmHg, while the pressure of 3 ATA allows the improved tissue oxygenation up to 10-15 times [6].The Hyperbaric Oxygentherapy increases tissue oxygenation and stimulates the formation of H₂O₂ as a secondary messenger for the phosphorylation of Nuclear Factor Kappa Beta (NF- κ B). The Nuclear Factor Kappa Beta/Rel transcription proteins play an important and quick role on a wide variety of genes in response to extracellular stimuli. Many aspects of the inflammatory response are also regulated by NF- κ B, including the production factors of complement cascade and the induction of proinflammatory cytokines such as IL1 and TNF α [7]

The Tumour Necrosis Factor (TNF) group consists of 19 cytokines which have an important role in regulating the development and functions of the immune system. Cytokines from the group of Tumour Necrosis Factor (TNF) regulates the inflammatory and immune processes. The lack of TNF can trigger cell deaths. Although TNF- α cytotoxicity affects some tumour cells, TNF α will only trigger apoptosis if the NF κ B signalling is inhibited. TNF α is an activator ofNF κ B, proinflammatory transcription factors that help cells survival through activation of pro-survival genes and suppression of pro-apoptotic genes.

Neovascularization holds a central function in wound healing. In vitro studies indicate that VEGF, the most potent angiogenic protein, was induced by NO. Barriers will disrupt NO synthesis of VEGF expression and angiogenesis, thus slowing down the process of wound healing.

The Hyperbaric Oxygen therapy suppresses the inflammatory response to prevent chronic wounds [8] Supplementary oxygen through the blood stream accelerate sangio genesis in ischemic wounds. Therefore, accelerates the healing time [6]. Numerous studies have proven the benefits of ATA 2.4 HBO five sessions and ten sessions [9]

Our previous studies result that HBO increases the expression of iNOS ($p = 0.001$), while the wound healing ($p = 0.002$) significant at 2.4 ATA HBO administration for 3x30 minutes in 5 sessions. Furthermore, the provision of 2.4 ATA HBO for 3x340 minutes in 10 sessions only enhance the expression of iNOS significantly ($p = 0.002$) but does not improve wound healing significantly ($p = 0.3$) compared with no HBO administration [10].

This study aims to determine the influence of the Hyperbaric Oxygen therapy (HBO) to increase the expression of TNF- α and its relationship with the increased expressions of e-NOS, TNF- α and VEGF in wound healing. This study tests the influence of hyperbaric oxygen therapy (HBO) to the

increase of eNOS, TNF α and VEGF expression by the immune histochemical examination and the macroscopic wound healing process by measuring the area of wound closure on Rattusnorvegicus Wistargallus. The results are expected to be a reference for drafting standard operational procedures (SOPs) of the use of hyperbaric oxygen therapy (HBO) for wound healing in health care centres.

2. Methods

This study was randomized by a controlled group of the pre test-post test designed in 28 Wistar rats. The samples were divided into 4 groups by random allocation, each consists of 7 rats. Treatment group 1 was given HBO 2.4 ATA for 3x30 minutes in 5 sessions; the treatment group 2 was given HBO 2.4 ATA for 3x30 minutes in 10 sessions, and each of the control groups were without HBO. All of the 28 rats were given a 1x1cm full-thickness excisional wound. An immunohistochemical examination of the expressions of eNOS, TNF α , VEGF and the wound size measurement were conducted to determine the wound healing. The immunohistochemical examination and the wound size measurement to the treatment group 1 and the control group 1 were conducted on day 0 (pre-HBO) and day 5 of the HBO therapy. The immunohistochemical examination and the wound size measurement on the treatment group 2 and the control group 2 were conducted on day 0 and day 10 of the HBO.

3. Results and discussion

The study was conducted after the experimental animals had been acclimatized for 7 days. Skin tissue sampling was done by making a full-thickness of 1x1 cm excisional wound on the dorsal part of the experimental animals. The skin tissue sampling on day 0 was performed on all animals (a total of 28 samples) in the treatment group and the control group. The treatment group was given Hyperbaric Oxygen treatment in 5 sessions and 10 sessions. Comprehensive measurement of a wound closure and tissue sampling were done after the treatment of Hyperbaric Oxygen on day 5 (a total of 14 samples) and on day 10 (a total of 14 samples). The slides were made and immunohistochemical staining was conducted.

Immunohistochemical examination for e-NOS, TNF- α and VEGF expressions was conducted under a microscope. This examination was intended to count the number of immunoreactive cells against e-NOS, TNF- α and VEGF. The quantitative data obtained by counting the number of cells in 10 different fields of view at 40x magnification.

3.1. Day 0 (Pre HBO) condition of variables of expressions of e-NOS, TNF- α , VEGF, and wound healing

The examination of e-NOS, TNF- α , VEGF expressions and wound healing done in groups with and without HBO while the extent of pre HBO wound receives the same treatment, i.e. full thickness of 1x1cm (100 mm²) excisional wound size. On day 0 the expressions of e-NOS, TNF- α and VEGF in inflammatory cells with low intensity were observed.




Figure 1. Average HBO on Day 0; Variables of expressions of eNOS, TNF α , VEGF and wound size

3.2. HBO condition of variables of expressions of e-NOS, TNF- α , VEGF and wound healing on day 5

The administration of HBO 2.4 ATA for 3x30 minutes in 5 sessions conducted in the treatment group, followed by an examination of e-NOS, TNF- α , VEGF expressions and the wound size to the treatment group and the control group. Figure 2 shows the average results of the variable expressions of e-NOS, TNF α , VEGF and the wound size. On day 5 there are increased of e-NOS, TNF- α , VEGF expressions with high intensity in the treatment group and the control group. The increased expressions of e-NOS, TNF- α , VEGF in the treatment group are higher than those in the control group. Meanwhile, the wound healing in the HBO treatment group is higher than that in the control group (smaller wound size and better healing).




Figure 2 Average HBO on day 5; Variables of Expressions of e-NOS, TNF- α , VEGF, and wound size

3.3. HBO condition on day 10; variables of expressions of e-NOS, TNF- α , VEGF, and wound size

The HBO administration of 2.4 ATA in 3x30 minute each session was conducted 10 times in the treatment group and followed by an examination of the expressions of e-NOS, TNF- α , VEGF and the wound size to the treatment group and the control group of HBO administration on day 10. Figure 3 shows that the intensity of the expressions of e-NOS, TNF α , VEGF were still high but lower than that of on day 5. There is no significant difference of the wound healing variable between the treatment group and the control group.




Figure 3. Average HBO on Day 10; e-NOS, TNF- α , VEGF Expressions and Wound size




Figure 4. TNF α Expressions using a light microscope Olympus CX31 objective magnification 40x. A: control group day 0, B: control group day-5, C: control group day 10, D: treatment group day 0, E: treatment group day-5, F: treatment group day 10. The intensity of expression of TNF- α is still low in the control group day 0 but the day-5 intensity increases the expression of TNF- α , then the 10th day intensity of the expression of TNF α declines again.




Figure 5. Expressions of e-NOS under a light microscope Olympus CX31 objective magnification 40x. A: control group day 0, B: control group day5, C: control group day 10, D: treatment group day0, E: treatment group day5, F: treatment group day 10. In the control group day 0, the e-NOS expression remains in low intensity; The intensity of e-NOS expression increases on day 5 but declines again on the 10th day.




Figure 6. VEGF Expressions under a light microscope Olympus CX31 objective magnification 40x. A: control group day 0, B: control group day5, C: control group day 10, D: treatment group day 0, E: treatment group day5, F: treatment group day 10. The intensity of VEGF expression is low in the control group day 0; The VEGF expression intensity increases on the 5th day but decreases again on the 10th day.

The results of examination on the expressions of e-NOS, TNF α and VEGF in the treatment group and the control groups on day0 show that the expressions of e-NOS, TNF- α and VEGF intensity are low (1-1). After the administration of HBO 2.4 ATA in five sessions, on the 5th day there seems to be increased expressions of e-NOS, TNF- α and VEGF (2-2) along with the increased number of inflammatory cells, in which the increased expressions of e-NOS, TNF- α and VEGF in the treatment group is significantly higher than those of the control groups. On the 10th day the intensity of the expressions of e-NOS, TNF- α , VEGF is still high. However,

the number of cells that express them reduces and the number of inflammatory cells remains high (2-1).

TNF α is a cytokine regulator and major effectors, mainly secreted by monocytes and macrophages that have been stimulated [11]. HBO administration increases the production of e-NOS in cultured endothelial cells [12]. Several biological processes and growth factors are stimulated and increased with Hypoxia and HBO. Such processes include angiogenesis, collagen synthesis, the activity of osteoclasts, and the release of VEGF and TNF- α . Oxygen stimulates biological processes under conditions of hypoxia and hyperoxia. It is called Oxygen Paradox. One mechanism for forming collagen fibroblasts is stimulated through peroxide. This occurs in HBO hypoxia wounds and during therapy. Peroxide formed in the HBO therapy resembles the stimulus that occurs during hypoxia. Another mechanism is cytokine stimulation which begins with hypoxia, and increases in cytokine regulation which is supported by the presence of hyperoxia occurring in the HBO therapy. VEGF, TNF- α TGF- β and PDGF β are released at wound hypoxia as an initial stimulus, but its activity increases in hyperoxia conditions, especially in the presence of lactic acid [13].

The evidence shows that HBO can accelerate wound healing through mechanisms expressions of e-NOS, TNF α and VEGF in the inflammatory phase and the proliferative phase. The administration of HBO can potentially be used for a post-surgical adjuvant therapy for certain cases, thus it is expected to reduce the duration of treatment (Length of Stay / LOS) in the hospital. Indeed, it is expected to reduce the cost and recovery time for patients after surgery so that patients can be immediately productive and return to their normal activities. Therefore, we need to do more research on the effects of hyperbaric oxygen therapy against various types of wounds, including wounds that are conventionally difficult to cure, post-trauma wounds and postoperative wounds.

The evidence also shows that the administration of HBO continuously for a long time does not provide significant benefits for wound healing. Therefore, time off (recovery time) needs to be given after administration of certain HBO in 5 sessions before continuing to the next session. Further research is needed to determine whether or not there are effects of the recovery period and the range of speeds of the wound healing/recovery time interval before the next continuing HBO therapy session. Further research with more dose variations (between 1-5 sessions and between 5-10 sessions) is needed to obtain the optimal dose of the Hyperbaric Oxygen therapy for acute wounds without infections. Further research needs to be conducted to determine the repetition time interval after the administration of the Hyperbaric Oxygen therapy in 5 sessions (recovery period) as well as the use of HBO as an adjuvant therapy for various types of injuries.

4. Conclusion

This study concludes that applying 2.4 ATA HBO for 3x30 minutes in 5 sessions increases the expressions of e-NOS, TNF- α , VEGF and accelerates wound healing, while giving HBO 2.4 ATA for 3x30 minutes in 10 sessions enhances the expression of e-NOS, TNF- α and VEGF but does not accelerate wound healing.

Furthermore, the evidence shows that giving HBO can accelerate wound healing through mechanisms of e NOS, TNF- α and, VEGF expression. The HBO administration can be potentially used for a post-surgical adjuvant therapy and specific cases as the administration is expected to shorten the duration of treatment in the hospitals, and reduce the cost and recovery time for patients after surgery so that they can return to normal activities.

References

- [1] Finochietto P V, Franco M C, Holod S, Gonzalez A S, Converso D P, Arciuch VGA, Maria P, Poderoso J J and Carreas MC 2009 Mitochondrial Nitric Oxide Synthase: A Masterpiece of Metabolic Adaptation, Cell Growth, Transformation, and Death *The Society for Experimental Biology and Medicine* **234** (9) 1020-8
- [2] Jiang Y, Huang S, Fu X, Liu H, Ran X, Lu S, Hu D, Li Q, Zhang H, Li Y, Wang R, Han C and Chen H 2011 Epidemiology of Chronic Cutaneous Wounds in China *Wound Repair Regeneration* **19** 181-188.
- [3] Sander A L, Henrich D, Muth C M, Marzi I, Barker J H and Frank J M 2006 In Vivo Effect of Hyperbaric Oxygen on Wound Angiogenesis and Epithelization, Wound Repair and Regeneration *Wound Healing Society* **17** 179-184.
- [4] Schreml S, Szeimies R M, Prantl L, Karrer S, Landthaler M and Babilas P 2010 Oxygen in Acute and Chronic Wound Healing, *British Journal of Dermatology, British Association of Dermatology* **163** (2):257-68.
- [5] Kumar S and Wong P F Leaper D J 2004 What is New in Wound Healing ? *Perspective in Medical Science, Turk J Med Sci* **34** 147-160
- [6] İşçimen A and Küçüktaş M 2008 Wound Healing and Hyperbaric Oxygen treatment, *Journal of Turkish Academy of Dermatology*.
- [7] Ghosh S 2007 *Handbook of Transcription Factor NF-kappaB*, CRC Press Taylor & Francis Group
- [8] Thom SR, 2010 Hyperbaric Oxygen: its Mechanisms and Efficacy *Wound Healing Plastic and Reconstructive Surgery* **127** 131-138
- [9] Lahat N, Bitterman H, Yanif N, Kinnarty A, and Bitterman N 1995 Exposure to hyperbaric oxygen induces tumour necrosis factor-alpha (TNF-alpha) secretion from rat macrophages, *Clinical and Experimental Immunology Clinical Sciences Ltd* **102** 655-659
- [10] Xu X, Wang Z, Li Q, Xiao X, Lian Q, Xu W, Sun X, Tao H, and Li R 2009 Endothelial nitric oxide synthase expression is progressively increased in primary cerebral microvascular endothelial cells during hyperbaric oxygen exposure.