

AJPCR

An Open Access Peer Reviewed Journal

Asian Journal of Pharmaceutical & Clinical Research

Volume 10(1) 2020
www.nnovare.com



[Home](#) | [About](#) | [Login](#) | [Register](#) | [Search](#) | [Current](#) | [Archives](#) | [Editorial Board](#)
[Submission](#) | [Instructions to Authors](#) | [Advertise](#) | [Contact Us](#)

USER

Search Articles

All


Search

Username

Password

Remember me

Login



Impact (Cites per doc)- 0.49
(SCImago, SJR 2016)

ICV 2016 = 104.84
Print ISSN- 0974-2441

Online ISSN- 2455-3891



SCOPUS™



Vol 10 Issue 2 February 2017

Table of Contents

Case Study(s)

BRIGHT ASTEROIDS IN THE RED SEA - AN EMERGING PATHOGEN CANDIDA HAEMOLUNII: ARE AZOLES THE TREATMENT OPTION?

Pooja Rao, Sanyuktha T, Keerthiraj B, Radhakrishna M

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.15775

Pages: 1-2 | [Share](#)

Review Article(s)

COMPARISON IN BENEFITS OF HERBAL MOUTHWASHES WITH CHLORHEXIDINE MOUTHWASH: A REVIEW

Renuka Selvakumar, Muralidharan Np

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.13304

Pages: 3-7 | [Share](#)

ADVANCEMENTS IN DRY POWDER INHALER

Ganga Srinivasan, Advait Shetty

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.14282

Pages: 8-12 | [Share](#)

PHYTOCHEMICAL SCREENING, CONTRIBUTION TO THE STUDY OF THE ANTIFUNGAL EFFECT OF FLAVONOIDS FROM DIFFERENT PARTS OF ZIZIPHUS LOTUS OF SOUTH-WEST ALGERIA

Alaa Slimani, Moussaoui Abdellah, Laazouni Hamadi

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.14595

Pages: 13-16 | [Share](#)

ERECTILE DYSFUNCTION AND ITS PHARMACOLOGY: AN OVERVIEW

Sunil Kumar Patnaik, Haritha Polimati, Rajeswara Rao Pragada

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.15183



JOURNAL METRICS

Source Normalized
Impact per Paper
(SNIP): 0.492

Impact per
Publication
(IPP): 0.588

SCImago Journal
Rank (SJR): 0.22

Cite Score: 0.49

Print ISSN: 0974–
2441

Online ISSN:
2455–3891

MOST DOWNLOADED ARTICLES

- PREGELATINIZED CASSAVA STARCH...
- ANTIBACTERIAL ACTIVITY OF HYDROLYZED...
- BUCCAL PENETRATION ENHANCERS-AN OVERVIEW
- Upper Thoracic spine(D2-D3)...
- ANTI-CANCER ACTIVITY OF DATURA METEL...
- ANTIBACTERIAL ACTIVITY OF THE...
- A REVIEW: LOVASTATIN PRODUCTION AND...
- DIPEPTIDYL PEPTIDASE-IV INHIBITORY...

Pages: 17-23 | Share

EFFECT OF ANTI-BIOFILM POTENTIAL OF DIFFERENT MEDICINAL PLANTS: REVIEW

M Gayathri, Abinaya Man

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.15334

Pages: 24-32 | Share

THE MAJOR HISTOCOMPATIBILITY COMPLEX: A REVIEW

Preeti Sharma, Pradeep Kumar, Rachna Sharma

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.15555

Pages: 33-36 | Share

LIFE, DEATH, AND PHARMA IN INDIA: A CASE STUDY

R.Gayathri Saravanan, C Vijayabanu

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.15566

Pages: 37-39 | Share

ROLE OF COMMUNITY PHARMACIST AND NUTRITIONISTS IN OBESITY MANAGEMENT: A LITERATURE REVIEW

Madeeha Malik, Natasha Kanwal, Azhar Hussain

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.15603

Pages: 40-42 | Share

CURRENT DEVELOPMENTS IN TRANSPORT MEDIA FOR AVULSED TEETH: AN UPDATE

Roma M, Shreya Hegde

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.15663

Pages: 43-46 | Share

A REVIEW: ANTIMICROBIAL AGENTS BASED ON NITROGEN AND SULFUR CONTAINING HETEROCYCLES

Praveen Kumar Sharma

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.15673

Pages: 47-49 | Share

EFFECT OF PHYTOESTROGENS ON LIPID PROFILE: MINI REVIEW

Takkella Nagamma, Akshay T Jagadeesh, Anjaneyulu K, Kumar Mr Bhat

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.15684

Pages: 50-53 | Share

PARENTERAL NUTRITION IN NEONATUS AND COMPATIBILITY ISSUES: A REVIEW ARTICLE

Yelly Oktavia S, Helmi Arifin, Rahmi Yosmar

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.15697

Pages: 54-64 | Share

APPLICABILITY OF NANOPARTICLES-HYDROGEL COMPOSITE IN TREATING PERIODONTAL DISEASES AND BEYOND

Nafiu Aminu, Seok Ming Toh

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.15709

Pages: 65-70 | Share

EDIBLE VACCINES: AN ADVANCEMENT IN ORAL IMMUNIZATION

• TURMERIC:
NATURE'S
PRECIOUS
MEDICINE.

• ETHNOBOTANICAL
STUDIES ON
SELECTED...

Srinivas Rajesham Bhairy, Rajashree Shreeram Hirlekar

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.15825

Pages: 71-77 | [Share](#)

CONVENTIONAL AND RECENT DIAGNOSTIC AIDS IN TUBERCULOUS LYMPHADENITIS: A BRIEF OVERVIEW

Shamimul Hasan, Shazina Saeed

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.16102

Pages: 78-81 | [Share](#)

Original Article(s)

ANTICANCER ACTIVITY OF CAMELLIA SINENSIS MEDIATED COPPER NANOPARTICLES AGAINST HT-29, MCF-7 AND MOLT-4 HUMAN CANCER CELL LINES

Vijay D. Mendhulkar, Anu Yadav

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.15710

Pages: 82-88 | [Share](#)

IS THERE TREATMENT EFFECT OF VINPOCETINE ON AUTONOMIC DYSFUNCTION IN RATS WITH ALZHEIMER

Azizuddin Khan, Nazan Dolu, Seda Gündüz, Ali Yucel Kara, Hale Acer, Selda Tasa

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.11367

Pages: 89-91 | [Share](#)

EFFECT OF YOHIMBINE ON CLOMIPRAMINE-INDUCED SEXUAL DYSFUNCTION IN MALE RATS

Devangam Sheshadri Shekar

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.12970

Pages: 92-96 | [Share](#)

THE DEVELOPMENT OF LEAN HEALTHCARE MANAGEMENT SYSTEM (LHMS) FOR HEALTHCARE INDUSTRY

Nurul Fadly Habidin

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.14193

Pages: 97-102 | [Share](#)

PHYTOCONSTITUENTS, PIGMENTS , G.C.M.S. ANALYSIS AND ALLELOPATHY EFFECT OF ALTERNANTHERA FICOIDEA (L.) P. BEAUV

Reshma Bhagawanrao Patil, Basavaraj A Kore

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.14498

Pages: 103-108 | [Share](#)

ANTICONVULSANT PROPERTIES OF SOME MEDICINAL PLANTS- A REVIEW

Nivedha Srinivasan, Anitha Roy

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.14501

Pages: 109-111 | [Share](#)

THE CHEMICAL COMPOSITION AND THE ANTIMICROBIAL PROPERTIES OF THE ESSENTIAL OIL EXTRACTED FROM THE LEAVES OF TEUCRIUM CAPITATUM L.

Jalila El Amri, Khalid El Badaoui, Zoubida Haloui

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.14504

"STUDY OF TOTAL CHOLESTEROL (TC), TRIACYLGLYCEROLS (TG), HIGH DENSITY LIPOPROTEIN CHOLESTEROL (HDL-C) IN TYPE II DIABETES MELLITUS."

Smita V Patil, Anita P Mandare, Gaikwad B Pandurang

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.14535

Pages: 116-118 | Share

DRUG COST ANALYSIS ON NATIONAL HEALTH INSURANCE ERA IN HEALTH SUPPORT FACILITIES

Satibi Satibi, Dewa Ayu Putu Satrya Dewi, Atika Dalili Akhmad, Novita Kaswindiarti, Dyah Ayu Puspandari

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.14577

Pages: 119-121 | Share

A STUDY OF DRUG UTILIZATION PATTERN OF THE GERIATRIC PATIENTS IN THE DEPARTMENT OF GERIATRIC MEDICINE IN A TERTIARY CARE HOSPITAL IN ASSAM, INDIA

Lakhimi Borah, Diptimayee Devi, Prasanjit Kumar Debnath, Dibyajyoti Deka

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.14762

Pages: 122-126 | Share

THE ROLE OF PHARMACISTS IN EVALUATING AND INTERVENING THE PATIENTS WITH DIABETIC NEUROPATHY

Sesilia Andriani Keban, Syamsudin Abdillah, Najuah Najuah

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.14796

Pages: 127-131 | Share

PHYSICAL, CHEMICAL CHARACTERISTICS AND RELEASE RATE OF MELOXICAM FROM NANOSTRUCTURED LIPID CARRIERS (NLC) SYSTEM USING SOME CONCENTRATION RATIOS OF MONOSTEARIN AND ALPHA TOCOPHEROL ACETATE LIPID MATRIX

Yulia Anggraeni, Isnaeni Isnaeni, Esti Hendradi

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.14806

Pages: 132-137 | Share

OBTENTION OF PROTEIN CONCENTRATE AND POLYPHENOLS FROM MACADAMIA (MACADAMIA INTEGRIFOLIA) WITH AQUEOUS EXTRACTION METHOD

Wilman Carrillo, David Lara, Edgar Vilcacundo, Cristian Carrillo, Monica Silva, Mario Alvarez, Cecilia Carpio

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.14808

Pages: 138-142 | Share

EVALUATION OF IN-VITRO CYTOTOXIC ACTIVITY OF CHLOROFORM EXTRACT OF SIDA ACUTA BURM.F.

Lakshmi Kanta Kanthal

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.14810

Pages: 143-145 | Share

EFFECTIVITY AND PHYSICOCHEMICAL STABILITY OF NANOSTRUCTURED LIPID CARRIER COENZYME Q10 IN DIFFERENT RATIO OF LIPID ALFA CETYL PALMITATE AND ALPHA TOCOPHERYL ACETATE AS CARRIER

Astridani Rizky Putranti, Esti Hendradi, Riesta Primaharinastiti

Abstract || [View PDF](#) || [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.14835

Pages: 146-152 | Share

Abstract || [View PDF](#) ||  [Download PDF](#) || DOI: 10.22159/ajpcr.2017.v10i2.17793
Pages: | [Share](#)

[Our Journals](#) || [Open Access Policy](#) || [Publication Ethics](#)

The publication is licensed under a Creative Commons License (CC BY)  [View Legal Code](#)

Copyright © 2017 All Rights Reserved: **Innovare Academic Sciences** || Powered By **CyberDairy**


USER

Username

Password

Remember me

Login



Impact (Cites per doc)- 0.49
(SCImago, SJR 2016)

ICV 2016 = 104.84

Print ISSN- 0974-2441

Online ISSN- 2455-3891



Editorial Board

Editorial Board

AJPCR is committed to have dynamic and potential advisory-editorial board. Those established in the field can directly send their resume. New people are first needed to serve as referee before being considered member of advisory-editorial board. Email your resume to editor@ajpcr.com

Editor-in-Chief

Dr. Anurekha Jain
Dept. of Pharmaceutical Sciences, Jyoti Mahila Vidyapeeth University, Jaipur, Rajasthan
Email: editor@ajpcr.com

Associate Editor

Dr. Neeraj Upmanyu
Peoples Institute of Pharmacy & Research Center, Bhopal, MP, India
Email: dmeerajupmanyu@gmail.com

Dr. Vikas Sharma
Shri Rawatpura Sarkar Institute of Pharmacy, Datiya, MP, India
Email: vikassharma15@gmail.com

Assistant Editor

Dr. Vimal Kumar Jain
Institute of Pharmacy, Nirmal University, Ahemdabad, Gujarat, India
Email: cognosy@gmail.com

Dr. Rupesh Kumar Gautam
ADINA Institute of Pharmaceutical Sciences, Sagar, MP, India
Email: drrupeshgautam@gmail.com

Editorial Board Members

- Dr. Debasish Maiti
Suryamaninagar, Tripura West, Agartala, Tripura, India
- Dr. Rashad Mohammed Musleh
University of Thamar, Yemen



JOURNAL METRICS

Source Normalized
Impact per Paper
(SNIP): 0.492

Impact per
Publication
(IPP): 0.588

SCImago Journal
Rank (SJR): 0.22

Cite Score: 0.49

Print ISSN: 0974-
2441

Online ISSN:
2455-3891

MOST DOWNLOADED ARTICLES

- PREGELATINIZED CASSAVA STARCH...
- ANTIBACTERIAL ACTIVITY OF HYDROLYZED...
- BUCCAL PENETRATION ENHANCERS-AN OVERVIEW
- Upper Thoracic spine(D2-D3)...
- ANTI-CANCER ACTIVITY OF DATURA METEL...
- ANTIBACTERIAL ACTIVITY OF THE...
- A REVIEW: LOVASTATIN PRODUCTION AND...
- DIPEPTIDYL PEPTIDASE-IV INHIBITORY...

- Dr. Shubhamoy Ghosh
Dept. of Pathology & Microbiology, Mahesh Bhattacharyya Homeopathic Medical College & Hospital, Govt. of West Bengal, India
- Mohd Abdul Hadi
Bhaskar Pharmacy College Affiliated To Jawaharlal Nehru Technological University, Hyderabad, India
- Dr. Amer A. Taqa
Department of Dental Basic Science, College of Dentistry, Mosul University, Iraq
- Mr. Atul Kabra
Department of Pharmacy, Manav Bharti University, Solan (H.P.), India
- Mr. Siddharth Kaushal Tripathi
Doctoral Research Associate National Center For Natural Products Research, School of Pharmacy, University of Mississippi, India
- Dr. Brajesh Kumar
Escuela Politécnica Del Ejército, Sangolqui, Ecuador, Latin America
- Dr. Farhan Ahmed Siddiqui
Faculty of Pharmacy, Federal Urdu University Arts, Science And Technology Karachi, Sindh, Pakistan
- Dr. Deepak Kumar Mittal
Iasca Department of Itm-Universe, Gwalior, India
- Dr. Jesse Joel Thathapudi
Karunya University (Deemed To Be University), Coimbatore, Tamil Nadu, India
- Dr. Kumaran Shanmugam
Analytical Chemistry, Academic Faculty, (Senior Scale) & International Co-Ordinator, Department of Biotechnology, Periyar Maniammai University, Vallam, Thanjavur-613 403, India
- Dr. Javad Sharifi Rad
Department of Pharmacognosy, Faculty of Pharmacy, Zabol University of Medical Sciences, P.O. Box 61615-585 Zabol, Iran
- Dr. Rajesh Mohanraj
Dept. of Pharmacology, CMHS, UAE
- Dr. P. Thillai Arasu
Department of Chemistry, Kalasalingam University, Srivilliputhur 626190, India
- Ms Vasundhra Saxena
Gautam Buddha Technical University Lucknow, India
- Ms Vinit Dattatray Chavhan
Sinhgad Technical Education Society'S Smt. Kashibai Navale College of Pharmacy, Kondhwa-Saswad Road, Kondhwa (BK), India
- Dr. Sami Saqf El Hait
Junior Executive - Quality Control At Jamjoom Pharmaceuticals Company Limited Jeddah, Saudi Arabia
- Md. Moklesur Rahman Sarker
Faculty of Medicine, University of Malaya, Malaysia

• TURMERIC:
NATURE'S
PRECIOUS
MEDICINE.

•
ETHNOBOTANICAL
STUDIES ON
SELECTED...

- Dr. Ahmed Hashim Mohaisen Al-Yasari
Department of Physics, College of Education For Pure Science, University of Babylon, Hilla,
Iraq
- Dr. Arun Kumar
Director & Dean, Faculty of Marine Sciences Faculty of Marine Sciences Cas In Marine
Biology Cas In Marine Biology, Parangipettai- 608502 Parangipettai- 608502 Tamil Nadu,
India
- Dr. Ashraf Ahmed
Dept. of Chemistry, Aligarh Muslim University, Uttar Pradesh 202001, India
- Dr. S. Bala Murugan
Dept. of Biotechnology, Bharathiar University, Coimbatore-641046, Tamil Nadu, India
- Dr. Sukhen Som
Department of Pharmaceutical Chemistry, M.M.U College of Pharmacy, K.K.Doddi,
Ramadevara Betta Road, Ramanagara- 562159 State- Karnataka India
- Dr. Sheikh Shoib
Department of Psychiatry, Institute of Mental Health And Neurosciences Kashmir (Imhans
K), Srinagar, India
- Dr. Hao Wu
Postdoctoral Fellow At Ngm Biopharmaceuticals, Inc, South San Francisco, CA 94080, USA
- Dr. Payal Bhaskar Joshi
Assistant Professor Department of Chemical Engineering, Mukesh Patel School of
Technology Management & Engineering (Mumbai Campus), Svkm'S Nmims (Deemed-To-
Be-University), India
- Dr. Nagarajan Kayalvizhi
Assistant Professor Department of Zoology, Periyar University, Salem, Tamilnadu, India
- Dr. Madhu Bala
Scientist 'F' And Joint Director, Institute of Nuclear Medicine And Allied Sciences (INMAS),
India
- Prof. Dr. Mamdouh Moawad Ali
Biochemistry Department, Genetic Engineering And Biotechnology, India
- Dr. Mohanraj Rathinavelu
Department of Pharmacy Practice, Raghavendra Institute of Pharmaceutical Education &
Research, Riper, India
- Dr. (Mrs.) Neeru Nathani
Dept. of Swasthavritta And Yoga, Faculty of Ayurveda, Institute of Medical Sciences,
Banaras Hindu University, India
- Dr. Rohini Karunakaran
Unit of Biochemistry, Faculty of Medicine, Aimst University Batu 3 1/2, Bukit Air Nasi, Jalan
Semeling, 08100 Bedong Kedah Darul Aman, Malaysia
- Dr. Imran Ahmad Khan
Royal Institute of Medical Sciences Multan, Pakistan
- Dr. Jitendra Gupta
Sper Timer (Publication Committee) Regional Head U.P. State, India

- Dr. Kamal A. Badr
Lecturer of Pharmaceutics And Industrial Pharmacy & Member of Quality Assurance Unit-
Faculty of Pharmacy- Delta University For Science And Technology, India
- Mr. Gurpreet Singh
Department of Pharmaceutical Sciences Guru Nanak Dev University, Amritsar, Punjab
(India) 143005
- Dr. Pranav Kumar Prabhakar
Lovely Faculty of Applied Medical Sciences, Lovely Professional University, Jalandhar-Delhi
G.T. Road
- Dr. Raj Mohan Raja Muthiah
Research Fellow (Harvard Medical School) 172 Hosmer Street, Apt 7. Marlborough, Ma
01752
- Dr. Sandip Narayan Chakraborty
Research Asst li, Translational Molecular Pathology, Ut Md Anderson Cancer Center, Life
Sciences Plaza, Houston, TX 77030
- Dr. Anup Naha
Dept. of Pharmaceutics "Swarna Kutir ", Ramnagar Road No.4, Mcops, Manipal-576 104,
Karnataka, India
- Dr. Tushar Treembak Shelke
Vice Principal , Head of Department of Pharmacology And Research Scholar, In
Jspms Charak College of Pharmacy & Research, Gat No. - 720(1&2), Pune, India
- Anindya Banerjee
Indian Pharmacopoeia Commission, Ministry of Health & Family Welfare
Govt of India
- Dr. Vijay Mishra
Lovely Institute of Technology (Pharmacy), Lovely Professional University, Phagwara,
Punjab, India
- Dr. Praveen Kumar Sharma
Department of Chemistry, Lovely Professional University, Punjab (India)-144411
- Deepansh Sharma
School of Bioengineering & Biosciences Lovely Professional University, Phagwara Punjab,
India
- Sai Prachetan Balguri
ORISE Research Fellow at U.S. FDA
- Dr. Mohd Abdul Hadi
Department of Pharmaceutics, Bhaskar Pharmacy college, Yenkapally (V), Moinabad (M),
R.R (Dt), Hyderabad-500 075, Telangana, India
- Tanay Pramanik
Department of Chemistry in Lovely Professional University, Punjab, India
- Dr. D. Nagsamy Venkatesh
Department of Pharmaceutics, JSS College of Pharmacy, Ooty, TN India

Editorial office

Asian Journal of Pharmaceutical and Clinical Research

B-11, In front of Beema Hospital, Nayi Awadi, Mandasaur 458001, MP, India
E-mail:editor@ajpcr.com

[Our Journals](#) || [Open Access Policy](#) || [Publication Ethics](#)

The publication is licensed under a Creative Commons License (CC BY)  [View Legal Code](#)

Copyright © 2017 All Rights Reserved: **Innovare Academic Sciences** || Powered By **CyberDairy**

PHYSICAL AND CHEMICAL CHARACTERISTICS OF MELOXICAM FROM NANOSTRUCTURED LIPID CARRIERS SYSTEM USING SOME CONCENTRATION RATIOS OF MONOSTEARIN AND ALPHA-TOCOPHEROL ACETATE LIPID MATRIX

YULIA ANGGRAENI¹, Isnaeni Yudi Haryanto², ESTI HENDRADI^{3*}

¹Post Graduated Student of Pharmaceutics Department, Faculty of Pharmacy, Airlangga University, Surabaya 60286, East Java, Indonesia.

²Department of Pharmaceutical Chemistry, Airlangga University, Surabaya 60286, East Java, Indonesia. ³Department of Pharmaceutics, Faculty of Pharmacy, Airlangga University, Surabaya 60286, East Java, Indonesia. Email: estihendradi@yahoo.com

Received: 21 August 2016, Revised and Accepted: 22 October 2016

ABSTRACT

Objective: The objective of this research is to develop nanostructured lipid carriers (NLC) of meloxicam (MLX) for topical application using monostearin and alpha-tocopherol as lipid matrix.

Methods: MLX-NLC was prepared by high-shear homogenization, and it was characterized for organoleptic, pH, viscosity, particle size and size distribution, zeta potential, crystallinity, morphology, and entrapment efficiency. The six different ratios of solid and liquid lipid matrix were 70:30, 75:25, 80:20, 85:15, 90:10, and 95:5, respectively.

Results: The pH value of all formulas met the pH range of topical dosage at 4.5-6.5. Viscosity test showed that a trend toward the decrease in viscosity with the increase in alpha-tocopherol content was observed. The results of particle size test proved that all six NLC system formulas have <1000 nm particle size with a quite narrow particle size distribution. Scanning electron microscope revealed nearly spherical shape NLC with negligible effect ratios of solid and liquid lipid on the particle morphology. The X-ray diffraction result showed the decreasing of peaks intensity of MLX-NLC system. It showed that MLX in NLC system is amorphous. The result showed that the increasing concentration of liquid lipid in NLC system caused the rising entrapment efficiency of MLX.

Conclusion: The result of this research indicates that the concentration ratio of monostearin and alpha-tocopherol acetate affects the physical and chemical characteristics of NLC system.

Keywords: Nanostructured lipid carriers, Meloxicam, Monostearin, Physical and chemical characteristics, Entrapment efficiency.

© 2017 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) DOI: <http://dx.doi.org/10.22159/ajpcr.2017.v10i2.14806>

INTRODUCTION

Meloxicam (MLX) is a non-steroidal anti-inflammatory drug, which inhibits cyclooxygenase-2 selectively, used orally to reduce the symptoms of rheumatoid arthritis. However, the use of oral MLX still showed gastrointestinal side effects, the risk of arterial thrombotic events, kidney malfunction, liver dysfunction at high doses, and long-term treatment. Topical route administration is an alternative to overcome these problems and also provides many advantages, including avoiding gastrointestinal irritation, systemic toxicity is minimal, avoiding hepatic metabolism, the plasma level is stable, and improving patient compliance [1].

MLX is classified as a BCS Class II drug (high permeability and low solubility), has a log p=3.42, and a poor wettability in water, causing difficulties in the pharmaceutical formulations design. Besides, the skin barrier properties also limit the permeability of various pharmaceutical active substances, requiring appropriate drug delivery systems to produce optimal therapeutic effect [2,3].

Nanostructured lipid carriers (NLC) have many advantages such as occlusion and skin hydration, absorption-increasing effects, active penetration enhancement, and controlled release properties. NLC was developed by exchanging the liquid lipid (oil) of oil-in-water (o/w) emulsions by a solid lipid, which can bring many advantages in comparison with a liquid core. The concept of NLC was developed by nanostructuring the lipid matrix, to give more flexibility for modulation

of drug release, increasing the drug loading, and preventing its leakage. The result is a less ordered lipid matrix with many imperfections, which can accommodate a higher amount of drug. The other advantages of using lipids as carrier systems for skin administration are related to their physiological and well-tolerated nature, which reduces the risk of toxicological problems and local irritancy. NLC is a lipid nanoparticle that can increase the penetration of lipophilic drug substance as it has occlusive properties and improves skin hydration. Therefore, MLX is encapsulated in the NLC system to increase its penetration into the skin [4].

METHODS

Materials

MLX was a gift sample from Apex Healthcare Limited; Monostearin (Cutina® GMS) was obtained from BASF; alpha-tocopherol acetate was obtained from Xinchang pharma; Kolliphor® P 188 (poloxamer 188) was obtained from BASF; Tween 80 was obtained from PT KAO; Ethanol p.a., sodium hydroxide, and NaH₂PO₄ (natrium dehydrogenate phosphate) p.a. were a product of Merck. Aquabidestilata was gift sample from PT. Widatra Bhakti, Pasuruan, Indonesia. All other ingredients used were analytical grade.

Methods

Preparation of MLX-NLC system

The method of this research is high-shear homogenization. MLX-NLC system was made by melting the lipid phase (monostearin and

α -tocopherol) with different lipid ratio (70:30, 75:25, 80:20, 85:15, 90:10, and 95:5), and MLX at 65°C temperature. At the same time, the surfactant solution (Kolliphor® P 188, Tween 80 and pH phosphate buffer 6.0±0.05) was prepared and heated at the same temperature. Then, this heated surfactant solution is dispersed into heated lipid phase using ultra-turrax at 5000 rpm speed for 15 minutes. Furthermore, this surfactant solution was added with phosphate buffer with 6.0±0.05 pH value until the volume reaches 100% b/b. Subsequently, the NLC dispersions were formed simply by cooling the warm pre-emulsion precursor to room temperature in the same container [5,6]. The composition of MLX-NLC system is shown in Table 1.

Physical and chemical characterization of MLX-NLC system

Fourier transform infrared (FTIR) spectroscopy

FTIR spectroscopy was conducted in the wave number range 4000-400/cm. The sample was combined with KBr and pressed into a pellet. The solid pellet was analyzed using FTIR spectroscopy [7].

pH

The electrode was inserted into 10 mL MLX-NLC system and then the number that is shown by pH meter must be written.

Viscosity

This viscosity evaluation has done to know the thickness of MLX-NLC system that is resulted from adding other materials, such as surfactants, and the effect of making technique. Viscosity measurement used cone and plate as the tools of the viscometer. 2.0 mL of MLX-NLC system placed in the sample cup, and then the viscometer is turned on and puts it down at a moment until the value is stable.

Particle size and size distribution

The examination of particle size and size distribution has done by Delsa™ Nano Submicron Particle Size Analyzer. Then, the tool is turned on and particle size menu is selected. The tool will measure it over 10 minutes. The observed data are the average droplet diameter and polydispersity index (PI). PI illustrates the variation on a sample. The small value of PI (<0.3) indicates that the sample is monodisperse.

Zeta potential

Zeta potential was measured using Malvern Instrument® with cell and the appropriate procedure. The instrument can measure on 0.001-40% concentration. The sample is dispersed into the water until it gets the concentration on the optimum intensity of the instrument. The suspension was sonicated for approximately 2 minutes to break agglomerate, and then it is measured.

Particle morphology

Particle morphology examination has done by scanning electron microscopy (SEM). This evaluation used to determine the form of particle contained in NLC. The MLX-NLC system was dispersed in CMC-Na gel and spread on glass plates to produce dry NLC system. Then, the dry NLC system was coated with Aurum before counted in vacuum condition.

Crystallinity

Differential scanning calorimetry (DSC)

Melting temperature measurement has done by DSC. The characterization using this instrument has done by weighing the sample of MLX and monostearin amount at 2-8 mg. Then, this powder inserted into an aluminum pan that is impermeable to air. This pan is heated to the temperature 30-300°C at calorimeter with the temperature rise 10°C every minute.

X-ray diffraction (XRD)

The sample is inserted to the holder glass, and the surface is trimmed by plate glass. This sample is placed in X-ray Philips X'Pert diffractometer. The measurement condition was: 40 kV voltage; 30 mA; Divergence slit size 1.0°. Then, it observed over a range of 2 θ angle from 5° to 50°; 0.15 mm receiving slit. Then, it observed over a range of 2 θ angle from 5° to 50°.

Entrapment efficiency of MLX

Drug E.E. was calculated by determining the amount of drug untrapped (Cf) after removal of NLC system by centrifugation. MLX which is untrapped in NLC system would be dispersed in phosphate buffer with pH 7.4±0.05 as a supernatant. The obtained supernatant was filtered using Millipore Whatman 0.2 μ m filter paper. This evaluation used UV spectrophotometer. Furthermore, it is calculated using the following formula:

$$EE(\%) = \left\{ \frac{C_t - C_f}{C_t} \right\} \times 100\%$$

Where Ct is the total amount of drug added in the formulation. The concentration of drug was determined by double-beam ultraviolet (UV)-vis spectrophotometer (Shimadzu UV-1800) with multiple wavelength methods [8].

Statistical analysis

Characterization evaluation (pH, viscosity, particle size, zeta potential, and entrapment efficiency of drug) of MLX-NLC system has statistically analyzed using one-way multivariate analysis of variance (MANOVA) method at 95% level. Then, significant difference of NLC system formula was obtained by honestly significant difference test.

RESULTS AND DISCUSSION

Physical and chemical characterization of MLX-NLC system

FTIR spectroscopy

The analysis of functional group change in MLX-NLC system was done by comparing the IR spectrum of MLX-NLC system with the material IR spectrum, as shown in Fig. 1.

MLX-NLC system with some ratios of the lipid matrix concentration has a similar FTIR spectrum profile. This condition shows that there is no difference in the interaction of some components when forming the NLC system. The characteristic peak of NH₃ and SO₂ group does not appear in the FTIR spectrum of MLX-NLC system. This condition shows that there was a possibility of intermolecular hydrogen bonds between N-H and S=O group with a hydroxyl group of some components. In

Table 1: Composition of MLX-NLC system

Formulation code	MLX (%b/b)	Monostearin (%b/b)	Alpha-copherol acetate (%b/b)	Tween 80 (%b/b)	Kolliphor 188 (%b/b)
F1	1	70	30	3	3
F2	1	75	25	3	3
F3	1	80	20	3	3
F4	1	85	15	3	3
F5	1	90	10	3	3
F6	1	95	5	3	3

NLC: Nanostructured lipid carriers, MLX: Meloxicam

addition, there is no new characteristic peak that appears in the FTIR spectrum of MLX-NLC system. Based on FTIR spectrum result, it could be seen that there was no chemical interaction between MLX with the constituent component of NLC system. There was only physical reaction in NLC system formation. The formation of MLX-NLC system with lipid matrix of monostearin and alpha-tocopherol acetate can be identified from NH_3 group and SO_2 group intensity lost of MLX [7].

pH

The pH parameter of MLX-NLC system has a connection with the pH stability of active ingredient and the convenience in topical administration. Topical dosage forms pH applied on the skin should have a value of 4.5-6.5 [9].

The obtained NLC system had a pH value in the range of 5.98-6.07. The histogram of average pH in some formulas of NLC system is shown in Fig. 2.

Statistical analysis using MANOVA showed that there was no significant difference between pH in each formula of NLC system which uses




Fig. 1: Fourier transform infrared spectrum of, (a) meloxicam, (b) monostearin, (c) alpha tocopherol acetate, (d) tween 80, (e) poloxamer 188, (f) formula 1 (70:30), (g) formula 2 (75:25), (h) formula 3 (80:20), (i) formula 4 (85:15), (j) formula 5 (90:10), (k) formula 6 (95:5)




Fig. 2: The histogram of average pH in some formulas of nanostructured lipid carriers system. The data obtained is the average of the pH value $\times 3$ replication formula $3 \pm$ standard deviation

multiple ratios of lipid matrix which is about 70:30; 75:25; 80:20; 85:15; 90:10; and 95:5 ($*p > 0.05$). Based on these results, it could be concluded that the ratios of solid and liquid lipid matrix did not affect the pH value of NLC system.

Viscosity test

MLX-NLC system which has been produced had viscosity values from 57.87 to 162.67 c.Ps. The histogram of average viscosity in some formulas of NLC system is shown in Fig. 3.

Statistical analysis using MANOVA showed that there were significant differences ($*p < 0.05$). From *post-hoc* test result, it could be known that F1 is not significantly different with F2, and F2 is not significantly different with F1 and F3, and then F4 is significantly different with all formulas; moreover, F5 is not significantly different with F6. From this result, it can be proven that the increased number of liquid lipid which is added to the formula is capable to decrease the viscosity of NLC system.

Viscosity is a barrier of liquid to flow, if the viscosity of the sample is higher and the barrier to flow is also higher [10]. The viscosity of NLC system is affected by the composition of NLC system formula.

Particle size and size distribution of NLC

From the particle examination result of all six NLC system formulas, the particle size was about < 1000 nm. The particle size of MLX-NLC system was about 431.1-515.3 nm. In addition, the histogram of average particle size in some NLC formulas is shown in Fig. 4.

Statistical analysis of particle size using MANOVA showed that there was no significant difference between the particle size in each formula of NLC system which uses multiple ratios of lipid matrix 70:30; 75:25;




Fig. 3: The histogram of average viscosity in some formulas of nanostructured lipid carriers system. The data obtained is the average of the pH value $\times 3$ replication formula $3 \pm$ standard deviation




Fig. 4: The histogram of average particle size in some nanostructured lipid carriers formulas in aqueous media using Delsa Nano®. The data obtained is the average of the pH value $\times 3$ replication formula $3 \pm$ standard deviation

80:20; 85:15; 90:10; and 95:5 (*p>0.05). In this study, the results are not in accordance with Khurana *et al.* who found that particle size of NLC prepared using cetyl palmitate (solid lipid) and caprylic acid (liquid lipid) depended on liquid lipid content. A trend toward the decrease in particle size with the increase in content was observed [8]. This is possible because the lipid matrix of different components with different characteristics may give different results. Cetyl palmitate is a wax lipid group, has a highly ordered crystal lattice and monostearin a class of glycerides having less ordered crystal lattice [11,12]. Differences in chemistry, physics, and crystallographic characteristics of the system will affect the characteristics of NLC [11].

Statistical analysis of PI using MANOVA showed that there were significant differences between PI in each formula of NLC system which uses multiple ratios of lipid matrix 70:30; 75:25; 80:20; 85:15; 90:10; and 95:5 (*p<0.05). From *post-hoc* test result, it could be known that F5 is significantly different with F1, F3, and F6. However, F6 is significantly different with F2. In addition, F1 and F6 have PI value which is larger than the other formulas. Moreover, F2, F3, F4, and F5 have more homogeneous particle size and they have particle size distribution which is narrower than the F1 and F6. F1 and F6 have broader particle size distribution than the others formula. This is possible because the F1 has a lower viscosity than other formula NLC systems. Stokes's law states that the value of the viscosity is inversely proportional to the speed of sedimentation [10]. Sedimentation velocity describes as the speed of formation of agglomerates. So that on systems with lower viscosity will be easier to occurrence of agglomeration. This resulted in the F1 is obtained PI value greater than other formulas. The high value of PI F6 is possible because most of the components of the lipid matrix is monostearin. Monostearin powder is generally in β -polymorph form and it will transform to α -polymorph after going through the process of melting lipid. α -polymorphic form monostearin is metastable, which will transform into β -polymorph at room temperature. The phenomenon of polymorph rapid change affects the instability of NLC system, such as the transformation of structure particle, agglomeration, and sedimentation in aqueous phase [13]. This condition can affect the formation of NLC system with polydisperse particle size distribution and higher tendency to agglomeration.

Zeta potential

Zeta potential measurement is an indirect measurement of NLC physical stability; it also affects the release kinetics and biological process of a nanoparticle. Zeta potential is the electrical potential on sliding area, which is defined as the electrical charge of outer particle surface where the opposite charge still able to strongly bound with other particles when they move in an electrical field.

Zeta potential measurement obtained 10.4-16.5 as a negative value. The histogram of average zeta potential in all NLC system is shown in Fig. 5.

Based on MANOVA, statistical analysis result showed that there are significant differences between zeta potential in each formula of NLC system that uses multiple ratios of lipid matrix 70:30; 75:25; 80:20; 85:15; 90:10; and 95:5 (*p<0.05). On *post-hoc* analysis, it could be known that there was a difference between F1 and F6 with F2, F3, F4, and F5. The result was almost equal to the particular size and PI results, where F1 and F6 had smaller zeta potential than the other formulas. On the other hand, the value of particle size and PI was larger than the other four formulas.

The examination of crystallinity

The X-ray diffraction data

X-ray diffraction analysis is used to know the crystallinity change of active substance and lipid component in NLC system. X-ray diffraction's pattern of MLX, monostearin, and some NLC formulas are shown in Fig. 6.

Based on the X-ray diffraction result, the diffractogram of MLX has some sharp peak at 25°, 15°, 18°, 19°, and it has some wide specific peak at

13° and 23°. Moreover, monostearin has some sharp specific peak at 5°, 19° and wide specific peak at 23°. At the results of all formulas with different ratio of solid lipid and liquid lipid, it was known that NLC system's peak intensity has decreased. The result showed that NLC system was amorphous. The condition indicated that MLX was entrapped in NLC system.

DSC

DSC analysis provides data such as thermogram to determine the melting point of a sample. The result of these data is shown in Fig. 7.

The melting point of NLC has decreased when it is compared with the melting point of the bulk lipid. This decrease occurs because there is a foreign molecule (molecule of the active substance or surfactant) which is dissolved in the lipid matrix. The melting point and crystallinity will




Fig. 5: The histogram of some nanostructured lipid carriers formulas average zeta potential in aqueous media using Malvern Instrument®. The data obtained is the average of the pH value ×3 replication formula 3±standard deviation




Fig. 6: The Examination result of meloxicam, monostearin, formula 1 (7:3), formula 2 (7.5:2.5), formula 3 (8:2), formula 4 (8.5:1.5), formula 5 (9:1), and formula 6 (9.5:0.5) X-ray diffraction




Fig. 7: The examination result of differential scanning calorimetry, (a) monostearin, (b) meloxicam, (c) formula 3 (8:2), (d) formula 5 (9:1), (e) formula 4 (8.5:1.5), (f) formula 6 (9.5:0.5), (g) formula 2 (7.5:2.5), (h) formula 1 (7:3)

change after they are formed in NLC system. The DSC of NLC system result shows that enthalpy result (ΔH) is lower than enthalpy (ΔH) bulk lipid [4]. Based on DSC result, the melting point of NLC system was bigger than the bulk lipid. This result is opposite with NLC system's theory. This condition is possible because MLX active substance (257.34°C) has higher melting point than the solid lipid (63.58°C); it affects the melting point of NLC system. This research used MLX as a drug model; it is because MLX has some specific sharp peak which show that MLX is a crystalline drug (based on the X-ray analysis results). In literature, it is mentioned that if there is a crystalline drug in particle or a dispersion, DSC method cannot be used to know the changes of melting characteristics of NLC system compared with bulk lipid. A small amount of crystalline active substance can produce a wide area [14]. Therefore, it is possible that the crystallinity of active substance affects melting characteristics of NLC system and influences the DSC result inappropriate with the NLC's theory. The DSC result showed that the melting point has decreased when it was compared with the melting point of active substance. In F1 and F6, the melting point was higher than the other formulas. This condition could happen because of particle size effect. The decrease of melting point in NLC system is determined by the particle size which is getting smaller. Therefore, this phenomenon is called the "effect of particle size." The decrease of melting point can be described by Gibbs-Thomson equation [5].

The examination of particle morphology in NLC system

The examination result of particle morphology in NLC using SEM is shown in Fig. 8.

Based on the observation of particle morphology using SEM with a magnification of 10,000 times, each formula showed that the particle is almost spherical shaped. This particle size could produce the controlled release and avoid the agglomeration. A particle with a sphere shape has a little contact with media which causes longer diffusion process than the non-spherical particle. The spherical surface particle has little contact between particles that causes smaller agglomeration risk.

MLX entrapment efficiency

In a number of drugs, the solubility in liquid lipid is greater than in solid lipid, and the addition of liquid lipid to the solid lipid can damage the crystal lattice in the matrix of lipid nanoparticle, and then it can increase the entrapment efficiency of active ingredient [15]. Entrapment efficiency increases by increasing the concentration of liquid lipid. In NLC system, high solubility of active ingredient in liquid lipid plays an important role in increasing the entrapment efficiency of active ingredient [8].

The measurement of entrapment efficiency in the sixth formula of NLC system showed the high result (>70%). The measurement result of entrapment efficiency of MLX in NLC system was 74.50-86.22. The histogram of average percentage entrapment efficiency of some NLC formulas is shown in Fig. 9.

Statistical analysis of particle size using MANOVA showed that there was a significant difference between the formulas of NLC system. From *post-hoc* test, it could be known that there were significant differences between the percentages of entrapment efficiency in F1 and F2 with F3, F4, F5, and F6. F1 had the highest percentage of entrapment efficiency, and F6 had the lowest percentage of entrapment efficiency. This result is consistent with the concentration theory of liquid lipid which can produce the big entrapment efficiency, because of the active substance solubility in liquid lipid increases [8].

CONCLUSION

The MLX-NLC system with monostearin and alpha-tocopherol acetate as lipid matrix is characterized by small particle size, low crystallinity, spherical morphology of particle, and high MLX entrapment efficiency. Increasing the liquid lipid concentration can reduce the viscosity of the NLC system and improve the entrapment efficiency while there is no effect on the particle size and morphology characteristics. Concentration




Fig. 8: The test result of particle morphology in nanostructured lipid carriers system with $\times 10,000$ magnification using scanning electron microscopy. (a) Formula 1 (7:3), (b) formula 2 (7.5:2.5), (c) formula 3 (8:2), (d) formula 4 (8.5:1.5), (e) formula 5 (9:1), (f) formula 6 (9.5:0.5)




Fig. 9: The histogram of average % entrapment efficiency of all nanostructured lipid carriers' system formulas. The data obtained is the average of the pH value $\times 3$ replication formula $3 \pm$ standard deviation

ratio also affects the pH, PI, and the zeta potential, but the value is not equal with the addition of a lipid component. The results were possible because of the nature of the solid lipid.

REFERENCES

- Sweetman SC. Martindale The Complete Drug Reference. 26th ed. USA: Pharmaceutical Press.
- Bachhav YG, Patravale VB. Formulation of meloxicam gel for topical application: *In vitro* and *in vivo* evaluation. *Acta Pharm* 2010;60(2):153-63.
- Khurana S, Jain NK, Bedi PM. Development and characterization of a novel controlled release drug delivery system based on nanostructured lipid carriers gel for meloxicam. *Life Sci* 2013;93(21):763-72.
- Souto EB, Muller RH. Lipid nanoparticles (solid lipid nanoparticles and nanostructured lipid carriers) for cosmetic, dermal, and transdermal applications. *Drug Pharm Sci* 2007;166:213-32.
- Han F, Li S, Yin R, Liu H, Xu L. Effect of surfactants on the formation and characterization of a new type of colloidal drug delivery system

- nanostructured lipid carriers. *Int J Pharm* 2008;315(1):210-6.
6. Yuan H, Wang LL, Du YZ, You J, Hu FQ, Zeng S. Preparation and characteristics of nanostructured lipid carriers for control-releasing progesterone by melt-emulsification. *Colloids Surf B Biointerfaces* 2007;60(2):174-9.
 7. Kumar SG, Mishra DN. Preparation, characterization and *in vivo* dissolution studies of solid dispersion of meloxicam with PEG 6000. *Pharm Soc Jpn* 2006;126(8):657-64.
 8. Khurana S, Jain NK, Bedi PM. Development of nanostructured lipid carriers for controlled delivery of mefenamic acid. *Int J Biomed Nanosci Nanotechnol* 2012;2:232-50.
 9. Simon P. Formulasi dan Uji Penetrasi Mikroemulsi Natrium Diklofenak Dengan Metode Sel Difusi Franz dan Metode Tape Stripping. Tesis. Indonesia: Universitas Indonesia; 2012.
 10. Sinko PJ, Singh Y. *Martin's Physical Pharmacy and Pharmaceutical Sciences: Physical Chemical and Biopharmaceutical Principles in the Pharmaceutical Sciences*. 6th ed. Philadelphia PA: Lippincott Williams & Wilkins, A Wolters Kluwer; 2011.
 11. Jennings V, Gohla S. Comparison of wax and glyceride solid lipid nanoparticles (SLN). *Int J Pharm* 2000;196(2):219-22.
 12. Rosita N, Setyawan D, Soeratri W, Mrtodihardjo S. Physical characterization of beeswax and glyceryl monostearat binary system to predict characteristics of solid lipid nanoparticle (SLN) loaded para methoxy cinnamic acid (PMCA). *Int J Pharm Pharm Sci* 2014;6 Suppl 2:939-45.
 13. Xia D, Cui F, Gan Y, Mu H, Yang M. Design of lipid matrix particles for fenofibrate: Effect of polymorphism of glycerol monostearate on drug incorporation and release. *J Pharm Sci* 2014;103(3):697-705.
 14. Nastruzi C. *Characterization of solid lipid nano and microparticles. Liposphere in Drug Targets and Delivery*. London: CRC Press; 2004. p. 41-66
 15. Müller RH, Mäder K, Gohla S. Solid lipid nanoparticles (SLN) for controlled drug delivery - A review of the state of the art. *Eur J Pharm Biopharm* 2000;50(1):161-77.