Unity in Diversity and the Standardisation of Clinical Pharmacy Services

Editors: Elida Zairina, Junaidi Khotib,
Chrismanwan Ardianto, Syed Azhar Syed Sulaiman,
Charles D. Sands III and Timothy E. Welty

CRC Press
Taylor & Francis Group
A BALKEMA BOOK

Unity in Diversity and the Standardisation of Clinical Pharmacy Services

Editors

Elida Zairina

Department of Pharmacy Practice, Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia

Junaidi Khotib & Chrismawan Ardianto

Department of Clinical Pharmacy, Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia

Syed Azhar Syed Sulaiman

Discipline of Clinical Pharmacy, School of Pharmaceutical Sciences, Universiti Sains Malaysia, Penang, Malaysia

Charles D. Sands III

(Formerly) McWhorter School of Pharmacy, College of Health Sciences, Samford University, Birmingham, Alabama, USA

Timothy E. Welty

Department of Clinical Sciences, College of Pharmacy and Health Sciences, Drake University, Iowa, USA

CRC Press is an imprint of the
Taylor & Francis Group, an **informa** business

A BALKEMA BOOK

Organizing committee

ADVISORS

Mohammad Nasih, *Rector of Universitas Airlangga, Indonesia*
Nurul Falah Eddy Pariang, *President of Indonesian Pharmacist Association, Indonesia*

INTERNATIONAL ADVISORY BOARD

Suharjono, *Faculty of Pharmacy, Universitas Airlangga, Indonesia*
Junaidi Khotib, *Faculty of Pharmacy, Universitas Airlangga, Indonesia*
Yunita Nita, *Faculty of Pharmacy, Universitas Airlangga, Indonesia*
Aris Widayati, *Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia*
Dyah Aryani Perwitasari, *Faculty of Pharmacy, Universitas Ahmad Dahlan, Indonesia*
Surakit Nathisuwan, *Faculty of Pharmacy, Mahidol University, Thailand*
Suphat Subongkot, *Faculty of Pharmaceutical Sciences, Khon Kaen University, Thailand*
Roger D. Lander, *McWorther School of Pharmacy, Samford University, USA*
Timothy E. Welty, *College of Pharmacy and Health Sciences, Drake University, USA*
Charles D. Sands III, *Former Dean and Professor (retired), McWhorter School of Pharmacy, Samford University, USA*
Alexandre Chan, *Department of Pharmacy, National University of Singapore, Singapore*
Priscilla How, *Department of Pharmacy, National University of Singapore, Singapore*
Syed Azhar Syed Sulaiman, *School of Pharmaceutical Sciences, Universiti Sains Malaysia*
Hiroyuki Kamei, *Faculty of Pharmacy, Meijo University, Japan*
Kwang-il Kwon, *College of Pharmacy, Chungnam National University, Korea*
Robert Sindelar, *Faculty of Pharmaceutical Sciences, The University of British Columbia, Canada*

STEERING COMMITTEE

Umi Athiyah, *Faculty of Pharmacy, Universitas Airlangga, Indonesia*
Riesta Primaharinastiti, *Faculty of Pharmacy, Universitas Airlangga, Indonesia*
Dwi Setiawan, *Faculty of Pharmacy, Universitas Airlangga, Indonesia*
Dewi Melani Haryadi, *Faculty of Pharmacy, Universitas Airlangga, Indonesia*
Wahyu Utami, *Faculty of Pharmacy, Universitas Airlangga, Indonesia*
Budi Suprpti, *Faculty of Pharmacy, Universitas Airlangga, Indonesia*
Agung Endro Nugroho, *Faculty of Pharmacy, Universitas Gadjah Mada, Indonesia*
Pinus Jumaryatno, *Faculty of Mathematics and Natural Sciences, Universitas Islam Indonesia*
Sabtanti Harimurti, *Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Indonesia*
Chao Zhang, *Peking University Third Hospital, China*
Helen Zhang, *United Family Health Care, China*
Zhu Zhu, *Peking Union Medical College Hospital, China*
Suo-Zhong Yuan, *Chinese Pharmaceutical Association, China*
Franco Cheng, *School of Pharmacy, The Chinese University of Hongkong, Hong Kong*

Vivian Lee Wing-Yan, *School of Pharmacy, The Chinese University of Hongkong, Hong Kong*
 Wan Gyoong Shin, *College of Pharmacy, Seoul National University, Korea*
 Kyung Eob Choi, *College of Pharmacy, CHA University, Korea*
 Jung Mi Oh, *College of Pharmacy, Seoul National University, Korea*
 Hyun-Take (Thomas) Shin, *College of Pharmacy, Sookmyung Women's University, Korea*
 Ramesh Adepu, *JSS College of Pharmacy, JSS University, India*
 Farshad Hashemian, *Pharmaceutical Sciences Branch, Islamic Azad University, Iran*
 Mehdi Rajabi, *Pharmaceutical Sciences Branch, Islamic Azad University, Iran*
 Takao Shimazoe, *Graduate School of Pharmaceutical Sciences, Kyushu University, Japan*
 Yolanda R. Robles, *College of Pharmacy, University of the Philippines, Philippines*
 Hazel Faye Ricaforte-Docuyan, *Philippines Society of Hospital Pharmacists, Philippines*
 Ng Hong Yen, *Pharmaceutical Society of Singapore, Singapore*
 Vivianne Shih Lee Chuen, *Pharmaceutical Society of Singapore, Singapore*
 Sutthiporn Pattarachayakul, *Faculty of Pharmaceutical Sciences, Prince of Songkla University, Thailand*
 Apornee Chaiyakum, *Faculty of Pharmaceutical Sciences, Khon Kaen University, Thailand*
 Nguyen Van Hung, *Faculty of Pharmacy, Haiphong University Medicine and Pharmacy, Vietnam*
 Pham Minh Hung, *Faculty of Pharmacy, Haiphong University Medicine and Pharmacy, Vietnam*
 Marshall E. Cates, *McWorther School of Pharmacy, Samford University, USA*
 Michael Hogue, *McWorther School of Pharmacy, Samford University, USA*
 James E. Tisdale, *College of Pharmacy, Purdue University, USA*
 Alan Lau, *College of Pharmacy, University of Illinois at Chicago, USA*
 Amrizal Marzuki, *Indonesian Society of Hospital Pharmacists, Indonesia*
 Saleh Rustandi, *Indonesian Society of Community Pharmacy, Indonesia*

SCIENTIFIC COMMITTEE

Elida Zairina, *Faculty of Pharmacy, Universitas Airlangga, Indonesia*
 Chrismawan Ardianto, *Faculty of Pharmacy, Universitas Airlangga, Indonesia*
 Arie Sulistyarini, *Faculty of Pharmacy, Universitas Airlangga, Indonesia*
 Mufarrihah, *Faculty of Pharmacy, Universitas Airlangga, Indonesia*
 Rita Suhadi, *Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia*
 Suci Hanifah, *Faculty of Mathematics and Natural Sciences, Universitas Islam Indonesia, Indonesia*
 Fita Rahmawati, *Faculty of Pharmacy, Universitas Gadjah Mada, Indonesia*
 M. Saiful Bachri, *Faculty of Pharmacy, Universitas Ahmad Dahlan, Indonesia*
 Aluwi Nirwana Sani, *Indonesian Pharmacist Association, Indonesia*
 Rizka Andalusia, *Indonesian Society of Hospital Pharmacists, Indonesia*
 Widayati, *Indonesian Society of Hospital Pharmacists, Indonesia*

CONFERENCE CHAIRMAN

Suharjono, *Faculty of Pharmacy, Universitas Airlangga, Indonesia*

CONFERENCE VICE-CHAIRMAN 1

Junaidi Khotib, *Faculty of Pharmacy, Universitas Airlangga, Indonesia*

CONFERENCE VICE-CHAIRMAN 2

Yunita Nita, *Faculty of Pharmacy, Universitas Airlangga, Indonesia*

SECRETARY

Anila Impian Sukorini, *Faculty of Pharmacy, Universitas Airlangga, Indonesia*
 Ana Hidayati, *Faculty of Pharmacy, Universitas Ahmad Dahlan, Indonesia*

FOOD AND BEVERAGE

Dewi Wara Shinta, *Faculty of Pharmacy, Universitas Airlangga, Indonesia*

Yosi Febrianti, *Faculty of Mathematics and Natural Sciences, Universitas Islam Indonesia, Indonesia*

Pinasti Utami, *Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Indonesia*

FACILITIES AND EQUIPMENT

Catur Dian Setiawan, *Faculty of Pharmacy, Universitas Airlangga, Indonesia*

Christianus Heru Setiawan, *Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia*

Yulianto, *Faculty of Mathematics and Natural Sciences, Universitas Islam Indonesia, Indonesia*

ACCOMMODATION AND TRANSPORTATION

Mutiara Herawati, *Faculty of Mathematics and Natural Sciences, Universitas Islam Indonesia, Indonesia*

Dita Maria Virginia, *Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia*

Mawardi Ihsan, *Faculty of Pharmacy, Universitas Gadjah Mada, Indonesia*

Table of contents

Preface	ix
Organizing committee	xi
Keynote speakers	xv
Plenary speakers	xvii
List of symposium speakers	xix
Medication management system in several care homes in Surabaya <i>G.N.V. Achmad, G. Nugraheni, W. Utami, S. Hardiyanti, S. Danutri, D.K. Lestari, Muhliseh & A.T. Mahardika</i>	1
Effectiveness of decision aid on knowledge, decision conflict, and outcome in diabetic patients <i>L. Aditama & F. Yulia</i>	7
Self-esteem scale: Translation and validation in Malaysian adults living with asthma <i>S. Ahmad, M. Qamar, F.A. Shaikh, N.E. Ismail, A.I. Ismail & M.A.M. Zim</i>	13
Factors affecting mortality among patients undergoing hemodialysis in Sudan <i>O. Amir, A. Sarrieff, A.H. Khan, M.B. Abdelraheem & B. Norsa'adah</i>	19
Evaluating the effectiveness of filgrastim in patients with solid cancer <i>E.N. Anggraeny, F. Rahmawati & K.W. Taroen-Hariadi</i>	27
Building care of hypertensive patients in reducing sodium intake in Banjarmasin <i>H. Ariyani, Akrom & R. Alfian</i>	33
Effect of glibenclamide on glycemic control in the presence of sodium diclofenac <i>T. Aryani, M.N. Zamzamah, Z. Izzah & M. Rahmadi</i>	41
A drug utilization study of antibiotics in patients with osteomyelitis <i>A.S. Budiadin, K.D. Kurdiana, B.S. Zulkarnain, H. Suroto & R. Diniya</i>	45
Factors influencing trastuzumab cardiotoxicity in breast cancer: A case-control study <i>C. Cherachat, C. Sukkasem, C. Somwangprasert, S. Pornbunjerd, N. Parinyanitikul & K. Tewthanom</i>	51
Transdermal patch loading diclofenac sodium for anti-inflammation therapy using a rat paw oedema model <i>P. Christanto, Isnaeni, A. Miatmoko & E. Hendradi</i>	55
Ethambutol-induced optic neuropathy in diabetic patients with tuberculosis <i>M. Djunaedi, U. Athiyah, Y. Priyandani & S.A.S. Sulaiman</i>	61
Identification of clinical specimens isolated from neonates <i>M. Djunaedi, S.A.S. Sulaiman, A. Sarrieff, N.B.A. Aziz & Habsah</i>	65
Effect of coenzyme Q10 on the malondialdehyde level and exercise performance of male runners in Jakarta <i>S. Gunawan, Purwastyastuti, F.D. Suyatna & E.I. Ilyas</i>	73
Compatibility of selected inotropic drugs in a syringe <i>S. Hanifah, R.A. Kennedy & P.A. Ball</i>	81

Ward pharmacist workload analysis in an Indonesian class A hospital <i>A.L. Hariadini, W. Utami & A. Rahem</i>	87
Effect of LD50 of ethanolic leaf extract from <i>Ipomoea reptans</i> Poir. in rats <i>F. Hayati, R. Istikharah, S. Arifah & D. Nurhasanah</i>	95
Oral indomethacin versus oral paracetamol for patent ductus arteriosus closure in neonates <i>N.A. Ibrahim, N.C. Umar, M.C.H. Chi'ing, P.E. Stephen & P. Anandakrishnan</i>	99
Clinical trial of Jamu X on blood glucosa and HbA1C in patients with type 2 diabetes mellitus <i>Z. Ikawati, M. Eko Cahyanto, N.R. Sholehah & N. Atikah</i>	107
Management of carbamate or organophosphate intoxication at a high care unit <i>Z. Izzah, T. Aryani, R. Rodhika & Lestiono</i>	113
Development of an antidepressant e-learning tool for pharmacology education <i>A. Karaksha, M. Dharmesti, A.K. Davey, G. Grant, S.A. Dukie, H. Budianto, E.V. Mutiara, V. Marina, I. Puspitaningrum & A. Shollina</i>	119
Effectiveness of nimodipine on non-traumatic subarachnoid hemorrhage based on computed tomography angiography <i>J. Khotib, S.S. Ganesen & A.F. Sani</i>	125
Study of pediatric compounded drug prescriptions in a health care facility in Bandung <i>F. Lestari, Y. Aryanti & U. Yuniarni</i>	133
Efficacy of honey vinegar in hyperlipidemic rats (<i>Rattus norvegicus</i>) <i>E.W. Lucia, K. Lidya & T. Annisa</i>	139
Improving the competence of pharmacist students through international lecturers <i>O.R. Mafruhah, S. Hanifah, C.P. Sari, P.A. Ball & H. Morrissey</i>	143
The effectiveness evaluation of antiretroviral therapy in Mangusada Hospital Bali <i>H. Meriyani, N.N.W. Udayani & K.A. Adrianta</i>	149
Drug therapy problems in pediatric and geriatric patients at Farmasi Airlangga Pharmacy <i>Mufarrihah, D.M. Machfud, V.D.A. Purworini, A. Yuda, Y. Priyandani & Y. Nita</i>	153
PCR primer design for detection of SNPs in SLC22A1 rs683369 encoding OCT1 as the main transporter of metformin <i>A.A. Mukminatin, V.D.A. Ningrum & R. Istikharah</i>	161
Pharmacist-patient communication: An observational study of characteristic information <i>I. Mulyono, S. Irawati, A. Pratidina & M. Claramita</i>	167
Multidrug resistance-1 gene variants in pediatric leukemia in Bali <i>R. Niruri, N.L. Ulandari, S.C. Yowani, I. Narayani & K. Ariawati</i>	171
Medication adherence in the elderly with chronic diseases using the Adherence to Refill and Medication Scale (ARMS) <i>Y. Nita, F.M. Saputra, S. Damayanti, P.I. Pratiwi, R. Zukhairah, A. Sulistyarini & Y. Priyandani</i>	175
Direct non-medical costs of patients with cervical cancer who underwent chemotherapy <i>R. Noviyani, P.A. Indrayathi, H. Thabrany, Andrijono, I.N.G. Budiana & K. Tunas</i>	179
Factors influencing correct measurements of liquid medicines by consumers <i>G. Nugraheni & G.N.V. Achmad</i>	185
The correlation of service quality and complaint handling with patient satisfaction <i>R.A. Oetari, M.E. Sariwatin, H. Basir & C. Wiedyaningsih</i>	189
Assessment of medication safety among Filipino pharmacists <i>R.C. Ongpoy Jr., P.P. David, R.C. Ongpoy, M.D.U. Dean & A.D. Atienza</i>	195
Effect of the combination of <i>Typhonium flagelliforme</i> Lodd. (Blume) and <i>Phyllanthus niruri</i> Linn. on the immune system <i>S.S. Pangestika, A.P. Gani, A. Yuswanto & R. Murwanti</i>	201

Organophosphate toxicity in red chili farmers, Ciamis, Indonesia <i>D.A. Perwitasari, D. Prasasti, I.W. Arsanti & I.A. Wiraagni</i>	207
Factors affecting the rational use of NSAIDs in self-medication <i>L. Pristianty, G.N.V. Achmad & A. Faturrohman</i>	213
The influence of adverse reactions of antituberculosis drugs to non-adherence in drug use <i>Y. Priyandani, C.D. Setiawan, A. Yuda, Y. Nita, U. Athiyah, M.B. Qomaruddin & Kuntoro</i>	217
(-)-Epigallocatechin gallate from green tea increases the level of a DNA repair enzyme <i>D.A. Purwanto</i>	223
Perceptions and practices of self-medication among healthcare students <i>M. Qamar, S. Norhazimah, F.A. Shaikh & S. Ahmad</i>	227
Smoking cessation counseling: Perceptions and barriers among community pharmacists <i>M. Qamar, S. Ahmad, K. Poobalan, F.A. Shaikh & M.A. Hammad</i>	233
Factors affecting medication noncompliance in patients with chronic diseases <i>A. Rahem</i>	241
The development and evaluation of a clinical pharmacy course at a pharmacy school in Indonesia <i>F. Rahmawati, D. Wahyono & M. Ihsan</i>	245
Relationship ejection fraction and segment ST-resolution in STEMI patients with streptokinase therapy <i>D.M.N. Ratri, S. Sjamsiah, H.P. Jaya & M. Aminuddin</i>	251
Management of hyponatremia in patients with heart failure: A retrospective study <i>S. Saepudin, P.A. Ball & H. Morrissey</i>	255
Continuous infusion versus intermittent bolus furosemide in heart failure NYHA III-IV <i>Samirah, S. Sjamsiah & M. Yogiarto</i>	261
The effectiveness of empirical and definitive antimicrobial therapy <i>I.P. Sari, R.H. Asdie, T. Nuryastuti, Sugiyono & Sumaryana</i>	265
<i>In vivo</i> analgesic effect of ethanolic extracts of exocarp, mesocarp, and seeds of <i>Carica pubescens</i> <i>H. Sasongko & Sugiyarto</i>	271
Evaluation of knowledge, attitude and perceived barriers towards adverse drug reaction reporting <i>F.A. Shaikh, S. Ahmad, E. Intra, M. Qamar & T.M. Khan</i>	275
Hydroxyethyl starch or gelatin, which is safer for the kidneys? <i>D.W. Shinta, J. Khotib, M. Rahmadi, B. Suprati, E. Rahardjo & J.K. Wijoyo</i>	281
In silico QSAR of 1-benzoyl-3-benzylurea lead and its analogue compounds as anticancer <i>F. Suhud, C. Effendi & Siswandono</i>	287
A study on antiemetics for postoperative nausea and vomiting at Dr. Soetomo Hospital <i>Suharjono, M.E.B.M.A. Nazim, B.P. Semedi & R. Diniya</i>	293
Ethanol extract of <i>Annona squamosa</i> L. improves the lipid profile in hyperlipidemia rats <i>R. Sumarny, Y. Sumiyati & D. Maulina</i>	297
The effect of Telmisartan on lipid levels and proinflammatory cytokines in ESRD patients undergoing hemodialysis <i>B. Suprati, W.P. Nilamsari, Z. Izzah, M. Dhrik & B. Dharma</i>	303
Risk assessment of ADEs: Patient safety incident reports at Ari Canti Hospital in 2016 <i>D.A. Swastini, N.W.S. Wahyuni & K. Widiyantara</i>	307
Medication-induced Adverse Drug Reaction (ADR) in the Malaysian elderly population <i>H.M. Taib, Z.A. Zainal, N.M. Ali & R. Hashim</i>	311

Cost-effectiveness analysis of patients with schizophrenia in Madani Hospital <i>M.R. Tandah, A. Mukaddas & W. Handayani</i>	317
Tacrolimus-induced symptomatic hyponatremia after kidney transplantation: A case study <i>T. Verayachankul & J. Tantivit</i>	323
Postoperative pain management in elderly patients: Evaluating the use of analgesics <i>A. Vonna, A. Apriani & Sadli</i>	327
Antimalarial activity and toxicological test of <i>Andrographis paniculata</i> tablets (AS202-01) <i>A. Widyawaruyanti, A.F. Hafid, D.A. Fitriiningtyas, L.S. Lestari, H. Ilmi & I.S. Tantular</i>	333
Socioeconomic status and obesity in an adult rural population in Indonesia <i>A. Widayati, Fenty, D.M. Virginia & P. Hendra</i>	339
A strategic approach to increase the compliance of patients with type 2 diabetes mellitus <i>N. Wulandari, D. Vivianhari & Nurhayati</i>	343
Disposal practices of unused medication among the public in Meradong, Sarawak, Malaysia <i>N.L.C. Yaacob, L.P. Wei, S. Ahmad, F. Naimat & A. Ahmad</i>	349
Drug use and potential drug interaction in the elderly <i>A. Yuda, E.C. Dewi, D.M. Fami, L. Jamila, M. Rakhmawati, K.P.P. Sari, K.P. Ningrum, G.N.V. Achmad & Y. Nita</i>	357
Effects of audiovisual education on the knowledge and adherence of patients with DMT1 <i>L.Y. Yusan, N. Rochmah, A. Rahem & A. Purnamayanti</i>	363
Root extract of <i>Imperata cylindrica</i> L. improves serum nitric oxide levels in diabetic mice <i>A. Zada, J.B. Dewanto, A. Dahlan, D. Dhianawaty, M.R.A.A. Syamsunarno, G.R. Mukarromah & N. Anggraeni</i>	367
Medication use during pregnancy in Surabaya: A cross-sectional study <i>E. Zairina, G. Nugraheni, G.N. Veronika Ahmad, A. Yuda, Y. Nita, M.P. Wardhana & K.E. Gumilar</i>	371
Eyedrops use perception during fasting <i>B.S. Zulkarnain, Sumarno, Y. Nita & R. Loebis</i>	375
Author index	381

Transdermal patch loading diclofenac sodium for anti-inflammation therapy using a rat paw oedema model

P. Christanto, Isnaeni, A. Miatmoko & E. Hendradi

Department of Pharmaceutics, Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia

Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia

ABSTRACT: The anti-inflammatory effect of transdermal delivery of a diclofenac sodium patch was evaluated. The patch matrix consists of ethyl cellulose N-20 and polyvinyl pyrrolidone K-30 to control the drug release. In this study, the patches were prepared with ethyl cellulose N-20 (EC-N20) and polyvinylpyrrolidone K-30 (PVP K-30) at weight ratio of 6:4 and 7:3 for EC/PVP-6/4 and EC/PVP-7/3 patches, respectively. The anti-inflammatory effect was determined by evaluating the swelling of rat's paw oedema that was induced with 1% carrageenan suspension. The results showed that the high concentration of PVP K-30 resulted in less rigid patch with pore structures. In addition, it improved the anti-inflammatory effect of diclofenac sodium resulted in higher efficacy of EC/PVP-6/4 than that of EC/PVP-7/3. There were no significant differences on drug stability observed for both formulations. It can be concluded that controlling diclofenac sodium released from patch using PVP K-30 could give benefits for anti-inflammation therapy.

1 INTRODUCTION

Diclofenac sodium is a Non-Steroidal Anti-Inflammatory Drug (NSAID) that is widely used to relieve pain and inflammation. Diclofenac sodium has fast absorption by oral administration, but only about 60% of the total drug amount reaches the systemic circulation. This phenomenon is caused by the first-pass metabolism that occurs in the liver (Chuasuan et al. 2008). The half-life of diclofenac sodium is very short, which is approximately two hours. Moreover, it produces severe side effects on the gastrointestinal tract i.e. stomach ulcers. Transdermal delivery may provide an alternative solution to overcome this problem.

Recently, commercial transdermal products of diclofenac sodium available on the market are mostly in topical liquids, gels or creams that are applied directly to the skin. However, these products provide only short-term pharmacological effects. The use of transdermal patch can provide many advantages for delivery of diclofenac sodium. The patch can control the drug release, so it can be used for long-term drug administration to achieve systemic dosing (Rathbone et al. 2002). The use of patch can also avoid gastrointestinal irritation, minimize pain, and bypass the hepatic first-pass metabolism. Moreover, due to its ability to control the release of drugs for extended and safe use, it reduces the occurrence of fluctuations and being beneficial for drugs that have very short half-life times and narrow therapeutic ranges. Thereby, it can increase the patient compliance (Kumar & Philip 2007).

It has been known that diclofenac sodium has low molecular weight, which is less 500 daltons, and partition coefficient or Log P of 1.1–1.3 (Kweon et al. 2004, Chuasuan et al. 2008). These properties have been reported to be ideal for drug delivery using transdermal patch (Rathbone et al. 2002).

Polymer has an important role in regulating the release of the drug in the matrix-patch type. Modification of polymer properties using combination of hydrophilic and lipophilic polymers, such as polyvinyl-pyrrolidone (PVP) and ethyl cellulose (EC) is useful for achieving good drug release rate (Kandavilli et al. 2002, Rathbone et al. 2002). To improve drug penetrated into the skin, the addition of penetration enhancers, such as menthol, can be considered.

In this study, we evaluated the anti-inflammatory effects of diclofenac sodium patch at a dose of 14.13 mg/7.065 cm². The patch was prepared with combination of EC N-20 and PVP K-30 at weight ratio of 6:4 and 7:3. These formulations also contain menthol and polyethylene glycol (PEG)-400 as they were previously reported as important components to prepare patch using the controlled matrix method (Hendradi et al. 2011). The anti-inflammatory effect was then evaluated by determine the thickness of oedema on hind paw of Wistar rats that was induced by injecting carrageenan suspension. Moreover, stability of patch was also determined as they likely affect the therapeutic effects of the preparation.

2 METHODS

2.1 Materials

Diclofenac sodium was purchased from Aarti Drug Ltd. (Tarapur, India). Ethyl cellulose (EC) N-20 was bought from Dow Chemical Company (Midland, USA). Polyvinylpyrrolidone (PVP) K-30 was a product of ISP Pte. Ltd. (Singapore). All other chemicals are the finest grade available.

2.2 Preparation of diclofenac sodium patch

Diclofenac sodium patch was prepared by using the controlled matrix method as previously reported (Hendradi et al. 2011). The patch composed of polymeric matrix, plasticizer, and penetration enhancer as shown in Table 1. The patches had surface area of 7.065 cm². The formulation was generated in triplicates.

2.3 Evaluation of physical characteristics of diclofenac sodium patch

2.3.1 Organoleptic properties

The organoleptic properties of diclofenac sodium patch were evaluated by visual observation of physical appearances and colour of patch, and smelling of its odour.

2.3.2 Moisture content

The Moisture Content was determined by weigh measurement of patch before and after desiccation. The patch was put in a safety cabinet and its mass was weighed. Then, this patch was put in a silica gel desiccator and allowed to stand for 24 hours. After 24 hours, the patch was removed from desiccator and measured for its weight. The moisture content (%) of patch was calculated as follows:

$$\% \text{Moisture Content} = \frac{w_1 - w_2}{w_2} \times 100 \%$$

Table 1. The formulation of diclofenac sodium patch.

Component	Use	Formulation	
		EC/PVP-6/4	EC/PVP-7/3
Diclofenac sodium	Active drug	14 mg	14 mg
EC N-20	Polymeric matrix	169 mg	197 mg
PVP K-30	Polymeric matrix	113 mg	85 mg
PEG-400	Plasticizer	73 mg	73 mg
Menthol	Penetration enhancer	4 mg	4 mg

which w_1 is diclofenac sodium patch's weight before being put in the desiccator, and w_2 is diclofenac sodium patch weight after being put in a desiccator for 24 hours (Patel et al. 2009, Hendradi et al. 2011).

2.4 Surface morphology and homogeneity of diclofenac sodium concentration of patch

Determination of surface homogeneity test of patch was evaluated using scanning electron microscopy. The homogeneity of diclofenac sodium concentration in patch was carried out by dividing the patch into four equal parts. Each part was then measured for its level of diclofenac sodium in phosphate buffer saline (PBS) pH 7.4 using UV spectrophotometer at λ of 276 nm. The measurement was in triplicates.

2.5 Evaluation of anti-inflammatory effects in rats

All animal experiments were performed through approval of ethical clearance committee of Universitas Airlangga, Indonesia. The Wistar rats with age of two months old were used as the experimental animals. They were divided into four groups, which each group consists of 4 animals. The negative control groups were treated with placebo patches prepared with the combination of EC N-20 and PVP K-30 at weight ratio of 7:3 and 6:4 for EC/PVP-6/4 and EC/PVP-7/3 groups, respectively, and do not contain diclofenac sodium. The treatments groups received diclofenac sodium patches with a dose of 14.13 mg/7.065 cm²/rat

and prepared with combination of EC N-20 and PVP K-30 in the weight ratio of 7:3 and 6:4 for EC/PVP-6/4 and EC/PVP-7/3 groups, respectively.

The anti-inflammatory effect of diclofenac sodium patches were evaluated using rat's paw oedema model as previously reported (Hendrati et al. 2003). The patches were applied onto the abdominal skin of the rats. At one hour later, the left rear foot plantar tissue of Wistar rat (two months old) was injected with suspension of carrageenan for oedema induction. The anti-inflammatory effect was determined by measuring the thickness of rat hind paw with the long sliding every 0.5 hours.

2.6 Stability of diclofenac sodium patch

Physical stability test of diclofenac sodium patch was conducted for three months, which were at day 7, 21, 54 and 84, by visual observation of consistency, physical appearances, and odour. In addition, the chemical stability tests were also carried out by determining the diclofenac sodium concentration

of patch during the same interval periods using spectrophotometric method at 276 nm. The measurements were in triplicates.

2.7 Data analysis

The anti-inflammatory effect of patch on reduction of inflammation was calculated using the following formula:

$$\% \text{ anti-inflammatory effect} = \frac{H_t - H_0}{H_0} \times 100 \%$$

which H_t is the thickness of swollen rat hind paw after carrageenan injection at determined measurement time, and H_0 is the thickness of swollen rat hind paw before rats were injected with carrageenan (Hendradi et al. 2003). The significance of differences on reduction of inflammation was statistically analyzed using one-way analysis of variance. A p value of 0.05 or less was considered significant.

3 RESULTS AND DISCUSSIONS

In this study, we prepared diclofenac sodium patch prepared with different weight ratio of PVP K-30 and EC N-20 to control the release of drug. Reducing the concentration of PVP K-30 showed enhanced anti-inflammatory effects on rat's hind paw oedema.

The organoleptic evaluation shows that diclofenac sodium patch EC/PVP-6/4 that was prepared with EC N-20 and PVP K-30 at weight ratio of 6:4, respectively, had more flexible consistency (Table 1). It may because of the low amount of PVP K-30. Since PVP K-30 is a hygroscopic and hydrophilic polymer, it may cause the patch absorbing water molecules from the air, thus becoming more flexible than EC/PVP-7/3.

Moreover, EC/PVP-6/4 had more transparent appearance than EC/PVP-7/3, as shown in Fig. 1 A,B. It is because the use of less amount of EC, which is a water insoluble hydrophobic polymer, than that of EC/PVP-6/4. Both formulations have the minty fresh smell (Table 1) due to the addition of menthol as penetration enhancer in patch.

Table 2. Organoleptic properties of diclofenac sodium patch at day 0.

Formulation	Consistency	Appearance	Odour
EC/PVP-6/4	Less rigid	Transparent	Minty fresh
EC/PVP-7/3	Rigid	Less transparent	Minty fresh

By using scanning electron microscopy (SEM), it can be seen that EC/PVP-6/4 patch have more pores and larger pore size than EC/PVP-7/3 patch as shown in Fig. 2. Since EC N-20 is a hydrophobic polymer, the low amount of this polymer in EC/PVP-6/4 resulted in hydrophilic patch with extensive pore structures (Fig. 2 A,B). On the other hand, the high amount of EC N-20 in EC/ PVP-7/3 produced tightened structures, thus having small amount of pore with relative small pore size (Fig. 2C,D). The addition of diclofenac sodium had no effect on the physical structures of patches.

The moisture content analysis indicates that EC/PVP-6/4 patch had slightly higher moisture content than EC/PVP-7/3 (Table 3). It is due to the PVP K-30 content, which is a hygroscopic polymer. However, the difference between these two patch formulations was negligible.

Figure 1. Physical appearances of diclofenac sodium patch prepared with EC N-20 and PVP K-30 at weight ratio of 6:4 and 7:3 for EC/PVP-6/4 (A), and EC/PVP-6/4 (B), respectively.

Figure 2. Scanning electron microscope (SEM) pictures of matrix type patch prepared with EC N-20 and PVP K-30 at weight ratio of 6:4 for EC/PVP-6/4 placebo patch (A), EC/PVP-6/4 loading diclofenac sodium (B), and 7:3 for EC/PVP-7/3 placebo patch (C), and EC/PVP-6/4 containing diclofenac sodium (D). Scale bar is 50 μm .

Table 3. Moisture content (%) and drug homogeneity of diclofenac sodium patches at day 0 (n = 3).

Formulation	Moisture content (%)	Drug homogeneity (%)
EC/PVP-6/4	12.81 ± 0.65	100.22 ± 0.61
EC/PVP-7/3	10.95 ± 0.11	98.56 ± 0.83

Figure 3. The anti-inflammatory effect of diclofenac sodium-loaded patch on the carrageenan-induced rat paw oedema (n = 4). Each value represents mean $\pm$ S.D. (n = 4). * $P < 0.05$.

Moreover, drug homogeneities of the two patch formulations were 98.56 ± 0.83 and $100.22 \pm 0.61\%$; these results show that both formulations produced homogeneous diclofenac sodium patches and meet product homogeneity requirements with percent of variation coefficient less than 2% (Departemen Kesehatan 2014).

In the in vivo study, there were significant differences on oedema reduction between diclofenac sodium-contained patch applications and placebo patches. Moreover, a significant higher increase in the anti-inflammatory effects was observed in the EC/PVP-6/4 treatment group than that of EC/PVP-7/3 (Fig. 3). It is possibly due to the large pores present in EC/PVP-6/4 patch enable drug to be dissolved, released, and readily penetrate into the skin, thus causing reduction of the inflammation. On the other hand, the tightened patch pore structure of EC/PVP-7/3 limits this process producing weaker anti-inflammatory effects than that of EC/PVP-6/4, although the concentration of diclofenac sodium released from this patch formulation was still able to cure the inflammation. In addition, according to the calculation of area under curve (AUC) of swelling thickness of rat's hind paw, it can be seen that administration of EC/PVP-6/4 patch into abdominal skin of rats produced the largest decrease on the paw oedema as shown in Fig. 4. Beside the pores, the use of menthol as penetration enhancer also plays important roles in enhancing anti-inflammatory effects of diclofenac sodium patch. It has been known that menthol affects the integrity of stratum cor-neum layers, thus improving the penetration of diclofenac sodium.

After storage at room temperature, the stability was evaluated at day 7, 21, 56 and 84. There were no changes observed in the organoleptic properties between patches determined at observation day and those of day 0 (Table 2). The consistency, physical appearances and odour of EC/PVP-6/4 and EC/PVP-7/3 patches were similar. Moreover, the moisture contents of patches, either EC/PVP-6/4 or EC/PVP-7/3, during observation periods had no significant differences from that of the day 0, as shown in Fig. 5. The moisture contents were about 11% and 13% for EC/PVP-6/4 and EC/PVP-7/3, respectively.

Figure 4. The areas under curve (AUC) of hind paw thickness of carrageenan suspension-induced rat oedema after topical administration of the placebo and diclofenac sodium-loaded patches onto abdominal skin (n = 4).

Figure 5. The stability evaluation for moisture content (%) of diclofenac sodium-loaded patches i.e. EC/PVP-6/4 and EC/PVP-7/3 stored at room temperature (25°C) for 84 days (n = 3).

Figure 6. The concentration of diclofenac sodium loaded in EC/PVP-6/4 and EC/PVP-7/3 patches during the storage at room temperature (25°C, n = 3).

In addition, it can be seen that the concentration of diclofenac sodium in the patches were stable until 84 days stored at room temperature, which were still within the range of 95–105% (Figure 6). This result suggests that the use of different ratio of EC N-20 and PVP K-30 in patch formulation have no different effects on chemical stability of the drug and both of them could maintain the stability of diclofenac sodium loaded in the patches.

The use of EC N-20 and PVP K-30 as polymeric matrix with menthol addition in patch could improve transdermal delivery of diclofenac sodium. It produced transdermal patch with good organoleptic properties, physically and chemically stable during storage at room temperature, and importantly, improved anti-inflammatory effects of diclofenac sodium on rat's paw edema by topical administration. Mostly, drug penetration into the skin is very limited, thus being the great problem for achieving good drug therapy. It is important to design a carrier for controlling drug release and enhancing skin drug penetration in an appropriate manner for achieving high drug efficacy. However, further investigation is still required to evaluate physicochemical interaction of EC N-20 and PVP K-30 for enhancing the therapeutic outcomes of patch loading diclofenac sodium.

4 CONCLUSIONS

In this study, to deliver diclofenac sodium for anti-inflammation therapy, we prepared patches using combination of EC N-20 and PVP K-30 as polymeric matrix and evaluated their anti-inflammatory effects using rat paw oedema model. High reduction on paw swelling was achieved in EC/PVP-6/4 patch that was prepared with the high amount of PVP K-30. This finding suggested that EC N-20 and PVP K-30 at weight ratio of 6:4, respectively, can act as an excellent polymeric matrix for transdermal patch.

ACKNOWLEDGEMENTS

We would like to thank Ministry of Research, Science, Technology and Higher Education (KEMENRISTEK-DIKTI) of Indonesia for providing funding for this research work.

REFERENCES

- Chuasuwana, B., Binjesoh, V., Polli, J.E., Zhang, H., Amidon, G.L., Junginger, H.E., Midha, K.K., Shah, V.P., Stavchansky, S., Dressman, J.B., Barends, D.M. 2008. Biowaiver monographs for immediate release solid oral dosage forms: diclofenac sodium and diclofenac potassium. *Journal of Pharmaceutical Sciences*: 1206–1219.
- Departemen Kesehatan. 2014. *Farmakope Indonesia V*. Departemen Kesehatan, Jakarta: 1526–1528.
- Hendrati, E., Obata, Y., Isowa, K., Nagai, T., Takayama, K. 2003. Effect of mixed micelle formulations including terpenes on the transdermal delivery of diclofenac. *Biological and Pharmaceutical Bulletin* 26(12): 1739–1743.
- Hendrati, E., Isnaeni, Fridayanti, A., Efrin, P. 2011. Optimasi sediaan transdermal patch natrium diclofenak tipe matriks. *Jurnal Farmasi Indonesia* 5(3): 112–119.
- Kandavilli, S., Nair, V., Panchagnula, R. 2002. Polymer In Transdermal drug delivery systems. *Pharmaceutical Technology*: 62–80.
- Kumar, R. and Philip, A. 2007. Modified transdermal technologies: breaking the barriers of drug permeation via the skin. *Tropical Journal of Pharmaceutical Research* 6 (1): 633–644.
- Kweon, J.H., Chi, S.C., Park, E.S. 2004. Transdermal delivery of diclofenac using microemulsions. *Archives of Pharmacal Research* 27(3): 351–356.
- Patel, R.P., Patel, G., Baria, A. 2009. Formulation and evaluation of transdermal patch of Aceclofenac. *International Journal of Drug Delivery* 1:41–51.
- Rathbone, M.J., Hadgraft, J., Robert, M.S. 2002. *Modified release drug delivery technology* 1–40 (90–92): 471–512. New York: Marcel Dekker.