New cis-ent-clerodanes from *Linaria japonica*

by Retno Widyowati

Submission date: 20-Sep-2018 10:04AM (UTC+0800)

Submission ID: 1005033738

File name: Bukti_C-1.pdf (1,012.26K)

Word count: 5894

Character count: 25320

Short communication

New *cis-ent*-clerodanes from *Linaria japonica*Retno Widyowati^{a,b}, Sachiko Sugimoto^a, Yoshi Yamano^a, Sukardiman^b, Hideaki Otsuka^c, Katsuyoshi Matsunami^{a,*}^a Graduate School of Biomedical & Health Sciences, Hiroshima University, Japan^b Faculty of Pharmacy, Airlangga University, Surabaya, Indonesia^c Graduate School of Pharmacy, Yasuda Women's University, Hiroshima, Japan

ARTICLE INFO

Article history:

Received 8 June 2015

Received in revised form 25 August 2015

Accepted 3 September 2015

Available online 19 September 2015

Keywords:

Linaria japonica

Linarenone

Cytotoxicity

*Leishmania**cis-ent*-Clerodane

ABSTRACT

Five new *cis-ent*-clerodanes, linarenones **1**–**5** and two known compounds (**6** and **7**) were isolated from whole plant of *Linaria japonica*. The structures of these compounds were determined by various spectroscopic analyses (UV, IR, HR-ESI-MS, 1D and 2D NMR). The absolute configuration of five new diterpenoids was confirmed by circular dichroism (CD) analysis and chemical conversion. Cytotoxicity of the isolated compounds against A549 cell lines and *Leishmania major* were evaluated. The new *cis-ent*-clerodane **3** was found to be moderately active against A549 cell lines, and new *cis-ent*-clerodanes **1,6** and desacetyl-linarenone (**7**) were active against *L. major*.

© 2015 Phytochemical Society of Europe. Published by Elsevier B.V. All rights reserved.

1. Introduction

Linaria japonica Miq. is a perennial plant that grows in seashore areas, and is used in folk medicine as a diuretic and a purgative. The isolation of flavonoid glycosides (Morita et al., 1974), iridoid glucosides (Kitagawa et al., 1972, 1973), and diterpenoids (Kitagawa et al., 1975, 1976, 1977, 1978, 1980) from *L. japonica* Mig. have reported earlier. On our investigation of the same plant, collected in sandy seashore areas of Tottori Prefecture, new flavonoids, phenylethanoids, iridoid and monoterpene glucosides have been isolated (Otsuka, 1992, 1993a,b, 1994a,b, 1995).

As a result of further investigation of the non-polar fraction, i.e., a mixture of hexane and ethyl acetate layers of the same plant, five new diterpenoids (**1**–**5**) along with two known diterpenoids (**6** and **7**) were isolated by various chromatographic techniques. The structures of these compounds were determined as follows by spectrometric analysis (Fig. 1). Herein, described are the isolation and structural elucidation of them as well as the cytotoxic activities against lung cancer cell lines (A549) and *Leishmania major*.

2. Results and discussion

2.1. Isolation and structural elucidation of new compounds

The mixture of hexane and ethyl acetate layers from methanol extract of aerial part of *L. japonica* was fractionated by silica gel and ODS column chromatography, then further purified by HPLC to afford seven compounds (Fig. 1). The known compounds were identified as linarenone (**6**) and desacetyl-linarenone (**7**) by comparing the physical and spectroscopic data in literature (Kitagawa et al., 1978). Desacetyl-linarenone (**7**) was isolated for the first time naturally.

Linarenone **1** was obtained as colorless powder with molecular formula of C₂₂H₃₄O₄ as determined by HR-ESI-MS at *m/z* 355.2348 [M+Na]⁺ (calcd. 385.2349). The IR spectrum had absorption bands at 3340, 1732 and 1237 cm^{−1} indicating the presence of hydroxy and ester carbonyl group. The ¹H NMR spectrum (Table 1) displayed signals due to tertiary methyls at δ_H 0.56 (s) and 1.23 (s), one secondary methyl at δ_H 0.84 (d, *J* = 6 Hz), two olefinic methyls at δ_H 1.64 (s) and 1.93 (s), eight methylene protons at δ_H 1.11 (br qd, *J* = 13, 3 Hz), 1.19 (ddd, *J* = 11, 3, 1.30 (dq, *J* = 13, 3 Hz), 1.61 (dd, *J* = 8.5, 2 Hz), 1.73 (dd, *J* = 16, 8 Hz), 2.06 (ddd, *J* = 14, 3, 3 Hz), 2.47 (br d, *J* = 18 Hz), and 2.71 (dd, *J* = 18, 6 Hz), two protons of oxygenated methylene at δ_H 4.15 (2H, *J* = 6 Hz), two methine protons at δ_H 1.42 (m) and 1.98 (br d, *J* = 6 Hz), and an oxygenated methine proton at δ_H 5.19 (dd, *J* = 8,

* Corresponding author.

E-mail address: matunami@hiroshima-u.ac.jp (K. Matsunami).

Fig. 1. Structures of compounds 1–7.

Table 1

The ^1H NMR spectroscopic data for compounds 1–5.

Position	1	2	3	4	5
4.8	2.47 br d (18)	2.46 br d (18)	2.25 br d (18)	2.28 br d (18)	2.48 br d (18)
β	2.71 dd (18, 6)	2.71 dd (18, 6)	2.66 dd (18, 6)	2.68 dd (18, 6)	2.72 dd (18, 6)
2	–	–	–	–	–
3	5.83 s	5.83 s	5.85 s	5.86 s	5.84 s
4	–	–	–	–	–
5	–	–	–	–	–
6- α	2.06 ddd (14, 3, 3)	2.07 ddd (14, 3, 3)	2.16 ddd (14, 3, 3)	2.16 ddd (14, 3, 3)	2.07 ddd (14, 3, 3)
β	1.19 ddd (14, 11, 3)	1.18 ddd (14, 11, 3)	1.29 ddd (14, 11, 3)	1.30 ddd (14, 11, 3)	1.22 ddd (14, 11, 3)
7- α	1.11 br qd (13, 3)	1.09 br qd (13, 3)	1.14 br qd (13, 3)	1.14 br qd (13, 3)	1.12 br qd (13, 3)
β	1.30 dq (13, 3)	1.33 dq (13, 3)	1.46 dq (13, 3)	1.46 dq (13, 3)	1.38 dq (13, 3)
8	1.42 m	1.43 m	1.49 m	1.48 m	1.55 m
9	–	–	–	–	–
10	1.98 br d (6)	1.97 br d (6)	1.86 br d (6)	1.86 br d (6)	1.81 br d (6)
11	1.61 dd (16, 2)	1.62 dd (16, 2)	6.44 d (16)	6.72 d (16)	2.16 ddt (16, 6, 2)
	1.73 dd (16, 8)	1.73 dd (16, 8)	–	–	2.32 dd (16, 8)
12	5.19 dd (8, 2)	5.21 dd (8, 2)	6.03 d (16)	5.75 d (16)	6.75 ddt (8, 6, 3)
13	–	–	–	–	–
14	5.59 t (6)	5.53 t (6)	2.24 s	–	2.86 br t (7)
15	4.15 d (6)	4.56 d (6)	–	–	4.37 t (7)
16	1.64 s	1.68 s	–	–	–
17	0.84 d (6)	0.84 d (6)	0.69 d (6)	0.70 d (6)	0.81 d (6)
18	0.56 s	0.56 s	0.75 s	0.76 s	0.66 s
19	1.23 s	1.23 s	1.24 s	1.24 s	1.20 s
20	1.93 s	1.93 s	1.95 s	1.95 s	1.93 s
12–OAc 2.01 s	2.01 s	–	–	–	–
1'	–	–	–	–	–
2'	–	2.52 septet (7)	–	–	–
3', 4'	–	1.14 d (7)	–	–	–

Recorded at 600 MHz in CDCl_3 . Chemical shifts (δ) are expressed in ppm and J values are presented in Hz in parenthesis.

73), one acetyl methyl proton signal at δ_{H} 18.01 (s), two olefinic protons at δ_{H} 5.59 (t, $J = 6$ Hz) and 5.83 (s). The ^{13}C NMR spectrum (Table 2) of **1** showed 22 carbon resonances that were classified by analysis of its chemical shift values and its HSQC spectrum as five methyls (δ_{C} 12.3, 16.1, 18.2, 20.3, 31.8), four methylenes (δ_{C} 28.1, 35.5, 36.5, 39.7), an oxygenated methylene (δ_{C} 58.7), two methines (δ_{C} 37.6, 47.5), an oxygenated methine (δ_{C} 74.5), two olefinic methines (δ_{C} 125.3, 128.3), four quaternary carbons (δ_{C} 39.2, 40.5, 137.7, 168.8), an acetoxy carbon signals at δ_{C} 21.0 and δ_{C} 169.9, and a ketone (δ_{C} 198.6).

The NMR spectroscopic data of **1** was closely resembled that of linarienone (**6**), except for some differences in the chemical shift values at C-13, 14 and 15 (δ_{C} 140.0, 121.0 and 60.2, respectively). The proton (δ_{H} 4.15) and carbon (δ_{C} 58.7) of **1** indicated that C-15 have hydroxy group. The acetate substituent was placed at C-12 on the basis of the HMBC correlation of H-12 (δ_{H} 5.19) to carbon signal at δ_{C} 169.9 (12–OAc) (Fig. 2). The A/B ring junction in

1 were deduced to be *cis* on the basis of the ^{13}C NMR chemical shifts of the angular methyl (δ_{C} 31.8 for C-19), which were found to be in the range δ_{C} 11–19 for *trans* and higher than δ_{C} 20 for *cis*-clerodane (**14**) (Manabe and Nishino, 1986).

Its relative configuration of **1** was established by NOESY analysis. The correlation $^2\text{H-15/Me-16}$ suggested as *E* configuration. Another correlations were observed between H-10/Me-19, H-10/H-12, H18/H-20 and Me-17/Me-18, which suggested the orientation of H-10, C-11 and Me-19 to be β -oriented, and those of Me-17 and Me-18 to be α -oriented (Fig. 3). Next, the absolute stereochemistry of the ring moiety of **1** was determined by an analysis of the CD spectrum. The significant positive cotton effect at 331 nm ($\Delta\epsilon = +1.02$) showed the same absolute stereochemistry with linarienone (**6**). In addition, from the biogenetic point of view, the remaining stereochemistry at C-12 of **1** should be identical with co-occurring *cis*-clerodane derivatives (**6** and **30**). This hypothesis was confirmed by the alkaline hydrolysis of **1** and **6**.

Table 2¹³C NMR spectroscopic data for compounds 1–5.

Position	1	2	3	4	5
1	35.5	35.6	36.6	36.3	36.2
2	198.6	198.5	198.3	198.4	198.7
3	128.3	128.4	128.7	128.5	128.8
4	168.8	168.1	167.1	167.2	168.6
5	39.2	39.2	38.3	38.1	39.7
6	36.5	36.5	36.5	36.2	36.7
7	28.1	28.1	27.5	27.2	28.4
8	37.6	37.6	40.5	40.1	36.7
9	40.5	40.5	45.7	45.7	42.2
10	47.5	47.5	49.8	49.5	48.5
11	39.7	39.5	156.3	159.9	37.8
12	74.5	74.3	130.1	120.5	136.7
13	137.7	139.9	197.9	169.8	127.4
14	125.3	120.7	28.2	–	25.6
15	58.7	60.2	–	–	65.6
16	12.3	12.5	–	–	171.3
17	16.1	16.2	17	16.7	16.5
18	18.2	18.2	13.4	13.1	18.6
19	31.8	31.7	31.9	31.7	31.9
20	20.3	20.3	20.5	20.3	20.8
12-OAc	169.9	169.8			
1'	21.0	177.0			
2'		33.7			
3', 4'		18.8			

Recorded at 125 MHz in CDCl₃. Chemical shifts (δ) are expressed in ppm.

The deacyl compounds (**1a** and **6b**) were revealed to be identical by spectroscopic and HPLC analysis. Hence, on the basis of above spectrum analysis, the structure of **1** was determined as (5S,8R,9S,10R,12R)-2-oxo-12,15-dihydroxy-*cis*-clerod-3Z,13(14)*E*-diene 12-acetate.

Linarenone B (**2**) was also a clerodane diterpenoid. It was obtained as colorless solid with molecular formula of C₂₆H₄₀O₅ as determined by HR-ESI-MS at *m/z* 455.6256 [M+Na]⁺ (calcd. 455.2768). The ¹H and ¹³C NMR spectrum (Tables 1 and 2) of **2** were similar to **1**, except for the appearance of isobutanoyl moiety, i.e., an ester carbonyl at δ_C 177.0 (C-1'), a methine at δ_C 33.7 (C-2'), two methyls at δ_C 18.8 (C-3' and C-4') and an esterified methylene at δ_H 4.56 (d, *J* = 6 Hz, H-15), δ_C 60.2 (C-15). It suggested the presence of an isobutanoyl moiety at C-15, which was also supported by HMBC correlations of H-15 to C-1' (Fig. 2). The relative and absolute stereochemistry of **2** was revealed to be the same as **1** by NOESY (Fig. 3), CD spectrum at 333 nm (Δε = +0.95) and a mild alkaline hydrolysis of **2** by which the deacyl derivative of **2** (**2a**) was

identical to **1a** and **6b**, thus the structure of **2** was deduced as (5S,8R,9S,10R,12R)-2-oxo-15-isobutanoyl-*cis*-clerod-3Z,13(14)*E*-diene 12-acetate.

Linarenone C (**3**) was also colorless amorphous powder and displayed an [M+Na]⁺ ion peak at *m/z* 297.1828 (calcd. 297.1825) corresponding to a molecular formula of C₁₈H₂₆O₂. The ¹H and ¹³C NMR spectrum (Tables 1 and 2) displayed two *trans* coupled doublets with *J* = 16 Hz at δ_H 6.03 (H-12) and δ_H 6.44 (H-11), a singlet methyl at δ_H 2.24 (Me-14), a carbonyl group at δ_C 197.9 (C-13), which is characteristic as a conjugated methyl ketone. The planar structure of **3** was determined by COSY and HMBC spectra as 2-oxo-14,15-bisnor-3,11*E*-kolavadien-13-one (Kabir et al., 2010) (Fig. 4), however, the chemical shift of C-19 has quite different (δ_C 31.9 for **3**, δ_C 18.9 for 2-oxo-14,15-bisnor-3,11*E*-kolavadien-13-one). It indicated that **3** is an epimer of C-19, i.e., a *cis*-clerodane diterpenoid. The relative configuration of **3** was deduced on the basis of NOESY correlations of H-10/H-11 and H-10/Me-19, which indicated β-orientation, and the correlation between Me-17/Me-18 for α-orientation (Fig. 5). The significant positive cotton effect of CD spectrum at 334 nm (Δε = +1.16) is similar to **1**. Thus the absolute stereochemistry of **3** should be (5S,8R,9S,10R,3Z,11*E*)-2-oxo-14,15-bisnor-3Z,11*E*-kolavadien-13-one.

Linarenone D (**4**), a colorless amorphous powder, was exhibited a HR-ESI-MS [M+Na]⁺ ion peak at *m/z* 299.1618 (calcd. 299.1618) suggesting to a molecular formula of C₁₇H₂₄O₃. The NMR spectroscopic data (Tables 1 and 2) of **3** and **4** indicated differences regarding the C-13 substituent. The carbon signal at δ_C 169.8 (C-13), the up-field shift to δ_H 5.75 (d, *J* = 16 Hz, H-12) and δ_C 120.5 (C-12) and the downfield shift to δ_H 6.72 (d, *J* = 16 Hz, H-11) and δ_C 159.9 (C-11) indicated a carboxylic acid at the C-13 position. The relative and absolute configuration of **4** was assigned to be the same as that of **3**, based on 1D and 2D NMR analysis including NOESY (Figs. 4 and 7) and CD spectrum at 325 nm (Δε = +1.22) as described for **3**. Thus, the structure of **4** was elucidated as (5S,8R,9S,10R)-2-oxo-14,15-bisnor-3Z,11*E*-kolavadien-13-carboxylic acid.

Linarenone E (**5**), isolated as a colorless powder, was assigned the molecular formula of C₂₀H₂₈O₃ as determined by HR-ESI-MS at *m/z* 339.1929 [M+Na]⁺ (calcd. 339.1931). The IR absorption band at 1754 cm⁻¹ was indicative of the presence of γ-lactone moiety. The ¹H and ¹³C NMR spectra (Tables 1 and 2) also revealed a *cis*-clerodane framework, with two tertiary methyls [δ_H/δ_C 0.66 (s)/18.6 (Me-18) and 1.20 (s)/31.9 (Me-19)], a secondary methyl [δ_H 0.81 (d, *J* = 6 Hz); δ_C 16.5 (Me-17)], an olefinic methyl [δ_H 1.93 (s); δ_C 20.8 (Me-20)], an olefinic methine [δ_H 6.75 (ddt, *J* = 8, 6, 3 Hz); δ_C 136.7 (C-12)], a typical γ-lactone ring [δ_C 127.4 (C-13), δ_H 2.86 (br t,

Fig. 2. HMBC and COSY correlations of **1** and **2**.

Fig. 3. Key NOESY correlations of **1** and **2**.

$J = 7$ Hz, H-14), δ_C 25.6 (C-14), δ_H 4.37 (t, $J = 7$ Hz, H-15), δ_C 65.6 (C-15) and δ_C 171.3 (C-16)] and a six membered ring ketone [δ_C 198.7 (C-2)]. The γ -lactone was connected to C-12 according to the HMBC correlations of H-12 to carbon at δ_C 171.3 (C-16) and 25.6 (C-14), together with the correlations of H-11, 14 and 15 to the carbon at δ_C 127.4 (C-13) (Fig. 6). The NOESY correlations of H-10/Me-19, H-10/H-12, H-3/H-18 and Me-17/Me-18 indicated that H-10, C-11 and Me-19 were β -oriented and Me-17 and Me-18 were α -oriented (Fig. 6), and the absolute stereochemistry of **5** was established by CD spectrum ($\Delta\epsilon = +1.13$ at 327 nm) that was similar to **6**. Accordingly,

7 the structure of **5** was determined as (5S,8R,9S,10R)-3Z,12E-2-oxodihydrofuran-2(3H)-one.

2.2. Cytotoxic and anti-Leishmania major activity of isolated compounds

Several clerodane type diterpenoids have been proven to be potential drug leads most notably with anti-cancer effects (Wang et al., 2013, 2012, 2010, and Yang et al., 2010) and anti-Leishmania activity (Andres et al., 1997 and Nikolas et al., 2006). Therefore, the

Fig. 4. HMBC and COSY correlations of **3** and **4**.

Fig. 5. Key NOESY correlations of **3** and **4**.

isolated compounds (**1–7**) were evaluated for their cytotoxicity against A549 (human lung cancer) cell lines and *L. major*. For A549, linarenone C (**3**) showed moderate inhibition ($IC_{50} = 51.2 \mu M$) and linarenone E (**5**) and linarienone (**6**) found to be active (IC_{50} 86.5 and $79.0 \mu M$, respectively). The presence of an acetyl group at C-12 of **3** may improve the activity compared with **4** (inactive) that have carboxylic acid at this position. For *L. major*, linarenone A (**1**), linarienone (**6**) and desacetyl-linarienone (**7**) showed moderate inhibition (IC_{50} 56.7, 50.3 and $52.7 \mu M$, respectively) and linarenone B (**2**) and linarenone E (**5**) found to be active (IC_{50} 89.4 and $97.3 \mu M$, respectively). It is noteworthy that **1** and **7** were relatively selective against *L. major* than cytotoxicity.

In summary, chemical investigation of non-polar fraction of *L. japonica* led to the isolation of seven compounds (**1–7**), including five new diterpenoids (**1–5**). These isolated compounds were evaluated for their cytotoxicity against A549 cell lines and *L. major*.

Linarenone C (**3**) showed the moderate inhibition toward A549 cell lines, and linarenone A (**1**), linarienone (**6**) and desacetyl-linarienone (**7**) had activity toward *L. major*.

3. Experimental

3.1. General experimental procedures

1H and ^{13}C NMR spectra were taken on a Bruker Ultrashield 600 spectrometer at 600 and 150 MHz, respectively, with TMS as an internal standard. IR and UV spectra were measured on a RIBA FT-720 and JASCO V-520 UV/vis spectrophotometer, respectively. Optical rotations and CD spectra were measured on a JASCO P-1030 digital polarimeter and a JASCO J-720 spectropolarimeter, respectively. Positive ion HR-ESI-MS was performed with an Applied Biosystems QSTAR XL NanoSpray™ System. Silica

Fig. 6. HMBC, COSY and key NOESY correlations of **5**.

gel open column chromatography (CC) a reversed phase [octadecyl] silylated silica gel (ODS)] CC were performed on silica gel 60 (E. Merck, Darmstadt, Germany), and Cosmosil 75C18-OPN (Nacal Tesque, Kyoto, Japan; Φ = 35 mm, L = 350 mm), respectively. HPLC was performed on an ODS column (Inertsil ODS-3, GL Science, Tokyo, Japan Φ = 6 mm, L = 250 mm) and the eluate was monitored with a JASCO RI-930 intelligent refractive index detector and a JASCO PU-1580 intelligent pump.

3.2. Plant material

Whole plants of *L. japonica* were collected in late July 1990 in seashore areas of Tottori Prefecture, and a voucher specimen (90-LJ-Tottori) was deposited at the Department of Pharmacognosy, Faculty of Pharmaceutical Sciences, Hiroshima University.

3.3. Extraction and isolation

The air-dried plants (2.30 kg) were extracted with MeOH (15 L $\times$ 2). The MeOH solution was concentrated and adjusted to 95% aqueous MeOH by the addition of H₂O and then partitioned with *n*-hexane (1.5 L $\times$ 2, 35.0 g). The remaining aqueous MeOH layer was evaporated and re-suspended in 1.5 L of H₂O and then partitioned with EtOAc (1.5 L $\times$ 2, 49.7 g) and 1-BuOH (1.5 L $\times$ 3, 151 g), successively.

The non-polar fraction (60.5 g, as a mixture of EtOAc and *n*-hexane layer) was proceeded on silica gel (300 g) CC with increasing polarity [Hexane–CHCl₃ (1:1), 4:1, CHCl₃–MeOH (50:1, 40:1, 30:1, 20:1, 15:1, 10:1, 7:1, 5:1, 3:1, 2:1, MeOH, each 24 L] yielding 12 fractions (Fr. Lj1–Lj12). The fraction Lj4 (10.1 g) was subjected to reversed phase (ODS) CC with 10% aq. MeOH to 100% MeOH, step gradient, led 19 fractions (Fr. Lj4-1–Lj4-19). The residue of fraction Lj4-10 (54.6 mg) was purified by HPLC (65% aq. MeOH) to give linarenone A (1, 7.58 mg). The fraction Lj4-12 (178 mg) was also purified by HPLC (67.5% aq. MeOH). Two peaks appeared at 18 and 35 min were collected to give desacetyl-linarinenone (7, 5.01 mg) and linarenone B (2, 3.07 mg). The other residue of fraction Lj4-9 (91.9 mg) was purified by prep. HPLC (45% aq. acetone) to give 3 (linarenone C, 2.83 mg) and 5 (linarenone E, 12.7 mg). The fraction Lj4-14 (238 mg) was purified by prep. HPLC (65% aq. acetone) to give 4 (linarenone D, 6.64 mg). Then the fraction Lj4-13 (491 mg) was purified by prep. HPLC (80% aq. MeOH) to give 6 (linarinenone, 20.6 mg).

3.3.1. Linarinone A (1)

Colorless amorphous solid; $[\alpha]_D^{25} + 10.0$ (c 0.50, CHCl₃); UV (EtOH) $\lambda_{\max}$ (log ϵ) nm: 230 (3.72); CD (c 2.76×10^{-5} M, MeOH) $\Delta\epsilon$ (nm): +1.02 (331); IR (film) $\nu_{\max}$ cm⁻¹: 3340, 2929, 1732, 1716, 1653, 1456, 1375, 1237, 1029; ¹H NMR and ¹³C NMR, see Tables 1 and 2; positive HR-ESI-MS m/z 385.2348 [M+Na]⁺ (calcd. for C₂₂H₃₄O₄Na: 385.2349).

3.3.2. Linarenone B (2)

Colorless amorphous solid; $[\alpha]_D^{25} + 5.7$ (c 0.20, CHCl₃); UV (EtOH) $\lambda_{\max}$ (log ϵ) nm: 224 (3.45); CD (c 1.94×10^{-5} M, MeOH) $\Delta\epsilon$ (nm): +0.95 (333); IR (film) $\nu_{\max}$ cm⁻¹: 2914, 1733, 1716, 1653, 1456, 1375, 1239, 1027; ¹H NMR and ¹³C NMR, see Tables 1 and 2; positive HR-ESI-MS m/z 455.2766 [M+Na]⁺ (calcd. for C₂₆H₄₀O₅Na: 455.2768).

3.3.3. Linarenone C (3)

Colorless amorphous solid; $[\alpha]_D^{25} + 15.5$ (c 0.21, CHCl₃); UV (EtOH) $\lambda_{\max}$ (log ϵ) nm: 232 (3.45); CD (c 2.33×10^{-5} M, MeOH) $\Delta\epsilon$ (nm): +1.16 (334), -2.10 (246); IR (film) $\nu_{\max}$ cm⁻¹: 2924, 1698, 1655, 1619, 1456, 1376, 1258, 1033; ¹H NMR and ¹³C NMR, see Tables 1 and 2; positive HR-ESI-MS m/z 297.1825 [M+Na]⁺ (calcd. for C₁₈H₂₆O₂Na: 297.1825).

3.3.4. Linarenone D (4)

Colorless amorphous solid; $[\alpha]_D^{25} + 12.6$ (c 0.44, CHCl₃); UV (EtOH) $\lambda_{\max}$ (log ϵ) nm: 218 (3.62); CD (c 3.21×10^{-5} M, MeOH) $\Delta\epsilon$ (nm): +1.22 (325), -3.32 (250); IR (film) $\nu_{\max}$ cm⁻¹: 2925, 1698, 1649, 1456, 1376, 1260, 1034; ¹H NMR and ¹³C NMR, see Tables 1 and 2; positive HR-ESI-MS m/z 299.1618 [M+Na]⁺ (calcd. for C₁₇H₂₄O₃Na: 299.1618).

3.3.5. Linarenone E (5)

Colorless amorphous solid; $[\alpha]_D^{27} + 48.8$ (c 0.82, MeOH); UV (EtOH) $\lambda_{\max}$ (log ϵ) nm: 239 (2.97); CD (c 5.21×10^{-5} M, MeOH) $\Delta\epsilon$ (nm): +1.31 (327); IR (film) $\nu_{\max}$ cm⁻¹: 2913, 1754, 1660, 1435, 1377, 1256, 1206, 1031; ¹H NMR and ¹³C NMR, see Tables 1 and 2; positive HR-ESI-MS m/z 339.1929 [M+Na]⁺ (calcd. for C₂₀H₂₈O₃Na: 339.1931).

3.3.6. Linarinenone (6)

Colorless amorphous solid; $[\alpha]_D^{26} + 23.5$ (c 0.17, MeOH); CD (c 2.77×10^{-5} M, MeOH) $\Delta\epsilon$ (nm): +0.76 (334), -0.05 (271); ¹H NMR (CDCl₃, δ): 0.55 (3H, s, H₃-18), 0.2 (3H, d, J = 6 Hz, H₃-17), 1.08 (1H, d, J = 13, 3 Hz, H-7 α), 1.18 (1H, ddd, J = 14, 11, 3 Hz, H-6 β), 1.21 (3H, s, H₃-19), 1.32 (1H, dq, J = 13, 3 Hz, H-7 β), 1.43 (1H, m, H-8), 1.62 (3H, m, H₂-11), 1.69 (3H, s, H₃-16), 1.71 (1H, dd, 16, 8 Hz, H-11), 1.85 (3H, s, H₃-5'), 1.92 (6H, br s, H₃-20), 1.95 (1H, br d, J = 6 Hz, H-10), 2.00 (3H, s, 12-OAc), 2.05 (1H, ddd, J = 28, 3, 3 Hz, H-6 α), 2.45 (1H, br d, J = 18 Hz, H-1 α) and 2.70 (1H, dd, J = 18, 6 Hz, H-1 β), 4.63 (2H, d, J = 6 Hz, H₂-15), 5.21 (1H, dd, J = 7, 2 Hz, H-12), 5.58 (1H, t, J = 6 Hz, H-14), 5.82 (1H, s, H-3), 6.02 (1H, dq, J = 7, 1 Hz, H-3') and ¹³C NMR (CDCl₃, δ): 12.7 (CH₃-16), 16.4 (CH₃-17), 18.4 (CH₃-18), 20.6 (CH₃-20), 20.6 (C-5'), 21.2 (C-6), 27.0 (CH₃CO), 28.3 (C-7), 31.9 (CH₃-4), 35.7 (C-1), 36.7 (C-6), 37.7 (C-8), 39.4 (C-11), 40.7 (C-5), 42.6 (C-9), 47.5 (C-10), 60.2 (C-15), 74.5 (C-12), 120.8 (C-14), 121.0 (C-3), 127.8 (C-3'), 128.5 (C-2'), 13.8 (C-4), 140.0 (C-13), 167.9 (CH₃CO), 170.0 (C-1'), 198.9 (C-2); positive HR-ESI-MS m/z 467.2773 [M+Na]⁺ (calcd. for C₂₇H₄₀O₅Na: 467.2768).

3.3.7. Desacetyl-linarinenone (6a = 7)

Colorless amorphous solid; $[\alpha]_D^{30.0} + 67.5$ (c = 0.04, MeOH); UV (EtOH) $\lambda_{\max}$ (log ϵ) nm: 247 (3.83); CD (c 1.24×10^{-5} M, MeOH) $\Delta\epsilon$ (nm): +8.61 (327), +0.10 (271); IR (film) $\nu_{\max}$ cm⁻¹: 3430, 2926, 1714, 1700, 1650, 1456, 1225, 1151, 1080; ¹H NMR (CDCl₃, δ): 0.56 (3H, s, H₃-18), 0.86 (3H, d, J = 6 Hz, H₃-17), 1.12 (1H, br qd, J = 13, 3 Hz, H-7 α), 1.22 (3H, s, H₃-19), 1.24 (1H, d, J = 14, 11, 3 Hz, H-6 β), 1.35 (1H, dq, J = 13, 3 Hz, H-7 β), 1.60 (2H, m, H-11), 1.72 (3H, s, H₃-16), 1.87 (3H, s, H₃-5'), 1.93 (3H, s, H₃-20), 1.96 (3H, dq, J = 7, 1 Hz, H₃-4), 2.02 (1H, br d, J = 6 Hz, H-10), 2.06 (1H, ddd, J = 14, 3, 3 Hz, H-6 α), 2.46 (1H, br d, J = 18 Hz, H-1 α), 2.69 (1H, dd, J = 18, 6 Hz, H-1 β), 4.16 (1H, br t, J = 6 Hz, H-12), 4.67 (2H, t, J = 7 Hz, H₂-15), 5.59 (1H, t, J = 7 Hz, H-14), 5.83 (1H, s, H-3), 6.04 (1H, dq, J = 7, 1 Hz, H-3'); positive HR-ESI-MS m/z 425.2661 [M+Na]⁺ (calcd. for C₂₅H₃₈O₄Na: 425.2662).

3.3.8. Desdiacyl-linarinenone (6b = 13 = 2a)

Colorless amorphous solid; $[\alpha]_D^{30} + 62.4$ (c 0.11, MeOH); UV (EtOH) $\lambda_{\max}$ (log ϵ) nm: 248 (4.50); CD (c 1.24×10^{-5} M, MeOH) $\Delta\epsilon$ (nm): +1.62 (330), +0.00 (292); IR (film) $\nu_{\max}$ cm⁻¹: 3382, 2927, 1730, 1717, 1649, 1456, 1379, 1258, 1004; ¹H NMR (CDCl₃, δ): 0.57 (3H, s, H₃-18), 0.86 (3H, d, J = 7 Hz, H₃-17), 1.10 (1H, br qd, J = 13, 3, H-7 α), 1.23 (3H, s, H₃-19), 1.26 (1H, ddd, J = 14, 11, 3 Hz, H-6 β), 1.35 (1H, dq, J = 13, 3, H-7 β), 1.54 (2H, m, H₂-11), 1.67 (3H, s, H₃-16), 1.94 (3H, s, H₃-20), 2.07 (1H, ddd, J = 14, 3, 3 Hz, H-6 α), 2.09 (1H, m, H-10), 2.40 (1H, br d, J = 18 Hz, H-1 α), 2.68 (1H, dd, J = 18, 6 Hz, H-1 β), 4.1 (1H, m, H-12), 4.17 (2H, m, H₂-15), 5.59 (1H, t, J = 7 Hz, H-14), 5.84 (1H, s, H-3) and ¹³C NMR (CDCl₃, δ): 12.6 (CH₃-16), 17.1 (C-17), 19.4 (C-18), 21.5 (CH₃-20), 28.9 (C-7), 32.6 (C-19), 36.4 (C-1), 37.3 (C-6), 38.2 (C-8), 39.9 (C-5), 41.2 (C-9), 42.0 (C-11), 47.9 (C-10), 59.5 (C-15), 74.3 (C-

12), 124.4 (C-14), 129.1 (C-3), 142.9 (C-13), 168.9 (C-4), 199.3 (C-2); positive HR-ESI-MS m/z 343.2246 $[M+Na]^+$ (calcd. for $C_{20}H_{32}O_3Na$: 343.2244).

3.4. Mild alkaline hydrolysis

A solution of linarienone (**6**) (11.5 mg) in MeOH-CH₂Cl₂ (1:1, 1.8 ml) was added 1 mol/l NaOMe solution (0.2 ml), stirred at rt for 3 h. The mixture was neutralized with ion exchange resin IR-120B GANO, H⁺-form) and filtered off. The filtrate was evaporated to dryness under reduced pressure. Purification of the reaction product by preparative TLC (*n*-hexane:EtOAc = 3:2) furnished desacetyl-linarienone (**6a**) (**7**) (0.3 mg, R_f = 0.45) and desdiacyl-linarienone (**6b**) (1.2 mg, R_f = 0.18). This procedure also performed for **1** and **2** to produce **1a** and **2a**, respectively, which were identical to **6b** by HPLC (65% aq. MeOH, ODS Φ = 6 mm $\times$ 250, UV-vis detector 248 nm, 1.5 ml/min, t_R = 16 min) and HR-ESI-MS analysis.

3.5. A549 growth inhibition assay

Human lung cancer cells (A549) were cultured in Dulbecco's modified Eagle's medium supplemented with 10% heat-inactivated fetal bovine serum, kanamycin (100 μ g/ml) and amphotericin B (0.5 μ g/ml). Into a 96-well plate, aliquots of the DMSO solution of the test compounds (1% final concentration) were incubated with A549 cells (5×10^3 cells/well) in CO₂ incubator at 37 °C for 72 h. MTT was added into each well and the plate was further incubated for 1.5 h. Absorbance was measured at 540 nm using a 300 EnSpire Multimode plate reader (Perkin Elmer). DMSO was used as a negative control and doxorubicin as a positive control (IC_{50} = 0.70 μ M). The viability was compared to that of control cells incubated in the same medium without the test compounds (Macahig et al., 2010).

$$\text{Inhibition (\%)} = [1 \times (A_{\text{sample}} - A_{\text{blank}}) / (A_{\text{control}} - A_{\text{blank}})] \times 100$$

3.6. Anti-Leishmania major assay

L. major promastigotes were cultured in M199 medium supplemented with 10% heat-inactivated fetal bovine serum and kanamycin (100 μ g/ml). Into a 96-well plate, aliquots of the DMSO solution of the test compounds (1% final concentration) were incubated with *L. major* cells (1×10^5 cells/well) at 27 °C for 72 h. MTT was added into each well and further incubated overnight. Absorbance was measured at 540 nm using a 2300 EnSpire Multimode plate reader (Perkin Elmer). DMSO was used as a negative and amphotericin B as positive control (IC_{50} = 0.58 μ M). The viability was compared to that of control cells incubated in the same medium without the test compounds (Macahig et al., 2010).

$$\text{Inhibition (\%)} = [1 \times (A_{\text{sample}} - A_{\text{blank}}) / (A_{\text{control}} - A_{\text{blank}})] \times 100$$

Acknowledgment

This work was supported by JSPS KAKENHI Grant Number 26460122, 00107385 and 60549012. The authors are grateful for access to the superconducting NMR instrument, UV and ESI-MS at

the Analytical Center of Molecular Medicine, the Analysis Center of Life Science and the Natural Science Center for Basic Research and Development of the Graduate School of Biomedical and Health Sciences, Hiroshima University. The authors would like to express their sincere thanks to Indonesian Directorate General of Higher Education (DIKTI) for doctoral scholarships.

References

- Andres, G., Emilio, L., Antonio, M., Francisco, R., Concepcion, M., Juan, J., Antonio, O., 1997. In vitro action of ent-manoyl oxides against *Leishmania donovani*. *J. Nat. Prod.* 60, 13–16.
- Kabir, S., Rahman, M.S., Chowdhury, A.M., Hasan, C.M., Rashid, M.A., 2010. An unusual bisnor-clerodane diterpenoid from *Polyalthia simirum*. *Nat. Prod. Commun.* 5 (10), 1543–1546.
- Kitagawa, I., Tani, T., Akita, K., Yosioka, I., 1972. Linarioside, a new chlorine containing iridoid glucoside, from *Linaria japonica*. *Tetrahedron Lett.* 5, 419–422.
- Kitagawa, I., Tani, T., Akita, K., Yosioka, I., 1973. On the constituents of *Linaria japonica* Mig. I. The structure of linarioside, a new chlorinated iridoid glucoside and identification of two related glucosides. *Chem. Pharm. Bull.* 21, 1978–1987.
- Kitagawa, I., Yoshihara, M., Tani, T., Yosioka, I., 1975. Linaridial, a new *cis*-clerodane-type diterpene dialdehyde from *Linaria japonica* Miq. *Tetrahedron Lett.* 1, 23–26.
- Kitagawa, I., Yoshihara, M., Tani, T., Yosioka, I., 1976. On the constituents of *Linaria japonica* Mig. II. The structure of linaridial, a new *cis*-clerodane-type diterpene dialdehyde. *Chem. Pharm. Bull.* 24, 294–302.
- Kitagawa, I., Yoshihara, M., Kamiguchi, T., 1977. Linarienone, a new *cis*-clerodane-type diterpene from the subterranean part of *Linaria japonica* Miq. *Tetrahedron Lett.* 14, 1221–1224.
- Kitagawa, I., Yoshihara, M.A., nd Kamiguchi, T., 1978. On the constituents of *Linaria japonica* Mig. III. The structure of linarienone, a new *cis*-clerodane-type diterpene. *Chem. Pharm. Bull.* 26, 79–87.
- Kitagawa, I., Kamiguchi, T., Yonetani, K., Yoshihara, M., 1980. On the constituents of *Linaria japonica* Mig. IV. Chemical correlation of *cis*-clerodane diterpenes with trans-clerodane diterpenes. *Chem. Pharm. Bull.* 28, 2403–2413.
- Macahig, R.A., Harinantenaina, L., Matsunami, K., Otsuka, H., Takeda, Y., Shinzato, T., 2010. Secoiridoid and iridoid glucosides from the leaves of *Fraxinus griffithii*. *J. Nat. Med.* 64, 1–8.
- Manabe, S., Nishino, C., 1986. Stereochemistry of *cis*-clerodane diterpenes. *Tetrahedron* 42, 3461–3470.
- Morita, N., Shimizu, M., Arisawa, M., Kobayashi, K., 1974. Studies on medicinal resources. XXXVII. The components of leaves of *Linaria japonica* Miq. and *L. vulgaris* Mill. (Scrophulariaceae). *Yakugaku Zasshi* 94, 913–916.
- Nikolas, F., Eleftherios, K., Babu, L.T., Alexios, L.S., Stephen, O.D., 2006. Antileishmanial activity of natural diterpenes from *Cistus* sp. and semisynthetic derivatives thereof. *Biol. Pharm. Bull.* 29, 1775–1778.
- Otsuka, H., 1992. Isolation of isolinariins A and B, new flavonoid glycosides from *Linaria japonica*. *J. Nat. Prod.* 55, 1252–1255.
- Otsuka, H., 1993a. Phenylethanoids from *Linaria japonica*. *Phytochemistry* 32, 979–981.
- Otsuka, H., 1993b. Iridoid glucosides from *Linaria japonica*. *Phytochemistry* 33, 617–622.
- Otsuka, H., 1994a. Iridolinarins A, B and C: Iridoid esters of an iridoid glucoside from *Linaria japonica*. *J. Nat. Prod.* 57, 357–362.
- Otsuka, H., 1994b. Linarionosides A–C and acyclic monoterpene diglucosides from *Linaria japonica*. *Phytochemistry* 37, 461–465.
- Otsuka, H., 1995. Iridoid mono and diesters of D-glucopyranose from *Linaria japonica*. *Phytochemistry* 39, 1111–1114.
- Wang, F., Ren, F.C., Li, Y.J., Liu, J.K., 2010. Scutellarbarbatines W–Z, new neo-clerodane diterpenoids from *Scutellaria barbata* and structure revision of a series of 13-spiro neo-clerodanes. *Chem. Pharm. Bull.* 58, 1267–1270.
- Wang, T., Wang, S., Xiao, D., 2012. A review of phytochemistry and antitumor activity of a valuable medicinal species: *Scutellaria barbata*. *J. Med. Plants Res.* 26, 4259–4275.
- Wang, B., Wang, X., Wang, S., Shen, T., Liu, Y., Yuan, H., Lou, H., Wang, X., 2013. Cytotoxic clerodane diterpenoids from the leaves and twigs of *Casearia balansae*. *J. Nat. Prod.* 76, 1573–1579.
- Yang, N., Liu, L., Tao, W., Duan, J., Tian, L., 2010. Diterpenoids from *Pinus massoniana* resin and their cytotoxicity against A431 and A549 cells. *Phytochemistry* 71, 1528–1533.

New cis-ent-clerodanes from *Linaria japonica*

ORIGINALITY REPORT

28%

SIMILARITY INDEX

13%

INTERNET SOURCES

28%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1

mycochem.cn

Internet Source

2%

2

M. Iqbal Choudhary, Muhammad Ismail, Khozirah Shaari, Ahmed Abbaskhan, Samina A. Sattar, Nordin H. Lajis. " Clerodane-Type Furanoditerpenoids from ", Journal of Natural Products, 2010

Publication

2%

3

Otsuka, Hideaki. "Isolation of Isolinariins A and B, New Flavonoid Glycosides from *Linaria japonica*", Journal of Natural Products, 1992.

Publication

1%

4

Koike, K.. "Quassinoids from *Quassia indica*", Phytochemistry, 199401

Publication

1%

5

Jiang, C.S.. "Synthesis and antitumor evaluation of methyl spongoate analogs", Steroids, 20101212

Publication

1%

6

Matsunami, Katsuyoshi, Ichiko Takamori,

Takakazu Shinzato, Mitsunori Aramoto, Kazunari Kondo, Hideaki Otsuka, and Yoshio Takeda. "Radical-Scavenging Activities of New Megastigmane Glucosides from *Macaranga tanarius* (L.) MœLL.-ARG.", CHEMICAL & PHARMACEUTICAL BULLETIN, 2006.

Publication

1%

7

Min Chen. "Four New Dibenzocyclooctene Lignans from *Kadsura renchangiana*", Helvetica Chimica Acta, 06/2004

Publication

1%

8

repository.library.du.ac.bd

Internet Source

1%

9

Kakuda, Rie, Koichi Machida, Yasunori Yaoita, Masafumi Kikuchi, and Masao Kikuchi. "Studies on the Constituents of *Gentiana* Species. II. A New Triterpenoid, and (S)-(+)- and (R)-($\hat{\sim}$)-Gentiolactones from *Gentiana lutea*", CHEMICAL & PHARMACEUTICAL BULLETIN, 2003.

Publication

1%

10

Dong-Xiao Guo, Rong-Xiu Zhu, Xiao-Ning Wang, Li-Ning Wang, Shu-Qi Wang, Zhao-Min Lin, Hong-Xiang Lou. " Scaparvin A, A Novel Caged -Clerodane with an Unprecedented C-6/C-11 Bond, and Related Diterpenoids from the Liverwort ", Organic Letters, 2010

Publication

1%

11

Hideaki Otsuka. "Phenylethanoids from *Linaria japonica*", *Phytochemistry*, 1993

Publication

1 %

12

Guo, Dong-Xiao, Xiao-Ning Wang, Rong-Xiu Zhu, Na Liu, Jin-Chuan Zhou, Wen-Tao Yu, and Hong-Xiang Lou. "cis-Clerodane diterpenoids from the Chinese liverwort *Scapania parva* Steph.", *Phytochemistry Letters*, 2012.

Publication

1 %

13

Taizong Wu, Qian Wang, Cheng Jiang, Susan L. Morris-Natschke, Hui Cui, Yan Wang, Yuan Yan, Jun Xu, Kuo-Hsiung Lee, Qiong Gu. " - Clerodane Diterpenoids from with Activity against Epstein–Barr Virus Lytic Replication ", *Journal of Natural Products*, 2015

Publication

1 %

14

J. Fausto Rivero-Cruz, Hee-Byung Chai, Leonardus B. S. Kardono, Fransisca M. Setyowati et al. " Cytotoxic Constituents of the Twigs and Leaves of ", *Journal of Natural Products*, 2004

Publication

1 %

15

www.thieme-connect.com

Internet Source

<1 %

16

Carpenter, Caitlyn D., Taryn O'Neill, Nadia

<1 %

Picot, John A. Johnson, Gilles A. Robichaud, Duncan Webster, and Christopher A. Gray. "Anti-mycobacterial natural products from the Canadian medicinal plant *Juniperus communis*", *Journal of Ethnopharmacology*, 2012.

Publication

17

James H. Wynne. "Synthesis of Novel 7 α -Thiol-Bridged Progesterone Derivatives", *Synthetic Communications*, 2003

<1 %

Publication

18

W. Zhang. "Two new flavanes and bioactive compounds from *Daphne tangutica* Maxim", *Natural Product Research*, 9/2007

<1 %

Publication

19

Ito, Masanobu, Keiichi Matsuzaki, Jue Wang, Akihiro Daikonya, Nai-Li Wang, Xin-Sheng Yao, and Susumu Kitanaka. "New Phenanthrenes and Stilbenes from *Dendrobium loddigesii*", *CHEMICAL & PHARMACEUTICAL BULLETIN*, 2010.

<1 %

Publication

20

Wenhui Ma. "Dibenzocyclooctane Lignans from the Stems of *Kadsura induta* and Their Antiviral Effect on Hepatitis B Virus", *Chemistry & Biodiversity*, 05/2007

<1 %

Publication

21

www.mdpi.com

Internet Source

<1 %

22

Cheng, L.. "Facile synthesis of cleistetroside-2, a partially acetylated oligorhamnoside from *Cleistopholis glauca* and *patens*", *Carbohydrate Research*, 20070813

Publication

<1 %

23

Su, J.H.. "Novel steroids from the soft coral *Nephthea chabrolii*", *Tetrahedron*, 20070115

Publication

<1 %

24

Lee, Chul, Jin Woo Lee, Qinghao Jin, Hak Ju Lee, Sung-Joon Lee, Dongho Lee, Myung Koo Lee, Chong Kil Lee, Jin Tae Hong, Mi Kyeong Lee, and Bang Yeon Hwang. "Anti-inflammatory Constituents from the Fruits of *Vitex rotundifolia*", *Bioorganic & Medicinal Chemistry Letters*, 2013.

Publication

<1 %

25

linknovate.com

Internet Source

<1 %

26

www.excli.de

Internet Source

<1 %

27

Jung-Rae Rho, Hyi-Seung Lee, Hee Jae Shin, Jong-Woong Ahn, Ji-Yun Kim, Chung J. Sim, Jongheon Shin. " New Sesterterpenes from the *Sponge* sp. ", *Journal of Natural Products*, 2004

<1 %

28

Oka, T.. "Synthesis of both enantiomers of the BC-ring part of ciguatoxin", Tetrahedron, 19960909

Publication

<1 %

29

Takaya, Y.. "Novel Guaianoids, Nardoguaianone E-I, from Nardostachys chinensis Roots", Tetrahedron, 20000922

Publication

<1 %

30

www.thieme-connect.de

Internet Source

<1 %

31

Shao-Jiang Song, Ling-Zhi Li, Pin-Yi Gao, Ying Peng, Jing-Yu Yang, Chun-Fu Wu. "Terpenoids and hexenes from the leaves of Crataegus pinnatifida", Food Chemistry, 2011

Publication

<1 %

32

Hammami, Saoussen, Zhenzhen Li, Mingjie Huang, Ridha El Mokni, Hatem Dhaouadi, and Sheng Yin. "New bioactive labdane diterpenoids from Marrubium aschersonii", Natural Product Research, 2016.

Publication

<1 %

33

Culioli, G.. "Trihydroxylated linear diterpenes from the brown alga Bifurcaria bifurcata", Phytochemistry, 200407

Publication

<1 %

34

Inada, A.. "Unusual cyclolanostanes from leaves of *Pandanus boninensis*", *Phytochemistry*, 200512

Publication

<1 %

35

academicjournals.org

Internet Source

<1 %

36

Varamini, P., M. Doroudchi, A. Mohagheghzadeh, M. Soltani, and A. Ghaderi. "Cytotoxic Evaluation of Four *Haplophyllum* Species with Various Tumor Cell Lines", *Pharmaceutical Biology*, 2007.

Publication

<1 %

37

Liu, L.. "Pestaloficiols A-E, bioactive cyclopropane derivatives from the plant endophytic fungus *Pestalotiopsis fici*", *Bioorganic & Medicinal Chemistry*, 20080601

Publication

<1 %

38

Eung Tae Yeon, Jin Woo Lee, Chul Lee, Qinghao Jin et al. "-Clerodane Diterpenoids from and Their Inhibitory Effects on LPS-Induced Nitric Oxide Production ", *Journal of Natural Products*, 2015

Publication

<1 %

39

ABE, Masako, Yoshio OZAWA, Yasushi UDA, Yoichi YAMADA, Yasujiro MORIMITSU, Yoshimasa NAKAMURA, and Toshihiko OSAWA. "Labdane-type Diterpene Dialdehyde,

<1 %

Pungent Principle of Myoga, Zingiber mioga Roscoe.", Bioscience Biotechnology and Biochemistry, 2002.

Publication

40

Takeda, Y.. "Phlorigidosides A-C, iridoid glucosides from Phlomis rigida", Phytochemistry, 200004

Publication

<1 %

41

Zhu, X.-H.. "Withanolide-type steroids from Solanum cilistum", Phytochemistry, 200104

Publication

<1 %

42

repositorium.sdum.uminho.pt

Internet Source

<1 %

43

Su, Yao, Jun-Long Bi, Yue-Hu Wang, Ying Tan, Jun Yang, Hong-Xin Liu, Wei Gu, Ge-Fen Yin, and Chun-Lin Long. "Chemical constituents from Chirita longgangensis var. hongyao with inhibitory activity against porcine respiratory and reproductive syndrome virus", Journal of the Brazilian Chemical Society, 2012.

Publication

<1 %

44

www.science.gov

Internet Source

<1 %

45

Zafer Asım Kaplancıklı. "Synthesis and Antimicrobial Activity of Some Thiazolyl-Pyrazoline Derivatives", Phosphorus Sulfur and

<1 %

Silicon and the Related Elements, 2007

Publication

46

de la Torre, M.C.. "Hemisynthesis of Oxetane-Containing Neoclerodane Diterpenoids", *Tetrahedron*, 20000929

Publication

<1 %

47

Qi, J.. "Linckosides C-E, three new neuritogenic steroid glycosides from the Okinawan starfish *Linckia laevigata*", *Bioorganic & Medicinal Chemistry*, 20040801

Publication

<1 %

48

www.devonfarms.co.uk

Internet Source

<1 %

49

Xue Gao. "Four new terpenoids from the roots of *Ligularia narynensis*", *Journal of Asian Natural Products Research*, 2/2008

Publication

<1 %

50

Jian-Hong Yang, Xue Du, Fei He, Hai-Bo Zhang, Xiao-Nian Li, Jia Su, Yan Li, Jian-Xin Pu, Han-Dong Sun. " Bioactive Abietane and -Kaurane Diterpenoids from ", *Journal of Natural Products*, 2013

Publication

<1 %

51

Evans, G.B.. "The synthesis and antibacterial activity of totarol derivatives. part 1: modifications of ring-C and pro-drugs", *Bioorganic & Medicinal Chemistry*, 199909

<1 %

52

Nasini, G.. "Structure and absolute configuration of acetosellin, a new polyketide from a phytotoxic strain of *Cercospora acetosella*", *Tetrahedron Letters*, 20020225

Publication

<1 %

53

cmjournal.biomedcentral.com

Internet Source

<1 %

54

Zhang, Y.. "Tagalsins A-H, dolabrane-type diterpenes from the mangrove plant, *Ceriops tagal*", *Phytochemistry*, 200506

Publication

<1 %

55

Tang, W.. "Novel NGF-potentiating diterpenoids from a Brazilian medicinal plant, *Ptychopetalum olacoides*", *Bioorganic & Medicinal Chemistry Letters*, 20090201

Publication

<1 %

56

Marcos, I.S.. "Synthesis of tri- and tetracyclic diterpenes. Cyclisations promoted by SmI^2 ", *Tetrahedron*, 20050124

Publication

<1 %

57

Jiang, Zhi-Yong, Wen-Juan Li, Li-Xiang Jiao, Jun-Ming Guo, Kai Tian, Chun-Tao Yang, and Xiang-Zhong Huang. "New Clerodane Diterpenes from *Tinospora sagittata* var. *yunnanensis*", *Planta Medica*, 2014.

Publication

<1 %

58 Gavagnin, M.. "Further chemical studies on the Antarctic nudibranch *Austrodoris kerguelensis*: new terpenoid acylglycerols and revision of the previous stereochemistry", *Tetrahedron*, 20030714 $<1\%$

Publication

59 mdpi.com $<1\%$

Internet Source

60 Akkol, Esra Küpeli, and Dilek Ercil. "Antinociceptive and anti-inflammatory activities of some *Linaria* species from Turkey", *Pharmaceutical Biology*, 2009. $<1\%$

Publication

61 Garcia, Abraham Ramirez-Apan, Teresa Ant. "Absolute configuration assignments by experimental and theoretical approaches of ent-labdane- and ci", *Canadian Journal of Chemistry*, Dec 2006 Issue $<1\%$

Publication

62 Ihsan Çalis. "Secondary metabolites from *Phlomis oppositiflora*", *Natural Product Research*, 7/2005 $<1\%$

Publication

63 acgpubs.org $<1\%$

Internet Source

64 Ma, D.. "Enantioselective synthesis of

functionalized @a-amino acids via a chiral guanidine catalyzed Michael addition reaction", Tetrahedron: Asymmetry, 19990226

Publication

<1 %

65

Armandodoriano Bianco, Marcella Guiso, Mauro Ballero, Sebastiano Foddai et al. "Glycosidic Monoterpenes from ", Natural Product Research, 2004

Publication

<1 %

66

Zhong-Mei Zou. "Steroidal alkaloids from roots of *Sarcococca vagans*", Phytochemistry, 199711

Publication

<1 %

67

Abdel waheb Fekih. "Hemisynthesis of new labdane derivatives from (-)-sclareol", Natural Product Research, 8/1/2006

Publication

<1 %

68

Julia Schuppan. "Apoptolidinone A: Synthesis of the Apoptolidin A Aglycone", Chemistry - A European Journal, 09/25/2006

Publication

<1 %

69

Otsuka, H.. "Iridoid mono- and diesters of d-glucopyranose from *Linaria japonica*", Phytochemistry, 199507

Publication

<1 %

70

P. N. Kotiyan. "Synthesis and characterization of an acrylate pressure sensitive adhesive for

<1 %

71

Shunji Ito. "Reaction of (η 4-cycloheptatriene)Fe(CO)₃ with haloketenes: structure and reactivity of the cycloadducts", Journal of Organometallic Chemistry, 20020401

Publication

<1 %

72

Huang, J.M.. "Structures of merrilactones B and C, novel anislactone-type sesquiterpenes from *Illicium merrillianum*, and chemical conversion of anislactone B to merrilactone A", Tetrahedron, 20010528

Publication

<1 %

73

Salim, A.A.. "Constituents of the leaves and stem bark of *Aglaia foveolata*", Tetrahedron, 20070813

Publication

<1 %

74

Ojika, M.. "Aplyronine A, a potent antitumor macrolide of marine origin, and the congeners aplyronines B and C: isolation, structures, and bioactivities", Tetrahedron, 20070409

Publication

<1 %

75

Suzuki, H.. "Synthesis of eleven-membered carbocycles via a homo-Cope type of five-carbon ring expansion reaction utilized @b-

<1 %

(hydroxymethyl)allylsilane", Tetrahedron,
20030421

Publication

76

Takahashi, H.. "Coumaroyl triterpenes from
Casuarina equisetifolia", Phytochemistry,
199906

Publication

<1 %

77

mscanner.stanford.edu

Internet Source

<1 %

78

Imuta, S.. "Total synthesis of actinobolin from
d-glucose by way of the stereoselective three-
component coupling reaction", Tetrahedron,
20060717

Publication

<1 %

79

Knight, J.G.. "Total synthesis of
deoxymannojirimycin and d-mannolactam via
carbonylation of 5-vinyloxazolidin-2-ones",
Tetrahedron, 20030113

Publication

<1 %

80

Veaceslav Kulcički, Nicon Ungur, Margherita
Gavagnin, Francesco Castelluccio, Guido
Cimino. "Ring B functionalization of scalarane
sesterterpenes by radical relay halogenation",
Tetrahedron, 2007

Publication

<1 %

81

Fleming, F.F.. "@b-Siloxy unsaturated nitriles:
stereodivergent cyclizations to cis- and trans-

<1 %

82

Asai, M.. "Stereocontrolled synthesis of (-)-5,11-dideoxytetrodotoxin", Tetrahedron, 20010521

Publication

<1 %

83

Imane Benabdelaziz, Santiago Gómez-Ruiz, Mohammed Benkhaled, Sandra Carralero et al. "New cycloartane-type ester triterpenes from Euphorbia pterococca and biological evaluation", Fitoterapia, 2018

Publication

<1 %

84

Nawwar, Mahmoud A., Noha F. Swilam, Amani N. Hashim, Ahmed M. Al-Abd, Ashraf B. Abdel-Naim, and Ulrike Lindequist. "Cytotoxic isoferulic acidamide from Myricaria germanica (Tamaricaceae)", Plant Signaling & Behavior, 2013.

Publication

<1 %

85

Evans, D.A.. "Enantioselective total synthesis of altohyrtin C (spongistatin 2)", Tetrahedron, 19990716

Publication

<1 %

86

Yukio Hitotsuyanagi, Kunihiro Nishimura, Hiroshi Ikuta, Koichi Takeya, Hideji Itokawa. "Syntheses of Antitumor Morphinane Alkaloids, Sinococuline and 6-epi-, 7-epi-, and 6-epi-7-

<1 %

epi-Sinococuline, from Sinomenine", The
Journal of Organic Chemistry, 1995

Publication

87

Guan-E Yang. "Cytotoxic Activities of Extracts
and Compounds from Viscum coloratum and
its Transformation Products by Rhodobacter
sphaeroides", Applied Biochemistry and
Biotechnology, 03/2009

Publication

<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography On