Skin Penetration of Ubiquinone (Co-Q10) In Nanoemulsi on Delivery System Using Virgin Cocunut Oil (Vco)

by Tristiana Erawati

Submission date: 01-Mar-2019 11:46AM (UTC+0800)

Submission ID: 1085775744

File name: of_Ubiquinone_Co-Q10_In_Nanoemulsi_on_Delivery_System_Using.pdf (338.19K)

Word count: 1318

Character count: 6877

Skin Penetration of Ubiquinone (Co-Q10) In Nanoemulsion Delivery System Using Virgin Coconut Oil (Vco)

Tristiana Erawati M^{*}, Widji Soeratri, Noorma Rosita

³
Pharmaceutics Department of Faculty of Pharmacy, Airlangga University, Kampus B UNAIR, Jl Dharmawangsa Dalam Surabaya, 60286

Received: 7th Dec, 17; Revised and Accepted: 21st Apr, 18; Available Online: 25th Jun, 2018

ABSTRACT

Ubiquinone (co-Q10) is a topical anti-aging with low solubility in water, so its penetration into the skin is also low. To increase its penetration ubiquinone (co-Q10) was loaded in the nanoemulsion delivery system using virgin coconut oil (VCO) as oil phase. Ubiquinone (co-Q10) in nanoemulsion formula consists of; ubiquinone (co-Q10), Tween 80, Span 80, ethanol 96%, VCO and acetate buffer solution pH 4.2 ± 0.5 with concentration; 1%; 18.66%; 1.92%; 3.42%; up to 100%, respectively. The characteristics of ubiquinone (co-Q10) nanoemulsion were observed in terms of droplet morphology by transmission electron microscope (TEM) and droplet size by particle analyzer, the pH and viscosity value. The ubiquinone (co-Q10) penetration test in the nanoemulsion system was compared with the emulsion on the rat skin.

Result of this research showed the droplet morphology of ubiquinone (co-Q10) in nanoemulsion was spherical. The droplet size was 93.2 ± 2.78 nm. The pH value was 4.12 ± 0.02 , and the viscosity was 8.5 cPas. Penetrasi of ubiquinone (co-Q10) in nanoemulsion was deeper (269 ± 48.12 and 993 ± 36.49 μ m) than in emulsion (26.33 ± 6.43 and 110.33 ± 9.124 μ m), after 2 and 6 hours sample application.

Keywords: Nanoemulsion, Skin Penetration, Ubiquinone (co-Q10), Virgin Coconut Oil (VCO).

INTRODUCTION

Nanoemulsion was known as a system to increase the solubility, and penetration of lipophilic substance^{1,2}. In this study nanoemulsion delivery system was used as vehicle of ubiquinone (co-Q10) which is known as anti-aging. Ubiquinone (co-Q10) can prevent collagen and elastine damage and help to avoid wrinkle of the skin³. Nanoemulsion is an emulsion of oil in water (O/W) or water in oil (W/O) with the droplet size between 50 - 500 nm. Smaller droplet size results in larger surface area it can increase the penetration of active ingredients⁴. For O/W nanoemulsion virgin coconut oil (VCO) used as oil phase in nanoemulsion because it can produced smallest droplet size than corn oil and soybean oil⁵.

MATERIALS AND METHODS

Research Material

Ubiquinone (Kangcare), virgin coconut oil (VCO), Tween 80 (Sigma Aldrich), Span 80 (Sigma Aldrich), ethanol 96% (E.Merck), acetic acid p.a (E.Merck), sodium acetate (E.Merck) p.a.

Research Instrument

Stirrer plate (Dragon Lab MS-Pro), ultrasonic (Branson 3510), Thermo shaker (Wine shake), pH meter (Eutech Instruments pH 700), Viscometer cone and plate, Transmission Electron Microscope (TEM) JEOL JEM 1400, Delsa™ Nano Submicron Particle Size and Zeta Potential Dynamic Light Scattering, Microscope Fluorecences and Brightfield Olympus FSX100.

Experimental Animals

Animals used were male Wistar rats with the inclusion criteria: 8-10 weeks of age, 100-250 grams of weight, healthy condition, with no defect or injury.

Ubiquinone (co-Q10) Nanoemulsion and Emulsion Preparation

The nanoemulsion formula in this study refers to the modified formula from Erawati et al, 2014⁴. In this experiment the concentration of ubiquinone (co-Q10), Tween 80, Span 80, ethanol 96%, VCO and acetate buffer solution pH 4.2 ± 0.5 were: 1%; 18.66%; 1.92%; 3.42%; up to 100%, respectively.

The emulsion formula consists of; 1% ubiquinone (co-Q10) was dissolved in 35% VCO and then added with 22% of surfactant mixture (Tween 80 and Span 80), it was stirred until the mixture became homogeneous. Subsequently, a solution of acetate buffer pH 4.2 ± 0.5 up to 100%.

RESULTS AND DISCUSSION

The characteristics of ubiquinone (co-Q10) nanoemulsion (QN) and ubiquinone (co-Q10) emulsion (QE) included pH, droplets size and viscosity were presented in Table 1. Ubiquinone (Co-Q10) nanoemulsion and ubiquinone (co-Q10) emulsion both have same pH value were 4.12 ± 0.02 . The droplets size of QN (93.2 ± 2.78 nm) was smaller than QE (10.55 ± 0.60 μ m). The viscosity of QN (8.5 cPas) lower than QE (14.2 cPas). The morphology of QN droplet

^{*}Author for Correspondence: era_ffua@yahoo.co.id

Figure 1: Droplet morphology of QN by Transmission Electron Microscope JEOL JEM 1400.

Figure 2: Histological preparation of skin penetration of co-Q10 in emulsion at 2 hours (A1), at 6 hours (A2) after sample application and in nanoemulsion at 2 hours (B1), at 6 hours (B2) after sample application was determined by Microscope Olympus FX-100 42x zoom.

by Transmission Electron Microscope JEOL JEM 1400 was presented in figure 1. Histological preparation of skin penetration of ubiquinone (co-Q10) in emulsion at 2 hours (A1), at 6 hours (A2) after

sample application and in nanoemulsion at 2 hours (B1), at 6 hours (B2) after sample application was determined by Microscope Olympus FX-100 42x zoom, presented in Figure 2. The result of in vivo skin penetration tests at two

Table 1: Characteristics of ubiquinone (co-Q10) nanoemulsion (QN) and emulsion (QE).

Characteristic	QN	QE
pH	4.12 ± 0.02	4.12 ± 0.02
Droplet size	93.2 ± 2.78 nm	10.55 ± 0.60 µm
Viscosity	8.5 cPas	14.2 cPas

different times (2 and 6 hours after sample application) was known that penetration depth of ubiquinone (co-Q10) in nanoemulsion were 269 ± 48.12 and 993 ± 36.49 µm, respectively. And than in emulsion were 26.33 ± 6.43 and 110.33 ± 9.124 µm, respectively. It shows that penetration of ubiquinone (co-Q10) in nanoemulsion delivery system into the rat skin faster and deeper than in emulsion.

Ubiquinone (co-Q10) penetration rate in nanoemulsion is faster than in the emulsion caused by three factors; the first is the lower nanoemulsion viscosity than the emulsion, the second is the nanoemulsion droplets size smaller than the emulsion, so the movement of the active ingredient is easier in nanoemulsion. The third is the nanoemulsion system which increases the release rate so that it's much more active ingredients are ready to be penetrated into the skin^{6,7}.

CONCLUSION

Penetrasi of ubiquinone (co-Q10) in nanoemulsion was deeper (269 ± 48.12 dan 993 ± 36.49 µm) than in emulsion (26.33 ± 6.43 and 110.33 ± 9.124 µm), after 2 and 6 hours sample application on Wistar rats skin.

ACKNOWLEDGEMENT

The research was funded by Faculty of Pharmacy Airlangga University

REFERENCES

1. Thakur N., Garg G., Sharma P.K. and Kumar N., 2012. Nanoemulsions: A Review on Various Pharmaceutical Application. *Global J. Pharmacol*, 6(3), pp. 222-225.
2. Bhat P., and Madhav S., 2011. A Detailed Review on Nanoemulsion Drug Delivery System. *International Journal of Pharmaceutical Sciences and Research*, Vol. 2 issue 9, pp. 2292- 2298.
3. Bank, Ginny., Kagan, Daniel., and Madhavi, Doddabele, 2010. Coenzyme Q10: Clinical Update and Bioavailability. *Journal of Evidence-Based Complementary & Alternative Medicine*, Vol. 16(2), pp. 129-137.
4. Gupta, A. and Kumar, R., 2010, Effect of geometry on droplet formation in the squeezing regime in a microfluidic T-junction. *Microfluidics and Nanofluidics*, 8(6), pp.799-812.
5. Erawati T. Hendradi E., Soeratri W., 2014, Praformulation Study of p-Methoxycinnamic Acid (APMS) Nanoemulsion Using Vegetable Oils (Soybean Oil, Corn Oil, VCO), *Int. J Pharm. Pharm. Sci*, Vol. 6, Issue 2, pp 99-101.
6. Tristiana Erawati M., Retnowati, Amalia Wardatul F., Widji Soeratri, 2017, Virgin Coconut Oil as Oil Phase in Tretinoin Nanoemulsion. *Internasional Journal of Drug Delivery Technology*, Volume 7, Issue 1, pp 42-44.
7. Mishra, Raj Kumar, Soni G.C., and Mishra, R.P., 2014, A Review Article: On Nanoemulsion. *World Journal of Pharmacy and Pharmaceutical Sciences*, Vol. 3, issue 9, p. 258-274.

Skin Penetration of Ubiquinone (Co-Q10) In Nanoemulsi on Delivery System Using Virgin Cocunut Oil (Vco)

ORIGINALITY REPORT

5%

SIMILARITY INDEX

5%

INTERNET SOURCES

4%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1

forddele.dk

Internet Source

2%

2

Gross-Heitfeld, Christoph, Jürgen Linders, Ralph Appel, Florian Selbach, and Christian Mayer. "Polyalkylcyanoacrylate Nanocapsules: Variation of Membrane Permeability by Chemical Cross-Linking", The Journal of Physical Chemistry B

Publication

1%

3

www.univmed.org

Internet Source

1%

4

Sergey Vladimirovich Moskvina, Oleg Ivanovich Apolikhin. "Effectiveness of low level laser therapy for treating male infertility", BioMedicine, 2018

Publication

1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Skin Penetration of Ubiquinone (Co-Q10) In Nanoemulsi on Delivery System Using Virgin Cocunut Oil (Vco)

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3