Synthesis of N'-(2-Methoxybenzylidene)-4-Hydroxy Benzohydrazide and N'-(4-Nitrobenzylidene)-4-Hydroxy Benzohydrazide, in Silico Study and Antibacterial Activity

by Isnaeni Isnaeni

Submission date: 27-Mar-2019 10:54AM (UTC+0800)

Submission ID: 1100520798

File name: y_Benzohydrazide,_in_Silico_Study_and_Antibacterial_Activity.pdf (490.82K)

Word count: 2942

Character count: 17504

Synthesis of N'-(2-Methoxybenzylidene)-4-Hydroxy Benzohydrazide and N'-(4-Nitrobenzylidene)-4-Hydroxy Benzohydrazide, in Silico Study and Antibacterial Activity

Suzana, Isnaeni, Tutuk Budiati*

Faculty of Pharmacy Airlangga University, Dharmawangsa Dalam Street, Surabaya 60286-Indonesia.

Available Online: 25th January, 2017

ABSTRACT

In this study, synthesized N'-(2-methoxybenzylidene)-4-hydroxybenzohydrazide and N'-(4-nitrobenzylidene)-4-hydroxybenzohydrazide in two step reaction by using methylparaben as starting material has been performed. Methylparaben was treated with hydrazine hydrate to obtain 4-hydroxybenzohydrazide. The reaction was carried out by microwave irradiation resulting 91 % yield. The obtained compound was then reacted with 2-methoxybenzaldehyde or 4-nitrobenzaldehyde to accomplish the target molecule, N'-(2-methoxybenzylidene)-4-hydroxybenzohydrazide and N'-(4-nitrobenzylidene)-4-hydroxybenzohydrazide in 55% and 72% yield respectively. Identification of N'-(2-methoxybenzylidene)-4-hydroxy benzo hydrazide and N'-(4-nitrobenzylidene)-4-hydroxybenzohidroksida was performed by FT-IR, MS, ¹H-NMR, and ¹³C-NMR spectroscopy. In silico study was done with receptor pdb 1C14. The N'-(4-nitrobenzylidene)-4-hydroxybenzohydrazide exhibited antimicrobial activity against *Escherichia coli* (MIC=31.3 ppm), *Bacillus subtilis* (MIC=500 ppm). Antimicrobial activity of N'-(2-methoxybenzylidene)-4-hydroxybenzohydrazide against *Bacillus subtilis* (MIC=31.3 ppm) and MIC= 500 ppm against *Escherichia coli*.

Keywords: Synthesis, N'-Benzylidene-4-hydroxybenzohydrazide derivatives, microwave, antibacterial.

INTRODUCTION

New drugs invention and development is crucial in the pharmaceutical industry. The application of conventional methods in developing new drugs such as through organic reactions which require high temperature has been widely practiced. Traditional heater equipment which are commonly used are sand-bath, oil-bath, and heating mantle. These techniques take much time to react the samples and may cause temperature differences on the samples. Moreover, heated reaction flask surface may cause localized overheating which results products, reagents, and substrate decomposition in prolonged heating. Therefore, new reaction methods which are expected to be able to perform quicker synthesis with higher product percentage and quality should be proposed. One of new synthesis methods is conducted by microwave irradiation. This technique is applied in new drugs development¹⁻³. Hydrazide derivatives are widely used in new drug development for its biological activity and various clinical applications such as anticancer⁴, antitumor⁵, antimicrobial⁶, antifungal^{7,8}, antiHIV⁹, and antituberculosis¹⁰. In some literature, it is mentioned that hydrazide derivatives have pharmacological activities (i.e. anticancer, anticonvulsant, antiinflammation, antibacterial, and antioxidant) are related to -CONHN=CN functional group in its molecular structure¹⁰. Although hydrazide derivatives can be synthesized with conventional¹¹ and microwave irradiation methods in this research, we attempt to conduct synthesis

reaction by applying microwave irradiation. This method is chosen because it takes shorter time and it's able to result higher percentage product. The products are more eco-friendly because no toxic solvents are used in this process. Moreover, optimization of 4-hydroxybenzohydrazide derivatives were using microwave irradiation reaction method has not been conducted. Based on the background explained above, the problem proposed in this research is whether microwave irradiation can be used to synthesize 4-hydroxybenzohydrazide compound and its derivatives. Synthesis of 4-hydroxybenzohydrazide derivatives used methyl paraben as starting material. This research is expected to be able to provide new understanding of condensation reaction through the application of microwave irradiation to synthesize 4-hydroxybenzohydrazide derivatives. The compounds are tested antibacterial activity against *Bacillus subtilis* (Gram positive) and *Escherichia coli* (Gram negative). The scheme of synthesis reaction is presented in Figure 1.

MATERIALS AND METHODS

All chemicals were used in this study from commercial sources. Methyl paraben, hydrazine hydrate, 2-methoxybenzaldehyde, 4-nitrobenzaldehyde ethanol 95%, chloroform, ethyl acetate, acetone, hexane, KBr, silica gel 60 GF₂₅₄ were purchased from Merck. Glassware commonly used in the chemical synthesis laboratories, Sanyo microwave EM-S 400 Watt, Spectrophotometer

*Author for Correspondence: suzanarushadi@yahoo.com

Buck Scientific IR M-500, FT-NMR spectrometer JEOL ECS-400.

Preparation of 4-hydroxybenzohydrazide (I) with microwave irradiation^{12,13}

Methyl paraben (10 mmol) and hydrazine hydrate (50 mmol) were stirred until homogeneous. The mixture was put in the microwave on 160 Watt power for 2-8 minutes while stirring every 2 minutes. The progress of the reaction was monitored by TLC. The mixture was cooled to room temperature and then added 20-30 ml of ice water, filtered, washed with ethanol. Crystals were recrystallized with absolute ethanol. The purity tests were done by melting point and thin layer chromatography (TLC) using three different eluents. Identifications were carried out by FT-IR, and ¹H-NMR spectroscopy.

Preparation of *N'*-(2-methoxybenzylidene)-4-hydroxybenzohydrazide and *N'*-(4-nitrobenzylidene)-4-hydroxybenzohydrazide^{12,13}

4-Hydroxybenzohydrazide (10 mmol) and benzaldehyde (20 mmol) were dissolved in ethanol (12 ml). Ethanol was evaporated until exhausted. The mixture was put in the microwave on 320 Watt power for 2 minutes (for synthesis IIb), on 160 Watt power for 4 minutes (for synthesis IIa). The mixture was cooled to room temperature, then added to 20-30 ml of ice water, filtered, washed with ethanol. Crystals were crystallized with absolute ethanol. The purity tests were done using melting point and thin layer chromatography using three different eluents. Identifications were carried out by FT-IR, MS, ¹H-NMR, and ¹³C-NMR spectroscopy.

Antibacterial Activity

Antibacterial activity was done by well diffusion method¹⁴⁻¹⁶. In this method, The wells are filled with test compounds. Petri plate was labeled with the name of the culture, sample and standard at the bottom of the plate. Inoculums were standardized by adjusting the turbidity of the culture according to McFarland standards. The standardized inoculums was prepared in the petri plates. Then sterile agar medium and suspension of microbial cultures were poured into a petri dish aseptically. Petri dishes are left at room temperature for 15 minutes and then incubated 37°C for 24 hours¹⁷. After incubation at 37°C for 24 hours, the diameter of the zone inhibition was measured. The inhibitory activity of DMSO was also employed as a negative control. Positive control used Nifuroxazide 31.3 ppm. The test compounds were evaluated for their antibacterial activity against *Escherichia coli* (ATCC 25922) and *Bacillus subtilis* (ATCC 6633)¹⁸.

RESULTS AND DISCUSSION

Synthesis of 4-hydroxybenzohydrazide derivatives using condensation reaction were carried in two reaction phases. The first phase was conducted reaction methyl paraben with hydrazine hydrate with microwave irradiation 160 Watt power for 2-8 minutes to produce 4-hydroxybenzohydrazide. It was done by the nucleophilic substitution reaction¹¹. The second phase was conducted to produce 4-hydroxybenzohydrazide derivatives using the condensation reaction method. This method was chosen for its practicality and not requiring toxic solvent. Besides,

the method took a relatively short time (about 8 minutes). The reaction was carried out without toxic solvent. Therefore, this reaction conformed to eco-chemistry because lacking of toxic solvent and energy saving^{19,20}. 4-Hydroxybenzohydrazide derivatives were synthesized by reacting 4-hydroxy benzohydrazide and 2-methoxybenzaldehyde or 4-nitrobenzaldehyde. Ethanol was used as solvent, then evaporated to run out.

Characterization of 4-hydroxybenzohydrazide (I)^{21,22}

Obtained in 91% yield as white needle-shaped crystals. The result of purity test was conducted using thin-layer chromatography indicated one stain. Three of eluents were chloroform:ethyl acetate (1:1), chloroform:aceton (1:2) and aceton:ethyl acetate (2:1), mp. 255-256°C. IR (KBr in cm⁻¹): 1620 (-C=O amide), 3318 (-OH phenolic), 1590 and 1467 (-C=C- aromatic), 1354 (C-N), 3005 (Csp²-H), 850 (para disubstitution on benzene), 3197 (-NH₂). ¹H-NMR (DMSO-d₆, δ, ppm): 6.74-6.72 (d, J=8.4 Hz, 2H, C₆H₄-), 7.65-7.62 (d, J=9.2 Hz, 2.H, C₆H₄-), 9.89 (s, 1H, OH), 9.44 (s, 1H, NH), 4.32 (s, 2H, NH₂).

Characterization of *N'*-(2-methoxybenzylidene)-4-hydroxybenzohydrazide (IIa)^{21,22}

Obtained in 55% yield as white needle-shaped crystals. The result of purity test was conducted using thin-layer chromatography indicated one stain, m.p. 229-230 °C. IR (KBr in cm⁻¹): 1606 (-C=O amide), 3090 (-OH phenolic), 1508 (-C=C- aromatic), 1557 (-NH), 1359 (C-N), 2941 (Csp³-H), 841 (para disubstitution on benzene), 1606 (C=N), 1254 (phenylalkyl ether). [M]⁺=270. ¹H-NMR (DMSO-d₆, δ, ppm): 11.61 (s, 1H, OH), 10.08 (s, 1H, NH), 8.74 (s, 1H, HC=N), 6.82-6.79 (d, J=8.4 Hz, 2H from aromatic ring), 6.99-6.95 (t, J=8 Hz, 1H from aromatic ring), 7.06-7.04 (d, J=8.8 Hz, 1H from aromatic ring), 7.18-7.34 (t, J=7.2Hz, 1H from aromatic ring), 7.78-7.76 (d, J=8.4Hz, 1H from aromatic ring), 7.82-7.80 (d, J=8 Hz, 2H from aromatic ring), 3.90 (s, 3H, OCH₃). ¹³C-NMR (DMSO-d₆, δ, ppm): 163.05, 161.16, 158.13, 142.71, 131.84, 130.18, 125.93 (2C), 124.34, 123.02, 121.25, 115.50 (2C), 112.31, 56.17.

Characterization of *N'*-(4-nitrobenzylidene)-4-hydroxybenzohydrazide (IIb)^{21,22}

Obtained in 72% yield as yellow needle-shaped crystals. The result of purity test was conducted using thin-layer chromatography indicated one stain, m.p. 289-290 °C. IR (KBr in cm⁻¹): 1657 (-C=O amide), 3467 (-OH phenolic), 1581 (-C=C- aromatic), 3168 (-NH), 3340 (NH₂), 1341 (C-N), 846 (para disubstitution on benzene), 1606 (C=N), 1512 (N=O). [M]⁺=285. ¹H-NMR (DMSO-d₆, δ, ppm): 11.90 (s, 1H, OH), 10.16 (s, 1H, NH), 8.47 (s, 1H, HC=N), 6.85-6.83 (d, J=8.8 Hz, 2H from aromatic ring), 7.80-7.78 (d, J=8.8 Hz, 2H from aromatic ring), 7.92-7.90 (d, J=8.4Hz, 2H from aromatic ring), 8.25-8.22 (d, J=8.8 Hz, 2H from aromatic ring). ¹³C-NMR (DMSO-d₆, δ, ppm): 163.47, 161.51, 148.12, 144.73, 141.42, 130.43 (2C), 128.33, 124.57 (2C), 123.99 (2C), 115.61 (2C).

In Silico Study

To estimate the antimicrobial activity of derivatives 4-hydroxybenzohydrazide, in silico study was done with receptor pdb 1C14 program MVD 5.0 (Mollegro Virtual Docker). 1C14 is a receptor model of inhibitor triclosan,

Figure 1: Synthesis pathway of N'-Benzylidene-4-hydroxybenzohydrazide derivatives.

Table 1: Result of docking molecule methyl paraben, N'-(2-methoxybenzylidene)-4-hydroxybenzohydrazide and N'-(4-nitrobenzylidene)-4-hydroxybenzohydrazide with receptor [pdb. 1C14](#) program MVD 5.0 (Mollegro Virtual Docker).

Compounds	Rerank (kcal/mol)	Score
<p>Methyl paraben</p>	-66.498
<p>4-hydroxybenzohydrazide (I)</p>	-65,510
<p>Nifuroxazide</p>	-106,971
<p>N-(2-methoxybenzylidene)-4-hydroxybenzohydrazide (IIa)</p>	-108.675
<p>N'-(4-nitrobenzylidene)-4-hydroxybenzohydrazide (IIb)</p>	-106.264	

with the ligand code NAD. Based on the rerank score it was found that the target compounds exhibited a greater antimicrobial activity than methyl paraben²⁴.

Antibacterial Activity

The synthetic compounds were evaluated for antibacterial activity against Gram positive (*Bacillus subtilis*) and Gram negative (*Escherichia coli*). Antibacterial activities against

Escherichia coli of the compounds as well as reference drug are summarized in Table 1. The results indicated that N'-(2-methoxybenzylidene)-4-hydroxybenzohydrazide and N'-(4-nitrobenzylidene)-4-hydroxybenzohydrazide have had activities against *E. coli*. Results of minimal inhibitory concentration (MIC) of N'-(2-methoxybenzylidene)-4-hydroxybenzohydrazide was 500 ppm and N'-(4-nitrobenzylidene)-4-hydroxybenzohydrazide was 31.3 ppm. N'-(4-nitrobenzylidene)-4-hydroxybenzohydrazide has MIC against *E. coli* (Gram negative) bigger than N'-(2-methoxybenzylidene)-4-hydroxybenzohydrazide.

Antibacterial activities against *Bacillus subtilis* of the compounds as well as reference drug are summarized in Table 2. The results indicated that N'-(2-methoxybenzylidene)-4-hydroxybenzohydrazide and N'-(4-nitrobenzylidene)-4-hydroxybenzohydrazide had activated against *Bacillus subtilis*. Results of minimal inhibitory concentration (MIC) of N'-(2-methoxybenzylidene)-4-hydroxybenzohydrazide was 31.3 ppm and N'-(4-nitrobenzylidene)-4-hydroxybenzohydrazide was 500 ppm. N'-(2-methoxybenzylidene)-4-hydroxybenzohydrazide has MIC against *B. subtilis* (Gram positive) bigger than N'-(4-nitrobenzylidene)-4-hydroxybenzohydrazide. The antibacterial activity of the target compounds due to the phenolic group ($\text{HO}-\text{C}_6\text{H}_5$)²⁵ and the azometin group ($-\text{HN}=\text{N}=\text{CH}-$)²⁶. Variations of the target compound antibacterial activity against different bacteria (Gram positive / Gram negative) is influenced by electronic, steric, and lipophilicity of compounds. In silico test lipophilicity factors not included, so it can give different results²⁴.

CONCLUSION

It can be concluded that N'-2-methoxybenzylidene-4-hydroxybenzohydrazide and its derivatives can be synthesized from the methyl paraben as starting material by microwave irradiation (160-320 Watt power for 2-8 minutes). The yields were obtained between 55-72%. The preliminary biological tests indicated that N'-(4-nitrobenzylidene)-4-hydroxybenzohydrazide has effective activity against *Escherichia coli* (MIC=31.3 ppm), and

Table 2: Antibacterial activity against *E.coli* of the test compounds.

Compound	Inhibition zone (mm)						Amoxi cillin	Control negative
	Concentration (ppm)							
	31,3	62,5	125	250	500	1000		
I	-	-	-	-	-	-	14.5±0.67	-
Ila	-	-	-	-	-	11.5±0.50	14.0±0.50	-
Ilb	-	10.0±0.33	12.0±0.50	14.5±0.67	16.5±0.67	13.0±0.50	14.0±0.67	-

(-) No inhibition zone, n=3 replications.

Table 3: Antibacterial activity against *Bacillus subtilis* of the test compounds.

Compound	Inhibition zone (mm)						Amoxi cillin	Control negative
	Concentration (ppm)							
	31,3	62,5	125	250	500	1000		
I	-	-	-	-	12.0±0.33	10.0±0.50	14.5±0.67	-
Ila	-	9.5±0.33	9.5±0.50	9.5±0.33	11.0±0.67	11.3±0.40	14.0±0.67	-
Ilb	-	-	-	-	-	9.5±0.33	14.0±0.50	-

(-) No inhibition zone, n=3 replications.

N'-(2-methoxybenzylidene)-4-hydroxybenzohydrazide has effective activity against *Bacillus subtilis* (MIC=31.3 ppm).

6. CKNOWLEDGMENT

This research was financially funded by Indonesian Directorate General of Higher Education (DGHE) or DIKTI through *Penelitian Unggulan Perguruan Tinggi*'s scheme of 2016.

REFERENCES

- Kappe O. Microwave-enhanced chemistry-anything technology revolutionizing organic synthesis and drug discovery 2004.
- Mavandadi F, Lidstrom P. Microwave-assisted chemistry in drug discovery. *Current Topics in Medicinal Chemistry* 2004; 4: 773-92.
- Suthakaran R, Kavimani S, Venkappayya D, Jayasree P, Deepthi V, Tehseen F, Suganthi K. Microwave assisted rapid synthesis and anti-microbial activity of 4-oxothiazolidine derivatives. *Rasayan J. Chem.* 2008; 1: 30-8.
- Al-Abdullah ES. Synthesis and anticancer activity of some novel tetralin-6-yl-pyrazoline, 2-thioxopyrimidine, 2-oxopyridine, 2-thioxo-pyridine and 2-iminopyridine derivatives. *Molecules* 2011; 16: 3410-9.
- Mohareb RM, Mohamed AA. The reaction of cyanoacetylhydrazine with w-bromo-(4-methyl)acetophenone: Synthesis of heterocyclic derivatives with antitumor activity. *Molecules* 2010; 15: 3602-17.
- Manivel P, Mohana RS, Kumar SR, Nawazkhan F. Synthesis of substituted isoquinolin-1-yl-2-(cycloalk-2-enylidene)hydrazines and their antimicrobial properties. *J. Chil. Chem. Soc.* 2009; 54: 183-5.
- Kostecka M. Synthesis of a new group of aliphatic hydrazide derivatives and the correlations between their molecular structure and biological activity. *Molecules* 2012; 17: 3560-73.
- Kaur R, Singh A, Budhlakoti P, Sanwal R. Synthesis, characterization and antifungal activity of certain (E)-1-(1-(substitutedphenyl)ethylidene)-2-(6-methylbenzo[d] thiazol-2-yl)hydrazine analogues. *Int. J. Pharm. Biol. Arch.* 2010; 1:56-61.
- Sechi M, Azeena U, Delussu MP, Dallochio R, Dessi A, Cosseddu A, Pala N, Neamati N. Design and synthesis of bis-amide and hydrazide containing derivatives of malonic acid as potential HIV-1 integrase inhibitors. *Molecules* 2008; 13: 2442-61.
- Raparti V, Chitre T, Bothara K, Kumar V, Dangre S, Khachane C, Gore S, Deshmane B. Novel 4-(morpholin-4-yl)-N'-(arylidene)benzohydrazides; synthesis, antimycobacterial activity and QSAR investigations. *Eur. J. Med. Chem.* 2009; 44: 3954-60.
- McMurry J. *Organic Chemistry*, 7th Ed. Thomson Learning Inc. USA 2008; 877-84.
- Habibi D, Marvi O. Montmorillonite KSF and montmorillonite K-10 clays as efficient catalysts for the solventless synthesis of bismaleimides and bisphthalimides using microwave irradiation. *General paper Arkivoc (xiii)* 2006: 8-15.
- Jain AK, Gupta PK, Ganesan K, Pande A, Malhotra RC. Rapid solvent free synthesis of aromatic hydrazides under microwave irradiation. *Defense Science Journal* 2007; 57 (2): 267-70.
- Balouri M., Sadiki M., Ibsouda SK. Methods for in vitro evaluating antimicrobial activity: a review. *Journal of Pharmaceutical Analysis*, 2016; 11 (5), pp. 1-10.
- Vellaiswamy G, Ramaswamy S. Synthesis, spectral characterization and antimicrobial screening of novel Schiff bases from sulfa drugs. *International Journal of Pharmacy and Pharmaceutical Sciences*, 2014; 6 (1): 487-91.
- Namasivayam N, Theivarasu C. Synthesis of substituted piperidine-4-ones with dichloro(cyclooctadiene) palladium (II) and antimicrobial activity, *International Journal of*

- Pharmacy and Pharmaceutical Sciences, 2014; 6 (1): 440-4.
17. Ramalingan C, Park YT, Kabilan S. Synthesis, stereochemistry and antimicrobial evaluation of substituted piperidin-4-one oxime ethers. *Eur J Med Chem*, 2006; 41: 683-96.
18. Kumar S, Goel N, Afzal O, Ali MR, Bawa S. *In vitro* antibacterial/antifungal screening of 2-chloroquinoline scaffold derivatives. *Journal of Antibiotics Research*, 2015; 1 (1):101. Kabalka GW, Wang I, Pagni RM. Potassium fluoride doped alumina: an effective reagent for ester hydrolysis under solvent free conditions. *Green Chemistry* 2001; 3: 261-2.
19. Paul S, Nanda P and Gupta R. PhCOCl-Py/BasicAlumina as a versatile reagent for benzoylation in solvent-free condition. *Molecule* 2003; 8: 374-80.
20. Pavia DL, Gary M, Lampman, George S, Kriz, James RV. *Introduction of Spectroscopy*, 4th Ed. Brooks/Cole USA 2009: 381-403.
21. Silverstein RM, Francis XW, David JK. *Spectrometric Identification of Organic Compounds*, 7th Ed. John Wiley and Sons Inc. USA 2005; 72-4.
22. Kapoor M., Mukhi PLS., Surolia N., Kusuga K., Surolia A. Kinetic and structural analysis of the increased affinity of enoyl-ACP (acyl-carrier protein) reductase for triclosan in presence of NAD⁺. *Biochemistry Journal*, 2004; 381:725-733.
23. Meng E.C., Shoichet B.K., Kuntz I.D. Automated docking with grid-based energy evaluation. *Journal of Computational Chemistry*, 2004; 13 (4), pp. 505-524.
24. Merkl R., Hradkova L., Filip V., Smidrkal J. Antimicrobial and antioxidant properties of phenolic acids alkyl esters. *Czech J. Food Sci.*, 2010; 28: 275-279.
25. Pangal A., Ahmed K., Shaikh S. Synthesis, Characterization and Study of Antimicrobial Activity of 2,6-Ditertiarybutyl-1,4-benzoquinone hydrazones. *International research Journal of Pharmacy*, 2013; 4 (8), pp. 172-176.

Synthesis of N'-(2-Methoxybenzylidene)-4-Hydroxy Benzohydrazide and N'-(4-Nitrobenzylidene)-4-Hydroxy Benzohydrazide, in Silico Study and Antibacterial Activity

ORIGINALITY REPORT

19%

SIMILARITY INDEX

11%

INTERNET SOURCES

17%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1

S Suzana, M I Sulistyowaty, Isnaeni, T Budiati. "Synthesis, docking molecule study and antibacterial activity of N'-(4-Fluorobenzylidene)-4- hydroxybenzohydrazide and N'-(4-Fluorobenzylidene)-4-hydroxybenzohydrazide)", Journal of Physics: Conference Series, 2019
Publication

10%
- 2

www.ijppsjournal.com
Internet Source

1%
- 3

Tatsumi Kimura. "Branched epoxy copolymers with oligo(benzoate) side chains carrying donor and acceptor functions in each end", Polymers for Advanced Technologies, 03/2001
Publication

1%
- 4

Kosuke Kuroda, Yumiko Shimada, Kenji Takahashi. "Hand-holding and releasing between the anion and cation to change their macroscopic behavior in water", Green Energy

1%

5	onlinelibrary.wiley.com Internet Source	1 %
6	Sanjaya, Ridwan, Agustine Eva Maria Soekesi, and A. Posmaria Setiasiwit Sitohang. "Technology entrepreneurship model development for teachers", International Journal of Technoentrepreneurship, 2015. Publication	1 %
7	"Handbook of Computational Chemistry", Springer Nature, 2017 Publication	1 %
8	ijpbs.net Internet Source	1 %
9	www.ajol.info Internet Source	1 %
10	lib.ui.ac.id Internet Source	<1 %
11	shodhganga.inflibnet.ac.in:8080 Internet Source	<1 %
12	Sébastien Moniot, Mariantonietta Forgione, Alessia Lucidi, Gebremedhin S. Hailu et al. "Development of 1,2,4-Oxadiazoles as Potent and Selective Inhibitors of the Human	<1 %

Deacetylase Sirtuin 2: Structure–Activity Relationship, X-ray Crystal Structure, and Anticancer Activity", Journal of Medicinal Chemistry, 2017

Publication

13

www.google.com

Internet Source

<1 %

14

www.imedpub.com

Internet Source

<1 %

15

www.faqs.org

Internet Source

<1 %

16

Jin, Hongjun, Jinda Fan, Xiang Zhang, Junfeng Li, Hubert P. Flores, Joel S. Perlmutter, Stanley M. Parsons, and Zhude Tu. "Radiosynthesis and in vivo evaluation of a novel σ 1 selective PET ligand", MedChemComm, 2014.

Publication

<1 %

17

worldwidescience.org

Internet Source

<1 %

18

Wang, C.. "Chlorotrimethylsilane-promoted one-pot synthesis of steroidal[17,16-d]pyrimidines", Steroids, 201012

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On