


IMRM



PROCEEDINGS OF INTERNATIONAL MEETING
ON REGENERATIVE MEDICINE
Surabaya, August 15th-15th 2017



From Foundational Bioscience
to Human Functioning

Editors

Sri Mardjati Muli Wulan, Hiroaki Kimura, Elana
Postema, Dharma Dwanandya and Apichana Kovalittha
Universitas Airlangga

Organized by

Faculty of Medicine, Universitas Airlangga



Proceedings

Proceedings of the International Meeting on Regenerative Medicine

August 15-16, 2017, in Surabaya, Indonesia



Editors: Sri Mardjati Mei Wulan¹; Hiroaki Kimura²; Klaas Postema³; Delvac Oceandy⁴ and Apichana Kovindha⁵

Affiliations: ¹ Airlangga University, Indonesia ; ² Hiroshima University Hospital, Japan ; ³ University of Groningen, Netherlands ; ⁴ University of Manchester, United Kingdom ; ⁵ University of Chiang Mai, Thailand

ISBN: 978-989-758-334-6

Conference Link: <http://conference.unair.ac.id/index.php/IMR/imurmua2017>

Foreword: As the world evolves rapidly, the field of science generally also has grown tremendously. This international scientific meeting is one of the real actions of Faculty of Medicine Universitas Airlangga to improve the quality and quantity of basic research and applied science and to develop institutional quality oriented to be able to compete at international level. Regenerative Medicine is an interesting-new field of medicine. This field holds the promise of engineering damaged tissues and organs by stimulating the body's own repair mechanisms to functionally heal previously irreparable tissues or organs. The International Meeting on Regenerative Medicine (IMRM) is held as collaboration among departments (PM&R, cardiology, orthopaedic, neurosurgery, plastic surgery & reconstruction and regenerative medicine). This event is attended by students, academician, researchers, undergraduate students, magister students, doctoral students and practioners. We

Volumes:
Show 50 papers

Vol. 1 - 978-989-758-334-6

Rehabilitation Programs for Total Hip Arthroplasty in Patients with Enclosed Hip Bilateral: A Case Report P. 5 - 8

Yuli Indah Kurnia and I Putu Pawana

DOI:10.5220/0007314400050008

Pyruvat Dehydrogenase Deficiency – Physical Medicine and Rehabilitation Approach: A Case Report P. 9 - 12

Yuli Indah Kurnia and S.M Mei Wulan

DOI:10.5220/0007314500090012

The Role of EMG Biofeedback and Comprehensive Rehabilitation in Brachial Plexus Injury after FFMT: A Case Report P. 13 - 17

Fitri Azizah Noor and Patricia Maria K

DOI:10.5220/0007314600130017

Rehabilitation Management in Late-Findings of Ankylosing Spondylitis Patient: A Case Report P. 18 - 22

Adesti Permasari and Reni Hendrarati Masduchi

DOI:10.5220/0007314700180022

Rehabilitation of Patients with Foville’s Syndrome after the First ICH Stroke: A Case Report P. 23 - 25

Bimo Sutomo and Nuniek Nugraheni

DOI:10.5220/0007314800230025

Rehabilitation Management of a Child with Scleroderma: A Case Report P. 26 - 30

Bayu Putra and Nurul Wardani

DOI:10.5220/0007314900260030

Patient with Multiple Non-contiguous Spondylitis (Spinal) Tuberculosis: A Case Report P. 31 - 34

Bayu Putra and Reni Masduchi

DOI:10.5220/0007315000310034

Correlation of ASIA Impairment Scale (AIS) Classification and Pain in Spinal Cord Injury Patients in Dr Soetomo Hospital Surabaya, Indonesia P. 35 - 38

Bona Pardede and Patricia Maria

DOI:10.5220/0007315100350038

Non-Traumatic Spinal Cord Injury – Tuberculous Spondylitis: A Case Report P. 39 - 42

Bona Pardede and Dewi Poerwandari

DOI:10.5220/0007315200390042

Correlation between Parity Status and Urinary Incontinence Among Elderly Women P. 43 - 45

DOI:10.5220/0007315300430045

Relation of Cognitive and Physical Functional Performance with Urinary Incontinence Problems in the Elderly P. 46 - 48

Dewi Ayub, Rwahita Satyawati and Nuniek Nugraheni

DOI:10.5220/0007315400460048

The Effect of Garlic Extract on Endothelial Progenitor Cells (EPCs) Quantification in Chronic Stable Angina Pectoris Patients P. 49 - 52

Alisia Putri, Yudi Her Oktoviono and Budi Susetyo Pikir

DOI:10.5220/0007315500490052

In Vitro Effect of Lisinopril on Endothelial Progenitor Cell (Epc) Proliferation from Peripheral Blood of Stable Coronary Artery Disease Patients P. 53 - 58

Yudi Her Oktoviono, Ragil Nur Rosyadi and Djoko Soemantri

DOI:10.5220/0007315600530058

Intraventricular Transplantation of Autologous Bone Marrow Mesenchymal Stem Cells for Stroke: Rationale, Technique and Early Clinical Experience P. 59 - 64

Asra Al Fauzi

DOI:10.5220/0007315700590064

Comparison of the Effects of Resistance Training with and without Vascular Occlusion on Creatine Kinase (CK) and C-Reactive Protein (CRP) P. 65 - 70

David Sugiarto, Hening Laswati, SM Mei Wulan and Hiroaki Kimura

DOI:10.5220/0007315800650070

Direct and Indirect Method of Sonographic Measurements of the Median Nerve in Carpal Tunnel Syndrome P. 71 - 75

Lydia Arfianti

DOI:10.5220/0007315900710075

Increasing Muscle Regeneration in Response to Exercise P. 76 - 80

S.M Mei Wulan

DOI:10.5220/0007316000760080

The Role of Comprehensive Rehabilitation Management in Noncontiguous Spinal Tuberculosis Involving Thoracic and Lumbar: A Case Report P. 81 - 84

Fitria Wardhani and Indrayuni Wardhani

DOI:10.5220/0007316100810084

Rehabilitation Program for Elderly with Neglected Elbow Dislocation with Coronary Heart Disease: A Case Report P. 85 - 89

Ni Made Maya Aprilia Sari and Rwahita Satyawati

DOI:10.5220/0007316200850089

Home-based Rehabilitation to Reduce Spasticity in Spastic Quadriplegic**Cerebral Palsy: A Case Report**

Mufidha Zulfia and SM Mei Wulan

DOI:10.5220/0007316300900092

92

Effectiveness of Aerobic Exercise Addition Using a Static Cycle in Improving Cardiorespiratory Fitness and Gait Speed after Sub-Acute Ischemic Stroke

Fifin Indraswari, Hening Laswati and Indrayuni Lukitra W.

DOI:10.5220/0007316400930100

P. 93 -
100**Incentive Spirometer as a Prognostic Factor for Post Cardiac Surgery Patients**

Bastianus Alfian Juatmadja and Dewi Poerwandari

DOI:10.5220/0007316501010103

P. 101 -
103**Rehabilitation Management of Chronic Inflammatory Demyelinating Polyneuropathy: A Case Report**

Sri Panca Lastiani and SM Mei Wulan

DOI:10.5220/0007316601040106

P. 104 -
106**Rehabilitation Management of Intracerebral Hemorrhage with Various Complications: A Case Report**

Gadiza Raizninha P and Ditaruni Asrina U.

DOI:10.5220/0007316701070110

P. 107 -
110**Congenital Scoliosis Caused by Hemivertebrae: A Case Report**

Masadah and Nurul Kusuma Wardani

DOI:10.5220/0007316801110114

P. 111 -
114**Anal Pressure Biofeedback Training in a Patient with Diverting Colostomy after Blunt and Penetrating Anorectal Injury: A Case Report**

Benni Azhari and Martha Kurnia Kusumawardani

DOI:10.5220/0007316901150119

P. 115 -
119**Comparison of the Immediate Effect between Radial Shock Wave Therapy and Ultrasound Diathermy on Ankle Range of Motion in Cerebral Palsy with Plantar Flexor Spasticity**

Nur Sulastri, Sri Mardjiati Mei Wulan and Noor Idha Handajani

DOI:10.5220/0007317001200124

P. 120 -
124**Rehabilitation Outcome of Mixed Transcortical Aphasia and Right Hand Function Disturbance in Intracerebral Hemorrhage: A Case Report**

Wahyu Sita Wardani and Subagyo

DOI:10.5220/0007317101250129

P. 125 -
129**An Obscure Case of Residual Poliomyelitis**

Anindya K Zahra and SM Mei Wulan

DOI:10.5220/0007317201300134

P. 130 -
134

**Benefits of Sport Rehabilitation for Patients with ACL and PCL Tears after P. 135 -
Reconstructive Surgery: A Case Report**

Bimo Sutomo and I Putu Alit Pawana

DOI:10.5220/0007317301350139

139

**Rehabilitation Management for Traumatic SCI During Pregnancy: A Case P. 140 -
Report**

Wahyu Sita Wardani and Patricia Maria

DOI:10.5220/0007317401400143

143

**Difficulty of Returning to Work After Traumatic Spinal Cord Injury: A Case P. 144 -
Report**

Astrina Nur Bahrun and Andriati

DOI:10.5220/0007317501440147

147

**Adolescent Idiopathic Scoliosis Rehabilitation in a Patient with P. 148 -
Tuberculosis-Destroyed Lung: A Case Report**

Benni Azhari and I Lukitra Wardhani

DOI:10.5220/0007317601480151

151

**Rehabilitation Intervention in Hypertensive Intracerebral Hemorrhage: A P. 152 -
Case Report**

Astrina Nur Bahrun and Dewi Poerwandari

DOI:10.5220/0007317701520154

154

**Correlation of Rehabilitation Management Initiation and Length of Stay P. 155 -
on Traumatic Brain Injury Patients in Dr. Soetomo Hospital Surabaya,
Indonesia**

Yohan Christian Suisan and Reni Hendrarati Masduchi

DOI:10.5220/0007317801550158

158

**The Role of Comprehensive Rehabilitation Management in Spina Bifida P. 159 -
Patient with CTEV: A Case Report**

Yohan Christian Suisan and S.M. Mei Wulan

DOI:10.5220/0007317901590162

162

**Urinary Incontinence Prevalence and Its Relation between Pelvic Floor P. 163 -
Muscle Strength and Type of Incontinence Urine Based on Incontinence
Questions (3IQ) among Elderly Men**

Gde Ganjar Oka Narasara , Rwahita Satyawati and Nuniek Nugraheni

DOI:10.5220/0007318001630166

166

Rehabilitation Challenges of Frequent Implant Failure of the Elbow P. 167 -

Alfin Windarjati Wyoso and I Putu Alit Pawana

DOI:10.5220/0007318101670170

170

**Comprehensive Rehabilitation Program for Cervical Spinal Cord Injury P. 171 -
Caused by Atlantoaxial Dislocation: A Case Series Report**

Meta Novantisari and I Lukitra Wardhani

176

Rehabilitation Program for Spinal Neurofibromatosis: A Case Report P. 177 - 180

Meta Novantisari and Damayanti Tinduh

DOI:10.5220/0007318301770180

Rehabilitation of Post Sciatic Nerve Repair with Mesenchymal Stem Cell Augmentation: A Case Report P. 181 - 185

Ikhwan Muhammad and Damayanti Tinduh

DOI:10.5220/0007318401810185

The Effect of Kinesio Taping on Lower Extremity Muscle Power in Adult Males P. 186 - 188

Stela Clara , Julia Windi Gunadi and Yenni Limyati

DOI:10.5220/0007318501860188

The Functional Capacity Difference between Below Elbow Prosthesis Three Jaw Chuck and Five Actuated Finger Terminal Devices on Able-Bodied Subjects P. 189 - 193

Rahmad , Fatchur Rachman , I Lukitra Wardhani and I Putu Alit P

DOI:10.5220/0007318601890193

Role of Allogeneic NK Cells Treatment in the Early Phase of Apoptosis on Poorly Differentiated Retinoblastoma Cells Culture P. 194 - 201

Hendrian Dwikoloso Soebagjo , Fitria Kusumastuti , Parma

Ratna Jaya , Susy Fatmariyanti , Endang Retnowati and Ugroseno S

Yudho Bintoro

DOI:10.5220/0007318701940201

The Relationship between Degree of COPD and Malnutrition in Patients in The Polyclinic of Meuraxa General Hospital P. 202 - 205

Husnah

DOI:10.5220/0007318802020205

Affect of Hatha Yoga Exercise in Executive Function of Diabetes Mellitus Patient P. 206 - 211

Ernie , Robby Tjandra and Suhartono

DOI:10.5220/0007318902060211

Tooth Mobility as an Impact of Psychological and Psycho-Physical Work Distress in a Rat Model P. 212 - 217

Zahreni Hamzah and Elyana Asnar

DOI:10.5220/0007319002120217

Comprehensive Program of Degenerative Scoliosis in Reducing Pain P. 218 - 221

Rwahita Satyawati and Ida Yuanita

DOI:10.5220/0007319102180221

Effects of Tai Chi Exercise on Cardiorespiration Function in Elderly P. 222 -

Women at Nursing Homes

D Kusumawardhana , PM Kurniawati and D Poerwandari

DOI:10.5220/0007319202220226

Functional Mobility and Pain Severity in Older Low Back Pain PatientsP. 227 -
232Dian Marta Sari , Pavankumar Balachandran , Ahmad Ramdan and
Marina A. Moeliono

DOI:10.5220/0007319302270232

Prev	1	2	Next
------	---	---	------

**SCITEPRESS - SCIENCE AND TECHNOLOGY PUBLICATIONS, LDA.****RESOURCES**

Proceedings

Papers

Authors

Ontology

CONTACTS

Science and Technology

Publications, Lda

Avenida de S. Francisco Xavier,

Lote 7 Cv. C,

2900-616 Setúbal, Portugal.

Phone: +351 265 520 185

Fax: +351 265 520 186

Email: info@scitepress.org

EXTERNAL LINKS

PRIMORIS

INSTICC

SCITEVENTS

CROSSREF

**PROCEEDINGS
SUBMITTED FOR
INDEXATION BY:**

Thomson Reuters

dblp

Ei Compendex

SCOPUS

Semantic Scholar

Role of Allogenic NK Cells Treatment in the Early Phase of Apoptosis on Poorly Differentiated Retinoblastoma Cells Culture

Hendrian Dwikoloso Soebago¹, Fitria Kusumastuti¹, Parma Ratna Jaya¹, Susy Fatmariyanti¹, Endang Retnowati² and Ugroseno S Yudho Bintoro³

¹Department/Medical Staff of Ophthalmology, Faculty of Medicine, Airlangga University, Dr. Soetomo Hospital

²Department of Clinical Pathology, Faculty of Medicine, Airlangga University, Dr. Soetomo Hospital

³Hematology and Medical Oncology Division, Department/Medical Staff of Internal Medicine, Faculty of Medicine, Airlangga University, Dr. Soetomo Hospital
hendriands@yahoo.com

Keywords: Allogeneic, Apoptosis, Cancer Immunotherapy, NK Cells, Poorly Differentiated Retinoblastoma.

Abstract: **Introduction:** Retinoblastoma is a retinal malignant tumor because of mutations on chromosome 13q14. This is associated with apoptosis deregulation processes. Both autologous or allogeneic NK cells play their roles in both innate and adaptive immune systems. They can stimulate apoptosis through several mechanisms. **Methods:** This study determined the role of allogeneic NK cells in retinoblastoma cell apoptosis. Poorly differentiated retinoblastoma tissues were tested with SDS-PAGE. Allogeneic NK cells were isolated from peripheral blood of the healthy family members of patients. There were two culture groups consisting of retinoblastoma cells only as the control and retinoblastoma treated with allogeneic NK cells as the treated group. Each group consisted of 10 well plates. Examinations of the expression of Bcl-2, Caspase-3, and Apoptosis were performed by the flowcytometry method. **Result:** The SDS PAGE electrophoresis tests showed strong expression of proteins with molecular weights of 14, 19, 28, 35, 53 and 85 kDa. The percentage of early-phase apoptosis is higher than late phase. There were significant correlations ($\alpha < 0.05$) between Bcl-2, Caspase-3, and the ratio of both on cell apoptosis. **Conclusion:** In conclusion, allogeneic NK cells play a role in retinoblastoma cell apoptosis, especially in the early phase. It is expected to be one of the new strategies in cancer immunotherapy.

1 INTRODUCTION

Retinoblastoma is the most common eye cancer occurring in children. Retinoblastoma is a malignant tumor of the retina that is derived from primitive neuroectodermal tissues. This tumor is caused by mutations on chromosome 13q14. The most common occurrence is diagnosed when children are aged less than 3 years old. The incidence rate reaches approximately 4% of all malignancies in children. In the United States, retinoblastoma affects 1 in 18,000 children aged less than 5, and the prevalence reached 250 to 300 cases per year, resulting in a 1% death rate in all cases. The figures are higher in developing countries and continue to increase annually, making the success of retinoblastoma therapy a constant challenge for developing countries. There were 44 cases from 2010 to 2012 in Dr. Soetomo General Hospital in Surabaya (Bunin & Orjuela, 2007;

Chantada & Leal-Leal, 2007; Chintagumpala *et al.*, 2007; Soebago *et al.*, 2013; Stewart & Wild, 2014).

Retinoblastoma is a tumor type that is known to be genetically influenced. The mutation of the tumor suppressor gene (RB1) which is located at chromosome 13q14 is closely related to tumor formation (Leiderman *et al.*, 2007). Due to the genetic mutations occurring in retinoblastoma, proliferation will increase as apoptosis decreases. Homeostasis is achieved when the proliferation rate on the tissue is balanced with apoptosis. When apoptosis deregulation takes place, the number of dividing cells will be higher than the dying cells and in the end, a tumor is formed (Cheng *et al.*, 2013).

The apoptosis process runs through two pathways, which are (1) the extrinsic pathway (cytoplasm) through the activity of the Fas death receptor by activating Fas ligand (FasL) interaction and (2) the intrinsic pathway (mitochondrial) that stimulates

cytochrome-c release, which depends on Bcl-2 protein regulation (B cell lymphoma) as an anti-apoptotic protein and Bax as a pro-apoptotic protein. Singh *et al.* (2015) reported that the Bcl-2 is positively correlated to the growth of invasive retinoblastoma cells even though it is not correlated with Bax expression. However, Sitorus *et al.* (2009) reported that Bcl-2 overexpression would not prevent apoptosis, even though the independent Caspase-3 pathway is the apoptosis main pathway in retinoblastoma (Singh *et al.*, 2015; Sitorus *et al.*, 2009).

One of the cell types which can influence apoptosis occurrence is the Natural Killer (NK) cell. This cell constitutes lymphocytes playing a role in both the innate and adaptive immune systems. These cells can induce apoptosis through several mechanisms. In the last decade, the knowledge of NK cells, both the autologous and allogeneic, has developed and become one of the new strategies in cancer immunotherapy. Davis & Rizzieri (2015) mention that applying NK cells in a therapy on malignancy showed success and the studies in this field continue to grow rapidly. Eguizabal *et al.* (2014) state that NK cells play an important role in building immunity against cancer. Soebagio *et al.* (2015) report that autologous NK cells play an important role in RB cells' aggressiveness through the expression of EZH2, Ki-67, and RB cell apoptosis (Davis & Rizzieri, 2015; Eguizabal *et al.*, 2014; Soebagio *et al.*, 2015).

According to the facts, NK cell-based immunotherapy acts as one of the promising alternative therapies. So, investigations into the role of Bcl-2 protein as a regulator and Caspase-3 as the apoptosis executor on retinoblastoma cells treated with allogeneic NK cell will be the basic treatment especially as an alternative therapy for retinoblastoma cases.

2 METHODS

2.1 Research Subjects

Poorly differentiated retinoblastoma tumor tissues were collected from the enucleated patients of Dr. Soetomo General Hospital, Faculty of Medicine, University of Airlangga, Indonesia after obtaining informed consent. Fresh tumor tissues were tested using the SDS-PAGE method to determine the molecular weight of proteins. There were two groups consisting of retinoblastoma cells only as the control group and retinoblastoma versus allogeneic NK cells

as the treated group. Each group consisted of 10 well plates of cells culture. Expression of Bcl-2, Caspase-3, and Apoptosis was examined with the flowcytometry test.

2.2 SDS-PAGE

The SDS-PAGE test is useful to describe the molecular weight of tissue. Basically, proteins in the tissue are extracted in a buffer gel. The gel is then divided into two parts, the stacking gel and separating gel. The stacking gel consisted of 830 μ L of UGB (upper gel buffer), 534 μ L of acrylamide (T-acryl), 1950 μ L of ddH₂O, 40 μ L of ammonium persulfate (APS), and 4 μ L of tetramethylethylene diamine (TEMED). The separating gel consisted of 2600 μ L of LGB (lower gel buffer), 4000 μ L of acrylamide (T-acryl), 3400 μ L of ddH₂O, 140 μ L of ammonium persulfate (APS), and 14 μ L of tetramethylethylene diamine (TEMED). As much as 3 μ L of tumor tissue was prepared with μ L Tris-cl + 15 μ L of RSB (Reducing Sample Buffer) and then it was separated by electrophoresis gel using with 100 V voltage. The gel was then dyed with Coomassie Brilliant Blue R-250 so that the strands of molecules became visible. The molecular weight of each strand could be measured with a standard marker strand. The weight calculation of protein molecules was conducted based on Rf (Retardation factor) strand value of each sample. The migration distance of the polypeptide is a proportional inverse of the logarithm (log) value of the polypeptide molecule weight.

$$Rf = \frac{\text{Migration distance}}{\text{Gel length}} \quad (1)$$

2.3 Culture of Retinoblastoma Cells

The retinoblastoma tissue was cleaned 3 times with sterile PBS. It was then finely chopped in serum-free media of type-I collagenase. Finally, it was incubated for 30 minutes at 37°C. After that, medium plus serum was added and then the tissue was filtered. Next, the tissue was put in a centrifuge for 10 minutes at the speed of 1600 rpm. The supernatant was discarded and the pellets were resuspended with medium plus serum. After that, the cell culture was conducted on plates so it became confluent. Multiple passages were conducted until cell lines formed. And then, the confluent culture was divided into two groups. The first was the control

group which was not treated with allogeneic NK cells and the second was treated with NK cells with the ratio of 1:1 (Soebagio *et al.*, 2015).

2.4 Peripheral Blood Collection for Allogeneic NK Cells

The isolation of peripheral blood mononuclear cells (PBMC) was based on Boyum's method (1968) with some modifications. Isolation of mononuclear cells is often used to analyze the cellular immune responses by reacting antibody with mononuclear cell-surface antigens (Bøyum, 1968; Rantam, 2003).

The peripheral blood collection for NK cell isolation was performed on healthy biological family members of the retinoblastoma patients. The blood samples were stored in a K₃EDTA vacuum tube for PBMC isolation. In addition, the collection of NK cells can also be conducted through stem cell growth by adding GM-CSF growth factor to the NK cell culture process (Bøyum, 1968).

2.5 Examination of the Number of NK Cells (CD3⁻CD56⁺/CD16⁺)

The NK cell examination was performed using flowcytometry by means of BD FACS Calibur™ reagents of BD TriTEST™ CD3, Fluorescein isothiocyanate (FITC)/CD16⁺CD56 phycoerythrin (PE)/CD45 and peridinin-chlorophyll protein (PerCP), which are the pigment reagents of immunofluorescein to determine the number of NK cells (CD3⁻CD56⁺/CD16⁺) (Hu *et al.*, 2012).

The NK cells expressing CD3⁻, CD56⁺, and CD16⁺ would experience fluorescence in accordance with the area and then gating was performed on NK cells in the expression areas of CD56⁺/CD16⁺ and areas without the expression of CD3 (CD3⁻). The number of NK cells (cells/μL) was obtained by calculating the ratio of cell event expressing CD3⁻CD56⁺/CD16⁺ with fluorescent bead event; the number was previously discovered on the BD TruCOUNT tube (Hu *et al.*, 2012).

2.6 Examination of Cells Expressing Bcl-2, Caspase-3, and Apoptotic Cells

The examination of cells expressing Bcl-2, Caspase-3, and apoptotic cells was conducted using flowcytometry by means of BD FACS Calibur™ reagents of the primary antibody of Anti-Bcl-2 (100) FITC and FITC Active Caspase-3 Apoptosis, and FITC Annexin V Apoptosis Detection Kit II

Cat. 556 570, fixation reagents of BD Cytofix/Cytoperm™, permeabilization reagents of BD FACS Permeabilizing Solution 2, washer reagents of BD Perm/Wash™ buffer, and lysis reagents of BD FACS Lysing Solution. Staining with Annexin V is typically used in conjunction with a vital dye such as propidium iodide (PI) for identification of early and late apoptotic cells. Meanwhile, the number of cells that expressed Bcl-2 and Caspase-3 (cells/μL) was obtained by calculating the ratio of cell-event expressing antibodies with fluorescent bead events that was previously discovered on the BD TruCOUNT tube.

2.7 Statistical Analysis

All of the results were collected in specific data collection sheets, grouped and presented in tabular forms and diagrams, and then analyzed by using SPSS 15.0 so as to analyze the amount of Bcl-2 and Caspase-3 and apoptosis between two groups using a T-Test and the Mann-Whitney Test. The correlation analysis between the variables was conducted by Spearman's Rank Correlation Test. The p value was <0.05 which statistically indicated a significance.

3 RESULTS

3.1 SDS-PAGE

The expression of protein fractions from poorly differentiated retinoblastoma samples was suggested from some arising proteins with the molecular weight range of 14-85 kDa. The SDS-PAGE electrophoresis test showed a strong expression of the protein with the molecular weights of 14, 19, 26, 35, 53 and 85 kDa (Figure 1). Some protein fraction expressions are in accordance with the theory on the possibility of some proteins with molecular weights contained in the poorly differentiated retinoblastoma samples.



Figure 1: Profile of retinoblastoma protein with SDS-PAGE test. Protein expression in retinoblastoma samples including proteins with molecular weights of 14, 19, 26, 35, 40, 53, and 85 kDa.

The characterization overview of the culture of retinoblastoma cells showed cell growth between the control group and the treated group during the three

days' incubation period. The control group showed retinoblastoma cell growth shown by the grouping and confluent cell layers (Figure 2B).

In the treated group, there was less growth of retinoblastoma cells and the culture showed the formation of remnant sand of NK cells with the dead (apoptotic) cells surrounding between (Figure 2D).




Figure 2. Pictures of culture retinoblastoma cells. A) Control group: culture of retinoblastoma cells (pre-seeding); B) Control group: culture of retinoblastoma cells 3 days post-seeding; C) Treated group: culture of retinoblastoma cells with NK cells (pre-seeding); D) Treated group: culture of retinoblastoma cells with NK cells 3 days post-seeding; (Yellow arrow: NK cells; White arrow: overview of remnants of cell group undergoing apoptosis; Green arrow: intact retinoblastoma cells) (magnification: 400x).

4 DISCUSSIONS

4.1 Expression Percentage and Ratio of Bcl-2, Caspase-3, and Apoptosis on Retinoblastoma Cells Culture

The results showed the percentage of apoptotic cells, necrotic cells and living cells as well as the expression of antibodies Bcl-2 and Caspase-3 and the expression ratio of Bcl-2 and Caspase-3 in the control group and treated group. The expression was calculated based on the mean percentage (\pm SD) of positive cells expressing antibodies through the flowcytometry test.

The mean percentage of the cells that expressed Bcl-2 in the treated group was $0.10 \pm 0.10\%$, while in

the control group the expression did not occur ($0.10 \pm 0.10\%$). Caspase-3 expression in the treated group was $90.27 \pm 1.70\%$. It was higher compared with the control group ($70.03 \pm 3.63\%$). The percentage of apoptotic cells in the treated group was $27.69\% \pm 2.36$. It was lower compared with the control group ($37.72\% \pm 2.01$).




Figure 3: Percentage and ratio expression of Bcl-2, Caspase-3, and Apoptosis between the two groups.

4.2 The Effect of NK Cell Treatment on the Total Expression of Bcl-2, Caspase-3, and Cultured Retinoblastoma Cell Apoptosis

The effect of NK cell treatment on retinoblastoma cell apoptosis was calculated by the pigmentation of Annexin V/PI and flowcytometry analysis (Figure 4). The number of cells undergoing apoptosis was calculated based on the increasing concentration of apoptotic cells at early or late phases. Flowcytometry worksheets showed four areas to evaluate cell condition. Living cells can be observed on the lower left, early phase of apoptotic cells on the lower right, late phase on the upper right, and necrotic cells on the upper left side. As the marker of the apoptotic cells, it was strong positively with Annexin V and negative with PI.

The number of early apoptosis in the treated group was $20.17\% \pm 1.81$, slightly higher than the control group (19.53 ± 2.36). In contrast, the percentage of the late phase apoptosis in the treated group of 7.51 ± 6.01 is significantly lower than the control group (18.2 ± 1.09).

The ratio of Bcl-2 and Caspase-3 was calculated to describe the difference between the expression of each parameter. There was no difference between the two groups (treated group was $0.11 \pm 0.11\%$ versus control group of $0.00 \pm 0.00\%$).

The different test result analysis of all variables using the Mann-Whitney test and Independent T-test between the treated group and control showed a significant result among them. In addition, Spearman's Rank correlation test between variables and apoptosis showed a significant result

Table 1: Different Test Result Analysis Control and Treatment in Phase Apoptosis.

Apoptosis Phases	Groups	Mean	P
Early	Rb	19.53±2.36	0.505
	Rb +NK	20.17±1.81	
Late	Rb	18.2±1.09	0.000 *
	Rb+NK	7.51±6.01	

Note: * significant at $\alpha = 0:05$

Table 2: Difference Test Results on all variables between the two groups.

Variables	Difference Test Results
Bcl-2	0.002*
Caspase-3	0.000*
Cell apoptosis	0.002*
The ratio of Bcl-2/Caspase-3	0.000*

Note: * significant at $\alpha = 0.05$

Table 3: Correlation Test Result between Variables and Apoptosis.

Variables	Apoptosis	
	r_s	p
Bcl-2	-0.605	0.005 *
Caspase-3	-0.657	0.002 *
The ratio of Bcl-2/ Caspase-3	-0.609	0.004 *

Note: * significant at $\alpha = 0.05$




Figure 4: Overview of flow cytometry on expression of Bcl-2, Caspase-3 and Apoptosis between the two groups.

A typical figure of retinoblastoma cell given NK cells is seen as the formation of remnant cells and NK cells surrounding cells that have an apoptotic process. Soebagjo *et al.* (2015) reported that NK cells have the ability to induce retinoblastoma cells thus inhibiting tumor cell growth and finally cause cell death. Therefore, NK cells are found surrounding the dead cells. In contrast, retinoblastoma cells grew normally in the control group (Soebagjo *et al.*, 2015).

The SDS-PAGE electrophoresis test on poorly differentiated retinoblastoma samples shows strong expression of proteins with the molecular weights of 14, 19, 26, 35, 53 and 85 kDa. The expressions show that there are proteins corresponding to Bcl-2 and Caspase-3 that play a role in retinoblastoma formation.

Raghupathi *et al.* (2003) measured the molecular weight of Bcl-2 (26 kDa) and Bax (21 kDa) by using immunoblot on HeLa cortex and hippocampus cells. In the cardiomyocytes, two types of Bcl-2 expressions with different molecular weights were obtained (26 and 32 kDa) (Raghupathi *et al.*, 2003; Cook *et al.*, 1999).

Caspase-3 with a molecular weight of 35 kDa is a form of fraction activated by T cells and is found in the cell cytoplasm. In addition, 35 kDa molecular weight is similar to the molecular weight of pro-caspase-7 and pro-caspase-9 (Pause *et al.*, 2008; Dean *et al.*, 2002; Kadirvel *et al.*, 2010).

The process of cell death or apoptosis is commonly used as the selective parameter of a proposed anti-cancer source. Apoptosis induction was indicated by an increase in the percentage of cells undergoing apoptosis. The average of early apoptosis in the treated group was higher compared with the control group, while late apoptosis in the treated group was lower compared with the control group. So, NK cells play a role in early phase apoptosis. Cheng *et al.* (2013) mentioned the role of NK cells especially in the early stage of a good immune system both on innate and adaptive immune systems. Poggi *et al.* (2005) mentioned that the apoptosis trigger induced by NK cells will bind phosphatidylserine exposed on the surface of apoptotic cells and is marked by Annexin V in the early apoptosis phase. In contrast, in late-phase apoptosis, the death rate of both retinoblastoma cells and NK cells cannot be distinguished. Poggi *et al.* (2005) mentioned that the role of NK cells is effective on 24-hour incubation and it will reach the peak at 48 hours of incubation together inside tumor cells (coculture) (Cheng *et al.*, 2013; Poggi *et al.*, 2005).

The flowcytometry analysis showed that Bcl-2 is relatively insignificant in the apoptosis process. However, caspase-3 showed significant cell expression but apoptotic cells were lower compared with the control group. The correlation of Bcl-2/Caspase-3 showed that Bcl-2 yielded less expression when compared with Caspase-3. Nevertheless, Caspase-3 turned out to be inversely proportional to the occurring cell apoptosis. This shows the apoptotic signaling process that occurred in allogeneic NK cells in retinoblastoma cells. In general, the process of NK cell apoptosis goes through two pathways, which are (1) the intrinsic pathway which goes through the NKG2D ligand or (2) the extrinsic pathway which goes through the TNF ligand/FasL (Dranoff, 2004; Krzewski & Coligan, 2012).

The NK cells will induce tumor cells through Killer Activation Receptors (KARs), i.e. the NKG2D ligand when the KIR inhibitory signal is hampered because the tumor cells are inhibited by the expression of MHC class I which cause cytotoxic granules (perforin and granzyme) to not be excreted. The cytotoxic granules will be excreted if NK cell receptors, FcγRIII (CD16) bind IgG tumor cells. Excreted perforin will open pores as the entrance of granzyme protein to mediate apoptosis (Dranoff, 2014; Abe *et al.*, 2008).

There are 5 types of granzyme known to exist in humans. These are granzyme A, B, H, K, and M. The difference of the types lies in the structure of the serine protease substrate (Grossman *et al.*, 2004). Hochegger *et al.* (2004) mention that the class of polymorphonuclear cells (PMN) excrete perforin and 2 types of granzyme which are granzyme A and B which contribute to the process of cytotoxic cells. Granzyme A induces cell death through the DNA characteristic of single chain cells and do not activate caspase cascade (Bots & Medema, 2006).

In NK cells, granzyme B plays a role. Granzyme B is capable of directly inducing apoptosis (bypass) via Caspase-3. This is consistent with the results of research and can occur because Bcl-2 is not sufficient to affect the apoptosis process. Overexpression of Bcl-2 in retinoblastoma itself is noted to not necessarily hinder the apoptosis process. The reason why Bcl-2 is not functional is supported by Sutton *et al.* (1997) who state that constraints on Bcl-2 will have an effect on increasing the ratio of the target and increase granzyme B. Granzyme B itself activates Bid pro-apoptotic proteins, along with Bax, which translocates to the mitochondria and affects the membrane permeability to release cytochrome-c after forming apoptosome by the Caspase cascade (Sitorus *et al.*, 2009; Bots & Medema, 2006; Sutton *et al.*, 1997; Pinkoski *et al.*, 2001).

In addition, Pinkoski *et al.* (2001) mentioned that in a direct path from granzyme B to caspase-3, apoptosis via the mitochondria can also occur via Smac/Diablo proteins that mediate apoptosis of granzyme B contributing to cell destruction by binding with XIAP. In addition, the release of Smac/Diablo in mitochondria will inhibit the Bcl-2 (Pinkoski *et al.*, 2001; Verhagen *et al.*, 2000; Adrain *et al.*, 2001).




Figure 5: A schematic possible pathway of the induction of NK Cells on apoptosis in Retinoblastoma Cells.

The test results indicate that the ratio of the expression of Caspase-3 increased, reversely proportional to cell apoptosis. Sitorus et al. (2009) stated that there is a possibility that apoptosis occurs through Caspase-3 -dependent or independent pathways. Apoptotic signals that have come to the Caspase-3 effector are possibly hindered by DNA enzyme of repair poly(ADP-ribose) polymerase-1 (PARP-1). Active Caspase-3 protein will divide various substrates, including PARP-1 which is the mitotic apparatus core, lamina nucleus, and also actin and endonucleases. PARP-1 plays an important role in DNA repair processes by participating in the initiation of base excision repair (BER), nucleotide excision repair, and base repair with single-stranded DNA ligase III mediation, as well as controlling 60-70% of cell metabolism, cell cycle and cell transcription. Cell death signals of Caspase-3 towards PARP-1 cleave and the PARP-1 fragments will activate cell apoptosis. The cleavage process of PARP-1 is conducted by cytokine of Matrix Metalloproteinase-2 (MMP-2). However, the cleavage process by MMP-2 is inhibited by Enhancer Zeste Homolog-2 (EZH2) that represses micro RNA 21 promoters and inhibits MMP-2 through TIMP-2 inhibitors. Barriers on PARP-1 will result in the increased expression of Caspase-3 and in opposite proportion to cell apoptosis (Lazebnik et al., 1994; Nicolescu et al., 2009).

5 CONCLUSIONS

It is concluded that the provision of allogeneic NK cells play a significant role in the apoptosis process of poorly differentiated retinoblastoma cells, especially at the early phase through the intrinsic pathway.

REFERENCES

- Abe, H., Akiyama S. & Okamoto, M., 2008, 'Clinical Cancer Immunotherapy: Molecular Targeting Immunotherapy', *J Integ Med*, 1(1), 38-46.
- Adrain, C, Creagh E.M. & Martin, S.J., 2001, 'Apoptosis-Associated Release of Smac/DIABLO From Mitochondria Requires Active Caspases and Is Blocked by Bcl-2', *EMBOJ*, 20(23), 6627-6636.
- Bots, M. & Medema, J.P., 2006, 'Granzymes at A Glance', *Journal of Cell Science*, 119, 5011-5014.
- Bøyum, A., 1968, 'Isolation of mononuclear cells and granulocytes from human blood' (Paper IV), *Scand. J. Clin. Lab. Invest*, 21 Suppl., 97, 77-89.
- Bunin, G.R. & Orjuela, M., 2007, *Clinical Ophthalmic Oncology*, Philadelphia: Saunders Elsevier, 410-414.
- Chantada, G.L. & Leal-Leal, 2007, *Geographic and Environmental Factors in Singh (Ed.) Clinical Ophthalmic Oncology*, Philadelphia: Saunders Elsevier, 2007. pp.417-420.
- Cheng, M., Chen, Y., Xiao, W., Sun, R. & Tian, Z., 2013, 'NK Cell-Based Immunotherapy for Malignant Diseases', *Cellular & Molecular Immunology*, 10, 230-252.
- Chintagumpala, M., Chevez-Barrios, P., Paysse, E.A, Plon, S.E. & Hurwitz, R., 2007, 'Retinoblastoma: Review of Current Management', *The Oncologist*, 10:12, 1237-1246.
- Cook, S.A., Sugden, P.H. & Clerk, A., 1999, 'Regulation of Bcl-2 Family Proteins During Development and in Response To Oxidative Stress in Cardiac Myocytes: Association with Changes in Mitochondrial Membrane Potential', *Circulation Research*, 85, 940-949.
- Davis, C.T. & Rizzieri, D., 2015, 'Immunotherapeutic Applications of NK Cells', *Pharmaceuticals*, 8, 250-256.
- Dean, A.L., Wu, Y., Huang, Z., Matsushita, K., Plesnila, N., Augustinack, J.C., et al., 2002, 'Caspase Activation and Neuroprotection in Caspase-3-Deficient Mice After In Vivo Cerebral Ischemia and In Vitro Oxygen Glucose Deprivation', *PNAS*, 23(99), 15188-15193.
- Dranoff, G., 2004, 'Cytokines in Cancer Pathogenesis and Cancer Therapy', *Nat Rev Cancer*, 4(1), 11-22.
- Eguizabal, C., Zenarruzabeitia, O., Monge, J., Santos, S., Vesga, M.A., Maruri, N., et al., 2014, 'Natural Killer Cells for Cancer Immunotherapy: Pluripotent Stem Cells-Derived NK Cells As An Immunotherapeutic Perspective', *Frontiers in Immunology*, 5, 1-10.
- Grossman, W.J., Verbsky, J.W., Tollefsen, B.L., Kemper, C., Atkinson, J.P. & Ley, T.J., 2004, 'Differential Expression of Granzymes A and B in Human Cytotoxic Lymphocyte Subsets and T Regulatory Cells', *Blood*, 104 (9), 2840-2848.
- Hochegger, K., Eller, P. & Rosenkranz, A.R., 2004, 'Granzyme A: An Additional Weapon of Human Polymorphonuclear Neutrophils (PMNs) in Innate Immunity?', *Blood*, 103 (3), 1176.
- Hu, P., Hegde, M. & Lennon, P.A., 2012, *Modern Clinical Molecular Techniques*, New York, Springer.

- Kadirvel, R., Ding, Y.H., Dai, D., Lewis, D.A. & Kallmes, D.F., 2010, 'Intrinsic Pathway Mediated Apoptosis In Elastase-Induced Aneurysms in Rabbits', *AJNR Am J Neuroradiol*, 31, 165-169.
- Krzewki, K. & Coligan, J.E., 2012, 'Human NK Cell Lytic Granules and Regulation of Their Exocytosis, Review Article', *Front Immunol.*, 3(335), 1-16.
- Lazebnik, Y.A., Kaufmann, S.H., Desnoyers, S., Poirier, G.G. & Earnshaw, W.C., 1994, 'Cleavage of Poly (ADP-Ribose) Polymerase by a Proteinase with Properties like ICE', *Nature*, 371, 346-347.
- Leiderman, Y.L., et al., 2007, 'Molecular Genetics of RB1- The Retinoblastoma Gene', *Seminars in Ophthalmology*, 22, 247-254.
- Nicolescu, A.C., Holt, A., Kandasamy, A.D., Pacher, P. & Schulz, R., 2009, 'Inhibition of Matrix Metalloproteinase-2 by PARP Inhibitors', *Biochem Biophys Res Commun.*, 387, 646-650.
- Paulsen, M., Ussat, S., Jakob, M., Scherer, G., Lepenies, I. & Schutze, S., et al., 2008, 'Interaction with XIAP Prevents Full Caspase-3/-7 Activation in Proliferating Human T Lymphocytes', *Eur J Immunol.*, 38, 1979-1987.
- Pinkoski, M.J., Waterhouse, N.J., Heibein, J.A., Wolf, B.B., Kuwana, T., Goldstein, J.C., Newmeyer, D.D., Bleackley, R.C. & Green, D.R., 2001, 'Granzyme B-mediated Apoptosis Proceeds Predominantly through a Bcl-2-Inhibitible Mitochondrial Pathway', *J Biol Chem.* 13, 276 (15), 12060-7.
- Poggi, A., Massaro, A., Negrini, S., Contini, P. & Zocchi, M.R., 2005, 'Tumor-Induced Apoptosis of Human IL-2-Activated NK Cells: Role of Natural Cytotoxicity Receptors', *J Immunol.*, 174, 2653-2660.
- Raghupathi, R., Strauss, K.I., Zhang, K., Krajewski, S., Reed, J.C. & McIntosh, T.K., 2003, 'Temporal Alterations in Cellular Bax:Bcl-2 Ratio Following Traumatic Brain Injury in The Rat', *J Neurotrauma*, 20(5), 421-435.
- Rantam, F.A., 2003, *Metode Immunologi*, Airlangga University Press, Surabaya.
- Singh, L., Pushker, N., Saini, N., Sen, S., Sharma, A., Bakhshi, S., et al., 2015, 'Expression of Pro-Apoptotic Bax and Anti-Apoptotic Bcl-2 Proteins in Human Retinoblastoma', *Clin Exp Ophthalmol.*, 43(3), 259-67.
- Sitorus, R.S., Gumay, S. & Van Der Valk, P., 2009, 'The Apoptotic Paradox in Retinoblastoma', *Ann N Y Acad Sci.*, 1171:77-86.
- Soebagio, H.D., Prastyani, R., Sujuti, H., Lyrawati, D. & Sumitro, S.B., 2013, 'Profile of Retinoblastoma in East Java, Indonesia', *World Journal of Medicine and Medical Science Research*, 3, 051-056.
- Soebagio, H.D., Fatmariyanti, S. & Lutfi, D., 2015, 'Effectiveness of Natural Killer (NK) Cells in Peripheral Blood Stem Cell Towards Expression of EZH2, Ki-67 and Apoptosis in Retinoblastoma (RB) Cells Culture', *Medicine Science*, 04, 1-17.
- Stewart, B.W. & Wild, C.P., 2014, 'Global Epidemiology of Cancer', Jones & Bartlett Learning LLC, 3, 45.
- Sutton, V.R., Vaux, D.L. & Trapani, J.A., 1997, 'Bcl-2 Prevents Apoptosis Induced by Perforin and Granzyme B, But Not That Mediated by Whole Cytotoxic Lymphocytes', *J Immunol.*, 158, 5783-5790.
- Verhagen, A.M., Ekert, P.G., Pakusch, M., Silke, J., Connolly, L.M., Reid, G.E., et al., 2000, 'Identification of DIABLO, a Mammalian Protein That Promotes Apoptosis by Binding to and Antagonizing IAP Proteins', *Cell*, 102(1), 43-5.