DAFTAR PUSTAKA

Ahmad Nisar, Fazal H, Abbasi1 BH, Farooq S, Ali1 M, Khan MA. (2012). Biological role of Piper nigrum L. (Black pepper): A review. Asian Pacific Journal of Tropical Biomedicine.

Arrington, L. (1972). Introductory Laboratory Animal. The Breeding, Care, and Management of Experimental Animal Science. New York: The Interstate Printers and Publishing, Inc.

Bai YF, Xu H. (2000). Protective action of piperine against experimental gastric ulcer. ActaPharmacolSin 21 : 357–359.

Bang JS, Oh DH, Choi HM, Sur BJ, Lim SJ, Kim JY, Yang HI, Yoo MC, Hahm DH, Kim KS. (2009). Anti-inflammatory and antiarthritic effects of piperine in human interleukin 1β-stimulated fibroblast-likesynoviocytes and in rat arthritis models. Arthritis Res Ther,11(2) : R49.

Bhardwaj RK, Glaeser H, Becquemont L, Klotz U, Gupta SK, Fromm MF. (2007) Piperine, a major constituent of black pepper, inhibits human P-glycoprotein and CYP3A4. J PharmacolExpTherapeut,302: 645–650.

D'Hooge R, Pei YQ, Raes A, Lebrun P, Bogaert PP, De Deyn PP. (1996). Anticonvulsant activity of piperine on seizures by excitatory amino acid receptor agonists. Arzneimittelforschung 46: 557– 560.

Darling, L.M. (2002). Spices as Aphrodisiacs. Dipetik february 4, 2015. Website: http://unitproj.library.ucla.edu/biomed/spice/index.cfm?spicefilename=aphr odisiacs.txt&itemsuppress=yes&displayswitch=0.

Daware MB, Mujumdar AM, Ghaskadbi S. (2000). Reproductive toxicity of piperine in Swiss albino mice. Planta Med,66: 231-236.

Ditjen POM, (1983). Tanaman Obat Keluarga (TOGA), Jakarta : Departemen Kesehatan RI

Ditjen POM. (2000). Parameter Standar Umum Ekstrak Tumbuhan Obat. Jakarta: Departemen Kesehatan RI.

Drewes, S.E., Horn, M.M., Munro, O.Q., Dhlamini, J.T.B., Meyer, J.J.M., Rakuambo, N.C., (2002). Pyrano-isoflavones with erectile dysfunction activity from Eriosema kraussianum. Phytochemistry 59, 739–747.

Gatkal, S., Punde, A., Balap, A., & Chaundari, P. (2012, October-january). Safety of herbal medicine. *International Journal Of Pharmaceutical and Chemical Sciences*, 1626.

Gibbs, R.D. History of Chemical Taxonomy, In Chemical Plant Taxonomy, Swain, T., ed. (1993), Academic Press: London. pp. 41-88.

Gupta SK, Bansal P, Bhardwaj RK, Velpandian T. (2000). Comparative anti-nociceptive, anti-inflammatory and toxicity profile of nimesulides vs nimesulide and piperine combination. Pharmacol Res 41: 657–662.

Lawrence, B.M. (1981) Major tropical spices—pepper (Piper nigrum L.). Essential oils 1979– 80; pp. 141–228, Allured Pub. Cor., USA

Lim, T. K. (2012). Piper cubeba. *Edible Medical And Non Medical Plants : Volume 4*.

Moeloek, N., (2006). Uji klinik ekstrak cabe jawa (Piper retrofractum Vahl) sebagai fitofarmaka androgenik pada pria hipogonad, Laporan akhir penelitian, PROM-Badan POM RI, Jakarta.

Nakatani, N., Inatani, R., Ohta, H., Nishioka, A., (1986), Chemical constituents of peppers (Piper spp.) and application to food preservation: Naturally occurring antioxidative compounds. Environmental Health Perspectives, 67: 135-42.

Parmar VS, Jain SC, Bisht KS, Jain R, Taneja P, Jha A, Tyagi OD, Prasad AK, Wengel J, Olsen CE, Boll PM (1997). Phytochemistry of the genus Piper. Phyto- chemistry 46:597–673

Patel DK, Kumar R, Prasad SK, Hemalatha S. (2011). Pharmacologically screened aphrodisiac plant-A review of current scientific literature. *Asian Pacific Journal of Tropical Biomedicine*.

Patil, S. A., M, S., & Patil, S. B. (2013). Aphrodisiac and phytochemical studies of Cocculus hirsutus extracts in albino rats . *Asian Pacific Journal of Reproduction*, 23-29.

Ravindran, C.S.Narayanan, & Roland, H. (2006). *Medicinal and Aromatic Plants—Industrial Profiles - Black Pepper.* Amsterdam: harwood academic.

Reynertson, K. A. (2007). Phytochemical Analysis of Bioactive Constituents From EdiblemMyrtaceae Fruits. *A dissertation submitted to the Graduate Faculty in Biology The City University of New York* .

Sandroni P. (2001). Aphrodisiacs past and present: a historical review. Clin Auton Res; 11: 303-307.

Sayre, L. E. (2013). *A Manual Of Organic Materia Medica and Pharmacognosy.*Philadelphia: Forgotten books.

Selven-diran K, Singh JP, Krishnan KB, Saktisekaran D. (2003). Cytoprotective effect of piperine against benzole [a]pyrene induced lung cancer with reference to lipid peroxidation and antioxidant system in Swiss albino mice. Fitoterapia74:109– 115.

Singh Amandeep R, Sarabjeet Singh, G. Jeyabalan, Ashraf Ali, Alok Semwal. (2012). Sexual dysfunction: An overview and medicinal plant used for treatment of sexual dysfunction. Crit Rev Pharm Sci; 1(2): 9-24.

Smith, J.B dan S. Mangkoewidjojo. 1988. Pemeliharaan, Pembiakan dan Penggunaan Hewan Percobaan di Daerah Tropis. Jakarta : UI Press.

Suwanphakdee C, Hodkinson T R, chantaranothai P. (2014). New records of Peperomia (Piperaceae) in Thailand. THAI FOR. BULL. (BOT.) 42: 61–67.

Tajuddin, Ahmad S, Latif A, Qasmi IA, Amin KMY. (2005). An experimental study of sexual function improving effect of

Myristica fragrans Houtt. (Nutmeg). BMC Complement Altern Med ; 5: 16.

Taryono R A, (2004). Cabe Jawa. Jakarta : Penebar Swadaya.

Thakur M, Bhargava S, Dixi VK. (2009) . Effect of Asparagus racemosus on sexual dysfunction in hyperglycemic male rats. Pharm Biol; 47: 390-395.

Tiwari P, Singh D. (2008). Antitrichomonas activity of sapindussaponins, a candidate for development as microbicidalcontraceptive. J. AntimicrobChemother,62: 526-534.

Usia, Tepy. 2012. Mengenal Cabe Jawa (Piper retrofractum Vahl.) sebagai Tanaman Obat Untuk Stamina Pria. InfoPOM, Badan Pengawas Obat dan Makanan Republik Indonesia, Jakarta.

Wahjoedi B, Pudjiastuti, Adjirni, Nuratmi B, Astuti Y. (2004). Efek androgenik ekstrak etanol cabe jawa (Piper retrofractum Vahl.) pada anak ayam. JBA Indonesia.;3:201-204.

Yakubu MT, Akanji MA, Oladiji A. (2007). Male sexual dysfunction and methods used in assessing medicinal plants with aphrodisiac potentials. Phcog Rev 2007; 11: 49-56.

Zamblé A, Sahpaz S, Brunet C, Bailleul F. (2008) Effects of Microdesmis keayana roots on sexual behavior of male rats. Phytomed; 15: 625-629.

Zanoli P, Zavatti M, Montanari C, Baraldi M (2009) Influence of *Eurycoma longifolia* on the copulary activity of sexually sluggish and impotence male rats.