

Journal of Physics: Conference Series


PAPER • OPEN ACCESS

Anti-inflammatory effect of ethyl acetate fraction of galing plant extract (*Cayratia trifolia*) on male wistar rats induced by carrageenan

D Santoso¹, I K Sudiana², A S Rahayu² and M Yunus³

Published under licence by IOP Publishing Ltd

Journal of Physics: Conference Series, Volume 1146, conference_1



References ▾

+ Article information

Abstract

Inflammation (inflammation) is a local reaction from living tissue or cells to an excitatory or injury. This study aims to determine the anti-inflammatory activity of ethyl acetate fraction of gallin plant extract (*Cayratia trifolia*) on one of the inflammatory parameters, namely swelling in the legs of rat with 1% carrageenan induction. In addition, this study also aims to determine the effective dose of ethyl acetate fraction of gallin plant extract (*Cayratia trifolia*) as anti-inflammatory. This study used 25 male wistar rats divided into 5 groups. Group 1 (negative control) Na CMIC 0.5%, group 2 (positive control) diclofenac sodium 0.0065 mg/gBW rats, and group 3, 4 and 5 suspension ethyl acetate fractions of gallin plant extract 0.0065; 0.013 and 0.026 mg/gBW rats. Inflammation in rats by inducing 1% carrageenan as much as 0.10 ml. The volume of edema every hour is known from the difference in foot volume at certain hours with normal foot volume. The AUC value of the edema volume was calculated by trapezoid method every one hour and calculated for anti-inflammatory power (IP). The results showed that ethyl acetate fraction of gallin plant extract (*Cayratia trifolia*) could reduce the volume of edema in the legs of male white wistar rats which was induced by caragenine with an effective dose of 0.0065 mg/gBW.

Export citation and abstract

BibTeX

RIS



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



References

- [1] Tjokronegoro A and Baziati A 1993 *Semiloka di Fakultas Kedokteran Universitas Indonesia* (Jakarta, 6-7 Juli 1992) *Etk Penelitian Obat Tradisional*
- [2] Chaitopadhyay Deprasad, Arunachalam, Ghosh L, Rajendran K, Mandal A B and Bhatta C S K 2005 *J. Pharm Pharmaceut Sci* **8** 558-564 www.cspscanada.org
- [3] Robbins SL and Kumar V 1995 *Teijemahan Staf Pengajar Laboratorium Patologi Fakultas Kedokteran UMAIR*, cetakan 2 (Jakarta: EGC) Buku Ajar Patologi I
- [4] Wirakusumah ES 2006 *Buah dan sayur untuk terapi* (Jakarta: Penebar Swadaya)
- [5] Nijveldt R J E, van Noord DEC, van Hoorn PG, Boelens K, van Norren PAM and van Leeuwen 2001 Flavonoids: a review of probable mechanisms of action and potential applications *American Journal of Clinical and Nutrition* **74** 418-425 [Crossref](#)
- [6] Nutritional Therapeutics 2003 NT Factor: Phosphoglycolipids-High Energy Potential. www.propas.com/FAO/soy_high_energy.htm [2 Desember 2005]
- [7] Katzung BG 2002 *Farmakologi Dasar dan Klinik* (Jakarta: Salemba Medika)

Export references:

BibTeX

RIS

37 Total downloads

Turn on MathJax

Share this article



brightrecruits.com jobs

University of Siegen – Institute of High Frequency and Quantum Electronics offers Now PhD Positions University of Siegen

KSETA Doctoral Fellow Karlsruhe Institute of Technology (KIT)

Teacher of Physics (possibility of 2 V/0) Caterham School

More jobs Post a job

Journal of Physics: Conference Series

Table of contents

Volume 1146

2018

◀ Previous issue Next issue ▶

The 5th ICAMBBE (International Conference on Advance Molecular Bioscience and Biomedical Engineering) 2018
3–4 September 2018, Brawijaya University, Malang, Indonesia

[View all abstracts](#)

Accepted papers received: 20 November 2018

Published online: 7 January 2019

Preface

OPEN ACCESS

The 5th ICAMBBE (International Conference on Advance Molecular Bioscience & Biomedical Engineering) 2018

011001

+ View abstract

[View article](#)



OPEN ACCESS

Peer review statement

011002

+ View abstract

[View article](#)



Papers

OPEN ACCESS

FAS and FASL polymorphisms and susceptibility to hepatitis B virus infection in Javanese individuals

012001

A A Prasetyo and D A Agustin

+ View abstract

[View article](#)



OPEN ACCESS

Effect of torbangun (*Caleus amboinicus* Loui) on blood pressure in women with hypercholesterolemia

012002

T Suryowati and M Gutom

+ View abstract

[View article](#)



OPEN ACCESS

The potential role of betatrophin in hepatocellular carcinoma and metabolic syndrome

012003

H Susanto, A Autennam, C H Wang, D K Wuragiti, A Y Handaya, M P Perther and S A Ruttatin Nisa

+ View abstract

[View article](#)



OPEN ACCESS

Interaction of acetaminophen and caffeine towards cyclooxygenase-2 (COX-2) in inhibition of prostaglandin (PGH₂) synthesis

012004

G C Kranamurti and F Fatchiyah

+ View abstract

[View article](#)



OPEN ACCESS

Effectivity of tomato (*Solanum lycopersicum*) and zinc combination to sperms of male white rats (*Rattus norvegicus*) exposed to monosodium glutamate

012005

I H Wirandoko, C Apriyani and D R Acriyanto

+ View abstract

[View article](#)



OPEN ACCESS

The morphological endogenous development of *Eimema necatrix* at primary and secondary infected chickens and its histopathological effect

012006

M Yunia, E Suprihati and A Wijaya

+ View abstract

[View article](#)



OPEN ACCESS

Effectiveness of betel leaves (*piper betel* L.) and breadfruit leaves (*Artocarpus altilis*) in reducing uric acid levels in hyperuricemic male white rats (*Rattus norvegicus*)

012007

F Farizal, D Nauphar, T M Pratamaewati and R S Brajawikaspa

+ View abstract

[View article](#)



OPEN ACCESS

The effects of specific inhibitor Lp-PLA2 on vasa vasorum angiogenesis through inhibition vascular endothelial growth factor expression: study *in vivo* using type 2 diabetes mellitus

012008

T A Winastuti, Hafsatuzzamzudah, F N Aini, K W Antia, S Kushardianti, S P W Aswiani and T Heriansyah

+ View abstract

[View article](#)



OPEN ACCESS

Antihyperglycemic effectiveness comparison of ethanol extract of soursop leaf (*Annona muricata* L.) against acarbose in streptozotocin-induced diabetic white rats

012009

R R Setiadi, A F M Z Zein and D Nauphar

+ View abstract

[View article](#)



OPEN ACCESS

The comparison on antihyperglycemic activity between *Acacia sinuata* leaf / *Mangifera indica* leaf, *Asplenium nidus* and

012010

JOURNAL LINKS

[Journal home](#)

[Information for organizers](#)

[Information for authors](#)

[Search for published proceedings](#)

[Contact us](#)

[Reprint services from Curran Associates](#)



The comparison of ameliorating activity between glibenclamide and metformin in streptozotocin-induced diabetic rats

Q Adyur, A F M Z Zein and V Melianawaty

[+ View abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

The effect of derapidib administration to inflammation marker in early development of atherosclerosis: *In vivo* study for dyslipidemia model

T Herianayah, F N Aim, Nafisauzzamrudin, P N B Saka, D Sargowo and T A Wihestuti

[+ View abstract](#) [View article](#) [PDF](#)

012011

OPEN ACCESS

Comparison of sport massage and combination of cold water immersion with sport massage on decrease of blood lactic acid level

N F Romadhona, G M Sari and D N Utomo

[+ View abstract](#) [View article](#) [PDF](#)

012012

OPEN ACCESS

Direct effect of short wave diathermy therapy and exercise with quadriceps bench against knee osteoarthritis patients with functional capabilities

D A Nugraha, R D Haryadi and S Cahyono

[+ View abstract](#) [View article](#) [PDF](#)

012013

OPEN ACCESS

Diabetes dance of persadia 1 effect on blood IL-6 level

W N Pratiwi, B Purwanto and Abdurachman

[+ View abstract](#) [View article](#) [PDF](#)

012014

OPEN ACCESS

Improved hypoglycemic effect of anthocyanin extract combination from red rice and black soybean

W M Rahayu, M Astuti and Y Marsono

[+ View abstract](#) [View article](#) [PDF](#)

012015

OPEN ACCESS

The activity of casein derived from goat milk yogurt as an antioxidant on histopathology of rat's liver exposure by 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD)

C Mahdi, A E P Haskito, M C Padaga and A Rosdiana

[+ View abstract](#) [View article](#) [PDF](#)

012016

OPEN ACCESS

The aquatic environmental quality of koi fish (*Cyprinus carpio*) pond infected by *Myxobolus* sp. based on the biological status of the phytoplankton

U Yanuar, N R Caesar, F Setiawan, M Sumsanto, M Musa and D K Wuragi

[+ View abstract](#) [View article](#) [PDF](#)

012017

OPEN ACCESS

Synthesis, docking molecule study and antibacterial activity of N'-(4-Fluorobenzylidene)-4-hydroxybenzohydrazide and N'-(4-Fluorobenzylidene)-4-hydroxybenzohydrazide

S Suzana, M I Sulistyowaty, Israeni and T Budiani

[+ View abstract](#) [View article](#) [PDF](#)

012018

OPEN ACCESS

Progress report on the rapid test kit development for early detection of systemic lupus erythematosus in Indonesia

K Handono, KN Bentia, FU Hakimah and ES Deei

[+ View abstract](#) [View article](#) [PDF](#)

012019

OPEN ACCESS

Hypoglycemic activity of hydroethanolic root extracts of *Ruellia tuberosa* L in diabetic rats

A Safrin, Sutrisno, A Rosdiana and C A Evindasari

[+ View abstract](#) [View article](#) [PDF](#)

012020

OPEN ACCESS

Anti-inflammatory effect of ethyl acetate fraction of galang plant extract (*Caryota trichilia*) on male wistar rats induced by carrageenan

D Santoso, I K Sudiana, A S Rahayu and M Yunus

[+ View abstract](#) [View article](#) [PDF](#)

012021 ✓

OPEN ACCESS

Isolation of lactic acid bacteria from cattle rumen as starter in silage manufacture

N D F K Foeh, N A Ndaong, R E M Mala, E Beribe, P L Pau, A Detha and F U Datta

[+ View abstract](#) [View article](#) [PDF](#)

012022

OPEN ACCESS

Individual mutations in Indonesian local etttawah goats based on the GDF9 gene

M Mudawahan, I D Ratnawingtyas, M Z Faedi and G Ciptadi

[+ View abstract](#) [View article](#) [PDF](#)

012023

OPEN ACCESS

Reproductive performance and fertility index of swamp buffalo (*Bubalus bubalis*) in ngawi regency, east java

A Budianto, G Ciptadi, L Hakim and A I Putri

[+ View abstract](#) [View article](#) [PDF](#)

012024

OPEN ACCESS

Cutaneous wound healing activity of herbal ointment containing the leaf extract of *Acalypha indica* L on mice (*Mus musculus*)

M Laut, N A Ndaong and T Utami

[+ View abstract](#) [View article](#) [PDF](#)

012025

OPEN ACCESS

Malondialdehyde (MDA) level and protein profile of serum after calving towards the provision of selenium-vitamin E™ on dairy cow fision holstein (FH)

F Darmawati, M F Sembiring, E E C Dikta, F Darmawati and C Chandra

012026

OPEN ACCESS

Antimicrobial activity of whey mare's milk against *Salmonella enteritidis*

012027

A Detha, A Saputra and A Ola

OPEN ACCESS

Cryoprotector combination ethylene glycol and propanediol on mice blastocyst viability post vitrification

012028

A Kresna, W Widjeki and T Damayanti

OPEN ACCESS

Potential polysaccharide gel from *Dunio zibethinus* var. Raja galuh rind extract towards *Klebsiella pneumoniae* bacteria

012029

A Amanah, T M Pratamasari, M Tsilimah and R Primanegara

OPEN ACCESS

The application of barcode DNA rbcL gene for identification of medicinal plants: red jaben and gofasa

012030

Sundari, Khadijah, A.M Jayasil and N.H Sulikanto

OPEN ACCESS

Prospects and problems the amplification of DNA metagenomic from clove plant collected from plantation community in temate island

012031

N Nurhesanah, S Sundari and N Papuanan

OPEN ACCESS

The immunogenicity of polyclonal antibody from induced *Vibrio parahaemolyticus* membrane by Elisa method

012032

M Puspitawaty and S Sundari

OPEN ACCESS

Reproductive characters of senduro goat at lumajang district east java

012033

G Ciptadi, M N Ihsan, A Budianto, M. Mudawarwan, A I Putri and M. N A Naufal

OPEN ACCESS

In vitro fertilization of cryopreserved goat oocytes in different cryoprotectants

012034

S Warjuningtih, M N Ihsan, G Ciptadi and H N Karima

OPEN ACCESS

Combination effect of core stability exercise and contract relax exercise on hamstring flexibility

012035

A L Na'ima, G M Sari and D N Utomo

OPEN ACCESS

Oviduct specific glycoprotein expression in goats (*Capra aegagrus*) testes and ovaries

012036

H Pratiwi, A Firmawati, H Herawati, N Ismaili, I A H Kurniawan and L 'Ainiyah

OPEN ACCESS

Altered profile of gut microbiota and the level short chain fatty acids in colorectal cancer patients

012037

F Yusuf, S Adewulan, A F Syam and F Fatchiyah

OPEN ACCESS

Protein profiling of coloring rice (*Oryza sativa* L) using SDS-PAGE and experion™ 260 analysis

012038

D R T Sari, A Salatin, J R K Cairns and F Patchiyan

OPEN ACCESS

Effect of supplementation of either powdered or encapsulated probiotic on carcass percentage, giblets and small intestinal morphometric of local duck

012039

O Sjorjan, I Adiansah and K Snolha

OPEN ACCESS

Classification of *Cytodactylus marmoratus* species complex based on the molecular approach

012040

M A Faizi and N Kurniawan

OPEN ACCESS

Albumin and enzyme profiles of dwarf snakehead, *Channa gachua* caught from River Brantas, East Java

012041

A M Hartati, A. Yuniarji, Endarismani, W.E. Kusuma and D.G.R. Widjaya

OPEN ACCESS

Candlenut leaf methanol extract induces re-endothelialization in high-fat diet/streptozotocin-induced rats

012042


D Nugrahenny, K Mardhoyan, R S Bekti, E Mayangsari, Nurdiana, U Kalsum, T I Winedar, A F Insantaqwa, M E D Prawira, A F Khasanah, K A Putri and Y L Puaptasari

PAPER • OPEN ACCESS

Anti-inflammatory effect of ethyl acetate fraction of galing plant extract (*Cayratia trifolia*) on male wistar rats induced by carrageenan

To cite this article: D Santoso *et al* 2019 *J. Phys.: Conf. Ser.* **1146** 012021

View the [article online](#) for updates and enhancements.



IOP | ebooksTM

Bringing you innovative digital publishing with leading voices
to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of
every title for free.

Anti-inflammatory effect of ethyl acetate fraction of galing plant extract (*Cayratia trifolia*) on male wistar rats induced by carrageenan

D Santoso^{1*}, I K Sudiana², A S Rahayu² and M Yunus³

¹Departments of Internal Medicine, Faculty of Medicine, Universitas Airlangga

²Departments of Anatomic Pathology, Faculty of Medicine, Universitas Airlangga

³Departments of Veterinary Parasitology, Faculty of Veterinary Medicine, Universitas Airlangga

*Corresponding author : drdjokosantoso@yahoo.com

Abstract. Inflammation (inflammation) is a local reaction from living tissue or cells to an excitatory or injury. This study aims to determine the anti-inflammatory activity of ethyl acetate fraction of galling plant extract (*Cayratia trifolia*) on one of the inflammatory parameters, namely swelling in the legs of rat with 1% carrageenan induction. In addition, this study also aims to determine the effective dose of ethyl acetate fraction of galing plant extract (*Cayratia trifolia*) as anti-inflammatory. This study used 25 male wistar rats divided into 5 groups. Group 1 (negative control) Na CMC 0.5%, group 2 (positive control) diclofenac sodium 0.0065 mg /gBW rats, and group 3, 4 and 5 suspension ethyl acetate fractions of galing plant extract 0.0065; 0.013 and 0.026 mg / gBB rats. Inflammation in rats by inducing 1% carrageenan as much as 0.10 ml. The volume of edema every hour is known from the difference in foot volume at certain hours with normal foot volume. The AUC value of the edema volume was calculated by trapezoid method every one hour and calculated for anti-inflammatory power (IP). The results showed that ethyl acetate fraction of galing plant extract (*Cayratia trifolia*) could reduce the volume of udema in the legs of male white wistar rats which was induced by caragenine with an effective dose of 0.0065 mg/gBW.

1. Introduction

Traditional medicine is a mixture of unrefined natural ingredients comes from plants or animals used for treatment traditional [1]. Traditional medicine is one of an alternative for the treatment of different diseases because the side effects are considered smaller and the price is cheaper than modern medicine. So that the use of traditional medicines can be accounted for, it needs to be done various studies both to look for active components and to assess its effectiveness and safety [1].

Inflammation is a local reaction of tissue or cells to a stimulus or injury. This study aims to determine the anti-inflammatory activity of the ethyl acetate fraction of galing plant extract (*Cayratia trifolia*) on one of the inflammatory parameters, namely swelling in the feet of wistar rat with 1% carrageenan induction. In addition, this study also aims to determine the effective dose of ethyl acetate fraction of galing plant extract as anti-inflammatory.

2. Materials and Methods

This study used 25 male wistar rats divided into 5 groups. Group 1 (negative control) Na CMC 0.5%, group 2 (positive control) diclofenac sodium 0.0065 mg /gBW rats, and group 3, 4 and 5 suspension ethyl acetate fraction of galing plant extract 0.0065; 0.013 and 0.026 mg / gBB rats. Inflammation in rats by inducing 1% carrageenan as much as 0.10 ml. The volume of edema every hour is known from the difference in foot volume at certain hours with normal foot volume. The AUC value of the edema volume was calculated by trapezoid method every one hour and calculated for anti-inflammatory power (IP). The AUC value of the edema volume obtained was analyzed statistically by the Kruskal-Wallis and Mann-Whitney tests using SPSS release 16.

3. Results and Discussion

The galing plant obtained is processed into simplicia on all parts of plant. Simplicia is pollinated and sieved with No. 30/40 sieve. Simplicia what is used is simplicia which passes on No. 30 sieve and does not pass on the sieve no.40. The simplicia powder is then extracted using a maceration method in 96% ethanol solvent for 5 days. The filtrate obtained was concentrated with rotary evaporator. The viscous extract obtained was fractionated using a separating funnel successively with n-hexane, chloroform, diethyl ether and ethyl acetate. This process done until the liquid is clear. The amount of solvent used for fractionation proportional to the amount of water-ethanol added to the ethanol extract (ratio of 1: 1). Ethyl acetate fraction is collected and concentrated by rotary evaporator until a viscous ethyl acetate fraction is obtained. Preliminary testing begins with phytochemical screening to find out the compounds contained in the ethyl acetate fraction of galing plant extract are thought to be anti-inflammatory. Phytochemical screening includes testing phenolic, alkaloids, flavonoids, saponins, tannins, and triterpenoid. If the results are positive, then the confirmation test is continued using TLC to ensure that there is a positive substance in the phytochemical screening test.

Table 1. Average% volume of edema, total AUC, and% DAI of various treatment groups

Group	% Volume of Edema							% Total of AUC (ml.hour)	% IP
	0 minute	30 minutes	60 minutes	90 minutes	120 minutes	150 minutes	180 minutes		
Negative Na-CMC 0.5%	126.31	138.65	132.42	137.24	143.15	141.53	144.36	88.75 ^b	0
Positive Na-diclofenac	122.18	114.28	113.17	110.42	105.44	103.67	102.35	67.08 ^a	21.5
Galing 0.0065mg per gram BB	123.05	111.64	111.25	109.32	106.52	105.44	102.25	68.43 ^a	22.6
Galing 0.013mg per gram BB	121.32	119.27	117.38	113.34	111.26	108.54	107.21	67.52 ^a	24.4
Galing 0.026mg per gram BB	119.25	118.27	110.32	105.29	101.63	100.45	100.33	66.84 ^a	23.8

Notes

a: significant different ($p<0.05$) on positive control group with Mann-Whitney test

b: significant different ($p<0.05$) on negative control group with Mann-Whitney test

The results obtained both phytochemical screening and assertion tests showed that the ethyl acetate fraction of galing plant extract contained secondary metabolites of alkaloids, flavonoids, saponins, and triterpenoids / steroids. Furthermore, the treatment of test compounds in male wistar rats at 2-3 months old to see the anti-inflammatory effect of galing plant ethyl acetate fraction. The inflammatory parameters observed were edema in the feet of rat with sub plantar 1% carrageenan administration. Carrageenan does not cause other tissue damage and does not cause scars, and provides a more sensitive response to anti-inflammatory drugs. Carrageenan latent time starts 1 hour and maximal edema formation occurs 3 hours after carrageenan administration [1]. The data observed were the percentage of edema volume, the Area Under Curve (AUC) value which illustrated the magnitude of the edema, and the percentage of anti-inflammatory power (IP). The results of this observation can be seen in Table 1.

Based on Table 1, the peak occurrence of edema caused by carrageenan occurs at 30 minutes. Furthermore, with the administration of galing both doses of 0.0065; 0.013; and 0.026 mg / gBW and diclofenac sodium, the percentage of edema volume decreases. This shows that galing can reduce the volume of edema or as anti-inflammatory. While the negative control group with the administration of ethyl acetate fraction carrier is CMC Na 0.5% the volume of edema remains even rising. Graph about the description of the percentage of edema volume of all treatment groups can be seen in Figure 1. Furthermore,% of the volume of the edema was used to calculate the AUC value. AUC describes the amount of inflammation that occurs. After that,% IP was also calculated to describe the percentage of anti-inflammatory power. The AUC value was inversely proportional to % IP. The smaller the AUC value means the amount of inflammation decreases so that the greater the percentage of anti-inflammatory power. Based on Table 1, the total AUC value of the negative control group was the largest compared to other groups. The positive control group of Diclofenac Na has a greater AUC value than the galing group of all doses. This is also shown % IP of galing greater than the drug Diclofenac Na. The higher the dose of galing, the higher than IP. This proves that galing can effect reducing edema which is one sign of inflammation or also called anti-inflammatory. A description of the total AUC value than the galing group of all doses. This is also shown % IP of galing greater than the drug Diclofenac Na. The higher the dose of galing, the higher than IP. This proves that galing can effect reducing edema which is one sign of inflammation or also called anti-inflammatory. A description of the total AUC value and % IP of all treatment groups can be seen in Figure 2.




Figure 1. Graph of relation between % volume of edema to time of various treatment groups

Statistical tests with SPSS 16.00 were also carried out to reinforce galing's anti-inflammatory effects. The results obtained were significant differences in AUC values between the negative control groups and the galing group of all doses. This proves that the galing plant extract ethyl acetate fraction can have anti-inflammatory effects. In addition the results of statistics obtained there was no significant difference in AUC values between groups of Diclofenac Na with the galing group all doses.

This shows that the ethyl acetate fraction has an anti-inflammatory effect comparable to Diclofenac Na. Based on the results of statistical tests, the effective dose of galing ethyl acetate fraction was 0.0065 mg / gBW.

Galing ethyl acetate fraction can have anti-inflammatory effects allegedly caused by secondary metabolites contained therein, namely flavonoids, saponins and steroids. Some flavonoids play a role in inhibiting lipoxygenase while other flavonoids play a role in inhibiting prostaglandin synthesis [3]. Flavonoids also have an influence on collagen metabolism in several ways, among others by crosslinking with collagen fibers so that the crosslinking of collagen becomes strong, and is able to stop the damage to collagen structure due to the presence of enzymes from white blood cells this process arises during inflammation [4]. The effects of flavonoids as an indirect antioxidant also support the anti-inflammatory effects of flavonoids. The presence of free radicals can attract various inflammatory mediators [5]. Flavonoids can stabilize Reactive Oxygen Species (ROS) by reacting with reactive compounds from radicals so that radicals become inactive [5]. Many saponins have been reported to have an anti-inflammatory effect, but the mechanism is not yet clearly known. Saponins consist of steroids or triterpenes (aglycones) which have actions such as detergents. The anti-inflammatory mechanism most likely is suspected saponins capable of interacting with many lipid membranes [6] such as phospholipids which are precursors of prostaglandins and other inflammatory mediators. Steroids in the body can inhibit the enzyme phospholipase A2 which is the enzyme responsible for the release of arachidonic acid which is then metabolized by the enzyme cyclooxygenase and lipo-oxygenase which will then release inflammatory mediators [7].




Figure 2. Total AUC (ml.hour) and % IP diagrams of all treatment groups

4. Conclusion

Ethyl acetate fraction of Galing (*Cayratia trifolia*) extract can have anti-inflammatory effects on male white rats induced by caragenine with an effective dose of 0.0065 mg / gBW.

Acknowledgements

This research was supported by Directorate of Higher Education, Ministry of Education and Culture of the Republic of Indonesia for funding through higher education excellence research grant for a research contract number Nomor: 122/SP2H/PTNBH/DRPM/2018

References

- [1] Tjokronegoro A and Baziat A 1993 *Etik Penelitian Obat Tradisional* Semiloka di Fakultas Kedokteran Universitas Indonesia 6-7 Juli 1992, Balai Penerbit Fakultas Kedokteran UI. Jakarta.
- [2] Chattopadyay, Deprasad, Arunachalam, Ghosh L, Rajendran K, Mandal A B and Bhatta C S K 2005 *J. Pharm Pharmaceut Sci* **8** (3): 558-564. www.cspsCanada.org.
- [3] Robbins SL and Kumar V 1995 *Buku Ajar Patologi I* Terjemahan Staf Pengajar Laboratorium Patologi Fakultas Kedokteran UNAIR , cetakan 2. Jakarta : EGC.
- [4] Wirakusumah ES 2006 *Buah dan sayur untuk terapi* Jakarta : Penebar Swadaya
- [5] Nijveldt R J E van Nood DEC van Hoorn PG, Boelens K, van Norren PAM and van Leeuwen. 2001 Flavonoids: a review of probable mechanisms of action and potential applications. *American Journal of Clinical and Nutrition* **74**: 418-425.
- [6] Nutritional Therapeutics 2003 NT Factor: Phosphoglycolipids-High Energy Potential. www.propax.com/FAQ/soy_high_energy.html [2 Desember 2005].
- [7] Katzung BG 2002 *Farmakologi Dasar dan Klinik*. Jakarta: Salemba Medika