




Knowledge E
Engaging minds



KnE Life Sciences

ISSN 2413-0877



www.KnEpublishing.com

Scientific and Editorial (S & E) Board International Conference on Natural Resources and Life Sciences the 1st NRLS - 2016

- Elieser Tarigan, Faculty of Electrical Engineering - University of Surabaya, IDN.
- Go Melisa Gunawan, Faculty of Biotechnology - University of Surabaya, IDN.
- Johan Sukweenadhi, Kyung Hee University - KOR; Faculty of Biotechnology - University of Surabaya, IDN.
- Juris Burlakovs, Faculty of Health and Life Sciences, Dept. of Biology and Environmental Science, Linnaeus University, SWD
- Lanny Sapei, Faculty of Chemical Engineering - University of Surabaya, IDN.
- Lieke Riadi, Faculty of Chemical Engineering - University of Surabaya, IDN.
- Maizirwan Mel, Department of Biotechnology Engineering, Faculty of Engineering, International Islamic University Malaysia, MYS.
- Maria Goretti Marianti Purwanto, Faculty of Biotechnology - University of Surabaya, IDN.
- Mariana Wahjudi, Faculty of Biotechnology - University of Surabaya, IDN.
- Nurul Azizah, Faculty of Biotechnology - University of Surabaya, IDN.
- Popy Hartatie Hardjo, Faculty of Biotechnology - University of Surabaya, IDN.
- Praptiningsih Gamawati Adinurani, Faculty of Agrotechnology, University of Merdeka, Madiun, IDN.
- Roy Hendroko Setyobudi, IABST, Jakarta and Ma Chung Research Center for Photosynthetic Pigments, Malang, IDN.
- Sharr Azni Harmin, Fakulti Sains & Bioteknologi, Universiti Selangor, MYS.
- Sulisty Emantoko Dwi Putra, Faculty of Science - University of Potsdam, DEU; Faculty of Biotechnology University of Surabaya, IDN.
- Theresia Desy Askitosari, RWTH Aachen University, Institute of Applied Microbiology, DEU.
- Tjie Kok, Faculty of Chemical Pharmaceutical Biology, University of Groningen, NLD.

VMIC—The Veterinary Medicine International Conference—is one of the world's leading conference focusing on a wide array of topics including Veterinary Medicine and Biomedical Science. It offers a stimulating venue for scientists, researchers, lecturers, general practitioners, and others to broaden their social scientific network. This conference aimed at improving human and public health by improving agricultural and food systems, advancing biomedical and comparative medical research, preventing and addressing zoonotic diseases, kit diagnostic, enhancing environmental and ecosystem health, and helping manage the 21st-century public health challenges.

Conference date: 12–14 July 2017

Location: Surabaya, East Java, Indonesia

Editors: Sri Agus Sudjarwo, Fedik A. Rantam, rer. nat. Gunawan Indrayanto, Muchammad Yunus, Rimyant, Wiwik Misaco, Tita Damayanti Lestari, Mustofa Helmi Effendi, Dikky Eka Mandala Putranto, and Shafia Khairani

Organizer: Faculty of Veterinary Medicine, Universitas Airlangga, Indonesia

Sponsors: Ministry of Research, Technology and Higher Education of the Republic of Indonesia, Universitas Airlangga, ROMINDO PRIMAVETCOM, Gajah Mada University, Bogor Agriculture Institute, ILRI (International Livestock Research Institute), Kagoshima University, Miyazaki University, Faculty of Veterinary Science, Chulalongkorn University, College of Veterinary Medicine, Tarlac Agricultural University, Yamaguchi University, Erasmus MC, Kasetsart University, Wyndham Veterinary Clinic, GERBU – Germany, USAID

Published: 29 November 2017

ISSN: 2413-0877

TABLE OF CONTENTS

Toll-Like Receptors (TLRs) Play Role in Adaptive Immunity in Rabbits Immunized by *Sarcoptes scabiei* Proteins

THE VETERINARY MEDICINE INTERNATIONAL CONFERENCE (VMIC)

Pages 1-9 Nunuk Dyah Retno Lastuti, Fedik Abdul Rantam, Poedji Hastutie, Dony Chrismanto

Preservation Effect of Grouper (*Epinephelus* sp) Fillet Against Survival of Anisakidae

THE VETERINARY MEDICINE INTERNATIONAL CONFERENCE (VMIC)

Pages 10-20 Hartanto M. Raharjo, Setiawan Koesdarto, A.T Soelih Estoepangestie, Kusuma Wardhani L.D

Acanthocephalan in *Xenochrophis piscator* Snake in Sidoarjo Indonesia

THE VETERINARY MEDICINE INTERNATIONAL CONFERENCE (VMIC)

Pages 21-27 Inggarsetya Syah Audini, Lucia Tri Suwanti, Setiawan Koesdarto, Emmanuel Djoko Poetranto

The Identification Blood Parasites On Pig (*Susdomesticus*) In Polewali Mandar District

THE VETERINARY MEDICINE INTERNATIONAL CONFERENCE (VMIC)

Pages 28-33 Silvana Arfin, Lucia Muslimin, Adryani ris

Spirometra in *Ptyas mucosus* Snake in Sidoarjo, Indonesia

THE VETERINARY MEDICINE INTERNATIONAL CONFERENCE (VMIC)

Pages 34-40 Garindra Tiara Pranashinta, Lucia Tri Suwanti, Setiawan Koesdarto, Emmanuel Djoko Poetranto

Toxicity of *Citrus mitis*, *Citrus aurantifolia*, and *Citrus maxima* leaf extract toward mortality of *Aedes aegypti* larvae (Diptera: Culicidae)

THE VETERINARY MEDICINE INTERNATIONAL CONFERENCE (VMIC)

Pages 41-47 Hamidah Hamidah, Hebert Adrianto

An In Vitro Antibacterial Activity Test of Meniran Herbs' (*Phyllanthus Niruri* L.) Ethanol Extract Against *Mycoplasma gallisepticum* causes Chronic Respiratory Disease (CRD) in Broiler Chickens

THE VETERINARY MEDICINE INTERNATIONAL CONFERENCE (VMIC)

Pages 48-61 Emy Koestanti Sabdongrum, Sri Hidanah, Retno Sri Wahjuni, Sri Chusniati, Arimbi Arimbi

Bioremediation of Mercury (II) Contaminated Seawater Using the Diatom *Skeletonema costatum*

THE VETERINARY MEDICINE INTERNATIONAL CONFERENCE (VMIC)

Pages 62-68 Thin Soedarti, Tini S., Sucipto H., Eko P. Kuncoro

BMP-2 Expression of Post Tooth Extraction that Catfish Oil Application

Utilization of Honey *Apis dorsata* as Antiosteoporosis on Requirements of Bone Calcium Ash Density on Ovariohysterectomized White Rat (*Ratus norvegicus*)

THE VETERINARY MEDICINE INTERNATIONAL CONFERENCE (VMIC)

Pages 627-632 Muhammad Huda Ramadhan Ibrahim, Abdullah Hasib Hasib, Siti Nur Rohmah, Salsabilla Abani, Samsi Yordan, Ira Sari Yudaniayanti

Increased Integrity of Plasma Membrane and Acrosome Cap Spermatozoa Limousin Cattle at Post Thawing in Frozen Media by adding Seawater Extract

THE VETERINARY MEDICINE INTERNATIONAL CONFERENCE (VMIC)

Pages 633-641 Nur Faidah, Tatik Hernawati, Mirni Lamid, Ismudiono Ismudiono, Tri Wahyu Suprayogi, Sri Mulyati

The Relation of Body Temperature and Vaginal Cytology Examination in Time Artificial Insemination Rate Fat-tailed Sheep (*Ovis Aries*) in The District Sidoarjo East Java

THE VETERINARY MEDICINE INTERNATIONAL CONFERENCE (VMIC)

Pages 642-649 Rhendyka Prasetya Anggriawan, Suzanita Utama, Hana Eliyani

Effect of Laser Acupuncture Shoot on Ova Point of Male Mojosari Duck (*Anas platyrhynchos*) on The Number of Sertoli and Leydig Cells

THE VETERINARY MEDICINE INTERNATIONAL CONFERENCE (VMIC)

Pages 650-657 Yuanara AR Adikara, Abdul Samik, Ira S Yudaniayanti, Tatang S Adikara, Eka P Hestianah, Suzanita Utama

Insulin-Like Growth Factor-I (IGF-I) from Crossbred Pregnant Mare Serum to Increase Follicle Number of Mice (*Mus musculus*)

THE VETERINARY MEDICINE INTERNATIONAL CONFERENCE (VMIC)

Pages 658-667 Abdullah Abdullah, Tjuk IRestiadi, Nunuk DR Lastuti, Tita Damayanti, Wurlina Wurlina, Erma Safitri

- Yoanes Maria Vianney, Faculty of Biotechnology - University of Surabaya, IDN.
- Zane Vincēviča-Gaile, Department of Environmental Science, University of Latvia, Riga - Latvia, European Union, LVA.

Conference Paper

Insulin-Like Growth Factor-I (IGF-I) from Crossbred Pregnant Mare Serum to Increase Follicle Number of Mice (*Mus musculus*)

Abdullah¹, Tjuk IRestiadi², Nunuk DR Lastuti³, Tita Damayanti², Wurlina², and Erma Safitri²

¹Student of Veterinary Medicine Faculty Surabaya, 60115, Indonesia

²Veterinary Reproduction Department of Veterinary Medicine Faculty, Surabaya, 60115, Indonesia

³Veterinary Parasitology Department of Veterinary Medicine Faculty, Surabaya, 60115, Indonesia

Abstract

The purpose of the research was to know the effect of Insulin-Like Growth Factor-I (IGF-I) derived from pregnant crossbred mare serum (PMS) in mice (*Mus musculus*) folliculogenesis. The subjects of this research were 20 female mice. The research was arranged by Completely Randomized Design (CRD) with four treatments and five replications. The treatments were C0 = 10 ng/ml of physiological NaCl, P1 = 10 ng/ml of IGF-I PMS, P2 = 20 ng/ml of IGF-I PMS, and P3 = 40 ng/ml of IGF-I PMS. Observed variables are the number of primary, secondary, tertiary and de Graff follicles. During the treatment, the estrous cycle was also observed. The data of follicles number were analyzed by Analysis of Variance (ANOVA) and followed by HSD (Honestly Significant Difference) test. The result showed that the addition of IGF-I PMS significantly affects ($p < 0.05$) on increasing the primary and secondary follicles number. The addition of IGF-I PMS 20 ng/ml and 40 ng/ml can increase the primary and secondary follicles significantly ($p < 0.05$).

Keywords: IGF-I crossbred mare serum pregnant; follicle; *Mus musculus*.

Corresponding Author:
Tjuk IRestiadi

Received: 03 October 2017
Accepted: 10 October 2017
Published: 29 November 2017

Publishing services provided
by Knowledge E

© Abdullah et al. This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Selection and Peer-review under the responsibility of the VMIC Conference Committee.

1. Introduction

The human need of animal protein is significantly increasing nowadays. It can be shown that Indonesian people consume 6.65% of meat during 2013-2014 [1] and is estimated that the number of consumption will keep rising up. On the contrary, the number of ranch industry is decreasing every year [2], especially in 2011 until 2014. At that time, the cattle ranch number in the slaughterhouse was significantly declining. In addition, the supplies of beef in some markets are decreasing as well. The government has attempted to solve this problem, like importing beef from Australia, but this has not

OPEN ACCESS

completely solved the problem. The people might not continually import the beef as the imported beef cost is much more expensive than the local one. Therefore, there was a thought to improve the productivity of livestock.

According to the regulations of the minister of agriculture no. 48 in 2016 about increasing the population of cows and buffaloes pregnancy rapidly. Article 5 verse 1 and 2 stated that the effort of increasing the population of cows and buffaloes pregnancy will be done by reproduction management system [3]. This system includes the examination for reproduction status and reproduction disorder, artificial insemination (AI) service and natural mating, fulfilling frozen semen and liquid N₂, controlling productive female individuals, and fulfilling forage and concentrate. Many problems, like reproduction, occur more often to stock farmers when they attempt to increase livestock population and productivity. The reproduction problems that usually occur are the cows are pregnant but there is not lust indication; reproduction hormone is abnormal (high progesterone), this causes false pregnancy and luteal cysts; the emergence of yellowish or greenery mucus with bad smell as there is inflammation inside the reproductive tract; unobserved lust; lack of feed which leads to ovarian hypofunction; repeated mating; and reproduction hormonal disorder (high estrogen) due to follicle cysts [4].

Reproduction or breeding is a process of forming a new individual whether it passes through natural mating or artificial mating [5]. Reproduction is a physiological process that is experienced by all living things in order to retain offspring and life sustainability [6]. This process requires reproduction hormone which has a role in reproduction system directly and indirectly. On the other hand, ovarium has also a significant role. It has two based functions: producing fertilizable oocyte that has faultless growth and secreting steroid hormone to set up the reproductive tract in a fertilization and implantation process [7] Thus when the ovary is in interference, it will disrupt reproduction system. A lot of previous research used bioactive materials which aimed to improve reproduction ability.

One of the bioactive materials that have a significant role in female animals reproduction system is Insulin-Like Growth Factor-I (IGF-I). IGF-I supplementation in maturation medium and oocyte culture can stimulate and increase the number of the mature oocytes. In addition, it also increases in vitro fertilization (IVF) [8] and the number of the embryo that reaches the stage of blastocyst stage for some animals, like pigs, cows [9], and buffaloes [10]. The result of this research [11] states that adding IGF-I during preantral cultures of pigs can intensify follicle diameter. Besides, it can increase mRNA Insulin-Like Growth Factor Receptor-I (IGFR-I) as well, by supplementing Follicle

Stimulating Hormone (FSH) during culture, and the oocyte will be processing meiosis stage after maturing. In addition [12], IGF can increase the proliferation of granulosa cells, steroidogenesis, and oocyte growth in mammals mainly. This research [13] also proves that identical events which happened on cows are related to IGF-I concentration enhancement toward serum and follicle itself. IGF-I from the ovary and/or from systemic circulation is an important thing in folliculogenesis regulation and is a mediator of the genetic component of ovulation on cows.

2. Materials and methods

This research applied laboratory experimental research with Completely Randomized Design (CRD) as the methods of the study. The subjects of this research were 20 Balb/C female mice in the age of 28-32 days which were got from Pusat Veterinaria Pharma (Pusvetma), Surabaya and were avowed by Research Ethics Commission of Faculty of Veterinary Medicine, Universitas Airlangga as ethically proper subjects. From 20 female mice, they were divided into four treatments groups. The mice in control group (C0) were injected by physiology NaCl 0.1 ml, treatment P1 were injected by IGF-I PMS 0.1 ml (10 ng/ml), treatment P2 were injected by IGF-I PMS 0.1 ml (20 ng/ml), and treatment P3 were injected by IGF-I PMS 0.1 ml (40 ng/ml). All injections were done subcutaneously and the IGF-I PMS injections were done 5 times in 5 days. Before injecting, there would be vaginal swab toward the mice in order to know the status of the reproduction.

2.1. Experimental Animals Surgery and Preparing Mice Ovary Histopathology

In the end of the research, the experimental animals would be anesthetized by ether to become stagnate and did surgery to get the left ovary. Afterwards, the ovary was separated from its fat and soaked in 10% formalin solution for 1-2 days to fix the tissue. Subsequently, the ovary was ready to go on the next stage for making mice ovary histopathology preparation. This process was using routine dyeing techniques Haematoxylin-Eosin in Veterinary Pathology Laboratory, Faculty of Veterinary Medicine, Universitas Airlangga.

2.2. Identify and Calculate Mice Follicles

Ovary preparation was observed by using a microscope with ocular lens magnification of 10x and objective lens magnification of 4x-40x. From this observation, the follicle was classified based on the oocyte size inside the follicles, the size of follicles which is based on the number of cells that cover them, and follicles morphology [14]. Primary follicles consist of a layer of flat granulosa cells and cubical granulosa cells, or it is only the cubical shape, while for secondary follicles, they consist of two layers or more of cubical granulosa cells [15]. Next, small antral follicles (tertiary follicles) have segmented cavity with 2 antrum or more, while big antral follicles (follicle de Graff) have one cavity, called antrum, which continuously enlarged [16]. These kinds of the follicle (both preantral and antral) in ovary was calculated by examining all viewing fields, then the data was conducted and analyzed.

3. Results

The result of counting the number of mice primary follicles after given Insulin-Like Growth Factor-I (IGF-I) derived from crossbreed pregnant mare serum (PMS) showed that there was escalation number of the primary follicles. Giving this serum showed significantly affect ($p < 0.05$) the treatment T1, T2, T3. The higher of the dose, the more increasing of the follicles. It can be seen that IGF-I PMS with a dose of 20 ng/ml (T2) and 40 ng/ml (P3) can increase the number of mice follicles into ($p < 0.05$).

Similar to the primary one, the result of counting the number of mice secondary follicles after injected IGF-I derived from crossbreed pregnant mare serum (PMS) also showed significantly affect ($P < 0.05$) the treatment T1, T2, and T3. IGF-I PMS with a dose of 20 ng/ml (P2) and 40 ng/ml (T3) can increase the number of mice secondary follicles, comparing to control group (C0) ($p < 0.05$). Besides, IGF-I PMS with a dose of 20 ng/ml and 40 ng/ml (T2 and T3) have significantly affected ($p < 0.05$) the treatment T1 in increasing mice secondary follicles number.

On the other hand, the result of mice tertiary follicles and de Graff follicles number after injected IGF-I derived from crossbreed pregnant mare serum (PMS) did not show any changes. Giving this treatment did not show significantly affect ($p > 0.05$) as well toward mice tertiary follicles and de Graff follicles for treatment T1, T2, and T3.

To understand more about the follicles, each follicle is presented in Figure 1, while the graphic for mice primary, secondary, tertiary, and de Graff follicles number is presented




Figure 1: The result of the ovarian follicles identification in mice (*Mus musculus*) (Haematoxylin-eosin, 400x). In the figure above, (A) is a primary follicles composed of a flat layer of granulosa cells, (B) is a secondary follicle consisting of two or more layers of both cubic and cylindrical granulosa cells, and (C) is a tertiary follicles which is a follicle with some cavities called call-exner bodies and (D) is the de Graaf with antral follicles containing liquor folliculi (Haematoxylin-Eosin, 100x).

in Figure 2. The results of counting primary, secondary, tertiary, and de Graaf follicles number after injected IGF-I PMS are provided in Table 1.

The function of the ovary is to supply germinal cells which are needed for living things sustainability. Besides, ovary produces hormones that are useful for female

Tabel 1: The average and standard deviation of the number of primaries, secondary, tertiary, and de Graff follicles in mice ovary (*Mus musculus*) after given IGF-I PMS.

Treatment	Follicle number (x ± SD)			
	Primary Follicle	Secondary Follicle	Tertiary Follicle	de Graff Follicle
CO	9.40 ^a ± 6.58	6.20 ^a ± 3.11	4.60 ^a ± 2.88	3.40 ^a ± 2.79
T1	14.00 ^{ab} ± 2.82	8.40 ^a ± 2.88	8.40 ^a ± 2.88	3.00 ^a ± 1.22
T2	23.80 ^b ± 7.19	21.60 ^b ± 6.34	8.60 ^a ± 4.27	3.60 ^a ± 1.14
T3	20.40 ^b ± 4.87	20.00 ^b ± 10.12	11.00 ^a ± 4.41	5.40 ^a ± 1.81

Note: The different superscript in one column shows significant differences (p<0.05)

T1 = injected with 10 ng/ml IGF-I PMS

T2 = injected with 20 ng/ml IGF-I PMS

T3 = injected with 40 ng/ml IGF-I PMS




Figure 2: The average and standard deviation of the number of primaries, secondary, tertiary, and de Graff follicles in mice ovary (*Mus musculus*) after given IGF-I PMS.

individuals growth [18-20]. Female individuals were born with limited primordial follicles in which when it runs out, it cannot be added again. This thing leads to folliculogenesis that can be called as irreversible process [21-23]. Ovary folliculogenesis comprises all follicle growth, differentiation, and ovulation. More than 99% of follicles will degenerate or will have atresia. In mammals, the follicles are controlled by endocrine factors, like FSH and LH, and paracrine factor. Among those factors, there is an element called Insulin-Like Growth Factor (IGF) which has an important role in modulating gonadotropin in proliferation and differentiation of follicular cells [24].

Based on the result of this research, giving Insulin-Like Growth Factor (IGF) derived from crossbreed pregnant mare serum can increase the number of primary and secondary follicles significantly (p<0.05) comparing to the control group. Moreover, the

doses of 20 ng/ml and 40 ng/ml in IGF also affects the escalation of primary and secondary follicles significantly ($p < 0.05$). On the other hand, the number of mice tertiary and de Graff follicles, injected with IGF, did not show significantly affect ($p > 0.05$) in control group (C0). This proves that IGF-I can stimulate receptor cells type 1 in granulosa cells and theca cells [25] and the reaction from the receptor in granulosa cells will be raised by estrogen and gonadotropin. Subsequently, IGF-I receptor reaction will increase in small antral follicles [26].

According to this research, Insulin-Like Growth Factor-I (IGF-I) derived from cross-breed pregnant mare serum can advance the number of primary and secondary follicles because the mice have not been in puberty. This is corresponding with the follicular development theory proposed by [27] in which the dominant mice ovary that belongs to pre-puberty mice like this research's subjects is primary and secondary follicular development.

There is also another theory that states ovary follicular development is divided into two stages: preantral and antral. Preantral encompasses primordial follicle, primer follicle, and secondary follicle. This phase, can be called as gonadotropin-independent phase, is marked by oocyte growth and differentiation which is affected by some growth factors through the autocrine/paracrine system, like Transforming Growth Factor β (TGF- β) superfamily and insulin-like Growth Factor I (IGF-I). Another phase, antral or gonadotropin-independent, is marked by a significant escalation of the follicle size. This phase encompasses de Graff follicle. In addition, antral is controlled by Follicle Stimulating Hormone (FSH) and luteinizing Hormone (LH) and another growth factors. These another factor will stimulate proliferation cell and affect gonadotropin activity [28-30]. Therefore, there were only primary and secondary follicles which increase significantly in this research, while tertiary and de Graff follicles were not increasing.

The working mechanism of protein complex IGF-I and IGFBP-3 toward immature and mature follicles was explained in [31]. It stated that immature follicle, LH, assisted with IGF-I and activin in granulosa cell, will stimulate theca cell to form androgen. IGF-I and IGFBP-3 were binded and disparted by TIMP (Tissue InhibitingMetalo Proteinase) with high concentration. However, the ability of disparting was not significant, so that the generated IGF-I would not be plentiful. Androgen was brought to granulosa cell, then it was changed into estradiol through aromatization process and was stimulated by IGF-I and FSH.

In mature follicle, LH, assisted with IGF-I and Inhibin in granulosa cell, would stimulate theca cell to form androgen. While IGFBP-3 would be disparted by TMP to produce IGF-I in large quantities. The androgen was brought to granulosa cell and changed

into estradiol through aromatization process. Afterwards, by controlled by aromatase inhibitor enzyme that was produced by granulosa cell, the androgen was stimulated by IGF-I and FSH to increase the secretion [31]. Based on this, the mice which have not been in estrous phase, they would get into the phase faster and it would be maintained during IGF-I injection.

Protein complex IGF-I and IGFBP-3 have an effect on forming FSH. This hormone, produced by the pituitary, will stimulate follicle growth in the ovary and form LH receptor in ovary follicles cells as well [32]. According to [33], injecting IGF-I and IGFBP-3 in diestrous phase can stimulate the anterior pituitary to produce LH and FSH to secrete estrogen in the blood. Estradiol secreting escalation by dominant follicle cells when ovulation happened will be causing lust, stimulate LH, and ovum excretion.

References

- [1] Direktorat Jendral Peternakan dan Kesehatan Hewan. 2015. Statistik Peternakan dan Kesehatan Hewan 2015. Kementerian Pertanian.
- [2] BPS. 2015. Peternakan. Badan Pusat Statistik Nasional.
- [3] Kementerian Pertanian. 2016. Peraturan Menteri Pertanian (Permentan) no. 48 tahun 2016 tentang Upaya Khusus Percepatan Peningkatan Populasi Sapi dan Kerbau Bunting. Kementerian Pertanian Republik Indonesia.
- [4] Ulum, F.M. dan Purwanta, B. 2015. Manual: Manajemen Kesehatan Reproduksi Ternak Sapi. Lembaga Penelitian dan Pengabdian Kepada Masyarakat. Institut Pertanian Bogor. Bogor.
- [5] Ismudiono, Srianto, P., Anwar, H., Madyawati, S.P., Samik, A., dan Safitri, E. 2010. Buku Ajar Fisiologi Reproduksi Pada Ternak. Airlangga University Press. Surabaya
- [6] Nursyah, D.A. 2012. Gambaran Siklus Estrus Tikus Putih (*Rattus norvegicus*) Ovariektomi Yang Diberi Tepung Daging Teripang (*Holothuria scabra*) [Skripsi]. Fakultas Kedokteran Hewan. Institut Pertanian Bogor. Bogor.
- [7] Palermo, R. 2007. Differential Actions of FSH and LH During Folliculogenesis. *Reproductive BioMedicine* Vol 15 3: 326-337.
- [8] Oberlender, G., Murgas, L.D.S., Zangeronimo, M.G., da Silva, A.C., Menezes, T.A., Pontelo, T.P. and Vieira, L.A. 2013. Role of Insulin-Like Growth Factor-I and Follicular Fluid from Ovarian Follicles with Different Diameters on Porcine Oocyte Maturation and Fertilization In Vitro. *Theriogenology*. 80: 319–327.
- [9] Neira, J.A., Tainturier, D., Pen, M.A. and Martal, J. 2010. Effect of the Association of IGF- I, IGF-II, bFGF, TGF-b1, GM-CSF, and LIF on the Development of Bovine Embryos

- Produced In Vitro. *Theriogenology* 73: 595–604.
- [10] Singhal, S., Prasad, S., Singh, B., Prasad, J.K. and Gupta, H.P. 2009. Effect of Including Growth Factors and Antioxidants in Maturation Medium Used for In Vitro Culture of Buffalo Oocytes Recovered In Vivo. *Anim Reprod Sci.* 113:44– 50.
- [11] Magalhaes-Padilha, D.M., Duarte, A.B.G., Araujo, V.R., Saraiva, M.V.A., Almeida, A.P., Rodrigues, G.Q., Matos, M.H.T., Campello, C.C., Silva, J.R. and Gastal, M.O. 2012. Steady-state Level of Insulin-Like Growth Factor-I (IGF-I) Receptor mRNA and The Effect of IGF-I on the In Vitro Culture of Caprine Preantral Follicles. *Theriogenology* 77:206–213.
- [12] Silva, J.R.V., Figueiredo, J.R., and vanden Hurk, R. 2009. Involvement of Growth Hormone (GH) and Insulin-Like Growth Factor (IGF) system in ovarian folliculogenesis. *Theriogenology* 71: 1193–1208.
- [13] Echternkamp, S.E., Spicer, L.J., Gregory, K.E., Canning, S.F., and Hammond, J.M. 1990. Concentrations of insulin-like growth factor-I in blood and ovarian follicular fluid of cattle selected for twins. *Biol Reprod.* 43:8–14.
- [14] Pedersen, T., and Peters, H. Proposal For A Classification of Oocytes and Follicles in The Mouse Ovary. *J Reprod Fertil.* 1968; 17:555–557.
- [15] Gougeon, A. 1996. Regulation of ovarian follicular development in primates: facts and hypotheses. *Endocr Rev.* 17:121–55.
- [16] Griffin, J., Benjamin, R., Emery, B.R., Huang, I., Peterson, C.M., Carrell, D.T. 2006. Comparative analysis of follicle morphology and oocyte diameter in four mammalian species (mouse, hamster, pig, and human). *J Exp Clin Assist Reprod.* 3:2.
- [17] Kusningrum, R.S. 2011. *Perancangan Percobaan.* Airlangga University Press. Surabaya. Hal: 165-212.
- [18] Hoyer, P.B. 2004. Can the clock be turned back on ovarian aging?. *Sci Aging Knowl Environ* 2004;p: 11.
- [19] Bhattacharya, P and Keating, A.F. 2012. Impact of environmental exposures on ovarian function and role of xenobiotic metabolism during ovotoxicity. *Toxicol Appl Pharmacol.* 261:227–235.
- [20] Anderson, L,D. and Hirshfield, A,N. 1992. An overview of follicular development in the ovary: from embryo to the fertilized ovum in vitro. *Md Med J;* 41:614–620.
- [21] Edson, M.A., Nagaraja, A.K., and Matzuk, M.M. 2009. The mammalian ovary from genesis to revelation. *Endocr Rev.* 30:624–712.
- [22] Hirshfield, A.N. 1991. Development of follicles in the mammalian ovary. *Int Rev Cytol.* 124:43–101.

- [23] Elvin, J.A., and Matzuk, M.M. 1998. Mouse models of ovarian failure. *Rev Reprod.* 3:183–195.
- [24] Monget, P., Besnard, N., Huet, C., Pisselet, C., and Monniaux, D. 1996. Insulin-like growth factor-binding proteins and ovarian folliculogenesis. *Horm Res.* 45:211–217.
- [25] Giudice, L.C. 1992. Insulin-like Growth Factor and Ovarian Follicular Development. *Endocrine Reviews.* 13: 641-669.
- [26] Wandji, S.A., Teresa, L.W., Jennifer, C., Steven, W.L. and Hammond, J.M. 1998. Expression of Mouse Ovarian Insulin-like Growth Factor System Component During Follicle Development and Atresia. *Endocrinology.* 134:5205-5214.
- [27] Partodiharjo, S. 1992. Ilmu Reproduksi Hewan. Ed. Ke-3. Jakarta: Mutiara Sumber Widya.
- [28] Webb, R., Garnsworthy, P.C., Gong, J.G., and Armstrong, D.G. 2004. Control of Follicular Growth: Local Interaction and Nutritional Influence. *Journal of Animal Science.* 82: E63-E74.
- [29] Anwar, R. 2005. Morfologi dan Fungsi Ovaarium. Subbagian Fertilitas dan Endokrinologi Reproduksi Bagian Obstetri dan Ginekologi Fakultas Kedokteran Unpad
- [30] Gannon, A.M. 2013. Exposure To Cigarette Smoke And Its Impact On The Ovarian Follicle Population Mechanisms Of Follicle Loss. Canada: McMaster University.
- [31] Roche, J.F. 1998. Controls of Folliculogenesis. *J Theriology.* 49: 457-467.
- [32] Peters, A.R. 1985. Hormonal Control of Bovine Oestrous Cycle 1. The Natural Cycle. *Br. Vet. J. Reprod. Fertil.* 14: 546-573
- [33] Rochler, M.M. 1993. Insulin-like Growth Factor Binding Protein, Vitamins and Hormone. *J. Reprod. Fertil.* 47: 1-114