ISSN : 2088-0197

Indonesian Journal of Cancer Chemoprevention

Vol.1, No.1, 2010





The Official Journal of
Indonesian Society of Cancer Chemoprevention

Indonesian Journal of Cancer Chemoprevention, 2010, 1(1)
ISSN : 2088-0197

ISCC

Indonesian Journal of Cancer Chemoprevention

Editor in Chief
Dr. Arief Nurrochmad, M.Si., M.Sc., Apt.
Vice
Dr. Agung Endro Nugroho, M.Si., Apt.
Secretary
Dr.rer.nat. Rr. Endang Lukitaningsih, M.Si., Apt.

Editors
Dr.Med.dr. Indwiani Astuti
Dr. Erna Prawita Setyowati, M.Si., Apt.
Dr. Muhammad Da'i, M.Si., Apt.

Board Reviewing Editors
Prof. Dr. Edy Meiyanto, M.Si., Apt.
Prof. drg. Ferry Sandra, D.D.S., Ph.D., L.F.I.B.A.
Prof. Dr. Sugiyanto, SU., Apt.
Prof. Dr. Sukardiman, SU., Apt.
Prof. Dr. Adek Zamrud Adnan, M.Sc., Apt.
Prof. Dr. Sismindari, SU., Apt.
Prof. Masashi Kawaichi, MD., PhD.
Prof. Dr. Hiroshi Itoh

Business Manager
Adam Hermawan, M.Sc., Apt.
Muthi Ikawati, M.Sc., Apt.

Lay outer and setting
Gatot Sadewo, AMd.

Editorial Office
Faculty of Pharmacy, Universitas Gadjah Mada Yogyakarta
Sekip Utara, Yogyakarta, Indonesia 55281
Telp. (+62-274) 6492662, Fax.(+62-274) 543120
Email : iscc2010@ymail.com
Bank : Bank BNI
Branch : Cabang UGM, Yogyakarta
Account No. : 0205640096
Account Name : Muthi Ikawati

Published by
Indonesian Society for Cancer Chemoprevention (ISCC)

Cover Picture :
"Molecular structure of **5** and **6**"
(see Maulana et al., p 1-11)

Indonesian Journal of Cancer Chemoprevention, 2010, **1**(1)
ISSN : 2088-0197

ISCC

INDONESIAN JOURNAL OF CANCER CHEMOPREVENTION

ARTICLES

Platinum Metal Complexes Of *Carbaboranylphophines*: Potential Anti Cancer Agents
(Ilham Maulana, Peter Loennecke, and Evamarie Hey-Hawkins) 1-11

Structure Modification of Ethyl *p*-methoxycinnamate Isolated from *Kaempferia galanga* Linn. and Citotoxicity Assay of The Products on WiDr cell
(Juni Ekowati, Marcellino Rudyanto, Shigeru Sasaki, Tutuk Budiati, Sukardiman, Adam Hermawan and Edy Meiyanto) 12-18

Kaempferia galanga L. Rhizome as a Potential Dental Plaque Preventive Agent
(Triana Hertiani, Sylvia Utami Tunjung Pratiwi, Iramie Duma Kencana Irianto and Aini Febriana) 19-25

Synergistic Combination Of Ciplukan (*Physalis angulata*) Herbs Ethanolic Extract and Doxorubicin on T47D Breast Cancer Cells
(Inna Armandari, Kartika Dyah Palupi, Sofa Farida, Adam Hermawan, Ratna Asmah Susidarti and Edy Meiyanto) 26-31

Leunca (*Solanum nigrum* L.) Herbs Ethanolic Extract Increase Cytotoxic Activity of Cisplatin on Hela Cervical Cancer Cells
(Raditya Prima Istiaji, Maya Fitria, Larasati, Fortunella Tjondro, Astrid Ayu Maruti, Erna Prawita Setyowati and Edy Meiyanto) 32-37

Antioxidant Activity Test Of 2,6-Bis-(2'-Furilidyn)-Cyclohexanone, ; 2,5-Bis-(2'-Furilidyn)-Cyclopentanone,; 1,5-Difuryl-1,4-Pentadien-3-One
(Ismi Rahmawati, Endang Sri Rejeki and Sardjiman) 38-42

Curcumin And Extract Of *Plantago major*,' L Increased SPF Value of Cold Cream Base
(Nining Sugihartini) 43-47

The Effect of Addition of Antioxidant to The Stability of Green Tea Water Extract As Anti Acne
(Nadya Fadila Saib) 48-53

Potency of Industrial Tea Waste: Comparison Between Green and Black Tea Industrial Wastes as UV Filter for Sunscreen
(Yohanes Martono) 54-59

The Influence of DMBA (7,12-dimethylbenz-[a]anthracene) Regimen in the Development of Mammae carcinogénesis on *Sprague dawley* Female Rat
(Agung Eru Wibowo, Sriningsih, Puspita Eka Wuyung, Raafqi Ranasasmita) 60-66

Indonesian Journal of Cancer Chemoprevention, 2010, 1(1):12-18
ISSN : 2088-0197

Structure Modification of Ethyl *p*-methoxycinnamate Isolated from *Kaempferia galanga* Linn. and Citotoxicity Assay of The Products on WiDr Cells

Juni Ekowati[1]*, Marcellino Rudyanto[1], Shigeru Sasaki[3], Tutuk Budiati[1], Sukardiman[2], Adam Hermawan[4], Edy Meiyanto[4]

[1]Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Airlangga University, Jl. Dharmawangsa Dalam, Surabaya 60286, Indonesia
[2]Department of Phytochemistry and Pharmacognosy, Faculty of Pharmacy, Airlangga University, Jl. Dharmawangsa Dalam, Surabaya 60286, Indonesia
[3]Institute of Medicinal Chemistry, Hoshi University, Ebara 2-4-41, Shinagawa, Tokyo 152-8501, Japan
[4]CCRC, Gadjah Mada University, Jl. Sekip Utara, Jogyakarta, 55281 Indonesia

Abstract :

Ethyl *p*-methoxycinnamate, major ingredient of *Kaempferia galanga* rhizome, have been reported not only has analgesic – anti inflammatory activities like NSAIDs which inhibited cyclooxygenase, but also inhibit tumor cell proliferation in specimen of mouse epidermis. Therefore, it will be interesting to carry out synthetic studies on the derivates of ethyl *p*-methoxycinnamate and searching their citotoxic activity on WiDr cell. We wish to report of structure modification on carboxyl moiety of ethyl *p*-methoxycinnamate and evaluation on their citotoxic activity on WiDr cell. Isolation of ethyl *p*-methoxycinnamate from *Kaempferia galanga* rhizome was carried out by percolation with ethanol 96% as solvent. Hydrolysis of ethyl *p*-methoxycinnamate in basic condition was performed to obtain *p*-methoxycinnamic acid. Preparation of some thiourea derivates of ethyl *p*-methoxycinnamate was carried out by microwave irradiation. Citotoxicity assay was carried out by MTT method for 48 h.

Modification of carboxyl group of ethyl *p*-methoxycinnamate to its thiourea form could be carried out by microwave irradiation gave; (*E*)-3-(4-methoxyphenyl)-*N*-(phenylcarbamothioyl)acrylamide (50%); (*E*)-3-(4-methoxyphenyl)-*N*-(4-methoxyphenylcarbamothioyl)acrylamide (26%) and (*E*)-3-(4-methoxyphenyl)-*N*-(4-methylphenylcarbamothioyl) acrylamide (54%), yield calculated for 2 step from the acid chloride. All compounds showed no citotoxic effect on WiDr cell at 48 h incubation.

Keywords : ethyl *p*-methoxycinnamate, microwave irradiation, *Kaempferia galanga*, citotoxicity, WiDr cell

INTODUCTION

Colon cancer or colorectal cancer, is one of type cancer that causes the highest mortality, in addition to beside of breast cancer in women, prostate cancer in men, lung and breast (NCI, 2007). Surgery therapy of colorectal cancer has surgery side effect : infection, anastomosis leakage, obstruction, and malabsorption syndrome. Chronic side effects also usually appear after several months of stopping radiotherapy, including persistent diarrhea, enteritis or proctitis, and wounds that never healed (Medina & Davis, 2005; Wells, 2006).

Generally, In addition chemotherapy agents have also some typical side effects, including hair loss because of the effects on hair follicles, nausea and vomiting due to stimulation of chemoreceptors prostrema area, diarrhea caused by the effects on the epithelium of the gastrointestinal tract, bone marrow depression in the back that cause neutropenia, thrombocytopenia, and anemia, and fertility disorders (Chabner et al, 2006; Lűllman et al., 2000).

The emergence of serious side effects of cancer treatment has encouraged continued development of research looking for new compounds from natural and synthetic materials as anti cancer.

*Corresponding author email : j_ekowati@yahoo.com

Ethyl *p*-methoxycinnamate (EPMS), the largest component of the rhizome essential oils kencur (Kaempferia galanga Linn.) rhizome has been reported to have analgesic and anti-inflammatory activity (Sadono & Hasmono, 2001), inhibitor for cyclooxygenase by docking study (Ekowati et al, 2010), inhibitor of tumor promoter teleocidin B-4-induced Epstein-Barr virus (EBV), inhibitor ornitin decarboxylase in the epidermis and skin papillomas of mice (Xue & Chen, 2001; Vimala et al, 1999) and cytotoxic to Artemia salina by brine shrimp lethality test (Tewtrakul et al., 2005). Those finding give an idea to the author to take advantage of EPMS of kencur rhizome as a starting material for synthesising of compounds that have a cytotoxic effect on colon cancer cells. For example, WiDr cells.

METHOD

Synthesis

Starting from ethyl *p*-methoxycinnamate **(1)** from *Kaempferia galangal* Linn., the *p*-methoxycinnamoyl isothiocyanates (**4**) were synthesized through the reaction of *p*-methoxy cinnamoyl chloride (**3**) and powder ammonium thiocyanate. The final compounds (**6a-6c**) were obtained by the reaction of some primary amines (**5a-5c**) and *p*-methoxycinnamoyl thiocyanates (**4**). (Fig. 1).

The reaction mixture is was irradiated under microwave 140W. The chemical structures of the synthesized compounds were confirmed by ^{1}H-NMR, ^{13}C-NMR, IR and HRMS spectral data, the purity were ascertained by melting point and TLC tests.

Ethyl *p*-methoxycinnamate was isolated from *Kaempferia galanga* Linn. under the known method [10]. *Kaempferia galanga* was collected from Purwodadi Botanical Garden. All reagents and solvent were purchased from standard commercial suppliers. Melting points were measured with a Electrothermal melting point apparatus without correction. IR spectra were recorded in KBr on Jasco FT-IR 5300, and major absorption was listed in cm^{-1}. ^{1}H-NMR and ^{13}C-NMR spectra were ^{1}obtained on BRUKER instrument, and chemical shift were reported in ppm on the δ-scale from internal Me_4Si. MS spectra were measured with a JEOL JMS 600 spectrometer by using the EI methods. TLC was carried out on glass plates coated with silica gel F_{254} (Merck). Spot detection was performed with UV 254 nm.

Structure Modification of ethyl *p*-methoxycinnamate Isolated from *Kaempferia galangal* Linn.

Ethyl p-methoxycinnamate in 5% KOH/ethanol solution was heated for 2 hours by using water bath, then and acidified with HCl to produce p-methoxycinnammic acid. The crude product was purified by recrystalization using certain solvent and determined its melting point. Mix of p-methoxycinnamic acid in dry benzene and one drop of pyridine, was then added a 5-fold excess of thionyl chloride was added. Care was taken to trap the formation of _HCl vapour formed. The mixture was refluxed overnight, the solvent and excess thionyl chloride were removed by rotary evaporation. The addition of benzene and evaporation process were repeated several times to remove the last traces of thionyl chloride to give a dark yellow solid. p-methoxycinnamoyl chloride was used in the next reaction without purification. After that following the process, powdered ammonium thiocyanate, appropriate p-methoxycinnamoyl chloride, PEG-400 and dichlo-rometane (12.5 ml) were placed in a dried Erlenmeyer flask and irradiated under microwave at 140 W. Then the appropriate amines (aniline, p-toluidine and p-anisidine) was then added and the mixture was irradiated under microwave 140 W. The mixture was filtered off to remove inorganic salts and the filtrate was concentrated under reduced pressure. The resulting solid was obtained after removing the dichlormethane, then

it was recrystalised using ethanol to give p-methoxycynnamoyl thiourea compounds (Fig.1).



Fig. 1. The schematic representation of compounds 1-6

Cytotoxicity Assay

The cytotoxic potencies of the prepared compounds prepared in this study were assessed using a standardized protocol (Reff). Tests were carried out by incubating 5x103 WiDr cells in 96 well plate for 24 hours for adaption, then followed by treatment with the serial concentration of the sample, and further incubatedtion again for 24 hours as described in the research procedure (Reff). Compounds were evaluated at concentrations of 5 µM, 10 µM, 20 µM, 50 µM, 100 µM dan 150 µM. The viability of cells obtained from the conversion of absorbance values formazan formed by MTT treatment as described in the research procedure (Reff). Profile of cell viability is presented of mean ± standard deviation (SD) of 3 experiments.

RESULTS AND DISCUSSION

Structure modification of ethyl *p*-methoxycinnamate

Transformation ethyl *p*-methoxycinnamate (1) to *p*-methoxycinnamic acid (2).

p-methoxycinnamic acid (yield 80%) as white crystal (m.p. 169°C). HRMS m/z EI, 178 (M^+). ^{1}H NMR (DMSO) 3.78 (3H, s), 6.60 (1H, d, J = 16 Hz), 6.96 (2H, d, J = 5.0 Hz), 7.53 (1H, d, J = 16 Hz), 7.62 (2H, d, J = 5.0 Hz). ^{13}C NMR (DMSO) 55.5 ppm, 114.54 ppm, 116.71 ppm, 127.02 ppm, 130.13 ppm, 143.92 ppm, 161.12 ppm, 168.03 ppm, IR (KBr) 2937, 2843, 2567, 1685, 1624, 1288, 1255 cm^{-1}. Calculated Mass $C_{10}H_{10}O_3$ 178.0630. Measured Mass 178.0617.

Transformation *p*-methoxycinnamic acid (2) to *p*-methoxycinnamoyl chloride (3)

p-methoxycinnamoyl chloride (yield 90%) as yellow solid (m.p. 51°C). HRMS m/z EI, 196 (M^+). ^{1}H NMR ($CDCl_3$) 3.87 (3H, s), 6.51 (1H, d, J = 15.2 Hz), 6.96 (2H, d, J = 5.0 Hz), 7.53 (1H, ^{2}d, J = 16 Hz), 7.62 (2H, d, J = 5.0 Hz). ^{13}C NMR (DMSO) 55.5 ppm, 114.75 ppm, 116.70 ppm, 127.02 ppm, 130.14 ppm, 143.94 ppm, 161.13 ppm, 168.02 ppm. Calc. Mass $C_{10}H_9O_2Cl$ 196.0291. Measured Mass 196.0276.

Synthesis of (*E*)-3-(4-methoxyphenyl)-*N*-(phenylcarbamothioyl)acrylamide (6a) (yield 50%) as pale green crystal (m.p. 200°C). MS m/z EI, 312 (M^+). ^{1}H NMR ($CDCl_3$) 3.87 (3H, s), 6.31 (1H, d, J = 15.60 Hz), 7.30 (1H, d, J = 15.60 Hz), 7.71 (2H, d, J = 9.60 Hz), 6.94 (2H, d, J = 9.60 Hz), 7.50 (2H, d, J = 9.60 Hz), 7.41 (2H, t, J = 6.80 Hz), 7.28 (1H, t, J = 10.80 Hz), 8.69 (1H, s), 12.63 (1H,s). IR (KBr) 3222, 3031, 1671, 1591, 1537, 1244, 1149, 826 cm^{-1}. ^{13}C NMR (DMSO) 55.46 ppm, 114.96 ppm, 124.16 ppm, 126.37 ppm, 126.79 ppm, 128.86 ppm, 130.41 ppm, 137.68 ppm, 146.45 ppm, 162.19 ppm, 166.02 ppm, 178.66 ppm. Calc. Mass $C_{16}H_{15}NO_3$ 312.0932. Measured Mass 312.0928.

Synthesis of (*E*)-3-(4-methoxyphenyl)-*N*-(4-methylphenylcarbamothioyl)acrylamide (6b) (yield 54%) as pale yellow crystal (m.p. 165°C) MS m/z EI, 326 (M^+). ^{1}H NMR ($CDCl_3$) 2.37 (3H, s), 3.87 (3H, s), 6.31 (1H, d, J = 15.40 Hz), 6.93 (2H, d, J = 9.60 Hz), 7.21 (2H, d, J = 8Hz), 7.78 (1H, d, J = 15.40 Hz), 7.54 (4H, m), 8.68 (1H, s), 12.51 (1H,s). IR (KBr) 3224, 3025, 1672, 1590, 1537, 1248, 1151, 827 cm^{-1}. ^{13}C NMR (DMSO) 55.43, 114.51, 115.53, 124.25, 126.42, 129.44, 130.39, 135.10, 146.27, 162.12, 166.08, 178.78 ppm. Calc. Mass $C_{16}H_{15}NO_3$ 326.1095. Measured Mass 326.1089.

Synthesis of (*E*)-3-(4-methoxyphenyl)-*N*-(4-methoxyphenylcarbamothioyl)acrylamide (6c).
(yield 26%) as pale yellow crystal (m.p. 181°C) MS m/z EI , 342 (M^+). ^{1}H NMR (DMSO) 3.76 (3H, s), 3.80 (3H,s), 6.86 (1H, d, J = 15.80 Hz), 6.95-7.03 (4H, m), 7.51-7.59 (4H, m), 7.70 (1H, d, J = 15.80 Hz), 11.43 (1H, s), 12.58 (1H,s). ^{13}C NMR (DMSO) 55.49, 55.61, 114.01, 114.84, 117.23, 125.92, 126.88, 130.33, 130.94, 144.60, 157.59, 161.65, 166.83, 179.23 ppm. IR(KBr) 3235, 3034, 1673, 1593, 1509, 1252, 1150, 825 cm^{-1}. Calc. Mass $C_{16}H_{15}NO_3$ 342.1038. Measured Mass. 342.1031.

Unlike reported reaction condition where some thiourea preparation was at room temperature (Xu et al.,2003), in this research Unlike reported the reaction condition to prepare some thiourea compounds which in room temperature (Xu et al, 2003), preparation of thiourea compounds from EPMS was done by on microwave irradiation. In 2004, Wei et al successfully developed methods of synthesis of *N*-phenyl-*N'*-benzoylthiourea with microwave irradiation method. In this process synthesis of compounds *N*-phenyl-*N'*-benzoylthiourea the reaction takes about 10 minutes with the percentage yield about results obtained around 98%.

Microwave irradiation increases the speed of reaction and increase the reaction with a small side reaction products. The situation can be explained by an influence of thermal heat and / or non-thermal effects or specific microwave effect. Heating by microwave irradiation involves two mechanisms namely dipole rotation and ionic conduction. Absorption of microwave energy occurs when a rotating dipole molecules align themselves with the electric field components or ions move back and forth because the same phenomenon. When molecules or ions rotate back and forth in the matrix, it will arise because the heat caused by friction (Mavandadi, 2004).

Preparation of some thiourea derivates of EPMS from kencur rhizome, i.e. compound 6a, 6b and 6c (Fig.1) takes place in two stages. First step is nucleophylic substitution reaction between p-metoksisinamoil ammonium chloride with thiocyanate. The nucleophilic substitution reaction is a substitution reaction of a nucleophile which has a pair of free electrons (atoms N in ammonium thiocyanate) on the C atom of *p*-methoxycinnamoyl chloride that binds chlorine atom. The reaction will release the halide ions of the acyl halide, the chloride ion, so that, it will be generated to become *p*-methoxycinnamoyl isothiocyanate. In the second phase step, occurs is the addition of nucleophiles, where the nucleophile is the N atom on aniline (5a), *p*-methylaniline (5b) and *p*-methoxyaniline (6c), this will attacks the C atom bound of *p*-methoxycinnamoyl isothiocyanate to atom N, produced (*E*)-3-(4-methoxyphenyl)-*N*-(phenylcarbamothioyl)acrylamide,(*E*)-3-(4-methoxyphenyl)-*N*-(4-methylphenylcarbamothioyl)acrylamide and (*E*)-3-(4-methoxyphenyl)-*N*-(4-methoxyphenylcarbamothioyl)acrylamide.

Differences of amine functional groups on compounds that are used, to can give different percentage results obtained. This was because of the differences in the nature of these functional groups, methyl substituents (*-p*) on aniline compound is an electron donating substituent allowing the nucleophilic addition of amine compounds on the *p*-methoxycinnamoyl isothiocyanate. Substituent methoxy (*-p*) on

aniline compound is also donating an electron, but in this study the lowest percentage of the results. This is probably due to the high solubility of this compound (6c) in ethanol, used during recrystallization.

Cytotoxic Assay for WiDr Cells

Cytotoxicity test was done to confirm the cytotoxic ability of 3 compounds derivates of ethyl *p*-methoxycinnamate against WiDr cells. Cytotoxic test can be carried out used by colorimetric method, which is based on the ability of mitochondrial dehydrogenase enzymes to convert MTT (3 - (4,5-dimetiltiazol-2-yl) -2.5 diphenyl tetrazolium-bromid), a yellow substrate that is not soluble in water, to a dark blue formazan that is not soluble in water and attached to the cells (Doyle and Griffiths, 2000).

Formazan formed is proportional to the number of living cells, since the reduction MTT into formazan can only be done by living cells. (Mosmann, 1983). The method is rapid, sensitive, accurate and can be used for test samples in large numbers automatically using spectrophotometer ELISA Reader (Doyle and Griffiths, 2000). This method also proved more reliable than cell calculations using hemositometer (Freimoser et al., 1999).



Fig. 2. Morphology WiDr cell after incubated with sample 6a, 6b and 6c (up to 150 µM for 24 h by inverted microscope (400x). No difference between control cell WiDr and those sample (6a, 6b and 6 c).



Fig. 3. Effect of sample treatment 6a, 6b and 6c on viability WiDr cell for 24 h (A) and 48h (B).

Table 1. IC $_{50}$ value of Samples

Compounds	$IC_{50}(\mu M)$
6a	-
6b	-
6c	-



Fig. 5. MTT reduction reaction to formazan by succinate dehydrogenase enzyme (Mosmann, 1983).

Reduction MTT into formazan can be done only by living cells. Thus, the absorbance of formazan formed is proportional to the population viability of cells (Mossman, 1983). There are variation of the enzymatic capability is expected to differ in each cell is still alive. Solvents used were DMSO. The highest DMSO concentration used in this study was 0.12% (v / v), has no effect on the test cell.

WiDr cell is one of cells that has a low sensitivity towards treatment with 5-fluorouracil (5-FU), antimetabolite class of chemotherapeutic agents. Transfection of _WiDr with normal p53 did not cause increasing sensitivity to 5-FU (Giovannetti et al., 2007). WiDr cell resistance against 5-FU one of which is mediated by the increased expression of the enzyme thymidylate synthetase, which is the main inhibitory target of 5-FU (Sigmond et al., 2003). However, P-glycoprotein (PGP) in WiDr cells did not too express high, so that there might be other mechanisms that facilitate resistance to 5-FU (Jansen, 1997).

The existence of the relationship between dose and percent of the living cell occurs if a longer incubation period was carried out, so that the activity was seem to be was carried out , so that the activity was seem to be (time-dependent). IC50 is the parameter used to demonstrate the anticancer potential of a test substance. IC50 of a test substance can be calculated if there is a correlation between levels with the percentage of living cells. IC50 of the test substance is still more likely to be developed as a cancer drug is equal to 100 ug / ml (Ueda et al, 2002).

CONCLUSION :

1. Modification of carboxyl group of ethyl *p*-methoxycinnamate to its thiourea form can be carried out by microwave irradiation to give ;
   a.(*E*)-3-(4-methoxyphenyl)-*N*-(phenylcarbamothioyl)acrylamide (6a) (50%)
   b.(*E*)-3-(4-methoxyphenyl)-*N*-(4-methoxyphenylcarbamothioyl)acrylamide (6b) (26%)
   c.(*E*)-3-(4-methoxyphenyl)-*N*-(4-methylphenylcarbamothioyl)acrylamide (6c) (54%)
2. All compounds showed no cytotoxic effect for WiDr cell after treatment for 24 and 48 h.
3. Substituent -phenyl, -methylphenyl *(-p)* and –methoxyphenyl *(-p)* on compounds (6a-c) have no influence on viability WiDr cells.

REFFERENCES

Chabner, B.A., Amrein, P.C., Druker, B.J., *et al.*, 2006. Chemotherapy of Neoplastic Diseases. *In* : Brunton, L.L., Lazo, J.S., Parker K.L., *Goodman and Gilman's the Pharmacological Basis of Therapeutics*, 11th Ed, New York : The McGraw- Hill Co., Inc. Chapter 51.

Doyle, A., *and* Griffiths, J.B., 2000, *Cell and Tissue Culture for Medical Research*, John Willey and Sons Ltd, New York.

Freimoser, F.M., Jakob, C.A., Aebi, M., *and* Tuor, U., 1999, The MTT {3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide} Assay Is a Fast and Reliable Method for Colorimetric Determination of Fungal Cell Densities, *Applied and Environmental Microbiology*, **65**, (8), 3727-3729.

Giovannetti, E., Backus, H.H.J., Wouters, D., Ferreira, C.G., van Houten, V.M.M. and Brakenhoff, R.H., 2007, Changes in the Status of p53 Affect Drug Sensitivity to Thymidylate Synthase (TS) Inhibitors by Altering TS Levels, *British J. Can.*, **96**:769-775.

Lullman, H., Mohr, K., Zielger, A., Bieger, D., 2002. *Color Atlas of Pharmacology*, 2nd Ed, New York : Thieme Stuttgard., p.296-299.

Mavandadi, F., Lidstrom, P., 2004. Microwave-Assisted Chemistry in Drug Discovery. *Medical Chemistry*. Vol.4, hal. 773-792.

Medina, P.J., Davis, L.E., 2005. *Colorectal Cancer. In* : Dipiro, J.T., Talbert, R.L., Matzke G.R., Wells, B.G., Posey, L.M (Eds). *Pharmacotherapy A Pathofisiologic Approach*, 6th Ed, New York : The McGraw-Hill Co., Inc. p.2383-2415.

Mosmann, T., 1983, Rapid Colorimetric Assay for Cellular Growth & Survival: Application to Proliferation & Cytotoxicity Assays, *Journal of Immunological Method*, **65**, 65-59.

NCI fact sheet 2007. What is cancer? accessed from http://www.nci.org October 2008

Sigmond, J., Backus, H.H., Wouters, D., Temmink, O.H., Jansen, G. and Peters, G.J., 2003, Induction of Resistance to the Multitargeted Antifolate Pemetrexed (ALIMTA) in WiDr Human Colon Cancer Cells is Associated with Thymidilate Synthase Overexpression, *Biochem. Pharmacol.* **25**, 12, p189-201.

Tewtrakul S., Yuenyongsawad S., Kummee S. and Atsawajaruwan L. 2005. Chemical component and biological activities of volatile oil of *kaempferia galanga* Linn. *Songklanakarin J.Sci.Technol.* **27** (Suppl.2) : Thai herbs.pp.504-507.

Ueda, J.Y., Tezuka, Y., Banskota., A.H., Tran, Q.L., Tran, Q.K., Harimaya, Y., Saiki, I., dan Kadota, S, 2002, Antiproliferative Activity of Vietnamese Medicinal Plants, *Biol. Pharm. Bull.* **25** (6): 753-760.

Vimala S, Norhanom AW, and Yadav, 1999. Antitumor promoter activity in Malaysian ginger rhizobia used in traditional medicine.*Br. J. Cancer.* **80** (1-2) : pp. 110-116.

Wells, B.G., DiPiro, J., Schwinghammer, T., Hamilton, C., 2006. *Pharmacotherapy Handbook*, 6th Ed, Boston : The McGraw-Hill Co. pp.614-621.

Wei, T.B., Lin, Q., Zhang, Y.M., and Wei, W., 2004. Microwave Promoted Efficient Synthesis of *N*-Aryl-N'-Aroyl thiourea under Solvent-Free and Phase Transfer Catalysis Conditions. *Journal of Synthesic Communication*, **34** (1), 181-186

Xue Y, Chen H. 2001. *Abstracts of the XVIth EUCARPIA Congress.Edinburg. Scotland.*

Xu, X., Qian, X., Li, Z., Huang, Q., Chen, G. 2003. *J. of Fluorine Chemistry*. **121** : pp. 51-54

Journal Profile

Indonesian Journal of Cancer Chemoprevention

eISSN : 23558989 | pISSN :

Health Science

Indonesian Society for Cancer Chemoprevention

sinta

S2

Sinta Score

GARUDA

Indexed by GARUDA

6

H-Index

6

H5-Index

242

Citations

214

5 Year Citations



Penerbit:

Indonesian Society for Cancer Chemoprevention

Website | Editor URL

Re: LoA P2

From: juni ekowaty (j_ekowati@yahoo.com)

To: iscc2010@ymail.com

Date: Wednesday, October 6, 2010, 12:49 PM GMT+7

Yth. Panitia The International Seminar on Chemoprevention for Health Promotion and Beauty,

Dengan hormat,

Bersama ini kami kirimkan bukti pembayaran keikutsertaan kami pada The International Seminar on Chemoprevention for Health Promotion and Beauty, sebesar Rp 200.000,- (kami adalah mhsw S3 MIPA Unair, KTM terlampir).

Terima kasih atas perhatian yg diberikan.

Hormat kami,

Juni Ekowati
Fakultas Farmasi Univ. Airlangga
(mhsw S3 MIPA Univ. Airlangga)
Surabaya

From: Iscc Indonesia <iscc2010@ymail.com>
To: j_ekowati@yahoo.com
Cc: muthi_ikawati@yahoo.co.id
Sent: Fri, September 24, 2010 2:31:15 PM
Subject: LoA P2

Yth. Peserta The International Seminar on Chemoprevention for Health Promotion and Beauty

Dengan hormat,
Bersama ini kami lampirkan "letter of acceptance" dari abstrak yang telah Bapak/Ibu kirimkan, untuk dapat dipresentasikan dalam seminar 9 Oktober 2010 di Fakultas Farmasi UGM, Jogjakarta.

Bapak/Ibu diharap dapat melakukan pengiriman full paper dari makalah yang akan dipresentasikan, oral maupun poster, paling lambat 29 September 2010, secara on line melalui website kami **http://farmasi.ugm.ac.id/iscc2010**. Makalah direncanakan akan diterbitkan dalam Indonesian Journal of Cancer Chemoprevention (dengan konfirmasi dan persetujuan pemakalah terlebih dulu).

Sekaligus pula melalui webiste tersebut dimohon melakukan **registrasi on line** dengan melengkapi data yang diperlukan, bagi yang belum.

Pembayaran dapat dilakukan melalui transfer ke Bank Mandiri Gedung Magister UGM Yogyakarta, nomor rek. 137-00-0633484-7 an. Dr. Ratna Asmah Susidarti, Apt., paling lambat tanggal 1 Oktober 2010.

Bukti pembayaran dikirimkan via email ke iscc2010@ymail.com atau muthi_ikawati@yahoo.co.id (cp: Muthi' Ikawati) atau melalui facsimile ke 0274-543120 (cp: Anita Ellyana).

Demikian yang disampaikan. Atas perhatian Bapak/Ibu kami ucapkan terima kasih, dan sampai ber temu di Jogjakarta!

Hormat kami,
Panitia

pembayaran ISCC-Juni.jpg
840.8kB

perbaikan naskah u paper ISCC

From: Iscc Indonesia (iscc2010@ymail.com)

To: j_ekowati@yahoo.com

Cc: info@iscc-indonesia.org

Date: Thursday, December 2, 2010, 02:31 PM GMT+7

Yth. Bu Juni Ekowati,

Ada beberapa hal yang reviewer tanyakan mengenai paper Ibu :
1. Mengenai metode pada metode no 2

2.3. Structure Modification of ethyl *p*-methoxycinnamate Isolated from *Kaempferia galangal* Linn.

Ethyl p-methoxycinnamate in 5% KOH/ethanol solution was heated for 2 hours ~~by~~ using water bath, ~~then~~ and acidified with HCl to produce p-methoxycinnammic acid. The crude product was purified by recrystalization using certain solvent and determined its melting point. Mix of p-methoxycinnamic acid in dry benzene and one drop of pyridine, was then added a 5-fold excess of thionyl chloride ~~was added~~. Care was taken to trap the formation of HCl vapour ~~formed~~. The mixture was refluxed overnight, the solvent and excess thionyl chloride were removed by rotary evaporation. The addition of benzene and evaporation process were repeated several times to remove the last traces of thionyl chloride to give a dark yellow solid. p-methoxycinnamoyl chloride was used in the next reaction without purification. ~~After that~~ following the process, powdered ammonium thiocyanate, appropriate p-methoxycinnamoyl chloride, PEG-400 and dichlo-rometane (12.5 ml) were placed in a dried Erlenmeyer flask and irradiated under microwave at 140 W. ~~Then~~ the appropriate amines (aniline, p-toluidine and p-anisidine) was then added and the mixture was irradiated under microwave 140 W. The mixture was filtered off to remove inorganic salts and the filtrate was concentrated under reduced pressure. The resulting solid was recrystalised ~~from~~ using (?)[s1] ethanol to give p-methoxycynnamoyl thiourea compounds (Fig.1).

[s1]mohon konfirmasi rekristalisasi menggunakan ethanol atau tadinya larut dalam ethanol lalu direkristalisasi dgn pelarut lain (?)

2. Bagian pembahasan :

~~Reduction MTT into formazan can be done only by living cells. Thus, the absorbance of formazan formed is proportional to the population viability of cells~~ (Mossman, 1983). ~~However,~~ There are variation of th e ~~is~~ enzymatic capability ~~is expected to differ~~ in each cell ~~is still alive~~. “There might be cells that are still normal, but there are also the cells exposed to toxic effects and not dead. Consideration then that is what underlies the conversion of absorbance data into percent of the cell, rather than as percent cell death. ” [s1] Solvents used were DMSO. The highest DMSO concentration used in this study was 0.12% (v / v), has no effect on the test cell.

WiDr cell morphology due to treatment did not provide significant changes in cell morphology control. Curve relationship with the percent concentration of the test sample proves that life WiDr cell percent WiDr cell growth does not decline as increasing of sample concentration. This is probably due to the resistant WiDr cells, so that the test compounds are not cytotoxic effect on WiDr cells. [s2]

[s1]perlu dikonfirmasi ke Authors maksudnya apa?

[s2]Perlu konfirmasi Authors

Kalimat tdk bunyi →Saya cenderung menghilangkan saja

Mohon dikirimkan revisiannya. Untuk koreksi misstyping akan dihandle oleh redaksi.
Kami tunggu kiriman revisiannya
Terima Kasih,

Salam,

Redaksi Jurnal ISCC

Naskah Jurnal ISCC

From: Iscc Indonesia (iscc2010@ymail.com)

To: ccrcfarmasiugm@yahoo.com; afadlan@chem.its.ac.id; anindita_adhika@yahoo.com; ardhi_quelle@yahoo.com; dianratih.ffup@gmail.com; diniwati@medis.fisika.net; elzariana@yahoo.com; erna_prawita@yahoo.com; fanie_mcganza@yahoo.com; hadna3ana@yahoo.com; hartatis2003@yahoo.com; irmaningsih75@yahoo.com; j_ekowati@yahoo.com; jamilahabbas@yahoo.com; khie_khiong@yahoo.com; kurnia_atini@yahoo.com; laela_uad@yahoo.com; lanhadi@yahoo.com; lindaweni@yahoo.com; mardisaad@gmail.com; Mariau.astra@yahoo.com; mariau.astra@yahoo.com; mo2y_pellucida@yahoo.com; nadyafadila@yahoo.com; noerkh2002@yahoo.com; pyunianto@gmail.com; rinandri@gmail.com; s.soeng@yahoo.com; sondaridewi@yahoo.com; sulaiman_zubair80@yahoo.co.id; syuliani@yahoo.com; tatat_04her@yahoo.com; wahyu_w60@yahoo.com; warsinah@rocketmal.com; yangie.dwi@gmail.com; yanifirda@yahoo.co.id; yohanes_mart@yahoo.co.id

Cc: info@iscc-indonesia.org; muthi_ikawati@yahoo.co.id; lukitaningsih_end@yahoo.com

Date: Wednesday, October 27, 2010, 03:13 PM GMT+7

Kepada
Yth. Bapak/Ibu Peserta Seminar International on Chemoprevention for Health Promotion and Beauty
(Fakultas Farmasi UGM, 9 Oktober 2010)

Dengan hormat,

Berdasarkan review Panitia, maka daftar naskah yang akan diterbitkan dalam

Indonesian Journal of Cancer Chemoprevention

(Indon J. Can. Chemoprev.)

volume 1, nomor 1, 2, 3 dan 4

adalah sebagaimana terlampir (tentatif).

Jurnal akan diterbitkan oleh Organisasi ISCC (Indonesian Society for Cancer Chemoprevention)/PAKKI (Perhimpunan Kemoprevensi Kanker Indonesia), dalam Bahasa Inggris dan volume 1 direncanakan akan didistribusikan Desember 2010.

Bagi peserta yang naskahnya belum diterbitkan dalam volume 1 akan diterbitkan dalam volume berikutnya. Sedangkan bagi peserta yang naskahnya direncanakan akan diterbitkan dalam volume 1 akan menjalani proses review lebih lanjut, yang telah dimulai sejak hari ini, 27 Oktober 2010, dan akan dikomunikasikan kepada masing-masing penulis mulai 5 November 2010. Daftar yang terlampir bersifat tentatif dan dapat berubah jika diperlukan (diutamakan adalah yang telah siap).

Demikian yang kami sampaikan, atas perhatiannya diucapkan terima kasih.

Hormat kami,
Panitia

ISCC journal-daftar naskah volume 1.doc
57kB

Letter of Acceptance of Indonesian Journal of Cancer Chemoprevention.

From: Iscc Indonesia (iscc2010@ymail.com)

To: j_ekowati@yahoo.com

Cc: info@iscc-indonesia.org; muthi_ikawati@yahoo.co.id

Date: Monday, March 7, 2011, 01:52 PM GMT+7

Dear Author,

Please find attachment file for Letter of Acceptance of Indonesian Journal of Cancer Chemoprevention.

Thank you

Yours sincerely,

Arief Nurrochmad, PhD

Editor in Chief

LoA Juni Ekowati.pdf
34.7kB

Indonesian Journal of Cancer Chemoprevention Vol 1 No 1

From: Iscc Indonesia (iscc2010@ymail.com)

To: bowoae@yahoo.com; Andita.Ayu@soho.co.id; sri.handayani_83@yahoo.com; ardhi_quelle@yahoo.com; arief_hf@yahoo.com; indwiani@gmail.com; abulathfi@gmail.com; nur_fauzi84@yahoo.co.id; jamilahabbas@yahoo.com; kurnia_atini@yahoo.com; ptlarissa@yahoo.com; anindita_adhika@yahoo.com; adek_adnan@yahoo.com; agungendronugroho@yahoo.com; daengta007@yahoo.com; witarto@yahoo.com; arry.yanuar@gmail.com; ilham.maulana@gmail.com; ag_yuswanto@yahoo.com; amin.subandrio@ui.ac.id; p.astuti@gmail.com; mo2y_pellucida@yahoo.com; saurriana@yahoo.com; rinandri@gmail.com; umar001@lipi.go.id

Cc: muthi_ikawati@yahoo.co.id; info@iscc-indonesia.org; lukitaningsih_end@yahoo.com; chan_ahsan@yahoo.com; yanifirda@yahoo.co.id; iim_halimah@yahoo.com; kurnia_atini@yahoo.com; pyunianto@gmail.com; wahono@bppt.go.id; chaidir@gmx.de; susidarti@yahoo.com; sriningsih_2202@yahoo.com; zalinarudin@yahoo.com; retnoaji@yahoo.com; ccrcfarmasiugm@yahoo.com; desakdewa2002@yahoo.com; sondaridewi@yahoo.com; dianratih.ffup@gmail.com; diniwati@medis.fisika.net; dwisningsih@yahoo.co.id; prayascittadevi@yahoo.com; yangie.dwi@gmail.com; dahlansugiyanto@yahoo.com; abulathfi@gmail.com

Date: Tuesday, March 8, 2011, 11:23 AM GMT+7

Dear Collegues,

We are from Indonesian Journal of Cancer Chemoprevention. We are pleased to inform you that the first edition of our journal (**Volume 1 No 1**), is now available. The member of Indonesian Society for Cancer Chemoprevention (ISCC) will get the **first free edition**.

Otherwise, you can order the journal by filling the attached order form and send it back.

For the next edition you can also get it by order.

We are looking forward to hear from you.

Fell free to contact us should you have any inquiries regarding to this journal.

Thank you

Best regards,

Arief Nurrochmad, Ph.D
Editor in Chief

Journal Order Form.doc
50kB

B. daftar Isi ISCC Vol 1 No 1.pdf
82.6kB