

NORM ESTIMATES FOR BESSEL-RIESZ OPERATORS ON GENERALIZED MORREY SPACES

by Eridani Eridani

Submission date: 14-Dec-2018 04:54PM (UTC+0800)

Submission ID: 1056964004

File name: mb-004517.pdf (147.15K)

Word count: 3366

Character count: 12483

NORM ESTIMATES FOR BESSEL-RIESZ OPERATORS ON GENERALIZED MORREY SPACES

MOCHAMMAD IDRIS, HENDRA GUNAWAN, Bandung,
ERIDANI, Surabaya

Received May 11, 2017. First published March 7, 2018.
Communicated by Dagmar Medková

Abstract. We revisit the properties of Bessel-Riesz operators and present a different proof of the boundedness of these operators on generalized Morrey spaces. We also obtain an estimate for the norm of these operators on generalized Morrey spaces in terms of the norm of their kernels on an associated Morrey space. As a consequence of our results, we reprove the boundedness of fractional integral operators on generalized Morrey spaces, especially of exponent 1, and obtain a new estimate for their norm.

Keywords: Bessel-Riesz operator; fractional integral operator; generalized Morrey space

MSC 2010: 42B20, 26A33, 42B25, 26D10

1. INTRODUCTION

Integral operators such as maximal operators and fractional integral operators have been studied extensively in the last four decades. Here we are interested in Bessel-Riesz operators, which are related to fractional integral operators. Let $0 < \alpha < n$ and $\gamma \geq 0$. The operator $I_{\alpha, \gamma}$ which maps every $f \in L_{\text{loc}}^p(\mathbb{R}^n)$, $1 \leq p < \infty$, to

$$I_{\alpha, \gamma} f(x) := \int_{\mathbb{R}^n} K_{\alpha, \gamma}(x - y) f(y) dy = K_{\alpha, \gamma} * f(x), \quad x \in \mathbb{R}^n,$$

where $K_{\alpha, \gamma}(x) := |x|^{\alpha-n}(1 + |x|)^{-\gamma}$ is called *Bessel-Riesz operator*, and the kernel $K_{\alpha, \gamma}$ is called *Bessel-Riesz kernel*. The boundedness of these operators on Morrey spaces and on generalized Morrey spaces was studied in [8] and [9].

The first and second authors are supported by ITB Research & Innovation Program 2016.

34

11

Let $1 \leq p < \infty$ and $\varphi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be of class \mathcal{G}_p , that is, φ is almost decreasing (there exists $C > 0$ such that $\varphi(r) \geq C\varphi(s)$ for $r \leq s$) and $\varphi^p(r)r^n$ is almost increasing (there exists $C > 0$ such that $\varphi^p(r)r^n \leq C\varphi^p(s)s^n$ for $r \leq s$). Clearly if φ is of class \mathcal{G}_p , then φ satisfies the *doubling condition*, that is, there exists $C > 0$ such that $C^{-1} \leq \varphi(r)/\varphi(s) \leq C$ whenever $1 \leq rs^{-1} \leq 2$. We define the *generalized Morrey space* $L^{p,\varphi}(\mathbb{R}^n)$ to be the set of all functions $f \in L^p_{\text{loc}}(\mathbb{R}^n)$ for which

$$\|f\|_{L^{p,\varphi}} := \sup_{B=B(a,r)} \frac{1}{\varphi(r)} \left(\frac{1}{|B|} \int_B |f(x)|^p dx \right)^{1/p} < \infty,$$

where $|B|$ denotes the Lebesgue measure of B . (Recall that the Lebesgue measure of $B = B(a, r)$ is $|B(a, r)| = C_n r^n$ for every $a \in \mathbb{R}^n$ and $r > 0$, where $C_n > 0$ depends only on n .)

If $1 \leq p \leq q < \infty$ and $\varphi(r) := C_n r^{-n/q}$, $r > 0$, then $L^{p,\varphi}(\mathbb{R}^n)$ is the classical Morrey space $L^{p,q}(\mathbb{R}^n)$, which is equipped with

$$\|f\|_{L^{p,q}} := \sup_{B=B(a,r)} |B|^{1/q-1/p} \left(\int_B |f(x)|^p dx \right)^{1/p}.$$

Particularly, for $p = q$, $L^{p,p}(\mathbb{R}^n)$ is the Lebesgue space $L^p(\mathbb{R}^n)$.

In [9], we know that for $\gamma > 0$, $K_{\alpha,\gamma}$ is a member of $L^t(\mathbb{R}^n)$ spaces for some values of t depending on α and γ . It follows from Young's inequality (see [3]) that

$$\|I_{\alpha,\gamma} f\|_{L^q} \leq \|K_{\alpha,\gamma}\|_{L^t} \|f\|_{L^p}, \quad f \in L^p(\mathbb{R}^n)$$

whenever $1 \leq p < t'$, $1/q = 1/p - 1/t'$ (where t' denotes the dual exponent of t) and $n/(n + \gamma - \alpha) < t < n/(n - \alpha)$. This tells us that $I_{\alpha,\gamma}$ is bounded from $L^p(\mathbb{R}^n)$ to $L^q(\mathbb{R}^n)$ with $\|I_{\alpha,\gamma}\|_{L^p \rightarrow L^q} \leq \|K_{\alpha,\gamma}\|_{L^t}$. In [8], it is also shown that $I_{\alpha,\gamma}$ is bounded on generalized Morrey spaces but without a good estimate for its norm as on Morrey spaces. We shall now refine the results by estimating the norms of the operators more carefully through the membership of K_{α} in Morrey spaces.

Note that for $\gamma = 0$, $I_{\alpha,0} = I_{\alpha}$ is the *fractional integral operator* with kernel $K_{\alpha}(x) := |x|^{\alpha-n}$. Hardy and Littlewood [6], [7] and Sobolev [13] proved the boundedness of I_{α} on Lebesgue spaces. The boundedness of I_{α} on Morrey spaces is proved by Peetre [12], and improved by Adams [1] and Chiarenza and Frasca [2]. Later, Nakai [11] obtained the boundedness of I_{α} on generalized Morrey spaces, which can be viewed as an extension of Spanne's result. In 2009, Gunawan and Eridani [4] proved the boundedness of I_{α} on generalized Morrey spaces which extends Adams' and Chiarenza-Frasca's results.

In this paper, we give a new proof of the boundedness of $I_{\alpha,\gamma}$ on generalized Morrey spaces. At the same time, an upper bound for the norm of the operators is obtained. As a consequence of our result, we have an estimate for the norm of I_α (from a generalized Morrey space to another) in terms of the norm of K_α on the associated Morrey space. A lower bound for the norm of the operators is discussed in Section 3.

2. THE BOUNDEDNESS OF $I_{\alpha,\gamma}$ ON GENERALIZED MORREY SPACES

We begin with a lemma about the membership of K_α in some Morrey spaces. Note that throughout this paper, the letters C and C_k denote constants which may change from line to line.

Lemma 2.1. *If $0 < \alpha < n$, then $K_\alpha \in L^{s,t}(\mathbb{R}^n)$, where $1 \leq s < t = n/(n - \alpha)$.*

Proof. Let $0 < \alpha < n$. Take an arbitrary $B = B(a, R)$, where $a \in \mathbb{R}^n$ and $R > 0$. For $1 \leq s < t = n/(n - \alpha)$ we observe that

$$\begin{aligned} |B|^{s/t-1} \int_B K_\alpha^s(x) dx &\leq |B(0, R)|^{s/t-1} \int_{B(0, R)} |x|^{(\alpha-n)s} dx \\ &\leq CR^{n(s/t-1)} R^{n(1-s/t)} = C. \end{aligned}$$

By taking the supremum over $B = B(a, R)$ we obtain $\|K_\alpha\|_{L^{s,t}}^s \leq C$. Hence $K_\alpha \in L^{s,t}(\mathbb{R}^n)$. \square

Remark 2.2. For $0 < \alpha < n$ and $\gamma > 0$ we know that $K_{\alpha,\gamma} \in L^t(\mathbb{R}^n)$ for $n/(n + \gamma - \alpha) < t < n/(n - \alpha)$, see [9]. By the inclusion property of Morrey spaces (see [5]) we have $K_{\alpha,\gamma} \in L^t(\mathbb{R}^n) = L^{t,t}(\mathbb{R}^n) \subseteq L^{s,t}(\mathbb{R}^n)$ for $1 \leq s \leq t$ and $n/(n + \gamma - \alpha) < t < n/(n - \alpha)$. Moreover, because $K_{\alpha,\gamma}(x) \leq K_\alpha(x)$ for every $x \in \mathbb{R}^n$, $K_{\alpha,\gamma}$ is also contained in $L^{s,t}(\mathbb{R}^n)$ for $1 \leq s < t = n/(n - \alpha)$.

As a counterpart of the results in [8] and [9], we have the following theorem on the boundedness of $I_{\alpha,\gamma}$ on Morrey spaces. Note particularly that the estimate holds for $p_1 = 1$.

Theorem 2.3. *If $0 < \alpha < n$ and $\gamma \geq 0$, then $I_{\alpha,\gamma}$ is bounded from $L^{p_1,q_1}(\mathbb{R}^n)$ to $L^{p_2,q_2}(\mathbb{R}^n)$ with*

$$\|I_{\alpha,\gamma} f\|_{L^{p_2,q_2}} \leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,q_1}}, \quad f \in L^{p_1,q_1}(\mathbb{R}^n)$$

whenever $1 \leq p_1 \leq q_1 < n/\alpha$, $1/p_2 = 1/p_1 - 1/s'$, and $1/q_2 = 1/q_1 - 1/t'$, with $1 \leq s < t = n/(n - \alpha)$ for $\gamma \geq 0$ or $1 \leq s \leq t$ and $n/(n + \gamma - \alpha) < t < n/(n - \alpha)$ for $\gamma > 0$.

Theorem 2.3 is in fact a special case of the boundedness of $I_{\alpha,\gamma}$ on generalized Morrey spaces, which is stated in the following theorem.

Theorem 2.4. Let $0 < \alpha < n$ and $\gamma \geq 0$. If $\varphi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is of class \mathcal{G}_{p_1} such that $\int_R^\infty \varphi(r)r^{n/t'}-1 dr \leq C\varphi(R)R^{n/t'}$ for every $R > 0$, then $I_{\alpha,\gamma}$ is bounded from $L^{p_1,\varphi}(\mathbb{R}^n)$ to $L^{p_2,\psi}(\mathbb{R}^n)$, where $\psi(r) := \varphi(r)r^{n/t'}$, with

$$\|I_{\alpha,\gamma}f\|_{L^{p_2,\psi}} \leq C\|K_{\alpha,\gamma}\|_{L^{s,t}}\|f\|_{L^{p_1,\varphi}}, \quad f \in L^{p_1,\varphi}(\mathbb{R}^n)$$

whenever $1 \leq p_1 < n/\alpha$ and $1/p_2 = 1/p_1 - 1/s'$, with $1 \leq s \leq t = n/(n - \alpha)$ for $\gamma \geq 0$ or $1 \leq s \leq t$ and $n/(n + \gamma - \alpha) < t < n/(n - \alpha)$ for $\gamma > 0$.

Proof. Suppose that $\gamma > 0$ and all the hypotheses hold. For $f \in L^{p_1,\varphi}(\mathbb{R}^n)$ and $B = B(a, R)$, where $a \in \mathbb{R}^n$ and $R > 0$, write

$$f := f_1 + f_2 := f_{\chi_{\tilde{B}}} + f_{\chi_{\tilde{B}^c}},$$

where $\tilde{B} = B(a, 2R)$ and \tilde{B}^c denotes its complement. To estimate $I_{\alpha,\gamma}f_1$, we observe that for every $x \in B$, Hölder's inequality gives

$$\begin{aligned} |I_{\alpha,\gamma}f_1(x)| &\leq \int_{\tilde{B}} K_{\alpha,\gamma}(x-y)|f(y)| dy \\ &= \int_{\tilde{B}} K_{\alpha,\gamma}^{s/p_2}(x-y)|f(y)|^{p_1/p_2} K_{\alpha,\gamma}^{(p_2-s)/p_2}(x-y)|f(y)|^{(p_2-p_1)/p_2} dy \\ &\leq \left(\int_{\tilde{B}} K_{\alpha,\gamma}^s(x-y)|f(y)|^{p_1} dy \right)^{1/p_2} \\ &\quad \times \left(\int_{\tilde{B}} K_{\alpha,\gamma}^{(p_2-s)/(p_2-1)}(x-y)|f(y)|^{(p_2-p_1)/(p_2-1)} dy \right)^{1/p_2'}. \end{aligned}$$

Meanwhile, we have

$$\begin{aligned} &\int_{\tilde{B}} K_{\alpha,\gamma}^{(p_2-s)/(p_2-1)}(x-y)|f(y)|^{(p_2-p_1)/(p_2-1)} dy \\ &\leq \left(\int_{\tilde{B}} K_{\alpha,\gamma}^s(x-y) dy \right)^{p_2'(1/s-1/p_2)} \left(\int_{\tilde{B}} |f(y)|^{p_1} dy \right)^{p_2'/s}. \end{aligned}$$

Therefore we obtain

$$\begin{aligned} |I_{\alpha,\gamma}f_1(x)| &\leq \left(\int_{\tilde{B}} K_{\alpha,\gamma}^s(x-y)|f(y)|^{p_1} dy \right)^{1/p_2} \\ &\quad \times \left(\int_{\tilde{B}} K_{\alpha,\gamma}^s(x-y) dy \right)^{1/s-1/p_2} \left(\int_{\tilde{B}} |f(y)|^{p_1} dy \right)^{1/s'}. \end{aligned}$$

$$\begin{aligned} &\leq \left(\int_{\tilde{B}} K_{\alpha,\gamma}^s(x-y) |f(y)|^{p_1} dy \right)^{1/p_2} \\ &\quad \times CR^{n(1-s/t)(1/s-1/p_2)+n/s'} \varphi^{p_1/s'}(2R) \|K_{\alpha,\gamma}\|_{L^{s,t}}^{1-s/p_2} \|f\|_{L^{p_1,\varphi}}^{p_1/s'}. \end{aligned}$$

We then take the p_2 th power and integrate both sides over B to get

$$\begin{aligned} &\int_B |I_{\alpha,\gamma} f_1(x)|^{p_2} dx \\ &\leq \int_B \int_{\tilde{B}} K_{\alpha,\gamma}^s(x-y) |f(y)|^{p_1} dy dx \\ &\quad \times (CR^{n(1-s/t)(1/s-1/p_2)+n/s'} \varphi^{p_1/s'}(2R) \|K_{\alpha,\gamma}\|_{L^{s,t}}^{1-s/p_2} \|f\|_{L^{p_1,\varphi}}^{p_1/s'})^{p_2}. \end{aligned}$$

By Fubini's theorem we have

$$\begin{aligned} &\int_B |I_{\alpha,\gamma} f_1(x)|^{p_2} dx \\ &\leq \int_{\tilde{B}} |f(y)|^{p_1} \left(\int_B K_{\alpha,\gamma}^s(x-y) dx \right) dy \\ &\quad \times (CR^{n(1-s/t)(1/s-1/p_2)+n/s'} \varphi^{p_1/s'}(2R) \|K_{\alpha,\gamma}\|_{L^{s,t}}^{1-s/p_2} \|f\|_{L^{p_1,\varphi}}^{p_1/s'})^{p_2} \\ &\leq CR^{n(1-s/t)} \|K_{\alpha,\gamma}\|_{L^{s,t}}^s \int_{\tilde{B}} |f(y)|^{p_1} dy \\ &\quad \times (R^{n(1-s/t)(1/s-1/p_2)+n/s'} \varphi^{p_1/s'}(2R) \|K_{\alpha,\gamma}\|_{L^{s,t}}^{1-s/p_2} \|f\|_{L^{p_1,\varphi}}^{p_1/s'})^{p_2} \\ &\leq CR^{n(1-s/t)+n} \varphi^{p_1} (2R) \|K_{\alpha,\gamma}\|_{L^{s,t}}^s \|f\|_{L^{p_1,\varphi}}^{p_1} \\ &\quad \times (R^{n(1-s/t)(1/s-1/p_2)+n/s'} \varphi^{p_1/s'}(2R) \|K_{\alpha,\gamma}\|_{L^{s,t}}^{1-s/p_2} \|f\|_{L^{p_1,\varphi}}^{p_1/s'})^{p_2} \\ &\leq C|B|\psi^{p_2}(R) \|K_{\alpha,\gamma}\|_{L^{s,t}}^{p_2} \|f\|_{L^{p_1,\varphi}}^{p_2}, \end{aligned}$$

whence

$$\frac{1}{\psi(R)} \left(\frac{1}{|B|} \int_B |I_{\alpha,\gamma} f_1(x)|^{p_2} dx \right)^{1/p_2} \leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\varphi}}.$$

Next, we estimate $I_{\alpha,\gamma} f_2$. For every $x \in B = B(a, R)$ we observe that

$$\begin{aligned} |I_{\alpha,\gamma} f_2(x)| &\leq \int_{\tilde{B}^c} K_{\alpha,\gamma}(x-y) |f(y)| dy \\ &\leq \int_{|x-y| \geq R} K_{\alpha,\gamma}(x-y) |f(y)| dy \\ &= \sum_{k=0}^{\infty} \int_{2^k R \leq |x-y| < 2^{k+1} R} K_{\alpha,\gamma}(x-y) |f(y)| dy \\ &\leq \sum_{k=0}^{\infty} K_{\alpha,\gamma}(2^k R) \int_{2^k R \leq |x-y| < 2^{k+1} R} |f(y)| dy \end{aligned}$$

$$\begin{aligned}
&\leq C \sum_{k=0}^{\infty} K_{\alpha,\gamma}(2^k R)(2^k R)^{n/p_1'} \left(\int_{2^k R \leq |x-y| < 2^{k+1} R} |f(y)|^{p_1} dy \right)^{1/p_1} \\
&\leq C \|f\|_{L^{p_1,\varphi}} \sum_{k=0}^{\infty} K_{\alpha,\gamma}(2^k R)(2^k R)^n \varphi(2^k R).
\end{aligned}$$

For every $k \in \mathbb{Z}$ we have

$$\begin{aligned}
K_{\alpha,\gamma}(2^k R) &\leq C(2^k R)^{-n/s} \left(\int_{2^k R \leq |x-y| < 2^{k+1} R} K_{\alpha,\gamma}^s(x-y) dy \right)^{1/s} \\
&\leq C(2^k R)^{-n/t} \|K_{\alpha,\gamma}\|_{L^{s,t}}.
\end{aligned}$$

Since $\int_R^\infty \varphi(r)r^{n/t'-1} dr \leq C\varphi(R)R^{n/t'}$, we get

$$\begin{aligned}
|I_{\alpha,\gamma}f_2(x)| &\leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\varphi}} \sum_{k=0}^{\infty} (2^k R)^{n/t'} \varphi(2^k R) \\
&\leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\varphi}} \int_R^\infty \varphi(r)r^{n/t'-1} dr \\
&\leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\varphi}} \varphi(R)R^{n/t'} \\
&= C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\varphi}} \psi(R).
\end{aligned}$$

Raising to the p_2 th power and integrating over B we obtain

$$\int_B |I_{\alpha,\gamma}f_2(x)|^{p_2} dx \leq C (\|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\varphi}})^{p_2} \psi^{p_2}(R) |B|,$$

whence

$$\frac{1}{\psi(R)} \left(\frac{1}{|B|} \int_B |I_{\alpha,\gamma}f_2(x)|^{p_2} dx \right)^{1/p_2} \leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\varphi}}.$$

Combining the two estimates for $I_{\alpha,\gamma}f_1$ and $I_{\alpha,\gamma}f_2$ we obtain

$$\frac{1}{\psi(R)} \left(\frac{1}{|B|} \int_B |I_{\alpha,\gamma}f(x)|^{p_2} dx \right)^{1/p_2} \leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\varphi}}.$$

Since this inequality holds for every $a \in \mathbb{R}^n$ and $R > 0$, it follows that

$$\|I_{\alpha,\gamma}f\|_{L^{p_2,\psi}} \leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\varphi}},$$

as desired.

We may repeat the same argument and use Lemma 2.1 to obtain the same inequality for the case where $\gamma = 0$ and $1 \leq s < t = n/(n - \alpha)$. \square

44

Remark 2.5. Theorems 2.3 and 2.4 give us upper estimates for the norm of the Bessel-Riesz operators (from one Morrey space to another). In particular, for $\gamma = 0$ we have an estimate for the norm of the fractional integral operator I_α in terms of the norm of its kernel (on the associated Morrey space), which follows from the inequality

$$\|I_\alpha f\|_{L^{p_2, \psi}} \leq C \|K_\alpha\|_{L^{s, t}} \|f\|_{L^{p_1, \varphi}}$$

15

for $1 \leq p_1 < n/\alpha$ and $1/p_2 = 1/p_1 - 1/s'$, with $1 \leq s < t = n/(n - \alpha)$.

In the following section, we discuss lower estimates for the norm of the operators in terms of the norm of the Bessel-Riesz kernel (on some Morrey spaces).

3. AN ESTIMATE FOR THE NORM OF THE OPERATORS

14

Recall that if $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ are normed spaces and if $T: (X, \|\cdot\|_X) \rightarrow (Y, \|\cdot\|_Y)$ is a linear operator, then the norm of T (from X to Y) is defined as

$$\|T\|_{X \rightarrow Y} := \sup_{f \neq 0} \frac{\|Tf\|_Y}{\|f\|_X}.$$

Knowing that the Bessel-Riesz operator $I_{\alpha, \gamma}$ is a linear operator on Morrey spaces, we would like to estimate the norm of $I_{\alpha, \gamma}$ from a (generalized) Morrey space to another. We obtain the following result.

Theorem 3.1. Let $0 < \alpha < n$, $\gamma \geq 0$, and φ be of class \mathcal{G}_{p_1} where $1 \leq p_1 < n/\alpha$.

If $\varphi(r)r^n$ is almost increasing and for every $R > 0$ we have

- (i) $\int_R^\infty \varphi(r)r^{n/t'-1} dr \leq C_1 \varphi(R)R^{n/t'}$,
- (ii) $\int_0^R \varphi^{p_1}(r)r^{n-1} dr \leq C_2 \varphi^{p_1}(R)R^n$, and
- (iii) $\int_0^R r^{n-1}/\varphi^{s'}(r)r^{n s'} dr \leq C_3 R^n / \varphi^{s'}(R)R^{n s'}$, where $1 \leq p_1 < t$ and $1 < s < t = n/(n - \alpha)$ for $\gamma \geq 0$ or $1 \leq p_1 \leq t$, $1 < s \leq t$, and $n/(n + \gamma - \alpha) < t < n/(n - \alpha)$ for $\gamma > 0$,

then we have

$$C_4 \|K_{\alpha, \gamma}\|_{L^{p_1, t}} \leq \|I_{\alpha, \gamma}\|_{L^{p_1, \varphi} \rightarrow L^{p_2, \psi}} \leq C_5 \|K_{\alpha, \gamma}\|_{L^{s, t}}$$

whenever $1/p_2 = 1/p_1 - 1/s'$ and $\psi(r) := \varphi(r)r^{n/t'}$. In particular, for $\gamma = 0$, $1 \leq p_1 < t$ and $1 < s < t = n/(n - \alpha)$ we have

$$C_4 \|K_\alpha\|_{L^{p_1, t}} \leq \|I_\alpha\|_{L^{p_1, \varphi} \rightarrow L^{p_2, \psi}} \leq C_5 \|K_\alpha\|_{L^{s, t}}$$

whenever $1/p_2 = 1/p_1 - 1/s'$ and $\psi(r) := \varphi(r)r^{n/t'}$.

43

Proof. Suppose that $\gamma > 0$ and all the hypotheses hold. By Theorem 2.4 we already have

$$\|I_{\alpha,\gamma}\|_{L^{p_1,\varphi} \rightarrow L^{p_2,\psi}} \leq C \|K_{\alpha,\gamma}\|_{L^{s,t}}.$$

To prove the lower estimate, put $\varrho(r) := \varphi(r)r^n$. Let $B = B(a, R)$, where $a \in \mathbb{R}^n$ and $R > 0$. By our assumptions on φ we have

$$|B|^{1/t} \psi(R) \left(\frac{1}{|B|} \int_B \varrho^{-s'}(|x|) dx \right)^{1/s'} \leq C \varphi(R) R^{n/s} \left(\int_0^R \frac{r^{n-1}}{\varphi^{s'}(r) r^{ns'}} dr \right)^{1/s'} \leq C.$$

Now take $f_0(x) := \varphi(|x|)$. Here $\|f_0\|_{L^{p_1,\varphi}} \approx 1$. Moreover, one may compute that

$$I_{\alpha,\gamma} f_0(x) \geq \int_{B(x, 2|x|)} K_{\alpha,\gamma}(x-y) f_0(y) dy \geq C K_{\alpha,\gamma}(2x) \varphi(|x|) |x|^n = C \varrho(|x|) K_{\alpha,\gamma}(x)$$

for every $x \in \mathbb{R}^n$. It follows that

$$\|\varrho(|\cdot|) K_{\alpha,\gamma}\|_{L^{p_2,\psi}} \leq C \|I_{\alpha,\gamma} f_0\|_{L^{p_2,\psi}} \leq C \|I_{\alpha,\gamma}\|_{L^{p_1,\varphi} \rightarrow L^{p_2,\psi}}.$$

23

Next, by Hölder's inequality we have

$$\left(\int_B K_{\alpha,\gamma}^{p_1}(x) dx \right)^{1/p_1} \leq \left(\int_B \varrho^{-s'}(|x|) dx \right)^{1/s'} \left(\int_B (\varrho(|x|) K_{\alpha,\gamma}(x))^{p_2} dx \right)^{1/p_2},$$

whence

$$\begin{aligned} |B|^{1/t-1/p_1} \left(\int_B K_{\alpha,\gamma}^{p_1}(x) dx \right)^{1/p_1} &\leq |B|^{1/t} \psi(R) \left(\frac{1}{|B|} \int_B \varrho^{-s'}(|x|) dx \right)^{1/s'} \\ &\quad \times \frac{1}{\psi(R)} \left(\frac{1}{|B|} \int_B (\varrho(|x|) K_{\alpha,\gamma}(x))^{p_2} dx \right)^{1/p_2} \\ &\leq C \|I_{\alpha,\gamma}\|_{L^{p_1,\varphi} \rightarrow L^{p_2,\psi}}. \end{aligned}$$

25

By taking the supremum over $B = B(a, R)$ we conclude that

$$C \|K_{\alpha,\gamma}\|_{L^{p_1,t}} \leq \|I_{\alpha,\gamma}\|_{L^{p_1,\varphi} \rightarrow L^{p_2,\psi}},$$

as desired. The same argument applies for the case where $\gamma = 0$ with $1 \leq p_1 < t$ and $1 < s < t = n/(n - \alpha)$. \square

Remark 3.2. One may observe that the constants C_4 and C_5 in Theorem 3.1 depend on φ , n , p_1 , s , and t , but not on α and γ . Although the lower and the upper bound are not comparable, we may still get useful information from these estimates,

especially for the norm of the operator I_α from $L^{p_1, \varphi}(\mathbb{R}^n)$ to $L^{p_2, \psi}(\mathbb{R}^n)$. Observe that for $1 \leq p_1 < t = n/(n - \alpha)$ we have $\|K_\alpha\|_{L^{p_1, t}}^{p_1} = C/((\alpha - n)p_1 + n) \geq C/\alpha$. Hence, if all the hypotheses in Theorem 3.1 hold for the case where $\gamma = 0$, then we obtain $\|I_\alpha\|_{L^{p_1, \varphi} \rightarrow L^{p_2, \psi}} \geq C/\alpha$, which blows up when $\alpha \rightarrow 0^+$. For $\varphi(r) := r^{-n/q_1}$ with $1 \leq p_1 < q_1 < \min\{s, n/\alpha\}$ and $1 < q_2 < n/(n - \alpha)$, our result reduces to the estimate $\|I_\alpha\|_{L^{p_1, q_1} \rightarrow L^{p_2, q_2}} \geq C/\alpha$, where $1/p_2 = 1/p_1 - 1/s'$ and $1/q_2 = 1/q_1 - \alpha/n$. A similar behavior of the norm of I_α from $L^{p_1}(\mathbb{R}^n)$ to $L^{p_2}(\mathbb{R}^n)$ for $1/p_2 = 1/p_1 - \alpha/n$ when $\alpha \rightarrow 0^+$ is observed in [10], Chapter 4.

Acknowledgement. All authors would like to thank the anonymous referee for his/her careful reading and useful comments on the earlier version of this paper.

References

- [1] *D. R. Adams*: A note on Riesz potentials. *Duke Math. J.* **42** (1975), 765–778. [zbl](#) [MR](#) [doi](#)
- [2] *F. Chiarenza, M. Frasca*: Morrey spaces and Hardy-Littlewood maximal function. *Rend. Mat. Appl.*, VII. Ser. **7** (1987), 679–693. [zbl](#) [MR](#)
- [3] *L. Grafakos*: *Classical Fourier Analysis*. Graduate Texts in Mathematics 249. Springer, New York, 2008. [zbl](#) [MR](#) [doi](#)
- [4] *H. Gunawan, Eridani*: Fractional integrals and generalized Olsen inequalities. *Kyungpook Math. J.* **49** (2009), 31–39. [zbl](#) [MR](#) [doi](#)
- [5] *H. Gunawan, D. I. Hakim, K. M. Limanta, A. A. Masta*: Inclusion properties of generalized Morrey spaces. *Math. Nachr.* **290** (2017), 332–340. [zbl](#) [MR](#) [doi](#)
- [6] *G. H. Hardy, J. E. Littlewood*: Some properties of fractional integrals I. *Math. Z.* **27** (1928), 565–606. [zbl](#) [MR](#) [doi](#)
- [7] *G. H. Hardy, J. E. Littlewood*: Some properties of fractional integrals II. *Math. Z.* **34** (1932), 403–439. [zbl](#) [MR](#) [doi](#)
- [8] *M. Idris, H. Gunawan, Eridani*: The boundedness of Bessel-Riesz operators on generalized Morrey spaces. *Aust. J. Math. Anal. Appl.* **13** (2016), Article ID 9, 10 pages. [zbl](#) [MR](#)
- [9] *M. Idris, H. Gunawan, J. Lindiarni, Eridani*: The boundedness of Bessel-Riesz operators on Morrey spaces. *Int. Symp. On Current Progress in Mathematics and Sciences 2015*. AIP Conf. Proc. vol. 1729, 2016, Article No. 020006. [doi](#)
- [10] *E. H. Lieb, M. Loss*: *Analysis*. Graduate Studies in Mathematics 14. AMS, Providence, 2001. [zbl](#) [MR](#) [doi](#)
- [11] *E. Nakai*: Hardy-Littlewood maximal operator, singular integral operators and the Riesz potentials on generalized Morrey spaces. *Math. Nachr.* **166** (1994), 95–103. [zbl](#) [MR](#) [doi](#)
- [12] *J. Peetre*: On the theory of $\mathcal{L}_{p, \lambda}$ spaces. *J. Functional Analysis* **4** (1969), 71–87. [zbl](#) [MR](#) [doi](#)
- [13] *S. L. Sobolev*: On a theorem of functional analysis. *Am. Math. Soc.*, Transl., II. Ser. **34** (1963), 39–68; Translated from *Mat. Sb.*, N. Ser. **4** (1938), 471–497. [zbl](#) [doi](#)

Authors' addresses: Mochammad Idris, Department of Mathematics, Institute of Technology Bandung, Bandung 40132, Indonesia, e-mail: idemath@gmail.com (*permanent address:* Department of Mathematics, Lambung Mangkurat University, Banjarbaru Campus, Banjarbaru 70714, Indonesia); Hendra Gunawan, Department of Mathematics, Institute of Technology Bandung, Bandung 40132, Indonesia, e-mail: hgunawan@math.itb.ac.id; Eridani, Department of Mathematics, Airlangga University, Campus C Mulyorejo, Surabaya 60115, Indonesia, e-mail: eridani.dinadewi@gmail.com.

NORM ESTIMATES FOR BESSEL-RIESZ OPERATORS ON GENERALIZED MORREY SPACES

ORIGINALITY REPORT

25%

SIMILARITY INDEX

14%

INTERNET SOURCES

23%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** personal.fmipa.itb.ac.id Internet Source **2%**
- 2** Hakim, Denny Ivanal, Eiichi Nakai, and Yoshihiro Sawano. "Generalized fractional maximal operators and vector-valued inequalities on generalized Orlicz–Morrey spaces", *Revista Matemática Complutense*, 2015. Publication **2%**
- 3** mathsoc.jp Internet Source **1%**
- 4** Mitsuru Sugimoto. "A remark on fractional integrals on modulation spaces", *Mathematische Nachrichten*, 10/2008. Publication **1%**
- 5** www.math.cas.cz Internet Source **1%**
- 6** C. Guidorzi, F. Frontera, E. Montanari, F. Rossi, L. Amati, A. Gomboc, C. G. Mundell. "The slope of the gamma-ray burst variability/peak luminosity correlation", **1%**

Monthly Notices of the Royal Astronomical Society, 2006

Publication

7 MAS, Editor. "Modern Applied Science, Vol. 3, No. 2, February 2009, all in one file", Modern Applied Science, 2009. **1%**
Publication

8 www.guliyev.com **1%**
Internet Source

9 Yong Zhong. "Modulation space estimates for the fractional integral operators", Science China Mathematics, 07/2011 **1%**
Publication

10 Yasuo Komori. "Factorization of functions in $H^1(\mathbb{R}^n)$ and generalized Morrey spaces", Mathematische Nachrichten, 04/2006 **1%**
Publication

11 Hendra Gunawan, Eder Kikianty, Christopher Schwanke. "Discrete Morrey spaces and their inclusion properties", Mathematische Nachrichten, 2017 **1%**
Publication

12 www.eudoxuspress.com **1%**
Internet Source

13 openaccess.city.ac.uk **1%**
Internet Source

14 www.grahamkendall.net **1%**
Internet Source

- 15 Zun Wei Fu. " λ -central BMO estimates for commutators of singular integral operators with rough kernels", Acta Mathematica Sinica English Series, 03/2008
Publication 1%
-
- 16 www.m-hikari.com
Internet Source 1%
-
- 17 Operator Theory Advances and Applications, 2016.
Publication 1%
-
- 18 Amjad Hussain, Gui-lian Gao. "Some new estimates for the commutators of n-dimensional Hausdorff operator", Applied Mathematics-A Journal of Chinese Universities, 2014
Publication <1%
-
- 19 Universitext, 2012.
Publication <1%
-
- 20 Tao, Xiangxing, and Sha He. "The boundedness of multilinear operators on generalized Morrey spaces over the quasi-metric space of non-homogeneous type", Journal of Inequalities and Applications, 2013.
Publication <1%
-
- 21 Adam Nowak. "The multi-dimensional pencil phenomenon for Laguerre heat-diffusion maximal operators", Mathematische Annalen, <1%

05/2009

Publication

22

Gunawan, Hendra, and Eridani Eridani. "Fractional Integrals and Generalized Olsen Inequalities", Kyungpook mathematical journal, 2009.

Publication

<1 %

23

Sawano, Yoshihiro, and Hidemitsu Wadade. "On the Gagliardo-Nirenberg type inequality in the critical Sobolev-Morrey space", Journal of Fourier Analysis and Applications, 2013.

Publication

<1 %

24

91.187.98.171

Internet Source

<1 %

25

Gunawan, Hendra, Denny I. Hakim, Kevin M. Limanta, and Al A. Masta. "Inclusion properties of generalized Morrey spaces : Inclusion properties of generalized Morrey spaces", Mathematische Nachrichten, 2016.

Publication

<1 %

26

L. P. Castro. "Finite Interval Convolution Operators with Transmission Property", Integral Equations and Operator Theory, 06/2005

Publication

<1 %

27

fsdona2011.uni-jena.de

Internet Source

<1 %

28

ejde.math.txstate.edu

Internet Source

<1 %

29

Shi, Yanlong, and Xiangxing Tao. "SOME MULTI-SUBLINEAR OPERATORS ON GENERALIZED MORREY SPACES WITH NON-DOUBLING MEASURES", Journal of the Korean Mathematical Society, 2012.

Publication

<1 %

30

Vagif S. Guliyev. "Boundedness of the Maximal, Potential and Singular Operators in the Generalized Morrey Spaces", Journal of Inequalities and Applications, 2009

Publication

<1 %

31

matwbn.icm.edu.pl

Internet Source

<1 %

32

www.acadsci.fi

Internet Source

<1 %

33

elib.mi.sanu.ac.rs

Internet Source

<1 %

34

Guliyev, Vagif S., and Lubomira G. Softova. "Global Regularity in Generalized Morrey Spaces of Solutions to Nondivergence Elliptic Equations with VMO Coefficients", Potential Analysis, 2013.

Publication

<1 %

35

link.springer.com

Internet Source

<1 %

36	kluedo.ub.uni-kl.de Internet Source	<1 %
37	"Computational Intelligence: Soft Computing and Fuzzy-Neuro Integration with Applications", Springer Nature America, Inc, 1998 Publication	<1 %
38	Yoshihiro Sawano. "A Note on Generalized Fractional Integral Operators on Generalized Morrey Spaces", Boundary Value Problems, 2009 Publication	<1 %
39	Advances in Harmonic Analysis and Operator Theory, 2013. Publication	<1 %
40	jsaer.com Internet Source	<1 %
41	www.blacklitterman.org Internet Source	<1 %
42	Guohua Liu. "Boundedness for the commutator of fractional integral on generalized morrey space in nonhomogeneous space", Analysis in Theory and Applications, 03/2011 Publication	<1 %
43	Byun, Sun-Sig, and Lubomira G. Softova. "Gradient estimates in generalized Morrey spaces for parabolic operators : Parabolic	<1 %

operators in generalized Morrey spaces",
Mathematische Nachrichten, 2015.

Publication

44

Seppo Heikkilä. "On extremal solutions of operator equations in ordered normed spaces", *Applicable Analysis*, 4/1992

Publication

45

Fan, Yun. "Boundedness of sublinear operators and their commutators on generalized central Morrey spaces", *Journal of Inequalities and Applications*, 2013.

Publication

46

Analysis Partial Differential Equations and Applications, 2009.

Publication

47

Sawano, Yoshihiro, Denny Ivanal Hakim, and Hendra Gunawan. "Non-smooth atomic decomposition for generalized Orlicz-Morrey spaces : Atomic decomposition for generalized Orlicz-Morrey spaces", *Mathematische Nachrichten*, 2015.

Publication

48

Lars Ehlers. "Strategy-proof allocation when preferences are single-plateaued", *Review of Economic Design*, 06/01/2002

Publication

49

Michael E. Taylor. "Function Space and Operator Theory for Nonlinear Analysis", *Applied Mathematical Sciences*, 2011

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

50

Vagif S. Guliyev. "On the limiting case for boundedness of the B-Riesz potential in B-Morrey spaces", Analysis Mathematica, 06/2009

Publication

<1%

51

V. S. Guliyev. "On the rearrangement estimates and the boundedness of the generalized fractional integrals associated with the Laplace-Bessel differential operator", Acta Mathematica Hungarica, 05/2008

Publication

<1%

52

www.jlimnol.it

Internet Source

<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

NORM ESTIMATES FOR BESSEL-RIESZ OPERATORS ON GENERALIZED MORREY SPACES

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9
