

Rare case of Primary Congenital Glaucoma with Hypoplasia Corpus Callosum

by Prastiya Indra Gunawan

Submission date: 30-Jun-2020 12:20PM (UTC+0800)

Submission ID: 1351644148

File name: case_of_Primary_Congenital_Glaucoma_Ayub_Med_Coll_Abbotabad.pdf (563.31K)

Word count: 1761

Character count: 9469

CASE REPORT

RARE CASE OF PRIMARY CONGENITAL GLAUCOMA WITH HYPOPLASIA CORPUS CALLOSUM

Prastiya Indra Gunawan, Evelyn Komarath*, Nurwasis*, Risa Etika**, Agus Harianto**, Darto Saharso

Division of Neurology, Department of Paediatrics, *Division of Glaucoma, Department of Ophthalmology, **Division of Perinatology, Department of Paediatrics, Airlangga University, College of Medicine, Dr Soetomo Hospital, Surabaya-Indonesia

Primary congenital glaucoma is a rare disease that causes elevated intraocular pressure within the first three years of life. Few studies have explored the association of primary congenital glaucoma with malformation of corpus callosum. We report on a six-month-old female presenting with unilateral primary congenital glaucoma associated with hypoplasia of corpus callosum in Indonesian infant. The patient had already undergone trabeculectomy surgery. However, there no obvious improvement following the procedure given the severity of the condition. **Conclusion:** The failure rate of surgery in severe primary congenital glaucoma conditions is still very high, and therapy can usually preserve vision if early identification of mild or moderate form is made.

Keywords: Primary congenital glaucoma; Hypoplasia of corpus callosum.

Citation: Gunawan PI, Komarath E, Nurwasis, Etika R, Harianto A, Saharso D. Rare case of primary congenital glaucoma with hypoplasia corpus callosum. J Ayub Med Coll Abbottabad 2018;30(2):286-8.

INTRODUCTION

Congenital glaucoma is defined as maldevelopment of some portions of the eye, which may or may not be associated with a systemic anomaly, and which causes elevated intraocular pressure (IOP) within the first three years of life.¹ However, when glaucoma develops in the absence of any secondary cause or related systemic problem, the term primary congenital glaucoma (PCG) is applied.^{2,3} The incidence of PCG is relatively rare, with overall prevalence at 1:10,000 births.¹ Most of the PCG is expressed bilaterally, and asymmetric expression should be suspected in clinically apparent unilateral cases.²

Hypoplasia of the corpus callosum (HCC) exists when all the components of the callosum are present, but the structure is thinner than normal.⁴ Some recent studies have reported some syndrome-associated congenital glaucoma with malformation of corpus callosum.⁵ The objective of this paper is to report a rare case of congenital glaucoma in an Indonesian infant with hypoplasia corpus callosum.

CASE REPORT

A six-month-old female was referred to the Paediatrics and Ophthalmology Department Soetomo Hospital, Surabaya, Indonesia due to protuding left eyeball. There was a bulge in the centre of the left eye since a week before admission (Figure 1-4). Patient was irritable and presenting photophobia as well as blepharospasm. Tears were coming out of her left eye. There was a previous history of trabeculectomy procedure in the left eye at the age of 40 days. The baby was born spontaneously with birth weight 1,000 grams at 28-29 weeks of gestational

age. The Apgar score was 7-8, meconal amniotic fluid was present, but no history of birth trauma. The mother was in healthy condition with good antenatal care. The USG anatomy of vitreoretina was in the normal limit (Figure-5). Examination under anaesthesia is presented in table-1.

The head MRI showed hypoplasia of the corpus callosum (Figure-6, 7). The patient was diagnosed as PCG with descemetocoele on left eye and seclutio of the pupil, with anterior synechia on the right eye. Enucleation or evisceration therapy on the left eye and pupiloplasty for the right eye were considered in order to let the light through the pupil into the retina, and allow the patient to learn to see with her right eye. Unfortunately, the patient's parent refused the procedure.

The patient was six months of age; otherwise, it was three months of corrected age. Evaluation of growth and development using the Denver II chart showed delay in the aspects of fine motor adaptive and social personal.

Table-1: Examination under anaesthesia

Right eye		Left eye
Light perception +	Visual acuity	Decreased of light perception
Normal	IOP	36 mmHg
Hyperaemia -	Conjunctiva	Hyperaemia -
Clear with Φ 10 mm	Cornea	Cloudy with Φ 16 mm, descemetocoele +
Deep	Anterior Chamber	Flat
Anterior synechiae	Iris	Anterior synechiae
Seclutio	Pupil	Seclutio
Difficult to evaluate	Lens	Difficult to evaluate
Fundus reflex +	Fundoscopy	Fundus reflex +
Normal cup and disc ratio		Difficult to evaluate cup and disc ratio




Figure-1: Right eye: seclusion of pupil, anterior synechiae, and normal cornea




Figure-2: Left eye: seclusion of pupil, anterior synechiae, enlargement of a cloudy cornea and descemetocoele



Figure-3: Right eye: examination with calliper revealed corneal diameter was 9 mm



Figure-4: Left eye: examination with calliper revealed corneal diameter was 24 mm




Figure-5: Left eye: The anatomy of vitreoretina was in the normal limit and axial length was 23 mm




Figure-6: The Head MRI coronal view showing hypoplasia of corpus callosum




Figure-7: The Head MRI sagittal view showing hypoplasia of corpus callosum

DISCUSSION

The manifestations of glaucoma in infants are different from those associated with open angle glaucoma in adults. These differences provide the key to early diagnosis. Photophobia, blepharospasm, and epiphora frequently result from the corneal changes that occur secondary to elevated intraocular pressure (IOP) in the

infant eye.⁶ When the condition reaches this stage, the eye is painful and tearing is present. In our case, the baby became sensitive to light and attempted to avoid it by covering the eye or burying the head in a blanket.

The size of the cornea is important in the diagnosis of PCG. Al-Umran KU found that the corneal diameter on preterm infant ranged from 7–7.5 mm to a maximum of 10 mm. The walls of neonatal eyes are distended by an elevated IOP because the corneal and scleral collagen has not yet sufficiently hardened. Before the age of six months, the rigidity of the cornea is less than the sclera; consequently, the cornea enlarges more rapidly under the influence of an increased IOP. The final diagnosis of congenital glaucoma can usually be made only when infant is under general anaesthesia (EUA) by evaluating (1) the IOP with Schiøtz, Perkins or Tono-Pen tonometer, (2) length of the horizontal cornea with calliper, (3) the angle of anterior chamber with gonioscopy, and (4) the fundi (1,2).^{1,2} In this patient, the enlargement of cornea was accompanied by the symptoms of tearing and blepharospasm. The EUA revealed left cornea was larger than normal value. The left cornea was so cloudy that it caused some difficulties in evaluating the Hab's striae and fundi. The cloudy cornea with corneal enlargement could be seen in the first week of life due to increasing of IOP, which reached 36 mmHg (normal IOP was less than 25 mmHg under anaesthesia). The left AXL was longer than normal value because immature collagen remained distensible due to elevated IOP, leading to eye enlargement. Paediatric glaucoma is classified into PCG when caused by congenital anomalies that affect aqueous outflow, and secondary glaucoma when caused by post-birth factors such as ocular trauma, surgery, and inflammation.³ Diagnosis of PCG in this case is made by exclusion of other causes of glaucoma. PCG is better treated with surgery than medication because of the pathophysiology of the disease.^{2,7} Surgical treatment includes goniotomy, trabeculotomy, or trabeculectomy, or a combination of these procedures.⁷ The surgery planned is usually trabeculo-trabeculectomy, due to the severity of the glaucoma, but in practice only trabeculectomy is performed, due to technical constraints in finding Schlemm's canal. This is consistent with research by Dietlein *et al.* who performed trabeculectomy in four patients who initially planned trabeculo-trabeculectomy because Schlemm's canal could not be localized.⁸ The trabeculectomy that had already been performed in this patient resulted no significant improvement. There is

more failure with trabeculectomy than goniotomy or trabeculotomy because of complications including scarring over the bleb.^{1,8,9} Corpus callosum is the largest connective structure in the brain. HCC exists when all the components of the callosum are present, but the structure is thinner than normal.¹⁰ HCC is a strong predictor of developmental delay and significantly associated with poor outcome in all of the developmental domains. Anderson *et al.* found that among very premature infants, corpus callosum grows at lower rate postnatal than in utero. Some recent studies have reported some syndrome-associated congenital glaucoma with malformation of corpus callosum.⁵ This patient suffered from HCC with developmental delay, but the other congenital anomaly associated with a syndrome has not been proven. In conclusion, PCG associated with HCC is a rare disease and presents challenging management. Recognizing the early presenting signs and symptoms of congenital glaucoma for medical practitioners is important. The failure rate of surgery in severe PCG conditions is still very high, and therapy can usually preserve vision if early identification of mild or moderate form is made.

Acknowledgement: The author would thank to Mrs Nur Fitriyah for English editing.

REFERENCES

- Walton D, Katsavounidou G. Newborn primary congenital glaucoma: 2005 Update. *J Pediatr Ophthalmol Strabismus* 2005;42(6):333–41.
- Ho CL, Walton DS. Primary congenital glaucoma. *J Pediatr Ophthalmol Strabismus* 2004;41(5):271–88.
- Yu Chan JY, Choy BN, Ng AL, Shum JW. Review on the management of primary congenital glaucoma. *J Curr Glaucoma Pract* 2015;9(3):92–9.
- Paul LK, Brown WS, Adolphs R, Tyszka JM, Richards LJ, Mukherjee P, *et al.* Agenesis of the corpus callosum: genetic, developmental and functional aspects of connectivity. *Nat Rev Neurosci* 2007;8(4):287–99.
- Abouzeid H, Youssef MA, ElShakankiri N, Hauser P, Munier FL, Schorderet DF. PAX6 aniridia and interhemispheric brain anomalies. *Mol Vis* 2009;15:2074–83.
- Girgis NM, Frantz KA. A case of primary congenital glaucoma: A diagnostic dilemma. *Optometry* 2007;78(4):167–75.
- Biglan AW. Glaucoma in Children: are we making progress? *J AAPOS* 2006;10(1):7–21.
- Dietlein TS, Jacobi PC, Kriegelstein GK. Prognosis of primary ab externo surgery for primary congenital glaucoma. *Br J Ophthalmol* 1999;83(3):317–22.
- Elder MJ. Congenital glaucoma in the West Bank and Gaza Strip. *Br J Ophthalmol* 1993;77(7):413–16.
- Anderson NG, Laurent I, Woodward LJ, Inder TE. Detection of impaired growth of the corpus callosum in premature infants. *Pediatrics* 2006;118(3):951–60.

Received: 10 September, 2016

Revised: 31 May, 2017

Accepted: 7 March, 2018

Address for Correspondence:

Prastiya Indra Gunawan, Jl. Delta Raya I No. 20, Perum Delta Sari Baru, Waru Sidoarjo-Indonesia

Tel: +62 81 13429476

Email: prastiya_ig@yahoo.co.id

Rare case of Primary Congenital Glaucoma with Hypoplasia Corpus Callosum

ORIGINALITY REPORT

22%

SIMILARITY INDEX

13%

INTERNET SOURCES

16%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Nadine M. Girgis, Kelly A. Frantz. "A case of primary congenital glaucoma: A diagnostic dilemma", Optometry - Journal of the American Optometric Association, 2007 4%

Publication
- 2 K U al-Umran. "Corneal diameter in premature infants.", British Journal of Ophthalmology, 5/1/1992 4%

Publication
- 3 www.ajol.info 2%

Internet Source
- 4 ayubmed.edu.pk 1%

Internet Source
- 5 webeye.opth.uiowa.edu 1%

Internet Source
- 6 www.ppjpi.unair.ac.id 1%

Internet Source
- 7 www.ayubmed.edu.pk

Internet Source

		1%
8	journals.plos.org Internet Source	1%
9	Biglan, A.W.. "Glaucoma in Children: Are We Making Progress?", Journal of AAPOS, 200602 Publication	1%
10	www.panafrican-med-journal.com Internet Source	1%
11	David S. Walton. "Chapter 22 The Childhood Glaucomas", Springer Science and Business Media LLC, 2014 Publication	1%
12	www.ahujaeyecentre.com Internet Source	1%
13	nrs.harvard.edu Internet Source	1%
14	journals.lww.com Internet Source	1%
15	www.ncbi.nlm.nih.gov Internet Source	1%
16	Morales-Fernandez, Laura, Jose M. Martinez-de-la-Casa, Javier Garcia-Bella, Carmen Mendez, Federico Saenz-Frances, Maite Garcia, Julio Escribano, and Julian Garcia-	1%

Feijoo. "Clinical Variability of Primary Congenital Glaucoma in a Spanish Family With Cyp1b1 Gene Mutations :", Journal of Glaucoma, 2015.

Publication

17

Hannah C. Glass. "Agenesis of the corpus callosum in California 1983-2003: A population-based study", American Journal of Medical Genetics Part A, 10/01/2008

<1%

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Rare case of Primary Congenital Glaucoma with Hypoplasia Corpus Callosum

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3
