

# Anal extrusion of VP Shunt Catheter

*by* Prastiya Indra Gunawan

---

**Submission date:** 30-Jun-2020 12:55PM (UTC+0800)

**Submission ID:** 1351656586

**File name:** 12.\_Anal\_extrusion\_of\_VP\_Shunt\_Personal\_Copy\_from\_Academia.pdf (223.77K)

**Word count:** 260

**Character count:** 2305

## Anal extrusion of the ventriculoperitoneal shunt catheter

Prastiya Indra Gunawan, IGM Reza Gunadi Ranuh, Alpha Fardah Atthiyah

Department of Pediatrics, Airlangga University College of Medicine, Dr Soetomo Hospital, Surabaya, Indonesia



An 11-month-old girl presented with a history of protrusion of a ventriculoperitoneal (VP) shunt through the anus (Panel A). The patient had a history of congenital hydrocephalus since birth and the VP shunt had already been inserted for 6 months. On examination, the child was found to be afebrile with no abdominal or meningeal symptoms. The shunt tube would protrude from the anus upon straining. The neurosurgeon decided to remove the VP shunt. The VP catheter was cut at the abdominal level and gen-

tly pulled out by the extruded anal portion. An incision at the head end was made to remove the rest of the shunt. After the procedure the patient was doing well. Abdominal ultrasound showed no evidence of free fluid collection, and there were no signs of peritonitis. VP shunting is most commonly indicated for hydrocephalus; however, there are some complications reported. Bowel perforation and anal extrusion of the distal shunt tube are rare and unusual complications of a VP shunt, and were first reported by Wilson and Bertrand in 1966. So far, to the best of our knowledge, 56 cases have been reported. The exact pathogenesis of shunt tube-related organ perforation is unclear, but various mechanisms have been suggested, including pressure necrosis, foreign body reaction, previous inflammations of the bowel wall, and poor general condition with weak bowel musculature, that may contribute to perforation. Careful attention is recommended so this complication is recognized and surgical removal of the shunt tube considered in the absence of infection, to avoid morbidity and mortality.

**Key words:** Anal extrusion • VP shunt • Hydrocephalus.

**Authors' contributions:** Conception and design: PIG and RGR; Acquisition, analysis and interpretation of data: PIG and AFA; Drafting the article: PIG, RGR and

AFA; Revising it critically for important intellectual content: RGR and AFA; Approved final version of the manuscript: PIG, RGR and AFA.

**Conflict of interest:** The authors declare that they have no conflict of interest.

Correspondence:

*prastiya\_ig@yahoo.co.id*

Tel.: + 62 811 342 9476; Fax.: + 62 315 501 748

Received: 18 October 2016

Accepted: 29 December 2016

ACTA MEDICA ACADEMICA  
PERSONAL COPY

# Anal extrusion of VP Shunt Catheter

---

## ORIGINALITY REPORT

---

9%

SIMILARITY INDEX

5%

INTERNET SOURCES

7%

PUBLICATIONS

0%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

1

[www.scilit.net](http://www.scilit.net)

Internet Source

5%

---

2

Susan Bhattarai, Ram Hari Chapagain, Deepak Mishra, Anil Kumar Shrestha, Sushan Man Shrestha. "Bacteriological Profile and Antibiotic Sensitivity Pattern of Neonatal Sepsis in Central Paediatric Referral Hospital in Nepal", Journal of Nepal Paediatric Society, 2019

Publication

4%

---

Exclude quotes On

Exclude matches Off

Exclude bibliography On

# Anal extrusion of VP Shunt Catheter

---

GRADEMARK REPORT

---

FINAL GRADE

**/100**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---