

Hepatitis B and C virus infection among hemodialysis patients in yogyakarta, Indonesia- Prevalence and molecular evidence for nosocomial transmission

by Maria Lusida

Submission date: 23-Jun-2020 10:16PM (UTC+0800)

Submission ID: 1348583546

File name: revalence_and_molecular_evidence_for_nosocomial_transmission.pdf (1.02M)

Word count: 10416

Character count: 54649

Hepatitis B and C Virus Infection Among Hemodialysis Patients in Yogyakarta, Indonesia: Prevalence and Molecular Evidence for Nosocomial Transmission

Hanggoro Tri Rinonce,^{1,2} Yoshihiko Yano,^{1,3*} Takako Utsumi,^{1,4} Didik Setyo Heriyanto,² Nungki Anggorowati,^{1,2} Dewiyani Indah Widasari,^{1,2} Maria Inge Lusida,⁴ Soetjipto,⁴ Heru Prasanto,⁵ Hak Hotta,¹ and Yoshitake Hayashi¹

¹Center for Infectious Diseases, Kobe University Graduate School of Medicine, Kobe, Japan

²Department of Anatomical Pathology, Faculty of Medicine, Dr. Sardjito Hospital, Gadjah Mada University, Yogyakarta, Indonesia

³Department of Gastroenterology, Kobe University Graduate School of Medicine, Kobe, Japan

⁴Indonesia-Japan Collaborative Research Centre for Emerging and Re-Emerging Infectious Diseases, Institute of Tropical Disease, Airlangga University, Surabaya, Indonesia

⁵Department of Internal Medicine, Faculty of Medicine, Dr. Sardjito Hospital, Gadjah Mada University, Yogyakarta, Indonesia

Hemodialysis patients are at an increased risk of acquiring hepatitis B virus (HBV) and hepatitis C virus (HCV) infection. However, the prevalence of hepatitis viral infection and its genotype distribution among hemodialysis patients in Indonesia are unclear. In order to investigate these issues and the possibility of nosocomial transmission, 161 hemodialysis patients and 35 staff members at one of the hemodialysis unit in Yogyakarta, Indonesia, were tested for serological and virological markers of both viruses. HBV surface antigen (HBsAg) was detected in 18 patients (11.2%) and in two staff members (5.7%). Anti-HCV was detected in 130 patients (80.7%) but not in any staff members. Occult HBV and HCV infection were detected in 21 (14.7%) and 4 (12.9%) patients, respectively. The overall prevalence rates of HBV and HCV infection among patients were 24.2% and 83.2%, respectively. HCV infection was independently associated with hemodialysis duration and the number of blood transfusions. Phylogenetic analysis revealed that 23 of 39 tested HBV strains (59%) were genotype B, 11 (28.2%) were genotype C, and 5 (12.8%) were genotype A. HCV genotype 1a was dominant (95%) among 100 tested HCV strains. Nosocomial transmission was suspected because the genotype distribution differed from that of the general population in Indonesia, and because the viral genomes of several strains were identical. These findings suggest that HBV and HCV infection is common among hemodialysis patients in Yogya-

karta, and probably occurs through nosocomial infection. Implementation of strict infection-control programs is necessary in hemodialysis units in Indonesia. *J. Med. Virol.* 85:1348–1361, 2013. © 2013 Wiley Periodicals, Inc.

KEY WORDS: hepatitis B; hepatitis C; hemodialysis; Yogyakarta; Indonesia; nosocomial transmission

INTRODUCTION

Hemodialysis patients are at increased risk of acquiring hepatitis B virus (HBV) and hepatitis C virus (HCV) infection because of frequent contact with blood, supplies, or surfaces contaminated with these viruses [Rahnavardi et al., 2008; Edey et al., 2010]. Consequently, the prevalence of both infections in hemodialysis patients is very high, although it varies among countries and among hemodialysis units within the same country. Nevertheless, the prevalence of infection is generally higher in

Grant sponsor: The Japanese Initiative for Global Research Network on Infectious Diseases (J_GRID) grant-in-aid from The Ministry of Education, Culture, Sports, Science, and Technology, Japan

*Correspondence to: Yoshihiko Yano, MD, PhD, Center for Infectious Diseases, Graduate School of Medicine, Kobe University, 7-5-1 Kusunoki-cho, Chuo-Ku, Kobe 650-0017, Japan.

E-mail: yanoyo@med-u.ac.jp

Accepted 30 January 2013

DOI 10.1002/jmv.23581

Published online in Wiley Online Library (wileyonlinelibrary.com).

developing countries than in developed countries, and might reflect its prevalence in the general population [Burdick et al., 2003; Rahnavardi et al., 2008; Johnson et al., 2009]. Unfortunately, few data are available for Indonesia. In a study conducted in West Java, the rates of HBV surface antigen (HBsAg) and anti-HCV seropositivity among hemodialysis patients were 6.8% and 73.5%, respectively [Saketi et al., 2003], whereas those in Yogyakarta were 7% and 81%, respectively [Hadiwandowo et al., 1994]. However, as the study by Hadiwandowo et al. [1994] was conducted in 1994, more recent data addressing the rates on infection in Yogyakarta are needed.

The number of blood transfusions and the duration of hemodialysis are established risk factors for HBV and HCV infection in hemodialysis patients [Vladutiu et al., 2000; Busek et al., 2002; Ferreira et al., 2006]. The implementation of blood product screening in blood banks and the use of erythropoietin treatment have decreased the prevalence of infection with these viruses in hemodialysis units. However, outbreaks of both viruses still intermittently occur [Center for Disease Control and Prevention, 2001; Finelli et al., 2005]. Nosocomial transmission might play an important role in such outbreaks, a hypothesis that is supported by the association between hemodialysis duration and risk of infection with these viruses [Soetjipto et al., 1996].

HBV and HCV genotypes show marked geographic distributions. Indonesia is a multiethnic country with as many as 24 HBV subgenotypes, of which HBV subgenotype B3 (HBV/B3) is dominant [Nurainy et al., 2008; Mulyanto et al., 2009, 2010, 2011, 2012; Utsumi et al., 2009]. At least 15 subgenotypes of HCV have been identified throughout Indonesia, of which HCV/1b is the most prevalent [Hotta et al., 1994; Widjaja et al., 1995; Soetjipto et al., 1996; Tokita et al., 1996; Inoue et al., 2000; Utama et al., 2008, 2010; Anggorowati et al., 2012]. Nevertheless, few data exist for hemodialysis patients. Therefore, it is important to gather further information on the prevalence of HBV and HCV infection in hemodialysis settings, as well as identify the prevalent genotypes by performing molecular epidemiological studies using phylogenetic analysis. This tool is very useful in epidemiology studies, and can be used to trace the possible routes of transmission, and provide evidence of nosocomial infection [Kramvis et al., 2005; Kondili et al., 2006].

The objectives of this study were to investigate the prevalence and genotype distribution of HBV and HCV among hemodialysis patients in Yogyakarta, Indonesia. The possibility of nosocomial transmission was also assessed using molecular analysis.

MATERIALS AND METHODS

Patients, Staff Members, and Hemodialysis Unit

A total of 161 patients with end-stage renal disease who were undergoing hemodialysis and 35 staff

members (doctors, nurses, technicians, and cleaning and administrative staff) at one of the hemodialysis unit in Yogyakarta, Indonesia, were enrolled in this study, during January and February 2010. After obtaining written informed consent, the demographic and risk factors associated with HBV and HCV infection data were collected using a standardized questionnaire. Risk factors evaluated in this study were hemodialysis duration, frequency of hemodialysis each week, history of blood transfusion, number of blood transfusions, history of renal transplantation, multiple sexual partners, sexually transmitted diseases, injection drug use, and hepatitis B vaccination.

The hemodialysis unit serves approximately 350 patients from Yogyakarta province and the southern region of Central Java province. It provides hemodialysis treatment in morning, afternoon, or evening shifts. The patients were randomly selected from each shift, regardless of age, sex, and race. Before starting their hemodialysis treatment, blood (5 ml) was collected in plain tubes, allowed to clot, and centrifuged at room temperature. The sera were separated and stored at -80°C until used.

Thirty machines were available at the hemodialysis unit. Two machines were dedicated for patients with HBsAg seropositivity, 27 were used for patients with HBsAg seronegativity, regardless of anti-HCV status, and one was reserved for emergencies. Separate rooms were only available for patients seropositive for HBsAg. The hemodialysis machine uses a hollow fiber dialyzer with standard acetate and bicarbonate solutions. The dialyzers were cleaned individually, rinsed with sterile saline solution, and disinfected with formaldehyde. Before use, the dialyzers were flushed with 2,500 ml of normal saline solution according to standard procedures. Dialyzers were reused up to a maximum of eight times for all patients.

The staff members did their best to follow the recommendations for HBV and HCV infection control issued by the Indonesian Society of Nephrology. However, because of resource limitation, not all of the patients with HBsAg seronegativity were vaccinated for HBV, and anti-HCV screening was not performed routinely. Assessment of HBV DNA and HCV RNA using PCR assays was not performed regularly because of its high cost. Furthermore, established precautions to avoid infection were not consistently implemented, particularly in emergencies. Some instruments and medications were shared among patients.

The study protocol was reviewed and approved by the Ethics Committees of Gadjah Mada University, Indonesia, and Kobe University, Japan.

Liver Enzymes and Serologic Markers

Serum alanine aminotransferase (ALT) and γ -glutamyl transpeptidase (GGT) levels were determined using methods recommended by the Japan Society of Clinical Chemistry. Elevated ALT levels were defined as values >40 IU/L while elevated GGT levels were

defined as values ≥ 70 IU/L for males and ≥ 30 IU/L for females.

All samples were screened and confirmed for HBsAg status using reverse passive hemagglutination assays (Mycell II HBsAg; Institute of Immunology, Tokyo, Japan) and chemiluminescent immunoassays (Architect; Abbott Japan, Tokyo, Japan). Patients with HBsAg seropositivity were then tested for hepatitis B virus e antigen (HBeAg). Only patients with HBeAg negativity were further tested for anti-HBe antibody using enzyme immunoassays (Immunis HBeAg and HBeAb EIA; Institute of Immunology). Testing for anti-HBs and anti-HBc was conducted using the passive hemagglutination method (Mycell II anti-HBs and Mycell II anti-rHBc; Institute of Immunology) for all samples. Anti-HCV antibodies were detected using the particle agglutination method (Ortho HCV Ab PA test II; Ortho-Clinical Diagnostics, Tokyo, Japan). All assays were conducted according to the manufacturers' instructions.

Detection of Hepatitis Viruses

HBV DNA was extracted from 200 μ l of sera using a DNA extraction kit (QIAamp DNA Blood Mini Kit, QIAGEN Sciences, Germantown, MD) according to the manufacturer's instructions. The HBV genome was amplified by nested PCR as described previously [Sugauchi et al., 2001; Cui et al., 2002; Wong et al., 2011]. The amplified products were visualized on a 2% agarose gel stained with ethidium bromide. Patients and staff members who were negative serologically for HBsAg but positive for HBV DNA, based on PCR assays of at least two regions of the HBV genome, were defined as having occult HBV infection [Raimondo et al., 2007].

HBV DNA was quantified by real-time detection PCR (RTD-PCR) with Taq Man chemistry, as previously described [Abe et al., 1999], using an Applied Biosystems 7500 RT-PCR System (Applied Biosystems, Foster City, CA). The detection limit was 1.8 log copies/ml. A viral load >4 log copies/ml was considered high, whereas a viral load ≤ 4 log copies/ml was considered low [Moutinho et al., 2006].

HCV RNA was extracted from 140 μ l of sera using an RNA extraction kit (QIAamp Viral RNA Mini Kit, QIAGEN GmbH, Hilden, Germany) according to the manufacturer's instructions. The NS5B region of the HCV genome was amplified by nested RT-PCR using a SuperScript III One-Step RT-PCR System (Invitrogen, Carlsbad, CA) with specific primers, as previously described [Akkarathamrongsin et al., 2010]. Patients or staff members seronegative for anti-HCV but positive for HCV RNA, based on PCR, were defined as having occult HCV infection [Jain and Nijhawan, 2008; Tu et al., 2009; Li and Wang, 2010].

To avoid cross-contamination between samples, standard precautions were used in all procedures. Reagents, samples, and amplified products were stored in separate areas.

Sequencing and Phylogenetic Analysis

The amplified fragments were directly sequenced using a BigDye Terminator v3.1. Cycle Sequencing Kit and an ABI PRISM 3100-Avant Genetic Analyzer (Applied Biosystems).

HBV genotypes and subgenotypes were determined by phylogenetic analysis based on the Pre-S2 and S region (nucleotides [nt] 1–459). Reference sequences were retrieved from GenBank. Multiple alignments were done using Clustal X software (<http://www.clustal.org>), and phylogenetic trees were constructed using the neighbor-joining method based on Kimura two-parameter distance estimation. To confirm the reliability of the phylogenetic tree topologies, bootstrap reconstruction was carried out 1,000 times, and bootstrap values $>70\%$ were considered significant. Analyses were conducted using Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0.2 (<http://megasoftware.net/>). Samples with a high viral load were subjected to further PCR amplification to obtain the entire genome sequences. Phylogenetic analysis of HCV was done based on the NS5B region (nt 8,244–8,615) through comparisons with confirmed reference sequences [Simmonds et al., 2005].

Detection of Amino Acid Substitutions in the “a” Determinant Region

Some of the amino acid substitutions in the “a” determinant region are thought to result in undetectable HBsAg in patients with detectable HBV DNA. Sequences of this region were aligned and analyzed to identify amino acid substitutions among HBV strains obtained from patients and staff members positive for HBV DNA. Additional primer pairs, as previously described [Juniausti et al., 2011], were used to amplify the DNA sequence spanning the “a” determinant region in samples that could not be detected using the primers pairs mentioned above.

Statistical Analysis

Statistical analysis was performed using χ^2 -tests or Fisher's exact test for categorical variables, and independent *t*-tests and the Mann-Whitney test for continuous variables. Risk factors associated with HBV and HCV infection were first evaluated in univariate analyses by determining odds ratios. Risk factors identified in univariate analyses were further analyzed by multiple logistic regression to control for possible confounders. Values of $P < 0.05$ were considered statistically significant. Patients or staff members were defined as HBV positive if they were HBV DNA positive, regardless of serological status. Patients or staff members who were either anti-HCV or HCV RNA positive were considered as HCV positive.

Nucleotide Sequence Accession Numbers

The HBV and HCV sequences described in this study are deposited in the DNA Data Bank of Japan

under accession numbers AB713527 to AB713532, AB713848 to AB713895, AB714147 to AB714242, and AB767304 to AB767307.

RESULTS

Characteristics of the Participants

The characteristics of the participants are shown in Table I. As shown in this table, most of the patients were male and were in their middle adulthood. The patients were mainly Javanese and came from Sleman Regency. Most of the staff members were female and they were generally younger than the patients ($P < 0.001$). They were all Javanese and mainly came from Sleman Regency. Most of them had been working in the hemodialysis unit for a long time (mean \pm SD, 14.1 ± 8.0 years; range, 3 months to 42 years).

Chronic glomerulonephritis and hypertensive kidney disease were the main causes of end-stage renal disease in the patients. The duration of hemodialysis treatment was ≥ 1 year for most patients, and the majority underwent hemodialysis twice a week. Blood transfusion was still commonly performed to treat renal anemia. However, only eight patients had received the HBV vaccine before starting hemodialysis treatment. The staff members were routinely checked for hepatitis markers and were vaccinated. Kidney transplantation was rarely performed to treat patient with end-stage renal disease.

Distribution of Hepatitis Viral Markers

Eighteen of 161 hemodialysis patients (11.2%) and 2 of 35 staff members (5.7%) were HBsAg positive. HBV DNA was detected in all of the patients and staff members who were positive for HBsAg. Twenty-one of 143 (14.7%) patients negative for HBsAg had detectable HBV DNA in at least two regions within the HBV genome, and were defined as having occult infection. In contrast, none of the staff members had occult infection. Thus, the overall rates of HBV infection among patients and staff members were 24.2% and 5.7%, respectively (Table II). Anti-HBs was more frequently detected among staff members than among patients.

Overall, 134 (83.2%) patients had HCV infection. In contrast, none of the staff members were positive for anti-HCV or HCV RNA (Table II). HCV RNA was detected in 96 of 130 seropositive patients (73.8%) and 4 of 31 seronegative patients (12.9%) were classified as having occult infection. Twenty-nine patients (18%) were co-infected with HBV and HCV (Table III).

The overall prevalence of HBV and HCV was significantly higher in hemodialysis patients than in staff members.

Risk Factors Associated With HBV and HCV Infection

The characteristics of patients and possible risk factors for HBV and HCV infection are summarized

TABLE I. Characteristics of the Hemodialysis Patients and Hemodialysis Unit Staff Members

Characteristics	Patients (n = 161)	Staff members (n = 35)
Gender, n (%)		
Male	93 (58)	13 (37)
Female	68 (42)	22 (63)
Age, mean \pm SD	48 \pm 13	40 \pm 7
Area of origin, n (%)		
Sleman Regency	70 (43)	21 (60)
Gunung Kidul Regency	17 (11)	0 (0)
Kulon Progo Regency	12 (7)	1 (3)
Bantul Regency	12 (7)	7 (20)
Yogyakarta City	22 (14)	3 (9)
Central Java Province	23 (14)	2 (6)
Other regions	5 (3)	1 (3)
Race, n (%)		
Javanese	157 (97)	35 (100)
Sundanese	1 (0.6)	0 (0)
Balinese	1 (0.6)	0 (0)
Batak	1 (0.6)	0 (0)
Gorontalo	1 (0.6)	0 (0)
Cause of end-stage renal diseases, n (%)		
Chronic glomerulonephritis	50 (31)	
Hypertensive kidney diseases	50 (31)	
Diabetic nephropathy	28 (17)	
Obstructive nephropathy	10 (6)	
Polycystic kidney disease	6 (4)	
Lupus nephropathy	1 (0.6)	
Unknown	16 (10)	
Duration of hemodialysis, n (%)		
<1 year	58 (36)	
≥ 1 year	103 (64)	
Number of dialyses per week, n (%)		
1 time	14 (9)	
2 times	141 (88)	
≥ 3 times	6 (4)	
History of blood transfusion, n (%)	153 (95)	
Number of blood transfusions, n (%)		
≤ 5	96 (60)	
> 5	65 (40)	
History of kidney transplantation, n (%)	1 (0.6)	
Multiple sexual partners, n (%)	2 (1.2)	
History of suffering sexually transmitted disease, n (%)	0 (0)	
History of injecting drug use, n (%)	1 (0.6)	
Hepatitis B vaccination before hemodialysis treatment, n (%)	8 (5)	

in Tables III and IV. None of the possible risk factors analyzed in this study were associated with HBV infection (some data are not shown). However, in univariate analysis, hemodialysis duration and the number of blood transfusions were significantly associated with HCV infection. After adjusting for age and gender in multivariate analysis, hemodialysis duration and the number of blood transfusions were independently associated with HCV infection (Table IV, some data are not shown).

TABLE II. Distribution of Hepatitis Viral Markers Among Hemodialysis Patients and Hemodialysis Unit Staff Members

Hepatitis viral markers	Number (%)		P-value*
	Patients (n = 161)	Staff members (n = 35)	
HBsAg positive	18 (11.2)	2 (5.7)	0.54
HBeAg positive ^a	5 (3.1)	0 (0.0)	1.00
Anti-HBe positive ^b	9 (5.6)	1 (2.8)	1.00
Anti-HBc positive	48 (29.8)	6 (17.1)	0.13
Anti-HBs positive	84 (52.5)	27 (77.1)	0.008
HBV DNA positive	39 (24.2)	2 (5.7)	0.01
Anti-HCV positive	130 (80.7)	0 (0.0)	< 0.001
HCV RNA positive	100 (62.1)	0 (0.0)	< 0.001
Anti-HCV and or HCV RNA positive	134 (83.2)	0 (0.0)	< 0.001

^aHBeAg was only tested for HBsAg positive samples.

^bAnti-HBe was only tested for HBeAg negative samples.

*P-values were determined by the χ^2 -test. Significant values are indicated in bold.

Elevated ALT levels were not associated with HBV infection. However, elevated GGT levels were more frequent in patients without HBV infection than in patients with HBV infection ($P = 0.03$), which might be due to the difference in the prevalence of HCV infection. The presence of HBV DNA was associated with anti-HBc positivity ($P < 0.001$; Table III). Elevated ALT levels were not associated with HCV infection. However, elevated GGT levels were more frequent in patients with HCV infection than in

TABLE III. Characteristics of Patients and Risk Factors Associated With HBV Infection

Characteristics and risk factors	HBV DNA positive			HBV DNA negative (n = 122)	Univariate analysis	
	HBsAg positive (n = 18)	HBsAg negative (n = 21)	Total (n = 39)		OR (95% CI)	P-value*
Gender						
Female	6	11	17	51	1	
Male	12	10	22	71	0.9 (0.4–1.9)	0.8
Age						
<45	5	11	16	49	1	
>45	13	10	23	73	0.96 (0.5–2.0)	0.92
Hemodialysis duration						
<1 year	5	11	16	42	1	
≥1 year	13	10	23	80	0.75 (0.4–1.6)	0.45
Number of blood transfusions						
≤5	11	14	25	70	1	
>5	7	7	14	52	0.75 (0.4–1.6)	0.46
ALT level ^a						
Not elevated	17	20	37	101	1	
Elevated	1	1	2	21	0.3 (0.1–1.2)	0.06
GGT level ^b						
Not elevated	10	10	20	39	1	
Elevated	8	11	19	83	0.4 (0.2–0.9)	0.03
Anti-HBs						
Negative	13	10	23	54	1	
Positive	5	11	16	68	0.5 (0.3–1.1)	0.10
Anti-HBc ^c						
Negative	3	14	17	96	1	
Positive	15	7	22	26	4.8 (2.2–10.3)	< 0.001
Anti-HCV and or HCV RNA						
Negative	3	7	10	17	1	
Positive	15	14	29	105	0.5 (0.2–1.1)	0.08
HBV viral load						
<4 log copies/ml	12	21	33	—	—	
>4 log copies/ml	6	0	6	—	—	0.006

^aElevated ALT levels were defined as values >40 IU/L.

^bElevated GGT levels were defined as values ≥70 IU/L for males and ≥30 IU/L for females.

^cThe difference was statistically significant if the HBsAg positive and HBsAg negative groups were compared ($P = 0.002$).

*P-values were determined by the χ^2 -test. Significant values are indicated in bold.

TABLE IV. Characteristics of Patients and Risk Factors Associated With HCV Infection

Characteristics and risk factors	HCV positive (n = 134)	HCV negative (n = 27)	Univariate analysis		Multivariate analysis	
			OR (95% CI)	P-value*	OR ^a (95% CI)	P-value*
Gender						
Female	56	12	1			
Male	78	15	0.9 (0.4–2.1)	0.79		
Age						
<45 years	55	10	1			
>45 years	79	17	0.8 (0.4–2.0)	0.69		
Hemodialysis duration						
<1 year	35	23	1		1	
≥1 year	99	4	16.2 (5.2–50.3)	<0.001	9.5 (3.0–30.5)	<0.001
Number of blood transfusions						
≤5	69	26	1		1	
>5	65	1	24.5 (3.2–185.7)	<0.001	11.0 (1.4–88.4)	0.02
ALT level^b						
Not elevated	113	25	1			
Elevated	21	2	0.4 (0.1–1.9)	0.37		
GGT level^c						
Not elevated	45	14	1			
Elevated	89	13	0.5 (0.2–1.1)	0.07		

^aAdjusted odds ratio; adjusted for age and gender.

^bElevated ALT levels were defined as values >40 IU/L.

^cElevated GGT levels were defined as values ≥70 IU/L for males and ≥30 IU/L for females.

*Significant values are indicated in bold.

patients without HCV infection, although the difference was not statistically significant. The mean GGT level was significantly higher in patients with HCV infection than in patients without HCV infection (mean ± SD, 158.0 ± 163.7 vs. 73.7 ± 80.4 IU/L, $P < 0.001$).

Thirty-three (84.6%) of 39 patients with HBV infection had a low (≤4 log copies/ml) or an undetectable viral load. The median titer was 2.8 log copies/ml (range, 1.8–5.5 log copies/ml). Significantly, more patients with overt infection (33.3%) than patients with occult infection (0.0%) had a high viral load ($P = 0.006$, Table III).

HBV and HCV Genotypes

The HBV strains obtained in this study were sequenced and compared with the sequences deposited in GenBank for the pre-S2 and S regions, and a phylogenetic tree was constructed (Fig. 1). Twenty-three of 39 strains (59.0%) isolated from patients belonged to HBV/B. Surprisingly, HBV/C and HBV/A were identified in 11 (28.2%) and 5 (12.8%) strains, respectively. All of the HBV/B strains were classified as HBV/B3. Ten (25.6%) HBV/C strains were classified as C2 and the remaining 1 strain (2.6%) as C7. All of the HBV/A strains were classified as HBV/A2. HBV/C2 was detected mainly in patients with occult infection. HBV/A2 was only detected in patients with occult infection. Two isolates from staff members were classified as HBV/B3. Some of the HBV/A2, HBV/B3, and HBV/C2 strains were identical to the fragment spanning nt 1–459 (YOGHhcv89 and 90;

YOGHhcv10, 29, 32, 49, 67, 68, 74, 75, 132, and 149; YOGHhcv54 and 72). These findings strongly suggest the occurrence of nosocomial transmission. Six HBV strains were fully sequenced, and all were HBV/B3.

One hundred HCV strains isolated in this study were compared with 25 strains from the GenBank database, and a phylogenetic tree was constructed (Fig. 2). HCV genotype 1 (HCV/1) was the most common genotype (98%), followed by HCV/3 (2%). HCV/1a, a rare subgenotype among blood donors in Indonesia, was dominant (95%), while HCV/1b, HCV/1c, and HCV/3a were identified in 1 (1%), 2 (2%), and 2 (2%) patients, respectively. The HCV/1a strains showed high similarity (mean ± SD, 96.1 ± 2.0; range, 89.2%–100%). Twenty of them, including two strains isolated from patients with occult infection, were identical to the fragment spanning nt 8,244–8,615 (YOGHhcv13, 46, 57, and 155; YOGHhcv21, 48, 85, 112, 134, and 144; YOGHhcv32, 68, 135, and 136; YOGHhcv42 and 137; YOGHhcv50 and 51; YOGHhcv90 and 93). These findings were also strongly suggestive of nosocomial transmission. To confirm these findings, 95 HCV/1a strains were compared with five local but unrelated HCV/1a strains (AB661787, AB661788, AB661792, AB661794, and AB661816) that were isolated in a previous study [Anggorowati et al., 2012], and with 14 HCV/1a strains isolated in other countries. The phylogenetic tree showed that most of the strains were clustered in the same distinct branches of the tree, but were separated from the five local strains and the strains from the other countries (Fig. 3).

113
Amino Acid Substitution in the “a” Determinant Region
 37

The clinical and molecular data of the patients and staff members with HBV infection, including the distribution of amino acid substitutions in the “a” determinant, are listed in Table V. Amino acid substitutions were more frequently detected in patients with HBsAg seropositivity (67.7%) than in patients with occult infection (47.6%), although this difference was not statistically significant. Four of the HBV/A2 strains were wild-type strains, and one had T126I, N131T, and T143S substitutions. Among the HBV/B3 strains, T131N substitutions were the most frequent. Several specific substitutions (Q129H, Q129R, M133T, C137G, and C137W) were also identified, but less frequently. Among the HBV/C2 strains, I126T substitutions were found in three strains, T131N in one strain, T131P in one strain, and S143T in three strains. The D144G and G145R mutations were both found in one strain. The I126T substitution was detected in the HBV/C7 strain. The T140I substitution was only identified in one HBV/B3 strain isolated from a staff member. None of those substitutions were associated with the occurrence of occult HBV infection. One HBV/A2 strain (YOGHhbv121) had the same amino acid sequence as HBV/C2, one HBV/B3 strain (YOGHhbv65) had the same amino acid sequence as HBV/C2, and one HBV/C2 strain (YOGHhbv36) had the same amino acid sequence as HBV/A2.

70
DISCUSSION

The prevalence of HBV and HCV infection among hemodialysis patients in Yogyakarta was consistently higher than that among local healthcare workers and blood donors [Hadiwandowo, 1991; Triwibowo, 1993; Hadiwandowo et al., 1994; Rahayujati et al., 2006], suggesting that hemodialysis patients are at greater risk of infection. The high prevalence of anti-HBc seropositivity among patients negative for HBV DNA clearly indicates that hemodialysis patients are susceptible to HBV infection.

This study also confirmed that hemodialysis patients are at greater risk of acquiring HBV and HCV infection compared with staff members at the hemodialysis unit. However, the staff members were still




Fig. 1. Phylogenetic tree analysis of the Pre-S2 and S gene sequences of 39 HBV strains isolated from patients, two strains isolated from staff members, and 58 sequences of reference HBV strains from different genotypes (A–J) obtained from GenBank. The GenBank HBV sequences are labeled with their genotype, accession number, and country of origin. The sequences determined in this study are indicated by isolate number, starting with YOGHhbv, and are labeled with black and white diamonds for strains obtained from patients seropositive for HBsAg and patients with occult infection, respectively. HBV strains isolated from staff members are labeled with black circles. Bootstrap values are given at the internal nodes.



at risk of HBV infection considering the high prevalence of anti-HBc seropositivity among staff members who were negative for HBsAg. To prevent HBV infection, all staff members in the hemodialysis unit are tested routinely for HBV markers and are vaccinated. Nevertheless, approximately one-fifth of the staff members had lost the vaccine's protective effect, as indicated by undetectable anti-HBs.

Conversely, the patients were not routinely vaccinated before starting hemodialysis because the national health insurance program in Indonesia does not yet cover HBV vaccines. Only eight patients in this study had been vaccinated before starting hemodialysis. However, 84 (52.2%) patients were anti-HBs positive, probably because of the onset of immunity after the resolution of HBV infection.

In this study, the prevalence rates of HBV and HCV infection among hemodialysis patients were similar to those in developing countries, and were higher than those in developed countries [Burdick et al., 2003; Rahnavardi et al., 2008; Johnson et al., 2009]. The differences in prevalence rates among these countries might depend on the differences in the prevalence rates of HBV and HCV among the general population. Indonesia is a country with moderate to high endemic rates of HBV infection [Hasan, 2005], and the prevalence of HCV was reported to be 3.9% [Akbar et al., 1997; Lavanchy, 2011]. Therefore, the prevalence rates of HBV and HCV among hemodialysis patients has tended to increase in recent years. Clearly, infection-control programs should be implemented in developing countries.

The prevalence of HBV infection in this study was higher than that in another hemodialysis unit in Jakarta, but the prevalence of HCV infection was lower than that in hemodialysis units in Jakarta and Surabaya, where the prevalence rates were 97.9% and 88%, respectively [Budihusodo et al., 1991; Santoso et al., 2010]. However, these findings may simply represent the high prevalence of HBV and HCV in Java, independent of hemodialysis units. In Yogyakarta, the prevalence rates of HBsAg and anti-HCV were reported to be 7% and 81%, respectively, in a study performed by Hadiwandowo et al. in 1994, who examined HBV and HCV infection among hemodialysis patients at the same hemodialysis unit [Hadiwandowo et al., 1994]. The current study

Fig. 2. Phylogenetic tree analysis of the NS5B gene sequences of 100 HCV strains isolated from patients, and 25 reference sequences of HCV strains belonging to genotypes (1–6) retrieved from GenBank. The GenBank HCV strains are labeled with their genotype, accession number, and country of origin. The sequences determined in this study are indicated by the isolate number, starting with YOGHhcv, and are labeled with black and white triangles for strains isolated from patients seropositive for anti-HCV and patients with occult infection, respectively. Bootstrap values are given at the internal nodes.



revealed a higher prevalence of HBsAg but a similar prevalence of anti-HCV to those reported in study by Hadiwandowo et al. [1994]. Although the results might be influenced by differences in the sensitivities and specificities of the techniques used, they do indicate that the current infection-control procedures did not decrease the prevalence of HBV and HCV infection. In these circumstances, nosocomial transmission might play an important role.

In multivariate analysis, multiple blood transfusions and duration of hemodialysis were independently associated with the prevalence of HCV infection among hemodialysis patients. In Indonesia, HCV screening was introduced at blood banks in 1995. However, HCV screening was only started in 2001 in Yogyakarta. Thus, hemodialysis patients that received blood transfusions before 2001 were at increased risk of acquiring HCV infection. Unfortunately, blood transfusion, rather than erythropoietin treatment, is still being used to correct renal anemia in the hemodialysis unit. Consistent with previous studies [Saketi et al., 2003; Santoso et al., 2010], the duration of hemodialysis was independently and positively associated with the prevalence of HCV infection, supporting the hypothesis that nosocomial transmission was responsible for HCV infection within the hemodialysis unit.

Serum aminotransferase levels are generally low in hemodialysis patients because of pyridoxine deficiencies, leading to impaired enzyme synthesis, and/or because of the effects of uremic serum [Yasuda et al., 1995; Arora et al., 2011]. Therefore, the ALT level was not a useful marker for detecting HBV and HCV infection among hemodialysis patients [Pujol et al., 1996; Arora et al., 2011]. To detect viral hepatitis among hemodialysis patients, the cutoff values for ALT should be reduced to 20 IU/L [Yasuda et al., 1995]. In this study, the mean ALT level was 25.6 IU/L in patients with HCV infection, which exceeded this revised cutoff value. Accordingly, this revised cutoff value might be useful for the early detection of HCV infection in Indonesia.

GTT levels were reported to be useful as an indirect marker for hepatitis infection [Fabrizi et al., 2007; Souza et al., 2008]. In this study, the mean GTT level was markedly higher in patients with HCV

Fig. 3. Phylogenetic tree analysis of NS5B gene sequences of 95 HCV/1a strains from hemodialysis patients, 18 unrelated sequences of HCV strains belonging to genotype 1 (1a-1c) retrieved from GenBank, and five unrelated sequences of HCV/1a strains from the same administrative region. GenBank HCV strains are labeled with their genotype, accession number, and country of origin. The sequences determined in this study are indicated by isolate number, starting with YOGHhc, and are labeled with black and white triangles for strains isolated from patients seropositive for anti-HCV and patients with occult infection, respectively. Unrelated strains from the same administrative region are labeled with black squares. Bootstrap values are given at the internal nodes.

TABLE V. Clinical and Molecular Data of Patients and Staff Members With HBV Infection

Name of strains or accession numbers	Age (year)	Sex	ALT level (IU/L)	GGT level (IU/L)	HBsAg	HBeAg ^a	Anti-HBe ^b	Anti-HBc	Anti-HBs	Viral load (Log copies/mL) ^c	Genotype	Amino acid substitutions in the "a" determinant region (amino acids 124-147)
Consensus sequence												
AJ012207											A2	CTTPAQQNSMFPSCCCTKPTDGNK
M54923											B3	-----T-----
D23684											C2	--I-----S-----
AF011104											C7	--I-----S-----
Patients with HBsAg seropositive												
YOGHhbv11	55	F	18	62	+	-	-	+		UD	B3	-----
YOGHhbv22	45	F	5	69	+	-	+	-		UD	B3	-----
YOGHhbv23	55	M	4	31	+	-	+	-		UD	B3	-----R-----
YOGHhbv30	63	F	14	433	+	-	+	-		4.9	B3	-----H-----
YOGHhbv44	34	M	8	37	+	+	NT	+		5.0	B3	-----M-----
YOGHhbv68	74	M	6	52	+	+	NT	+		4.6	B3	-----N-----
YOGHhbv74	60	F	5	104	+	-	+	+		UD	B3	-----H-----G-----
YOGHhbv75	56	M	10	38	+	-	+	+		UD	B3	-----N-----
YOGHhbv94	57	M	11	39	+	+	NT	+		4.9	B3	-----N-----W-----
YOGHhbv98	40	M	5	64	+	+	NT	+		5.5	B3	-----
YOGHhbv123	60	M	11	29	+	-	+	+		1.8	B3	-----
YOGHhbv126	57	M	9	61	+	-	+	+		UD	B3	-----H-----G-----
YOGHhbv132	23	M	13	73	+	+	NT	+		4.8	B3	-----
YOGHhbv149	60	M	31	124	+	-	+	+		UD	B3	-----
YOGHhbv162	49	M	9	12	+	-	+	+		1.8	B3	-----N-----
YOGHhbv72	73	F	8	27	+	-	+	+		2.9	C2	-----GR-----
YOGHhbv96	50	F	37	117	+	-	+	+		2.4	C2	-----
YOGHhbv21	38	M	62	190	+	-	+	+		UD	C7	--T-----
Patients with occult infection												
YOGHhbv89	42	F	11	20	-	NT	NT	+	+	2.9	A2	-----
YOGHhbv90	54	M	26	192	-	NT	NT	-	-	2.6	A2	-----
YOGHhbv99	53	M	39	36	-	NT	NT	-	-	2.6	A2	-----
YOGHhbv109	12	F	8	42	-	NT	NT	-	-	2.7	A2	-----
YOGHhbv121	51	F	5	19	-	NT	NT	-	-	3.1	A2	--I-----S-----
YOGHhbv10	25	M	71	346	-	NT	NT	-	-	UD	B3	-----
YOGHhbv14	41	M	4	61	-	NT	NT	-	-	UD	B3	-----
YOGHhbv29	79	F	6	27	-	NT	NT	-	-	2.7	B3	-----N-----
YOGHhbv45	29	M	5	31	-	NT	NT	+	+	UD	B3	-----H-----
YOGHhbv46	49	F	10	86	-	NT	NT	-	+	2.6	B3	-----
YOGHhbv49	38	F	4	15	-	NT	NT	-	-	2.3	B3	-----N-----
YOGHhbv65	46	F	5	54	-	NT	NT	-	-	2.7	B3	--I-----S-----
YOGHhbv67	50	F	40	80	-	NT	NT	+	+	2.8	B3	-----
YOGHhbv36	33	M	6	48	-	NT	NT	-	+	UD	C2	--T-----T-----
YOGHhbv54	36	M	8	199	-	NT	NT	+	+	2.8	C2	--T-----T-----
YOGHhbv58	49	M	26	581	-	NT	NT	-	+	3.1	C2	-----
YOGHhbv84	36	M	15	60	-	NT	NT	+	-	2.6	C2	-----
YOGHhbv105	40	F	30	81	-	NT	NT	-	-	2.9	C2	-----T-----
YOGHhbv107	29	M	21	434	-	NT	NT	+	-	2.6	C2	-----
YOGHhbv118	59	F	3	17	-	NT	NT	-	-	2.8	C2	--T-----
YOGHhbv129	55	M	5	36	-	NT	NT	-	+	UD	C2	-----P-----
Staff members												
YOGHhbv2s	33	F	6	12	+	-	-	+	-	2.4	B3	-----I-----
YOGHhbv3s	33	M	17	305	+	-	+	+	-	UD	B3	-----

^{a,b}NT, not tested.
^cUD, undetectable (<1.8 log copies/ml).

infection (158 IU/L) than in patients without HCV infection (73 IU/L). In addition, elevated GTT levels were more frequent than elevated ALT levels among patients with HCV infection. These data suggest that GGT is more applicable than ALT as a marker for HCV infection in hemodialysis patients. Clinical measurement of ALT and GGT levels might be useful, providing the cutoff value for the ALT level is reduced.

Occult HBV infection was detected in 14.7% of hemodialysis patients in Yogyakarta. The prevalence of occult HBV infection ranges from 0% to 26.6% among hemodialysis units in different countries [Minuk et al., 2004; Fabrizi et al., 2005; Gwak et al., 2008; Di Stefano et al., 2009; Mina et al., 2010; Motta et al., 2010]. The sensitivities and specificities of the tests used, vaccination rates, healthcare programs, and isolation of hemodialysis for HBV-infected patients could explain these differences in prevalence rates [Hollinger et al., 2010]. Another study showed that occult HBV infection was associated with the presence of isolated anti-HBc and anti-HCV [Di Stefano et al., 2009]. In this study, occult HBV infection was not associated with anti-HBc seropositivity or anti-HCV seropositivity. The small

number of samples possibly affected the results of this study.

The prevalence of occult HBV infection in the general population of Indonesia is not well documented. In blood donors and school children, the prevalence rates of occult HBV infection were reported to be 8.1% and 2.2%, respectively [Theджа et al., 2010; Utsumi et al., 2010], which were much lower than the prevalence rate of occult HBV infection determined in the present study. These results suggest that hemodialysis patients are more susceptible to occult HBV infection compared with the general population.

Some amino acid substitutions in the "a" determinant region may affect the antigenicity of HBsAg, resulting in diagnostic failure. The M133L and T143M amino acid substitutions within the "a" determinant region possible altered HBsAg antigenicity among Indonesian blood donors [Theджа et al., 2010], whereas the T126I and T143S substitutions were more frequent among school children [Utsumi et al., 2010]. However, none of the substitutions associated with the disappearance of HBsAg were identified in patients with occult HBV infection in this study. Other mechanisms, such as low viral load and

nucleotide mutations in other regions, might explain the high prevalence of occult HBV infection. The amino acid sequence similarity in the “a” determinant region between genotypes supports the possibility of mixed-genotype infection or inter-genotype recombination in hemodialysis patients. Further studies are needed to address these issues.

The differences in the definitions, methods, hemodialysis unit conditions, and geographical areas resulted in marked differences in the prevalence rates of occult HCV infection, which ranged from 0% to 90% [Barril et al., 2008; Jain and Nijhawan, 2008; Tu et al., 2009; Li and Wang, 2010]. The present study revealed a high prevalence of occult HCV infection (12.9%). However, this rate might be lower than the actual rate of occult HCV infection based on the new definition comprising the presence of HCV RNA in hepatocytes or peripheral blood mononuclear cells (PBMCs) in the absence of serological or virological evidence of infection [Fabrizi and Martin, 2008]. Further studies are needed to address this issue by using PBMCs and liver tissue obtained from biopsy, because occult HCV infection was reported to be associated with increased mortality and nosocomial transmission within the hemodialysis unit [Li and Wang, 2010].

A nationwide study reported that HBV/B was the most common genotype in Indonesia, where the prevalence in some cities on Java, including in Yogyakarta, ranges from 81% to 98% [Haryanto et al., 2008; Mulyanto et al., 2009]. HBV/C is the second most common genotype and is mainly distributed in the northern part of Sulawesi, North Moluccas, and Papua, and is rarely found in Java. HBV/D is the third most common genotype and HBV/A is the fourth most common genotype place with a prevalence of 0.8%. HBV/A shows a very restricted distribution, being found only in Kalimantan, and has never been identified in Java [Mulyanto et al., 2009; Thedja et al., 2011]. Surprisingly, this study showed significant differences in genotype distribution, as the prevalence of HBV/C and HBV/A has increased since earlier studies. The increased prevalence of HBV/C and HBV/A suggests that cross-infection occurs among patients. To our knowledge, this is the first report to identify HBV/A in patients in Java.

HBV/C2 and HBV/A2 were particularly common in patients with occult infection, and the partial HBV genome (nt 1–459) of several strains was identical (YOGHhbv54 and YOGHhbv72; YOGHhbv89 and YOGHhbv90). Thus, occult hepatitis B cases might play an important role in HBV transmission in our hemodialysis unit. To prevent HBV transmission, HBV-DNA should be examined before hemodialysis treatment, although implementing such programs may be hindered by their cost. Alternatively, based on other studies, screening of anti-HBc and anti-HCV might be useful.

In Indonesia, few studies have documented the distribution of HCV genotypes among hemodialysis

patients. In this study, the most prevalent HCV genotype was 1a, differing from that observed in HCV-infected blood donors or patients with liver diseases, where genotype 1b prevailed [Hotta et al., 1994; Widjaja et al., 1995; Soetjipto et al., 1996; Tokita et al., 1996; Inoue et al., 2000; Utama et al., 2008, 2010]. A study conducted in the same hemodialysis unit 16 years before this study showed similar findings, as HCV/1a was dominant (89%) among hemodialysis patients, but was not detected in patients with liver diseases [Hadiwandowo et al., 1994]. The HCV/1a predominance might reflect an outbreak of HCV infection from a common source. To test this hypothesis, the HCV genome of strains obtained from hemodialysis patients was compared, which revealed that many of the strains were similar, or identical. Phylogenetic tree analysis of strains isolated from patients and unrelated strains isolated from the same geographic region and other countries supported this finding. Taken together, the present findings provide convincing evidence for nosocomial transmission of HCV within this hemodialysis unit. As genotype 1 is less responsive to standard therapy than genotypes 2 or 3 [Poynard et al., 1998], clinicians should acknowledge that it will be difficult to treat 98% of the HCV-infected patients, meaning special management approaches are needed.

Several mechanisms may explain the predominance of HCV/1a. This genotype might be adapted for patients with impaired immunity, and may be easily transmitted between patients in a hemodialysis unit. Furthermore, frequent exposure to HCV-contaminated blood, supplies, or surfaces, may favor the emergence of mixed genotype infection. Some evidence suggests that HCV/1a selection occurs following a period of infection. When mixed-genotype infection involving HCV/1a occurs, this subtype tends to prevail and persists as the only subtype during the course of the disease. This phenomenon has been demonstrated in experimental studies using chimpanzees [Okamoto et al., 1994], and has been observed in hemodialysis patients [Qian et al., 2000].

There are many possible routes of viral transmission that may be responsible for nosocomial infection. HBV and HCV cannot pass through the dialyzer membrane to cause cross-infection between patients [de Jong et al., 1992; Hubmann et al., 1995]. However, contaminated blood could be transferred if the dialyzers are not effectively sterilized between patients [Hardy et al., 2000]. Infection should not occur if the dialyzer reuse procedures are followed correctly. Moreover, a large-scale study showed that dialyzer reuse was not a risk factor for HCV seroconversion [Jadoul et al., 1993; Finelli et al., 2005]. The reuse of dialyzers is unlikely to provide a route of nosocomial transmission in the hemodialysis unit. It is suspected that the most likely cause of cross-infection between patients treated in the same hemodialysis unit is cross-contamination of supplies and surfaces (including gloves) through a failure to follow established

infection-control procedures within the hemodialysis unit [KDIGO, 2008]. Unfortunately, the main route of nosocomial transmission could not be determined in the present study. Further studies are therefore needed to assess the possible routes of transmission in this hemodialysis unit.

In conclusion, the prevalence of HBV and HCV infection among hemodialysis patients in Yogyakarta, Indonesia, remains high. Nosocomial transmission might play an important role in infection through a failure to follow infection-control procedures within the hemodialysis unit. Hemodialysis units should implement and adhere to strict infection-control procedures designed to prevent the transmission of HBV and HCV. Such procedures should include hygienic precautions aimed at preventing the transfer of blood or fluids contaminated with blood between patients, either directly or via contaminated equipment or surfaces.

ACKNOWLEDGMENTS

We thank Totok Utoro, Ahmad Hamim Sadewa, Abu Tholib Aman, and Suhardi Darmo Atmodjo for their encouragement and for facilitating sample collection, Teguh Triyono and Umi Solekhah Intansari for their advice, Lini Sunaryo for helping with data collection, and Niken Terate Sekar for reviewing the grammar in the manuscript.

REFERENCES

- Abe A, Inoue K, Tanaka T, Kato J, Kajiyama N, Kawaguchi R, Tanaka S, Yoshida M, Kohara M. 1999. Quantitation of hepatitis B virus genomic DNA by real-time detection PCR. *J Clin Microbiol* 37:2899–2903.
- Akbar N, Basuki B, Mulyanto, Garabrant DH, Sulaiman A, Noer HM. 1997. Ethnicity, socioeconomic status, transfusions and risk of hepatitis B and hepatitis C infection. *J Gastroenterol Hepatol* 12:752–757.
- Akkarathamrongsin S, Praianantathavorn K, Hacharoen N, Theamboonlers A, Tangkijvanich P, Tanaka Y, Mizokami M, Poovorawan Y. 2010. Geographic distribution of hepatitis C virus genotype 6 subtypes in Thailand. *J Med Virol* 82:257–262.
- Anggorowati N, Yano Y, Heriyanto DS, Rinonce HT, Utsumi T, Mulya DS, Subronto YW, Hayashi Y. 2012. Clinical and virological characteristics of hepatitis B or C virus co-infection with HIV in Indonesian patients. *J Med Virol* 84:857–865.
- Arora S, Siddiqui M, Ali M, Barzani Y, Ali S, Kitchloo K, Aggarwal P, Bhatt P, Bhandari R, Maini A, Duffoo F, Arya V. 2011. Aspartate aminotransferase and alanine aminotransferase levels in hemodialysis patients with positive hepatitis B/hepatitis C serology: Are they good predictors of liver injury? *Clin Gastroenterol Hepatol* 9:185.
- Barril G, Castillo I, Arenas MD, Espinosa M, Garcia-Valdecasas J, Garcia-Fernandez N, Gonzalez-Parra E, Alcazar JM, Sanchez C, Diez-Baylon JC, Martinez P, Bartolome J, Carreno V. 2008. Occult hepatitis C virus infection among hemodialysis patients. *J Am Soc Nephrol* 19:2288–2292.
- Budihusodo U, Sulaiman HA, Akbar HN, Lesmana LA, Wasposito AS, Noer HM, Akahane Y, Suzuki H. 1991. Seroepidemiology of HBV and HCV infection in Jakarta, Indonesia. *Gastroenterol Jpn* 26:196–201.
- Burdick RA, Bragg-Gresham JL, Woods JD, Hedderwick SA, Kurokawa K, Combe C, Saito A, LaBrecque J, Port FK, Young EW. 2003. Patterns of hepatitis B prevalence and seroconversion in hemodialysis units from three continents: The DOPPS. *Kidney Int* 63:2222–2229.
- Busek SU, Baba EH, Tavares Filho HA, Pimenta L, Salomao A, Correa-Oliveira R, Oliveira GC. 2012. Hepatitis C and hepatitis B virus infection in different hemodialysis units in Belo Horizonte, Minas Gerais, Brazil. *Mem Inst Oswaldo Cruz* 97:775–778.
- Center for Disease Control and Prevention. 2001. Recommendations for preventing transmission of infections among chronic hemodialysis patients. *MMWR Recomm Rep* 50:1–43.
- Cui C, Shi J, Hui L, Xi H, Zhuoma, Quni, Tsedan, Hu G. 2002. The dominant hepatitis B virus genotype identified in Tibet is a C/D hybrid. *J Gen Virol* 83:2773–2777.
- de Jong GM, de Bruin W, Verresen L, Moshage H, Desmyter J, Yap SH. 1992. High-flux membranes are not permeable to hepatitis B virus-DNA. *Nephron* 60:368.
- Di Stefano M, Volpe A, Stallone G, Tartaglia L, Prato R, Martinelli D, Pastore G, Gesualdo L, Fiore JR. 2009. Occult HBV infection in hemodialysis setting is marked by presence of isolated antibodies to HBcAg and HCV. *J Nephrol* 22:381–386.
- Edey M, Barraclough K, Johnson DW. 2010. Review article: Hepatitis B and dialysis. *Nephrology (Carlton)* 15:137–145.
- Fabrizi F, Martin P. 2008. Occult hepatitis C virus infection in hemodialysis. *J Am Soc Nephrol* 19:2248–2250.
- Fabrizi F, Messa PG, Lunghi G, Aucella F, Bisegna S, Mangano S, Villa M, Barbisoni F, Rusconi E, Martin P. 2005. Occult hepatitis B virus infection in dialysis patients: A multicentre survey. *Aliment Pharmacol Ther* 21:1341–1347.
- Fabrizi F, De Vecchi AF, Qureshi AR, Aucella F, Lunghi G, Bruchfeld A, Bisegna S, Mangano S, Limido A, Vigilante D, Forcella M, Delli Carri P, Martin P. 2007. Gamma glutamyl-transpeptidase activity and viral hepatitis in dialysis population. *Int J Artif Organs* 30:6–15.
- Ferreira RC, Teles SA, Dias MA, Tavares VR, Silva SA, Gomes SA, Yoshida CF, Martins RM. 2006. Hepatitis B virus infection profile in hemodialysis patients in Central Brazil: Prevalence, risk factors, and genotypes. *Mem Inst Oswaldo Cruz* 101:689–692.
- Finelli L, Miller JT, Tokars JI, Alter MJ, Arduino MJ. 2005. National surveillance of dialysis-associated diseases in the United States, 2002. *Semin Dial* 18:52–61.
- Gwak GY, Huh W, Lee DH, Min BH, Koh KC, Kim JJ, Oh HY. 2008. Occult hepatitis B virus infection in chronic hemodialysis patients in Korea. *Hepatogastroenterology* 55:1721–1724.
- Hadiwandowo S. 1991. Infeksi virus hepatitis B pada karyawan rumah sakit di Yogyakarta. *Berita Kedokteran Masyarakat* 7:54–59.
- Hadiwandowo S, Tsuda F, Okamoto H, Tokita H, Wang Y, Tanaka T, Miyakawa Y, Mayumi M. 1994. Hepatitis B virus subtypes and hepatitis C virus genotypes in patients with chronic liver disease or on maintenance hemodialysis in Indonesia. *J Med Virol* 43:182–186.
- Hardy NM, Chiao J, Arora N, Mars R, Jenkins SG. 2000. Hepatitis C virus in the hemodialysis setting: Detecting viral RNA from blood port caps by reverse transcription-polymerase chain reaction. *Clin Nephrol* 54:143–146.
- Haryanto A, Mulyani NS, Widowati T, Wijayanti N, Hadi P. 2008. Molecular genotyping of HBV by using nested PCR-RFLP among hepatitis B patients in Daerah Istimewa Yogyakarta Province and surrounding area. *Indonesian J Biotechnol* 13:1098–1104.
- Hasan I. 2005. Epidemiology of hepatitis B. *Acta Med Indones* 37:231–234.
- Hollinger FB, Habibollahi P, Daneshmand A, Alavian SM. 2010. Occult hepatitis B infection in chronic hemodialysis patients: Current concepts and strategy. *Hepat Mon* 10:199–204.
- Hotta H, Handajani R, Lusida MI, Soemarto W, Doi H, Miyajima H, Homma M. 1994. Subtype analysis of hepatitis C virus in Indonesia on the basis of NS5b region sequences. *J Clin Microbiol* 32:3049–3051.
- Hubmann R, Zazgornik J, Gabriel C, Garbeis B, Blauhut B. 1995. Hepatitis C virus—Does it penetrate the haemodialysis membrane? PCR analysis of haemodialysis ultrafiltrate and whole blood. *Nephrol Dial Transplant* 10:541–542.
- Inoue Y, Sulaiman HA, Matsubayashi K, Julitasari, Iinuma K, Ansari A, Laras K, Corwin AL. 2000. Genotypic analysis of hepatitis C virus in blood donors in Indonesia. *Am J Trop Med Hyg* 62:92–98.
- Jadoul M, Cornu C, van Ypersele de Strihou C. 1993. Incidence and risk factors for hepatitis C seroconversion in hemodialysis: A

- prospective study. The UCL Collaborative Group. *Kidney Int* 44:1322–1326.
- Jain P, Nijhawan S. 2008. Occult hepatitis C virus infection is more common than hepatitis B infection in maintenance hemodialysis patients. *World J Gastroenterol* 14:2288–2289.
- Johnson DW, Dent H, Yao Q, Tranaeus A, Huang CC, Han DS, Jha V, Wang T, Kawaguchi Y, Qian J. 2009. Frequencies of hepatitis B and C infections among haemodialysis and peritoneal dialysis patients in Asia-Pacific countries: Analysis of registry data. *Nephrol Dial Transplant* 24:1598–1603.
- Juniastuti, Utsumi T, Nugrahaputra VE, Amin M, Hayashi Y, Hotta H, Lusida MI. 2011. Another novel subgenotype of hepatitis B virus genotype C from papuans of Highland origin. *J Med Virol* 83:225–234.
- KDIGO. 2008. KDIGO clinical practice guidelines for the prevention, diagnosis, evaluation, and treatment of hepatitis C in chronic kidney disease. *Kidney Int Suppl* 109:S1–S99.
- Kondili LA, Genovese D, Argentini C, Chionne P, Toscani P, Fabro R, Cocconi R, Rapicetta M. 2006. Nosocomial transmission in simultaneous outbreaks of hepatitis C and B virus infections in a hemodialysis center. *Eur J Clin Microbiol Infect Dis* 25:527–531.
- Kramvis A, Kew M, Francois G. 2005. Hepatitis B virus genotypes. *Vaccine* 23:2409–2423.
- Lavanchy D. 2011. Evolving epidemiology of hepatitis C virus. *Clin Microbiol Infect* 17:107–115.
- Li H, Wang SX. 2010. Hepatitis C viral infection in a Chinese hemodialysis unit. *Chin Med J (Engl)* 123:3574–3577.
- Mina P, Georgiadou SP, Rizo C, Dalekos GN, Rigopoulou EI. 2010. Prevalence of occult hepatitis B virus infection in haemodialysis patients from central Greece. *World J Gastroenterol* 16:225–231.
- Minuk GY, Sun DF, Greenberg R, Zhang M, Hawkins K, Uhanova J, Gutkin A, Bernstein K, Giulivi A, Osioy C. 2004. Occult hepatitis B virus infection in a North American adult hemodialysis patient population. *Hepatology* 40:1072–1077.
- Motta JS, Mello FC, Lago BV, Perez RM, Gomes SA, Figueiredo FF. 2010. Occult hepatitis B virus infection and lamivudine-resistant mutations in isolates from renal patients undergoing hemodialysis. *J Gastroenterol Hepatol* 25:101–106.
- Moutinho RS, Perez RM, Medina-Pestana JO, Figueiredo MS, Koide S, Alberto FL, Silva AE, Ferraz ML. 2006. Low HBV-DNA levels in end-stage renal disease patients with HBeAg-negative chronic hepatitis B. *J Med Virol* 78:1284–1288.
- Mulyanto, Depamede SN, Surayah K, Tsuda F, Ichiyama K, Takahashi M, Okamoto H. 2009. A nationwide molecular epidemiological study on hepatitis B virus in Indonesia: Identification of two novel subgenotypes, B8 and C7. *Arch Virol* 154:1047–1059.
- Mulyanto, Depamede SN, Surayah K, Tjahyono AA, Jirintai, Nagashima S, Takahashi M, Okamoto H. 2010. Identification and characterization of novel hepatitis B virus subgenotype C10 in Nusa Tenggara, Indonesia. *Arch Virol* 155:705–715.
- Mulyanto, Depamede SN, Wahyono A, Jirintai, Nagashima S, Takahashi M, Okamoto H. 2011. Analysis of the full-length genomes of novel hepatitis B virus subgenotypes C11 and C12 in Papua, Indonesia. *J Med Virol* 83:54–64.
- Mulyanto, Pancawardani P, Depamede SN, Wahyono A, Jirintai S, Nagashima S, Takahashi M, Nishizawa T, Okamoto H. 2012. Identification of four novel subgenotypes (C13–C16) and two inter-genotypic recombinants (C12/G and C13/B3) of hepatitis B virus in Papua province, Indonesia. *Virus Res* 163:129–140.
- Nurainy N, Muljono DH, Sudoyo H, Marzuki S. 2008. Genetic study of hepatitis B virus in Indonesia reveals a new subgenotype of genotype B in east Nusa Tenggara. *Arch Virol* 153:1057–1065.
- Okamoto H, Mishiro S, Tokita H, Tsuda F, Miyakawa Y, Mayumi M. 1994. Superinfection of chimpanzees carrying hepatitis C virus of genotype II/1b with that of genotype III/2a or I/1a. *Hepatology* 20:1131–1136.
- Poynard T, Marcellin P, Lee SS, Niederau C, Minuk GS, Ideo G, Bain V, Heathcote J, Zeuzem S, Trepo C, Albrecht J. 1998. Randomised trial of interferon alpha2b plus ribavirin for 48 weeks or for 24 weeks versus interferon alpha2b plus placebo for 48 weeks for treatment of chronic infection with hepatitis C virus. *International Hepatitis Interventional Therapy Group (IHIT). Lancet* 352:1426–1432.
- Pujol FH, Ponce JG, Lema MG, Capriles F, Devesa M, Sirit F, Salazar M, Vasquez G, Monsalve F, Blitz-Dorfman L. 1996. High incidence of hepatitis C virus infection in hemodialysis patients in units with high prevalence. *J Clin Microbiol* 34:1633–1636.
- Qian KP, Natov SN, Pereira BJ, Lau JY. 2000. Hepatitis C virus mixed genotype infection in patients on haemodialysis. *J Viral Hepat* 7:153–160.
- Rahayujati TB, Nurdjanah S, Ng N. 2006. Beberapa faktor yang berhubungan dengan kejadian hepatitis B dan C pada pendonor darah. *Berita Kedokteran Masyarakat* 22:33–39.
- Rahnayardi M, Hosseini Moghaddam SM, Alavian SM. 2008. Hepatitis C in hemodialysis patients: Current global magnitude, natural history, diagnostic difficulties, and preventive measures. *Am J Nephrol* 28:628–640.
- Raimondo G, Pollicino T, Cacciola I, Squadrito G. 2007. Occult hepatitis B virus infection. *J Hepatol* 46:160–170.
- Saketi JR, Boland GJ, van Loon AM, van Hattum J, Abdurachman SA, Sukandar E. 2003. Prevalence of hepatitis C virus infection among haemodialysis patients in West Java, Indonesia. *Adv Exp Med Biol* 531:201–209.
- Santoso D, Pranawa, Yogiantoro M, Widodo, Wardana A, Mardiana N, Irwanadi C, Soewanto, Shou I, Maeda K, Hamada C, Fukui M, Horikoshi S, Tomino Y. 2010. Hepatitis C virus infection in hemodialysis patients: Comparison of The Surabaya Dialysis Center and Juntendo University Hospital Dialysis Center. *Indones J Trop Infect Dis* 1:105–109.
- Simmonds P, Bukh J, Combet C, Deleage G, Enomoto N, Feinstone S, Halfon P, Inchauste G, Kuiken C, Maertens G, Mizokami M, Murphy DG, Okamoto H, Pawlotsky JM, Penin F, Sablon E, Shin IT, Stuyver LJ, Thiel HJ, Viazov S, Weiner AJ, Widell A. 2005. Consensus proposals for a unified system of nomenclature of hepatitis C virus genotypes. *Hepatology* 42:962–973.
- Soetjipto, Handajani R, Lusida MI, Darmadi S, Adi P, Soemarto, Ishido S, Katayama Y, Hotta H. 1996. Differential prevalence of hepatitis C virus subtypes in healthy blood donors, patients on maintenance hemodialysis, and patients with hepatocellular carcinoma in Surabaya, Indonesia. *J Clin Microbiol* 34:2875–2880.
- Souza JF, Longui CA, Miorin LA, Sens YA. 2008. Gamma-glutamyltransferase activity in chronic dialysis patients and renal transplant recipients with hepatitis C virus infection. *Transplant Proc* 40:1319–1323.
- Sugauchi F, Mizokami M, Orito E, Ohno T, Kato H, Suzuki S, Kimura Y, Ueda R, Butterworth LA, Cooksley WG. 2001. A novel variant genotype C of hepatitis B virus identified in isolates from Australian Aborigines: Complete genome sequence and phylogenetic relatedness. *J Gen Virol* 82:883–892.
- Theджа MD, Roni M, Harahap AR, Siregar NC, Ie SI, Muljono DH. 2010. Occult hepatitis B in blood donors in Indonesia: Altered antigenicity of the hepatitis B virus surface protein. *Hepatol Int* 4:608–614.
- Theджа MD, Muljono DH, Nurainy N, Sukowati CH, Verhoef J, Marzuki S. 2011. Ethnogeographical structure of hepatitis B virus genotype distribution in Indonesia and discovery of a new subgenotype, B9. *Arch Virol* 156:855–868.
- Tokita H, Okamoto H, Iizuka H, Kishimoto J, Tsuda F, Lesmana LA, Miyakawa Y, Mayumi M. 1996. Hepatitis C virus variants from Jakarta, Indonesia classifiable into novel genotypes in the second (2e and 2f), tenth (10a) and eleventh (11a) genetic groups. *J Gen Virol* 77:293–301.
- Triwibowo. 1993. Anti-HIV, anti-HCV, syphilis, HBsAg serologic tests among high-risk groups and blood donors in Yogyakarta, Indonesia. *Southeast Asian J Trop Med Public Health* 24:275–277.
- Tu AW, Buxton JA, Whitlock M, Djurdjev O, Chong M, Kraiden M, Beaulieu M, Levin A. 2009. Prevalence and incidence of hepatitis C virus in hemodialysis patients in British Columbia: Follow-up after a possible breach in hemodialysis machines. *Can J Infect Dis Med Microbiol* 20:e19–e23.
- Utama A, Budiarto BR, Monasari D, Octavia TI, Chandra IS, Gani RA, Hasan I, Sanityoso A, Miskand UA, Yusuf I, Lesmana LA, Sulaiman A, Tai S. 2008. Hepatitis C virus genotype in blood donors and associated liver disease in Indonesia. *Intervirol* 51:410–416.
- Utama A, Tania NP, Dhenni R, Gani RA, Hasan I, Sanityoso A, Lelosutan SA, Martamala R, Lesmana LA, Sulaiman A, Tai S. 2010. Genotype diversity of hepatitis C virus (HCV) in HCV-

- associated liver disease patients in Indonesia. *Liver Int* 30:1152–1160.
- Utsumi T, Lusida MI, Yano Y, Nugrahaputra VE, Amin M, Juniastuti, Soetjipto, Hayashi Y, Hotta H. 2009. Complete genome sequence and phylogenetic relatedness of hepatitis B virus isolates in Papua, Indonesia. *J Clin Microbiol* 47:1842–1847.
- Utsumi T, Yano Y, Lusida MI, Amin M, Soetjipto, Hotta H, Hayashi Y. 2010. Serologic and molecular characteristics of hepatitis B virus among school children in East Java, Indonesia. *Am J Trop Med Hyg* 83:189–193.
- Vladutiu DS, Cosa A, Neamtu A, State D, Braila M, Gherman M, Patiu IM, Dulau-Florea I. 2000. Infections with hepatitis B and C viruses in patients on maintenance dialysis in Romania and in former communist countries: Yellow spots on a blank map? *J Viral Hepat* 7:313–319.
- Widjaja S, Li S, Ali S, Simon S, Sulaiman A, Lesmana LA, Yap SH. 1995. Hepatitis C virus RNA detection and HCV genotype in patients with chronic non-A, non-B hepatitis in Jakarta. *J Virol Methods* 51:169–175.
- Wong DK, Huang FY, Lai CL, Poon RT, Seto WK, Fung J, Hung IF, Yuen MF. 2011. Occult hepatitis B infection and HBV replicative activity in patients with cryptogenic cause of hepatocellular carcinoma. *Hepatology* 54:829–836.
- Yasuda K, Okuda K, Endo N, Ishiwatari Y, Ikeda R, Hayashi H, Yokozeki K, Kobayashi S, Irie Y. 1995. Hypoaminotransferase-mia in patients undergoing long-term hemodialysis: Clinical and biochemical appraisal. *Gastroenterology* 109:1295–1300.

Hepatitis B and C virus infection among hemodialysis patients in Yogyakarta, Indonesia- Prevalence and molecular evidence for nosocomial transmission

ORIGINALITY REPORT

25%

SIMILARITY INDEX

17%

INTERNET SOURCES

22%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1

"The 21st Conference of the Asian Pacific Association for the Study of the Liver", Hepatology International, 2011

Publication

1%

2

"Guideline 3: Preventing HCV transmission in hemodialysis units", Kidney International, 2008

Publication

1%

3

Quang Duy Trinh. "Sequence analysis of the VP7 gene of human rotaviruses G2 and G4 isolated in Japan, China, Thailand, and Vietnam during 2001-2003", Journal of Medical Virology, 05/2010

Publication

1%

4

Replacement of Renal Function by Dialysis, 2004.

Publication

1%

5

Ian Baudi, Sayuki Iijima, Nyasha Chin'ombe, Sekesai Mtapuri-Zinyowera et al. "Molecular

1%

epidemiology of co-infection with hepatitis B virus and human immunodeficiency virus (HIV) among adult patients in Harare, Zimbabwe", *Journal of Medical Virology*, 2017

Publication

6	hcvets.com Internet Source	1%
7	www.e-cmh.org Internet Source	<1%
8	spandidos-publications.com Internet Source	<1%
9	sintadev.ristekdikti.go.id Internet Source	<1%
10	jasn.asnjournals.org Internet Source	<1%
11	www.scitechnol.com Internet Source	<1%
12	Abdelrahim, Soha S., Rasha M. M. Khiry, Mona Abdel-Monem Esmail, Mahmoud Ragab, Mohamed Abdel-Hamid, and Sayed F. Abdelwahab. "Occult hepatitis C virus infection among Egyptian hemodialysis patients : Occult HCV among Egyptian hemodialysis patients", <i>Journal of Medical Virology</i> , 2016. Publication	<1%

13	www.jourlib.org Internet Source	<1%
14	Eva Mortensen, Amanda Kamali, Patricia L. Schirmer, Cynthia Lucero-Obusan et al. "Are current screening protocols for chronic hepatitis B virus infection adequate?", Diagnostic Microbiology and Infectious Disease, 2016 Publication	<1%
15	www.jidc.org Internet Source	<1%
16	pdfs.semanticscholar.org Internet Source	<1%
17	repositorio.ufpa.br Internet Source	<1%
18	www.koreascience.or.kr Internet Source	<1%
19	repositorio.bc.ufg.br Internet Source	<1%
20	www.ijpmonline.org Internet Source	<1%
21	wnt-signaling.com Internet Source	<1%
22	VIPA THANACHARTWET. "Viral hepatitis infections among dialysis patients: Thailand	<1%

registry report", Nephrology, 8/2007

Publication

23	academic.oup.com Internet Source	<1%
24	Choong Keun Cha. "Association of lamivudine-resistant mutational patterns with the antiviral effect of adefovir in patients with chronic hepatitis B", Journal of Medical Virology, 03/2009 Publication	<1%
25	www.mdpi.com Internet Source	<1%
26	s3.amazonaws.com Internet Source	<1%
27	article.sapub.org Internet Source	<1%
28	"Posters", Clinical Microbiology and Infection, 4/2007 Publication	<1%
29	zombiedoc.com Internet Source	<1%
30	www.mrinstitute.org Internet Source	<1%
31	dro.dur.ac.uk Internet Source	<1%

32

renal.org

Internet Source

<1%

33

Emad Abd-Allah. "Incidence of Hepatitis C Virus (HCV), Hepatitis B Virus (HBV) and Dual Infection in Egyptian Patients on Haemodialysis", *Kidney*, 02/23/2010

Publication

<1%

34

onlinelibrary.wiley.com

Internet Source

<1%

35

biosains.mipa.uns.ac.id

Internet Source

<1%

36

Seyed-Moayed Alavian, Seyed-Vahid Tabatabaei, Nastaran Mahboobi. "Epidemiology and risk factors of HCV infection among hemodialysis patients in countries of the Eastern Mediterranean Regional Office of WHO (EMRO): a quantitative review of literature", *Journal of Public Health*, 2010

Publication

<1%

37

www.cdc.gov

Internet Source

<1%

38

docplayer.com.br

Internet Source

<1%

39

www.kdigo.org

Internet Source

<1%

40

www.researchsquare.com

Internet Source

<1%

41

Kaoru Inokuchi, Keisuke Nakata, Keisuke Hamasaki, Manabu Daikoku et al. "Prevalence of hepatitis B or C virus infection in patients with fulminant viral hepatitis. An analysis using polymerase chain reaction", *Journal of Hepatology*, 1996

Publication

<1%

42

Aghakhani, Arezoo, Mohammad Banifazl, Ali Akbar Velayati, Ali Eslamifar, and Amitis Ramezani. "Occult Hepatitis B Virus Infection in Hemodialysis Patients: A Concept for Consideration : Occult HBV in Hemodialysis Patients", *Therapeutic Apheresis and Dialysis*, 2012.

Publication

<1%

43

Livia Melo Villar, Helena Medina Cruz, Moyra Machado Portilho, Jakeline Ribeiro Barbosa et al. "Chapter 1 Emergence of New Epidemiological Hepatitis B and C Profiles in High Risk Groups in Latin America", *IntechOpen*, 2018

Publication

<1%

44

link.springer.com

Internet Source

<1%

45	repub.eur.nl Internet Source	<1%
46	archive.org Internet Source	<1%
47	ousar.lib.okayama-u.ac.jp Internet Source	<1%
48	clinics.org.br Internet Source	<1%
49	preview-breast-cancer-research.biomedcentral.com Internet Source	<1%
50	journals.plos.org Internet Source	<1%
51	Abeer Elkady. "Virological and clinical implication of core promoter C1752/V1753 and T1764/G1766 mutations in hepatitis B virus genotype D infection in Mongolia", <i>Journal of Gastroenterology and Hepatology</i> , 3/2008 Publication	<1%
52	vactruth.com Internet Source	<1%
53	Fumio Tsuda, Soeliadi Hadiwandowo, Naoto Sawada, Masako Fukuda, Takeshi Tanaka, Hiroaki Okamoto, Yuzo Miyakawa, Makoto Mayumi. "Infection with GB virus C (GBV-C) in	<1%

patients with chronic liver disease or on maintenance hemodialysis in Indonesia", *Journal of Medical Virology*, 1996

Publication

54

unair.ac.id

Internet Source

<1%

55

budilukmanto.org

Internet Source

<1%

56

jcm.asm.org

Internet Source

<1%

57

K Takaguchi. "Detection of hepatitis B virus DNA in the liver and serum of patients with hepatitis B surface antigen and hepatitis C virus antibody negative chronic liver disease", *Hepatology Research*, 2002

Publication

<1%

58

T. G. Wreghitt. "Blood-borne virus infections in dialysis units—A review", *Reviews in Medical Virology*, 04/1999

Publication

<1%

59

Atsunori Kusakabe. "Virological features of hepatitis B virus-associated nephropathy in Japan", *Journal of Medical Virology*, 09/2007

Publication

<1%

60

www.wvdhhr.org

Internet Source

<1%

61	jcancer.org Internet Source	<1%
62	Renata C. Ferreira. "Prevalence of hepatitis B virus and risk factors in Brazilian non-injecting drug users", <i>Journal of Medical Virology</i> , 04/2009 Publication	<1%
63	www.cari.org.au Internet Source	<1%
64	www.spandidos-publications.com Internet Source	<1%
65	Zhifeng Li, Lunbiao Cui, Minghao Zhou, Xian Qi et al. "Development and application of a one-step real-time RT-PCR using a minor-groove-binding probe for the detection of a novel bunyavirus in clinical specimens", <i>Journal of Medical Virology</i> , 2013 Publication	<1%
66	M Izuma. "In vitro cytokine production of peripheral blood mononuclear cells in response to HCV core antigen stimulation during interferon- \hat{I}^2 treatment and its relevance to sCD8 and sCD30", <i>Hepatology Research</i> , 2000 Publication	<1%
67	www.wjgnet.com Internet Source	<1%

68

de Jesus Rodrigues de Freitas, Maria, Amanda Alves Fecury, Marcella Kelly Costa de Almeida, Andrei Silva Freitas, Vanessa de Souza Guimarães, Andrea Marinho da Silva, Ygor Fernando Ferreira da Costa, Renata Aparecida Andrade da Costa, Patrícia Ferreira, and Luisa Caricio Martins. "Prevalence of hepatitis C virus infection and genotypes in patient with chronic kidney disease undergoing hemodialysis : HCV in Hemodialysis Patients", *Journal of Medical Virology*, 2013.

Publication

<1%

69

Nagakawa, Osamu, Yoshiho Miyatomi, Yasuhiro Shigeta, Emi Inayama, Kouichi Murakami, Takehiko Sakai, Takashi Kouno, Motoyuki Masai, Kohji Shirai, and Toyohiko Yoshida. "Occult Hepatitis B Virus Infection in Japanese Chronic Hemodialysis Patients : Occult HBV in Japanese HD Patients", *Therapeutic Apheresis and Dialysis*, 2013.

Publication

<1%

70

www.apocpcontrol.org

Internet Source

<1%

71

hrcak.srce.hr

Internet Source

<1%

72

appswl.elsevier.es

Internet Source

<1%

73 Hui-Lan Zhu, Xu Li, Jun Li, Zhen-Hua Zhang. <1%
"Genetic variation of occult hepatitis B virus
infection", World Journal of Gastroenterology,
2016
Publication

74 numonthly.com <1%
Internet Source

75 naosite.lb.nagasaki-u.ac.jp <1%
Internet Source

76 journals.sagepub.com <1%
Internet Source

77 journals.lww.com <1%
Internet Source

78 Leroux-Roels, Geert, and Annemieke Dhondt. <1%
"Hepatitis and Dialysis", Complications of
Dialysis, 2000.
Publication

79 Hui, C.k.. "Occult hepatitis B virus infection in
hematopoietic stem cell donors in a hepatitis B
virus endemic area", Journal of Hepatology,
200506
Publication

80 www.oncotarget.com <1%
Internet Source

81 Serwah , Abdel Hamid A. | Mohamed , Waleed

S. | Serwah , Mohamed | Edreis , Awateif | ElZaydi , Ahmad. "Occult Hepatitis C Virus Infection in Haemodialysis Unit : A Single-Center Experience \ The Egyptian Journal of Hospital Medicine .- 2014 , Vol. 56 , pp. 280 - 288.", The Egyptian Journal of Hospital Medicine, 2014

<1%

Publication

82

www.bjamp.org

Internet Source

<1%

83

www.medica.ro

Internet Source

<1%

84

jcspk.pk

Internet Source

<1%

85

Yuko Iwasa, Shigeru Otsubo, Oriie Sugi, Keitaro Sato et al. "Patterns in the prevalence of hepatitis C virus infection at the start of hemodialysis in Japan", Clinical and Experimental Nephrology, 2008

<1%

Publication

86

Mederacke, I., M. Meier, J. B. Luth, H. Schmidt-Gurtler, R. Raupach, R. Horn-Wichmann, K. Wursthorn, A. Potthoff, G. Colucci, M. P. Manns, H. Wedemeyer, and H. L. Tillmann. "Different kinetics of HBV and HCV during haemodialysis and absence of seronegative viral hepatitis in patients with end-stage renal disease",

<1%

Nephrology Dialysis Transplantation, 2011.

Publication

-
- | | | |
|----|--|-----|
| 87 | Hiroshi Kamitsukasa, Masanobu Iri, Akihisa Tanaka, Shigeo Nagashima, Masaharu Takahashi, Tsutomu Nishizawa, Hiroaki Okamoto. "Spontaneous reactivation of hepatitis B virus (HBV) infection in patients with resolved or occult HBV infection", Journal of Medical Virology, 2015
Publication | <1% |
| 88 | www.scialert.net
Internet Source | <1% |
| 89 | creativecommons.org
Internet Source | <1% |
| 90 | asbmr.onlinelibrary.wiley.com
Internet Source | <1% |
| 91 | Sanchez-Tapias, JoseM.. "Nosocomial transmission of hepatitis C virus", Journal of Hepatology, 1999
Publication | <1% |
| 92 | pure.uva.nl
Internet Source | <1% |
| 93 | dokter-medik.blogspot.com
Internet Source | <1% |
| 94 | Lara B. Lemos. "Clinical and Laboratory | <1% |

Characteristics of Acute Hepatitis C in Patients With End-stage Renal Disease on Hemodialysis", Journal of Clinical Gastroenterology, 02/2008

Publication

95

Su, Yingying, Jessie L. Norris, Chunpeng Zang, Zhihang Peng, and Ning Wang. "Incidence of hepatitis C virus infection in patients on hemodialysis: A systematic review and meta-analysis : Meta-analysis of HCV in hemodialysis", Hemodialysis International, 2012.

<1%

Publication

96

Cibele Franz, Renata de Mello Perez, Mariano Gustavo Zalis, Ana Carolina Jonard Zalona et al. "Prevalence of occult hepatitis B virus infection in kidney transplant recipients", Memórias do Instituto Oswaldo Cruz, 2013

<1%

Publication

97

Francesco Vitale. "Can the serological status of "anti-HBc alone" be considered a sentinel marker for detection of "occult" HBV infection?", Journal of Medical Virology, 04/2008

<1%

Publication

98

scholarbank.nus.edu.sg

Internet Source

<1%

99

Manoochehr Makvandi. "Update on occult hepatitis B virus infection", World Journal of

<1%

Gastroenterology, 2016

Publication

-
- 100** Andi Utama. "Low prevalence of hepatitis B virus pre-s deletion mutation in Indonesia", *Journal of Medical Virology*, 10/2011
Publication <1%
-
- 101** Bin Zhou. "Geographical and Ethnic Distribution of the HBV C/D Recombinant on the Qinghai-Tibet Plateau", *PLoS ONE*, 04/11/2011
Publication <1%
-
- 102** Shiv K. Sarin. "Epidemiology, Screening, and Natural History of Chronic Hepatitis B Infection", *Chronic Viral Hepatitis*, 2009
Publication <1%
-
- 103** "APASL Seoul 2008 Meeting", *Hepatology International*, 2008
Publication <1%
-
- 104** keyvanlab.com
Internet Source <1%
-
- 105** Dzekova Vidimliski, Pavlina, Igor Nikolov, Nadica Matevska Geshkovska, Aleksandar Dimovski, Lionel Rostaing, and Aleksandar Sikole. "Review: Occult hepatitis C virus infection: Still remains a controversy : Occult Hepatitis C Remains a Controversy", *Journal of Medical Virology*, 2014.
Publication <1%
-

106 "Guideline 2: Treatment of HCV infection in patients with CKD", *Kidney International*, 2008 <1%
Publication

107 Fabrizio Fabrizi. "Management of hepatitis B and C virus infection before and after renal transplantation", *Current Opinion in Organ Transplantation*, 12/2006 <1%
Publication

108 Bira Tsatsralt-Od, Masaharu Takahashi, Tsutomu Nishizawa, Kazunori Endo, Jun Inoue, Hiroaki Okamoto. "High prevalence of dual or triple infection of hepatitis B, C, and delta viruses among patients with chronic liver disease in Mongolia", *Journal of Medical Virology*, 2005 <1%
Publication

109 Hiroaki Okamoto, Satoyuki Kobata, Hajime Tokita, Taisuke Inoue et al. "A second-generation method of genotyping hepatitis C virus by the polymerase chain reaction with sense and antisense primers deduced from the core gene", *Journal of Virological Methods*, 1996 <1%
Publication

110 F. Fabrizi, A.F. De Vecchi, A.R. Qureshi, F. Aucella et al. "Gamma Glutamyltranspeptidase Activity and Viral Hepatitis in Dialysis" <1%

Population", The International Journal of Artificial Organs, 2018

Publication

- 111** Nashwan M.R. Ibrahim, Zana Sidiq Mohammed Saleem, Nawfal R Hussein. "The Prevalence of HIV, HCV, and HBV Among Hemodialysis Patients Attending Duhok Hemodialysis Center", International Journal of Infection, 2017 <1%
- Publication
-

- 112** Mahmoud Reza Pourkarim, Valentijn Vergote, Samad Amini-Bavil-Olyaei, Zohre Sharifi et al. "Molecular characterization of hepatitis B virus (HBV) strains circulating in the northern coast of the Persian Gulf and its comparison with worldwide distribution of HBV subgenotype D1", Journal of Medical Virology, 2014 <1%
- Publication
-

- 113** M Takatori. "Mutations in the non-structural protein 5A gene in patients with chronic hepatitis C virus 1b infection during repeated interferon treatment", Hepatology Research, 2000 <1%
- Publication
-

- 114** S Sainokami. "Pretreatment hepatitis C virus dynamics for predicting virological response to interferon- α 2b monotherapy in patients with chronic hepatitis C virus infection", Hepatology Research, 2003 <1%
- Publication

115 Solange Z Freitas, Rivaldo V da Cunha, Regina MB Martins, Sheila A Teles, Mary L Ibanhes, Ana RC Motta-Castro. "Prevalence, genotypes and risk factors associated with hepatitis C virus infection in hemodialysis patients in Campo Grande, MS, Brazil", Memórias do Instituto Oswaldo Cruz, 2008

Publication

116 ELISABETH LAMPE. "Molecular analysis and patterns of ALT and hepatitis C virus seroconversion in haemodialysis patients with acute hepatitis", Nephrology, 4/2008

Publication

117 Selma A., Francisco C., Natalia M.. "Chapter 2 Hepatitis B Virus (HBV) Variants in Hemodialysis Patients", IntechOpen, 2011

Publication

118 J. Samal. "Molecular Mechanisms Underlying Occult Hepatitis B Virus Infection", Clinical Microbiology Reviews, 01/01/2012

Publication

119 Haldun Selcuk, Mehmet Kanbay, Murat Korkmaz, Gurden Gur, Ali Akcay, Hande Arslan, Nurhan Ozdemir, Ugur Yilmaz, Sedat Boyacioglu. "Distribution of HCV Genotypes in Patients with End-Stage Renal Disease According to Type of Dialysis Treatment",

120 Brian J.G. Pereira. "Hepatitis C Virus Infection in Dialysis: A Continuing Problem", Artificial Organs, 1/1999 $<1\%$
Publication

121 Masato Komori, Nobukazu Yuki, Takayuki Nagaoka, Masatoshi Yamashiro et al. "Long-term clinical impact of occult hepatitis B virus infection in chronic hepatitis B patients", Journal of Hepatology, 2001 $<1\%$
Publication

122 Mulyanto, Pingki Pancawardani, Sulaiman Ngongu Depamede, Arif Wahyono et al. "Identification of four novel subgenotypes (C13–C16) and two inter-genotypic recombinants (C12/G and C13/B3) of hepatitis B virus in Papua Province, Indonesia", Virus Research, 2012 $<1\%$
Publication

123 Añteneli, Añzcan. "Hepatitis B-related events in autologous hematopoietic stem cell transplantation recipients", World Journal of Gastroenterology, 2010. $<1\%$
Publication

124 Hartono Gunardi, Melanie Y. Iskandar, Turyadi, Susan I. Ie, Pramita G. Dwipoerwantoro, Rino A. $<1\%$

Gani, David H. Muljono. "Hepatitis B virus infection in children of HBV-related chronic liver disease patients: a study of intra-familial HBV transmission", *Hepatology International*, 2016

Publication

125

Gloria M. Calderón. "Prevalence and risk factors of hepatitis C virus, hepatitis B virus, and human immunodeficiency virus in multiply transfused recipients in Mexico", *Transfusion*, 10/2009

<1%

Publication

126

I Gede Putu Surya. "Serological markers of hepatitis B, C, and E viruses and human immunodeficiency virus type-1 infections in pregnant women in Bali, Indonesia", *Journal of Medical Virology*, 04/2005

<1%

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Hepatitis B and C virus infection among hemodialysis patients in Yogyakarta, Indonesia- Prevalence and molecular evidence for nosocomial transmission

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14
