

High rate of seronegative HCV infection in HIV-positive patients

by Maria Lusida

Submission date: 23-Jun-2020 09:27PM (UTC+0800)

Submission ID: 1348568386

File name: rate_of_seronegative_HCV_infection_in_HIV-positive_patients.pdf (362.35K)

Word count: 5159

Character count: 27231

High rate of seronegative HCV infection in HIV-positive patients

JUNIASTUTI^{1,2}, TAKAKO UTSUMI^{2,5}, NASRONUDIN^{2,3,6}, LINDAWATI ALIMASARDJONO¹, MOCHAMAD AMIN^{2,3}, MYRNA ADIANTI³, YOSHIHIKO YANO⁵, SOETJIPTO^{2,4}, YOSHITAKE HAYASHI⁵, HAK HOTTA⁵ and MARIA INGE LUSIDA^{1,2,3}

¹Department of Microbiology, School of Medicine, Airlangga University, Surabaya, East Java 60131;

²Indonesia-Japan Collaborative Research Center for Emerging and Re-emerging Infectious Diseases;

³Institute of Tropical Disease, Airlangga University, Surabaya, East Java 60115; ⁴Department of Biochemistry, School of Medicine, Airlangga University, Surabaya, East Java 60131, Indonesia; ⁵Center for Infectious Diseases, Kobe University Graduate School of Medicine, Kobe, Hyogo 6500017, Japan; ⁶Department of Internal Medicine, Dr Soetomo General Hospital, Surabaya, East Java 60131, Indonesia

Received September 25, 2013; Accepted October 18, 2013

DOI: 10.3892/br.2013.188

15

Abstract. Co-infection with human immunodeficiency virus (HIV) and hepatitis C virus (HCV) is a significant global health problem. The two viruses are transmitted with high efficacy via blood-to-blood contact, mainly intravenous drug use (IVDU), whereas HCV is less easily transmitted sexually. Antibody testing is the main screening method for HCV infection, although it may not be the optimal option for HIV infection. The aim of this study was to investigate HCV infection in HIV-positive patients, with and without a detectable anti-HCV antibody response. A total of 187 plasma samples were obtained from HIV-positive patients in Surabaya, Indonesia and examined for anti-HCV [HCV enzyme immunoassay (EIA) 3.0], HCV genotype/subtype [reverse transcription-polymerase chain reaction (RT-PCR) using primers targeting a part of NS5B/5'UTR followed by sequencing] and HCV viral load (quantitative RT-PCR). A total of 119 patients (63.6%) were found to be anti-HCV-positive and, among these, HCV RNA was detected in 73 (61.3%), with HCV-1a as the predominant subtype (31.5%). Of the 68 anti-HCV-negative samples, HCV RNA was detected in 26/68 (38.2%) mostly as the HCV-3a subtype (50%). High HCV viral loads were more common among the HCV-seropositive patients. The HCV-seropositive samples with detected HCV RNA were mostly obtained from HIV-positive patients with parenteral transmission (IVDU) (76.7%); however, the HCV-seronegative samples with detected HCV RNA were mostly from patients

who had acquired HCV through heterosexual transmission (61.5%). In conclusion, HIV-positive patients were at high risk of becoming co-infected with HCV and several remained HCV-seronegative. Furthermore, there may exist differences in HCV seropositivity and subtypes between HIV-positive patients who acquired HCV sexually and those who acquired HCV parenterally.

Introduction

The epidemic of human immunodeficiency virus (HIV) infection in Asia, including Indonesia, is rapidly expanding (1). The introduction of highly active antiretroviral therapy (HAART) has markedly reduced HIV-related morbidity and mortality. However, non-HIV-related conditions, particularly liver disease, currently constitute an increasingly high proportion of the causes of mortality among HIV-infected individuals (2). Hepatitis C virus (HCV) has emerged as an important cause of morbidity and mortality among HIV-positive individuals (3).

As the majority of individuals who acquire HCV are asymptomatic, it is difficult to determine some of the characteristics of acute infection (4). Early diagnosis is rare and the extent of this epidemic is unknown, since the majority of at-risk individuals are not tested for acute HCV infection (5). These and several other aspects of HCV infection may be further complicated by co-infection with HIV-1. In HIV-infected individuals, untreated acute HCV infection typically progresses to chronic HCV infection, a leading cause of non-AIDS-related morbidity and mortality among HIV-infected individuals in the HAART era (2).

HIV and HCV share common transmission pathways, which may explain the high rate of co-infection with the two viruses. Of the 33.4 million HIV-infected individuals worldwide in 2008, it is estimated that ≥5 million have concomitant HCV infection. Whereas the two viruses are transmitted with high efficacy via blood-to-blood contact [particularly in intra-

45

Correspondence to: Professor Maria Inge Lusida, Institute of Tropical Disease, Airlangga University, Campus C Jalan Muljorejo, Surabaya, East Java 60115, Indonesia
E-mail: ingelusida@yahoo.com

35

Key words: hepatitis C virus, subtypes, anti-hepatitis C virus, human immunodeficiency virus co-infection

venous drug users (IVDUs), HCV is less easily transmitted sexually and its risk remains controversial (6).

Antibody testing is the main screening method for HCV infection (7). However, serological screening in HIV-infected patients may not be the optimal screening method, possibly as a result of immunosuppression (8). Therefore, HCV RNA testing is recommended for the diagnosis of HCV infection (8,9).

The aim of this study was to investigate HCV infection in anti-HCV-positive and -negative HIV patients in Surabaya, Indonesia.

⁶⁸ Materials and methods

Collection of field samples. Plasma samples were obtained from HIV-positive patients, who visited the Institute of Tropical Disease (ITD), Airlangga University, Surabaya, Indonesia, for an HIV viral load examination requested by a clinician. The majority of the patients (176/187, 94%) were on HAART with activity against AIDS (lamivudine+zidovudine+efavirenz or lamivudine+zidovudine+nevirapine) and exhibited no symptoms of acute hepatitis. The plasma samples were stored at -80°C prior to examination. The study protocol was reviewed and approved by the Ethics Committees of Kobe University, Japan and Airlangga University, Indonesia and informed consent was obtained from all the patients. The HIV viral load data were retrieved from the patient database maintained at ITD, Airlangga University, Indonesia.

Anti-HCV tests. All the plasma samples were subjected to HCV enzyme immunoassay (EIA) 3.0 (Hepalisa Anti HCV; PT Indec Diagnostics, Jakarta, Indonesia) to detect anti-HCV, according to the manufacturer's instructions. The third-generation anti-HCV test which detects multiple antigenic determinants (core, NS3, NS4 and NS5) was used to increase sensitivity.

Viral RNA extraction, reverse transcription-polymerase chain reaction (RT-PCR) amplification and sequencing. HCV RNA was extracted from 140 µl plasma using a commercially available kit (QIAamp Viral RNA kit; Qiagen, Tokyo, Japan).

For the amplification of the NS5B region of the HCV genome, the extracted RNA was reverse-transcribed and amplified using SuperScript One-Step RT-PCR (Invitrogen, Tokyo, Japan) and a set of primers. The reaction was initially performed at 45°C for 30 min for RT and at 94°C for 2 min, followed by the first-round PCR over 40 cycles, with each cycle consisting of denaturation at 94°C for 1 min, annealing at 45°C for 1.5 min and extension at 72°C for 2 min, using outer primers 166 (5'-TGGGGATCCCGTATGATACCCGCTGCTTTGA-3', nt 8230-8260, +) and 167R (5'-GGCGGAATTCCTGGTCATAGCCTCCGTGAA-3', nt 8601-8630, -). All the PCRs were performed using Platinum Taq DNA polymerase (Invitrogen Life Technologies, Carlsbad, CA, USA). The second-round PCRs were performed under the same conditions to amplify the HCV genome, using different sets of primers as follows: i) HC23 (5'-TTTGACTCAACCGTCACTGA-3', nt 8256-8275, +), HC24 (5'-CTCAGGCTCGCCGCATCCTC-3', nt 8577-8596, -) and HC26 (5'-CTCAGGTTCCGCTCGTCTC-3', nt 8577-8596, -); ii) HC15 (5'-ACT

GTCCTGAACAGGACAT-3', nt 8265-8284, +) and HC16 (5'-GCTCTATCCTCATCGACGCC-3', nt 8568-8587, -); iii) HC23 and HC28 (5'-CACGAGCATGGTGCAGTCCGGAGC-3', nt 8507-8531, -); iv) HC23 and HC32 (5'-AGGTAGCACGTCAGCGTGTTC-3', nt 8454-8476, -); and v) HC23 and HC34 (5'-TAGCACGTCATGGTGTTC-3', nt 8451-8473, -) (10). If the result of amplification of the NS5B region was negative, the extracted RNA was reverse transcribed and amplified using the same condition but using a different set of primers as follows: i) UTR1 (5'-CCGGAGAGCCATAGTGGTC-3', +) and UTR2 (5'-AGTACCACAAGGCCTTTCGC-3', -) (first-round PCR); ii) UTR3 (5'-TGGTCTGCGGAACCGGTGAG-3', +) and UTR4 (5'-ACCCAACACTACTCGGCTAG-3', -) (second-round PCR) (11).

The PCR products were electrophoresed in a 2% agarose gel containing ethidium bromide and were visualized under UV illumination. Amplified cDNA fragments were sequenced by a direct sequencing method with the BigDye Terminator v1.1 Cycle Sequencing kit and an ABI Prism 310 sequencer (Applied Biosystems, Foster City, CA, USA).

Sequence analysis. Each obtained sequence was compared to those of the reported subtypes and on the basis of the percent homologies on the NS5B region, each isolate was assigned a subtype (10,12). When a subtype assignment was not possible due to the lack of NS5B amplification, the nucleotide sequences of the 5'UTR region were determined and compared with the consensus sequence motifs for each of the major genotypes previously reported (13).

Quantification of plasma HCV RNA titers. The quantification of plasma HCV RNA titers was performed with TaqMan® Gene Expression Master Mix (Applied Biosystems) using Applied Biosystems 7300 Real-Time PCR system.

Results

Samples. A total of 187 plasma samples were obtained from 187 HIV-positive patients (153 men and 34 women; mean age, 29.4 years; age range, 3 months-58 years). The risk factors for infection were IVDU (62.6%), heterosexual and homosexual intercourse (35.8 and 1.1%, respectively) and vertical transmission (0.5%) (Table I).

A total of 119 patients (63.6%) were anti-HCV-positive and the remaining 68 (36.4%) were anti-HCV-negative. The group of HCV-seropositive patients was most likely (73.1%) to have acquired HCV through parenteral transmission (IVDU), followed by heterosexual transmission (26.9%) (Table I).

HCV RNA detection. HCV RNA was detected in 73 of the 119 anti-HCV-positive samples (61.3%); among these, HCV-1a (31.5%) was the predominant subtype, followed by 3a (23.3%), 1 (16.4%), 1c (10.9%), 1b (9.6%), 3k (4.1%) and 2, 2a and 4a (1.4% each). Of the 68 anti-HCV-negative samples, HCV RNA was detected in 26 (38.2%), with HCV-3a being the most prevalent subtype (50%), followed by 3k (23.1%), 1c (15.4%) and 1b (11.5%) (Table II). An HCV viral load of >100,000 IU/ml was more commonly observed among the group of HCV-seropositive and HCV RNA-positive patients (33/73,

Table I. Modes of transmission according to the anti-HCV status.

Anti-HCV	Modes of transmission (%)				Total (%)
	IVDU	Heterosexual	Homosexual	Vertical	
Positive	87 (73.1)	32 (26.9)	-	-	119 (63.6)
Negative	30 (44.1)	35 (51.5)	2 (2.9)	1 (1.5)	68 (36.4)
Total	117 (62.6)	67 (35.8)	2 (1.1)	1 (0.5)	187 (100)

⁶³ HCV, hepatitis C virus; IVDU, intravenous drug use.

⁷² Table II. Distribution of HCV genotypes/subtypes among samples with detected HCV RNA according to the anti-HCV status.

Anti-HCV	HCV genotypes/subtypes (%)								
	1	1a	1b	1c	2	2a	3a	3k	4a
Positive (n=73)	12 (16.4)	23 (31.5)	7 (9.6)	8 (10.9)	1 (1.4)	1 (1.4)	17 (23.3)	3 (4.1)	1 (1.4)
Negative (n=26)	-	-	3 (11.5)	4 (15.4)	-	-	13 (50)	6 (23.1)	-

HCV, hepatitis C virus.

Table III. Modes of transmission among samples with detected HCV RNA according to the anti-HCV status.

Anti-HCV	Modes of HCV transmission (%)		
	IVDU	Heterosexual	Homosexual
Positive (n=73)	56 (76.7)	17 (23.3)	-
Negative (n=26)	9 (34.6)	16 (61.5)	1 (3.9)

⁶² HCV, hepatitis C virus; IVDU, intravenous drug use.

¹⁰ Table IV. HIV viral load among anti-HCV-positive and HCV RNA-positive patients vs. anti-HCV-negative and HCV RNA-positive patients.

HIV viral load (copies/ml)	Anti-HCV (%)	
	Negative	Positive
≥100,000	14 (53.9)	41 (56.2)
<100,000	7 (26.9)	27 (37.0)
Undetected	5 (19.2)	5 (6.8)
Total	26	73

⁶¹ HIV, human immunodeficiency virus; HCV, hepatitis C virus.

45.2%) compared to the group of HCV-seronegative and HCV RNA-positive patients (1/26, 3.9%).

The majority of HCV-seropositive samples with detected HCV RNA were obtained from HIV-positive patients who had acquired HCV through parenteral transmission (IVDU)

(56/73, 76.7%), followed by those with a history of heterosexual transmission (17/73, 23.3%). However, HCV-seronegative samples with detected HCV RNA were mostly obtained from HIV-positive patients with a history of heterosexual transmission (16/26, 61.5%), followed by those with parenteral (IVDU) (9/26, 34.6%) and homosexual transmission (1/26, 3.9%) (Table III).

⁶⁰ HIV viral load. High HIV viral loads (≥100,000 copies/ml) were mostly detected among the group of HCV-seropositive and HCV RNA-positive patients (41/73, 56.2%) and the group of HCV-seronegative and HCV RNA-positive patients (14/26, 53.9%) (Table IV).

Discussion

⁵⁹ HIV/HCV co-infection affects over one-third of HIV-infected individuals worldwide (14). Following the introduction of HAART, HCV infection has been considered as the principal cause of morbidity and mortality among HIV-infected individuals (15). Co-infected patients exhibit a higher mortality rate compared to singly HIV-infected and HCV infection is considered a predictor of mortality (16). Despite these emerging trends, screening for HCV in HIV-infected patients is not routinely performed, since HCV is perceived as a slowly progressive disease, which would be unlikely to affect the natural history of HIV and associated opportunistic infections (4). In Indonesia, the rapidly increasing number of new HIV infections makes the epidemic one of the fastest growing in Asia (1). However, the incidence rate of HCV infection among HIV-positive patients has not yet been determined in Indonesia. We examined repository blood specimens obtained from HIV-positive patients in Surabaya to identify and describe cases of newly acquired HCV infection.

Antibody testing is the main screening method for HCV infection in HIV-infected individuals (7). Tests available for the diagnosis of HCV infection in the acute phase vary in sensitivity and the third-generation antibody ELISA tests were developed to detect multiple antigenic determinants (core, NS3, NS4 and NS5) to increase the sensitivity. In this study, the result of anti-HCV tests using EIA 3.0 demonstrated that, of the 187 plasma samples obtained from HIV-positive patients, 119 (63.6%) were anti-HCV-positive and the remaining 68 (36.4%) were anti-HCV-negative (Table I). However, serological screening in HIV-infected patients may not be the optimal screening method, possibly as a result of immunosuppression (8).

In our study, the group of HCV-seropositive patients was more likely (73.1%) to have acquired HCV through parenteral transmission (IVDUs), followed by heterosexual transmission (26.9%) (Table I). Individuals infected with HIV are frequently co-infected with HCV due to the shared modes of transmission (9). The risk of HCV transmission is significantly higher for patients who acquire HIV infection parenterally compared to those who acquire it sexually. The role of homosexual and heterosexual transmission of HCV remains controversial; it is considered to occur, although with a low efficiency. The sexual route is a common mode of HIV transmission, although it is not as effective for HCV (17). A previous study reported that $\leq 90\%$ of HIV-positive IVDUs tested positive for HCV antibodies (18), although only 4-8% of the individuals who acquire HIV through sexual contact have detectable HCV antibodies (19,20). In our study, the heterosexual transmission history in HCV-seropositive patients (26.9%) was significantly higher compared to previously reported findings, which may be due to certain mechanisms involved in HCV transmission between sexual partners, particularly those who engage in habits that are associated with a high risk of virus transmission. Transmission may also result from exposure to unreported parenteral risk factors or from sharing certain personal items, such as toothbrushes or razors, which may result in accidental exposure to the partner's blood (21).

In HIV/HCV co-infection, abnormal antibody and cellular immune responses to HCV have been described (22-24). Immunosuppression by HIV infection may impair antibody formation and false-negative HCV antibody tests have been reported in individuals co-infected with HIV (22,25). Loss of HCV antibodies is observed in rare cases with advanced immune deficiency in HIV/HCV co-infection and does not necessarily indicate viral clearance (23). Identifying HCV seroconversion in serial samples is suboptimal, since the antibody development may be delayed in HIV-infected individuals to >1 year after the initial infection (26), compared to HIV-negative patients, who generally produce antibodies to HCV within 6 weeks of infection (27). This delay in the formation of antibodies may result in a significant delay in the diagnosis of HCV in HIV-positive individuals and the patients may miss the opportunity to receive effective treatment for their infection (59 vs. 40% success rate) (28,29). The European AIDS Clinical Society co-infection guidelines endorse an approach that should be considered, with serological testing for HCV for all patients upon entry into HIV care and annually thereafter for HCV-uninfected individuals, with HCV RNA testing for all HCV antibody-negative patients exhibiting

an unexplained increase in alanine transaminase levels and at high risk for HCV infection (IVDU, mucosal trauma during intercourse) (30). Matthews and Dore (31) suggested that consideration should be given to HCV RNA testing, despite a negative HCV antibody status, in cases of unexplained transaminase elevation in patients with CD4 cell counts $<200/\text{mm}^3$ when acute hepatitis C is suspected. Frequent serum HCV RNA testing is thus a possible screening strategy (32). Of note, the HCV antibody status may be negative, despite active HCV viremia, in 10-15% of immunosuppressed patients (31).

Our study demonstrated that HCV RNA was detected in 73 (61.3%) of the 119 anti-HCV-positive samples; of the 68 anti-HCV-negative samples, HCV RNA was detected in 26 (38.2%) (Table II). In the HCV-seropositive and HCV RNA-positive group of patients, a high HCV viral load ($>100,000$ IU/ml) was more frequently detected (33/73, 45.2%) compared to the HCV-seronegative and HCV RNA-positive group of patients (1/26, 3.9%). Higher HCV viral loads may also be responsible for the increased transmissibility, as has been noted in HIV-co-infected patients (17); in this study, they were mostly found among anti-HCV-positive patients. A previous study demonstrated that HCV RNA was detected more frequently among the 31 HCV-seropositive patients (84.4% of tests) compared to the 20 HCV viremic HCV-seronegative patients (51.5% of tests) (9). Another study reported that HCV RNA was detected in the blood of $>80\%$ of HIV-positive individuals who were positive for HCV antibodies (33). Those findings are consistent with those of our study, according to which HCV RNA was highly detected among HIV-positive and HCV-seropositive patients, but was also relatively highly detected among HIV-positive and HCV-seronegative patients. There was a predominantly high HIV viral load ($\geq 100,000$ copies/ml) among the HCV-seropositive and HCV RNA-positive patients (41/73, 56.2%) and also among HCV-seronegative and HCV RNA-positive patients (14/26, 53.9%) (Table IV), which may have contributed to these results. As previously reported, patients with higher HIV RNA titers tend to have lower CD4 cell counts. The impaired T-helper type 1 immune response in turn may alter the response of immune cells to HCV, permitting greater HCV replication (34,35).

IVDU continues to be the primary risk factor for acquisition of HCV/HIV co-infection, with HCV usually acquired within the first year of IVDU (36); however, patients with a sexual transmission history generally acquired HIV prior to HCV (3,37). It was previously suggested that the presence of HIV may increase the heterosexual transmission of HCV (20). HCV infection appears to occur more frequently among HIV-infected HCV-seronegative individuals than appreciated, particularly if HIV acquisition was through sexual as opposed to parenteral risk factors (9). Those findings confirmed the results of our study, demonstrating that the majority of the HCV-seronegative samples with detected HCV RNA were obtained from HIV-positive patients with heterosexual transmission history (61.5%), differed from HCV-seropositive samples with detected HCV RNA, which were mostly obtained from HIV-positive patients with parenteral transmission (IVDU) history (76.7%) (Table III). However, men engaging in sexual intercourse with other men did not appear to exhibit an overall increased risk for co-infection (38-40), although

epidemics of acute HCV were previously described among HIV-infected men engaging in sexual intercourse with other men exhibiting high-risk behaviors (41). The two HIV-infected homosexual males included in this study were anti-HCV-negative, although HCV RNA was detected in one of them (data not shown). HCV/HIV co-infection among homosexual men requires further investigation.

The distribution of HCV genotypes/subtypes in HIV-positive patients reflects the route of transmission. HCV RNA was detected in 73 (61.3%) of the 119 anti-HCV-positive samples and, among these, HCV-1a (31.5%) was the predominant subtype, followed by 3a (23.3%), 1 (16.4%), 1c (10.9%), 1b (9.6%), 3k (4.1%) and 2, 2a and 4a (1.4% each). Of the 68 anti-HCV-negative samples, HCV RNA was detected in 26 (38.2%), with HCV-3a being the most prevalent subtype (50%), followed by 3k (23.1%), 1c (15.4%) and 1b (11.6%) (Table II). HCV subtypes 1a and 3a were more commonly reported among IVDUs (42). The high incidence of HCV-3a in the group with a history of heterosexual transmission may be attributed to the sexual partners, who were IVDUs; however, this hypothesis requires further investigation. Considering the differences in the predominant transmission mode between HCV-seropositive and -seronegative individuals, it is likely that the mode of HCV transmission is associated with the HCV subtypes.

In conclusion, HIV-positive patients are at a high risk of becoming co-infected with HCV, several of whom may remain HCV-seronegative, partly due to their immunodeficiency status. These data may also suggest that there may exist differences in HCV seropositivity and subtypes between HIV-positive patients who acquired HCV sexually and those who acquired HCV parenterally. It seems prudent to consider HCV infection in HIV-positive patients who test negative for HCV antibodies, as delayed HCV seroconversion in HIV-positive patients may result in delayed diagnosis and treatment.

Acknowledgements

The authors are grateful to all the participants in this study. This study was supported by a grant from the Japan Initiative for Global Research Network on Infectious Diseases (J-GRID) and the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan.

References

- World Health Organization (WHO): Review of the health sector response to HIV and AIDS in Indonesia 2007. WHO, Regional Office for South-East Asia, p2, 2007. (http://209.61.208.233/catalogue/2005-2011/pdf/aids/review_of_the_health_sector_response_to_HIV_and_AIDS_in_Indonesia.pdf). Accessed December 13, 2012.
- Weber R, Sabin CA, Friis-Moller N, *et al*: Liver-related deaths in persons infected with the human immunodeficiency virus: the D:A:D study. *Arch Intern Med* 166: 1632-1641, 2006.
- Low E, Vogel M, Rockstroh J and Nelson M: Acute hepatitis C in HIV-positive individuals. *AIDS Rev* 10: 245-253, 2008.
- Sherman KE, Rouster SD, Chung RT and Rajicic N: Hepatitis C Virus prevalence among patients infected with human immunodeficiency virus: a cross-sectional analysis of the US adult AIDS Clinical Trials Group. *Clin Infect Dis* 34: 831-837, 2002.
- Taylor LE, DeLong AK, Maynard MA, *et al*: Acute hepatitis C virus in an HIV clinic: a screening strategy, risk factors, and perception of risk. *AIDS Patient Care STDS* 25: 571-577, 2011.
- Rockstroh JK and Spengler U: HIV and hepatitis C virus co-infection. *Lancet Infect Dis* 4: 437-444, 2004.
- Nelson M, Matthews G, Brook MG and Main J; BHIVA Coinfection Guideline Committee; British HIV Association: BHIVA guidelines on HIV and chronic hepatitis: coinfection with HIV and hepatitis C virus infection (2005). *HIV Med* 6 (Suppl 2): 96-106, 2005.
- Chamie G, Bonacini M, Bangsberg DR, *et al*: Factors associated with seronegative chronic hepatitis C virus infection in HIV infection. *Clin Infect Dis* 44: 577-583, 2007.
- George SL, Gebhardt J, Klinzman D, *et al*: Hepatitis C virus viremia in HIV-infected individuals with negative HCV antibody tests. *J Acquir Immune Defic Syndr* 31: 154-162, 2002.
- Apichartpiyakul C, Chittivudikarn C, Miyajima H, Homma M and Hotta H: Analysis of hepatitis C virus isolates among healthy blood donors and drug addicts in Chiang Mai, Thailand. *J Clin Microbiol* 32: 2276-2279, 1994.
- Doi H, Apichartpiyakul C, Ohba KI, Mizokami M and Hotta H: Hepatitis C virus (HCV) subtype prevalence in Chiang Mai, Thailand, and identification of novel subtypes of HCV major type 6. *J Clin Microbiol* 34: 569-574, 1996.
- Simmonds P, Bukh J, Combet C, *et al*: Consensus proposals for a unified system of nomenclature of hepatitis C virus genotypes. *Hepatology* 42: 962-973, 2005.
- Klefer GE, van Doorn LJ, Brouwer JT, Schalm SW, Heijntink RA and Quint WG: Sequence analysis of the 5' untranslated region in isolates of at least four genotypes of hepatitis C virus in The Netherlands. *J Clin Microbiol* 32: 306-310, 1994.
- Bruno R, Sacchi P, Puoti M, Soriano V and Filice G: HCV chronic hepatitis in patients with HIV: clinical management issues. *Am J Gastroenterol* 97: 1598-1606, 2002.
- Camino N, Sheldon J and Soriano V: Update on hepatitis C treatment in HIV-coinfected patients. *Minerva Gastroenterol Dietol* 50: 67-77, 2004.
- Braitein P, Yip B, Montessori V, Moore D, Montaner JS and Hogg RS: Effect of serostatus for hepatitis C virus on mortality among antiretrovirally naive HIV-positive patients. *CMAJ* 173: 160-164, 2005.
- Poles MA and Dieterich DT: Hepatitis C virus/human immunodeficiency virus coinfection: clinical management issues. *Clin Infect Dis* 31: 154-161, 2000.
- Thio CL, Nolt KR, Astemborski J, Vlahov D, Nelson KE and Thomas DL: Screening for hepatitis C virus in human immunodeficiency virus-infected individuals. *J Clin Microbiol* 38: 575-577, 2000.
- Tor J, Libre JM, Carbonell M, *et al*: Sexual transmission of hepatitis C and its relation with hepatitis B virus and HIV. *BMJ* 301: 1130-1133, 1990.
- Eyster ME, Alter HJ, Aledort LM, Quan S, Hatzakis A and Goedert JJ: Heterosexual co-transmission of hepatitis C virus (HCV) and human immunodeficiency virus (HIV). *Ann Intern Med* 115: 764-768, 1991.
- Mendes-Correa MC, Barone AA and Gianini RJ: Risk factors associated with hepatitis C among patients co-infected with human immunodeficiency virus: a case-control study. *Am J Trop Med Hyg* 72: 762-767, 2005.
- Chamot E, Hirschel B, Wintch J, *et al*: Loss of antibodies against hepatitis C virus in HIV-seropositive intravenous drug users. *AIDS* 4: 1275-1277, 1990.
- Cribier B, Rey D, Schmitt C, Lang JM, Kim A and Stoll-Keller F: High hepatitis C viremia and impaired antibody response in patients coinfecting with HIV. *AIDS* 9: 1131-1136, 1995.
- Valdez H, Anthony D, Farukhi F, *et al*: Immune responses to hepatitis C and non-hepatitis C antigens in hepatitis C virus infected and HIV-1 coinfecting patients. *AIDS* 14: 2239-2246, 2000.
- Sorbi D, Shen D and Lake-Bakaar G: Influence of HIV disease on serum anti-HCV antibody titers: a study of intravenous drug users. *J Acquir Immune Defic Syndr Hum Retrovirol* 13: 295-296, 1996.
- Thomson EC, Nastouli E, Main J, Karayiannis P, Eliahoo J, Muir D and McClure MO: Delayed anti-HCV antibody response in HIV-positive men acutely infected with HCV. *AIDS* 23: 89-93, 2009.
- Netski DM, Mosbrugger T, Depla E, *et al*: Humoral immune response in acute hepatitis C virus infection. *Clin Infect Dis* 41: 667-675, 2005.
- Torriani FJ, Rodriguez-Torres M, Rockstroh JK, *et al*: APRICOT Study Group: Peginterferon Alfa-2a plus ribavirin for chronic hepatitis C virus infection in HIV-infected patients. *N Engl J Med* 351: 438-450, 2004.

29. Gilleece YC, Browne RE, Asboe D, *et al.*: Transmission of hepatitis C virus among HIV-positive homosexual men and response to a 24-week course of pegylated interferon and ribavirin. *J Acquir Immune Defic Syndr* 40: 41-46, 2005.
30. Rockstroh JK, Bhagani S, Benhamou Y, *et al.*; EACS Executive Committee: European AIDS Clinical Society (EACS) guidelines for the clinical management and treatment of chronic hepatitis B and C coinfection in HIV-infected adults. *HIV Med* 9: 82-88, 2008.
31. Matthews GV and Dore GJ: HIV and hepatitis C co-infection. *Asian Biomed* 3: 29-37, 2009.
32. European Association for the Study of the Liver: EASL Clinical Practice Guidelines: management of hepatitis C virus infection. *J Hepatol* 55: 245-264, 2011.
33. Perez-Olmeda M, Rios P, Nunez M, Garcia-Samaniego J, Romero M and Soriano V: Virological characteristics of hepatitis C virus infection in HIV-infected individuals with chronic hepatitis C: implications for treatment. *AIDS* 16: 493-495, 2002.
34. Chung RT, Evans SR, Yang Y, *et al.*: Immune recovery is associated with persistent rise in hepatitis C virus RNA, infrequent liver test flares, and is not impaired by hepatitis C virus in co-infected subjects. *AIDS* 16: 1915-1923, 2002.
35. Chung R: Management of HIV/HCV coinfection. *The PRN Notebook* 9: 14-19, 2004.
36. Garten RJ, Zhang J, Lai S, Liu W, Chen J and Yu XF: Coinfection with HIV and hepatitis C virus among injection drug users in southern China. *Clin Infect Dis* 41: S18-S24, 2005.
37. Filippini P, Coppola N, Scolastico C, Rossi G, Onofrio M, Sagnelli E and Piccinino F: Does HIV infection favor the sexual transmission of hepatitis C? *Sex Transm Dis* 28: 725-729, 2001.
38. Tedaldi EM, Hullsiek KH, Malvestutto CD, *et al.*; Terry Bein Community Programs for Clinical Research on AIDS: Prevalence and characteristics of hepatitis C virus coinfection in a human immunodeficiency virus clinical trials group: the Terry Bein Community Programs for Clinical Research on AIDS. *Clin Infect Dis* 36: 1313-1317, 2003.
39. Bollepalli S, Mathieson K, Bay C, Hillier A, Post J, Van Thiel DH and Nadir A: Prevalence of risk factors for hepatitis C virus in HIV-infected and HIV/hepatitis C virus-coinfected patients. *Sex Transm Dis* 34: 367-370, 2007.
40. Larsen C, Pialoux G, Salmon D, *et al.*: Prevalence of hepatitis C and hepatitis B infection in the HIV-infected population of France, 2004. *Euro Surveill* 13: 18888, 2008.
41. Danta M, Brown D, Bhagani S, *et al.*: Recent epidemic of acute hepatitis C virus in HIV-positive men who have sex with men linked to high-risk sexual behaviours. *AIDS* 21: 983-991, 2007.
42. Pol S, Thiers V, Nousbaum JB, Legendre C, Berthelot P, Kreis H and Brechot C: Changing distribution of HCV genotype in Europe in the last decades. *J Hepatol* 21: S13, 1994.

High rate of seronegative HCV infection in HIV-positive patients

ORIGINALITY REPORT

24%

SIMILARITY INDEX

10%

INTERNET SOURCES

22%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1

Sherman, K. E., S. D. Rouster, R. T. Chung, and N. Rajicic. "Hepatitis C Virus Prevalence among Patients Infected with Human Immunodeficiency Virus: A Cross-Sectional Analysis of the US Adult AIDS Clinical Trials Group", *Clinical Infectious Diseases*, 2002.

Publication

1%

2

Shin-ichi Miyoshi, Jiyou Wang, Keizo Katoh, Mitsutoshi Senoh, Tamaki Mizuno, Yoko Maehara. "An extracellular serine protease produced by *Vibrio vulnificus* NCIMB 2137, a metalloprotease-gene negative strain isolated from a diseased eel", *World Journal of Microbiology and Biotechnology*, 2011

Publication

1%

3

Zylberberg, H., and S. Pol. "Reciprocal Interactions Between Human Immunodeficiency Virus and Hepatitis C Virus Infections", *Clinical Infectious Diseases*, 1996.

Publication

1%

Elizabeth Freda O. Telan, Genesis May J.

4

Samonte, Ilya P. Abellanosa-Tac-An, Evelyn T. Alesna et al. "The early phase of an HIV epidemic in a population exposed previously to HCV in the Philippines", Journal of Medical Virology, 2011

Publication

1%

5

Fabrice Merien, Denis Portnoi, Pascale Bourhy, Françoise Charavay, Alain Berlioz-Arthaud, Guy Baranton. " A rapid and quantitative method for the detection of species in human leptospirosis ", FEMS Microbiology Letters, 2005

Publication

1%

6

Assisted Reproductive Technologies and Infectious Diseases, 2016.

Publication

1%

7

Nevot, M., C. Boesecke, M. Parera, C. Andrés, S. Franco, B. Revollo, P. Ingiliz, C. Tural, B. Clotet, J. K. Rockstroh, and M. A. Martinez. "Hepatitis C virus NS3/4A quasispecies diversity in acute hepatitis C infection in HIV-1 co-infected patients", Journal of Viral Hepatitis, 2014.

Publication

1%

8

M. Arif. "Risk of hepatitis C virus infection among household contacts of Saudi patients with chronic liver disease", Journal of Viral

1%

Hepatitis, 3/1996

Publication

9

John S. Wayne, Lynda Walker, Margaret Patterson, David H. K. Chui. "IDENTIFICATION OF TWO NEW β -THALASSEMIA SPLICE MUTATIONS: IVS-I-1 (G \rightarrow C) AND IVS-I (-2) (A \rightarrow C)", Hemoglobin, 2009

Publication

<1%

10

Steven Shev, Svante Hermodsson, Annika Lindholm, Ewa Malm, Anders Widell, Gunnar Norkrans. "Risk Factor Exposure Among Hepatitis C Virus RNA Positive Swedish Blood Donors - The Role of Parenteral and Sexual Transmission", Scandinavian Journal of Infectious Diseases, 2009

Publication

<1%

11

"Poster", Respiriology, 11/2006

Publication

<1%

12

Bradshaw, Daniel, Gail Matthews, and Mark Danta. "Sexually transmitted hepatitis C infection : the new epidemic in MSM?", Current Opinion in Infectious Diseases, 2012.

Publication

<1%

13

hdl.handle.net

Internet Source

<1%

14

El-Shamy, Ahmed, Michiko Shindo, Ikuo Shoji, Lin Deng, Tadao Okuno, and Hak Hotta.

<1%

"Polymorphisms of the core, NS3 and NS5A proteins of hepatitis C 1 virus genotype 1b associate with development of hepatocellular carcinoma", Hepatology, 2012.

Publication

15

www.highbeam.com

Internet Source

<1%

16

Yamanaka, A.. "A simple method for evaluating dengue vaccine effectiveness in mice based on levels of viremia caused by intraperitoneal injection of infected culture cells", Vaccine, 20090608

Publication

<1%

17

Rezaianzadeh, Abbas, Jafar Hasanzadeh, Abbas Alipour, Mohamed Ali Davarpanah, Abdorreza Rajaeifard, and Seyed Hamid Reza Tabatabaee. "Impact of Hepatitis C on Survival of HIV-Infected Individuals in Shiraz; South of Iran", Hepatitis Monthly, 2012.

Publication

<1%

18

Jin Yang, Mei-xin Fang, Jie Li, Guo-qiang Lou, Hang-jun Lu, Nan-ping Wu. "Detection of hepatitis C virus by an improved loop-mediated isothermal amplification assay", Archives of Virology, 2011

Publication

<1%

19

hepatmon.com

<1%

20

A H Mohsen. "Hepatitis C and HIV-1 coinfection", Gut, 2002

Publication

<1%

21

www.nexusediciones.com

Internet Source

<1%

22

Denise P.C. Chan, Hsin-Yun Sun, Horas T.H. Wong, Shui-Shan Lee, Chien-Ching Hung. "Sexually acquired hepatitis C virus infection: a review", International Journal of Infectious Diseases, 2016

Publication

<1%

23

Tomoaki Tanimoto, Nguyen Hung Cuong, Azumi Ishizaki, Phan Thi Thu Chung et al. "Multiple routes of hepatitis C virus transmission among injection drug users in Hai Phong, Northern Vietnam", Journal of Medical Virology, 2010

Publication

<1%

24

Francesc X. López-Labrador, Sergi Ampurdanés, Xavier Forns, Antoni Castells et al. "Hepatitis C virus (HCV) genotypes in Spanish patients with HCV infection: relationship between HCV genotype 1b, cirrhosis and hepatocellular carcinoma", Journal of Hepatology, 1997

Publication

<1%

25

William A. Bower. "Changes in Hepatitis C Virus (HCV) Viral Load and Interferon-?? Levels in HIV/HCV-Coinfected Patients Treated With Highly Active Antiretroviral Therapy", JAIDS Journal of Acquired Immune Deficiency Syndromes, 07/2006

Publication

<1%

26

Deepa G Gamage, Tim RH Read, Catriona S Bradshaw, Jane S Hocking, Kerry Howley, Marcus Y Chen, Christopher K Fairley. "Incidence of Hepatitis-C among HIV infected men who have sex with men (MSM) attending a sexual health service: a cohort study", BMC Infectious Diseases, 2011

Publication

<1%

27

"13th European Congress of Clinical Microbiology and Infectious Diseases", Clinical Microbiology and Infection, 2003

Publication

<1%

28

Everton Hadlich. "Hepatitis C virus (HCV) viremia in HIV-infected patients without HCV antibodies detectable by third-generation enzyme immunoassay", Journal of Gastroenterology and Hepatology, 9/2007

Publication

<1%

29

Haines, B.P.. "Expression of the Lingo/LERN gene family during mouse embryogenesis",

<1%

Gene Expression Patterns, 200801

Publication

30

M. Furione, R. Maserati, M. Gatti, F. Baldanti, A. Cividini, R. Bruno, G. Gerna, M. U. Mondelli. "Dissociation of Serum and Liver Hepatitis C Virus RNA Levels in Patients Coinfected with Human Immunodeficiency Virus and Treated with Antiretroviral Drugs", Journal of Clinical Microbiology, 2004

Publication

31

Brouwer, J.T.. "Early prediction of response in interferon monotherapy and in interferon-ribavirin combination therapy for chronic hepatitis C: HCV RNA at 4 weeks versus ALT", Journal of Hepatology, 199902

Publication

32

S. Ramia, S. Koussa, A. Taher, S. Haraki, S. Klayme, D. Sarkis, R. Naman. "Hepatitis-C-virus genotypes and hepatitis-G-virus infection in Lebanese thalassaemics", Annals of Tropical Medicine & Parasitology, 2013

Publication

33

aem.asm.org

Internet Source

34

Danta, Mark, and Alison J. Rodger. "Transmission of HCV in HIV-positive populations :", Current Opinion in HIV and AIDS,

<1%

<1%

<1%

<1%

<1%

2011.
Publication

35 Deepak Joshi. "Role of liver transplantation in human immunodeficiency virus positive patients", World Journal of Gastroenterology, 2015
Publication

36 vir.sgmjournals.org
Internet Source

37 citeseerx.ist.psu.edu
Internet Source

38 onlinelibrary.wiley.com
Internet Source

39 Norma de Paula Cavalheiro, Abel de La Rosa, Slava Elagin, Fatima Mitiko Tengan, Antonio Alci Barone. "Hepatitis C virus: molecular and epidemiological evidence of male-to-female transmission", Brazilian Journal of Infectious Diseases, 2010
Publication

40 Thomas Berg, Uwe Hopf, Klaus Stark, Renate Baumgarten, Hartmut Lobeck, Eckart Schreier. "Distribution of hepatitis C virus genotypes in German patients with chronic hepatitis C: correlation with clinical and virological parameters", Journal of Hepatology, 1997
Publication

41 Isao Nakano, Yoshihide Fukuda, Yoshiaki Katano, Satoshi Nakano, Takashi Kumada, Tetsuo Hayakawa. "Why is the interferon sensitivity-determining region (ISDR) system useful in Japan?", *Journal of Hepatology*, 1999
Publication <1%

42 "Parallel Session : Abstracts", *Hepatology Research*, 07/14/2010
Publication <1%

43 Marina Nunez, Angel Mariño, Celia Miralles, Miguel A Berdún et al. "Baseline Serum Hepatitis C Virus (HCV) RNA Level and Response at Week 4 Are the Best Predictors of Relapse After Treatment With Pegylated Interferon Plus Ribavirin in HIV/HCV-Coinfected Patients", *JAIDS Journal of Acquired Immune Deficiency Syndromes*, 2007
Publication <1%

44 www.omicsonline.org
Internet Source <1%

45 doaj.org
Internet Source <1%

46 uregina.ca
Internet Source <1%

47 Meghan D. Morris, Jennifer Evans, Martha Montgomery, Michelle Yu, Alya Briceno, <1%

Kimberly Page, Judith A. Hahn. "Intimate Injection Partnerships Are at Elevated Risk of High-Risk Injecting: A Multi-Level Longitudinal Study of HCV-Serodiscordant Injection Partnerships in San Francisco, CA", PLoS ONE, 2014

Publication

48

Sulkowski, M.S.. "Viral hepatitis and HIV coinfection", Journal of Hepatology, 200802

Publication

49

doctiktak.com

Internet Source

50

jvi.asm.org

Internet Source

51

Ashwani K Singal, Bhupinderjit S Anand. "Management of hepatitis C virus infection in HIV/HCV co-infected patients: Clinical review", World Journal of Gastroenterology, 2009

Publication

52

Shepard, C.W.. "Global epidemiology of hepatitis C virus infection", The Lancet Infectious Diseases, 200509

Publication

53

www.hiv.uw.edu

Internet Source

54

Catherine Delamare, Bruno Carbonne,

Nazbanou Heim, Nadia Berkane, Jean Claude Petit, Serge Uzan, Jean-Didier Grangé.
"Detection of hepatitis C virus RNA (HCV RNA) in amniotic fluid: a prospective study", Journal of Hepatology, 1999

Publication

<1%

55

Jain, M. K., B. Adams-Huet, D. Terekhova, L. E. Kushner, R. Bedimo, X. Li, and M. Holodniy.
"Acute and chronic immune biomarker changes during interferon/ribavirin treatment in HIV/HCV co-infected patients", Journal of Viral Hepatitis, 2014.

Publication

<1%

56

www.e-cmh.org
Internet Source

<1%

57

www.hivguidelines.org
Internet Source

<1%

58

urg.prn.org
Internet Source

<1%

59

E. Herrero Martínez. "Hepatitis B and hepatitis C co-infection in patients with HIV", Reviews in Medical Virology, 07/2001

Publication

<1%

60

www.aidsetc.com
Internet Source

<1%

61

medcraveonline.com

<1%

62

R. S. Ross. "Changes in the epidemiology of hepatitis C infection in Germany: Shift in the predominance of hepatitis C subtypes", Journal of Medical Virology, 02/2000

Publication

<1%

63

sti.bmj.com

Internet Source

<1%

64

E. Sikuler, S. Elsana, A. Yaari, Y. Shemer-Avni, L. Naggan, M. Margalith. "Salivary anti-HCV as a marker of serum anti-HCV. A followup cohort study", Journal of Hepatology, 2001

Publication

<1%

65

Giannini, C.. "Comparative analysis of two assays for genotyping hepatitis C virus based on genotype-specific primers or probes", Journal of Hepatology, 199509

Publication

<1%

66

Katayama, Yuko, Nina G. Barzaga, Arlene Alipio, Soetjipto, Hisaya Doi, Satoshi Ishido, and Hak Hotta. "Genotype Analysis of Hepatitis C Virus among Blood Donors and Inmates in Metro Manila, The Philippines", Microbiology and Immunology, 1996.

Publication

<1%

67

cid.oxfordjournals.org

Internet Source

<1%

68

Rachel Trowbridge, Theo P. Sloots, Philip Buda, Joan Faoagali, Catherine Hyland, Ian Young, Eric J. Gowans. "An ELISA for the detection of antibody to the core antigen of hepatitis C virus: use in diagnosis and analysis of indeterminate samples", *Journal of Hepatology*, 1996

Publication

<1%

69

Tian, Di, Lin Li, Yongjian Liu, Hanping Li, Xiaoyuan Xu, and Jingyun Li. "Different HCV Genotype Distributions of HIV-Infected Individuals in Henan and Guangxi, China", *PLoS ONE*, 2012.

Publication

<1%

70

Robério Amorim de Almeida Pondé. "Hidden hazards of HCV transmission", *Medical Microbiology and Immunology*, 2010

Publication

<1%

71

J. T. Blackard. "HCV/ HIV co-infection: time to re-evaluate the role of HIV in the liver?", *Journal of Viral Hepatitis*, 5/2008

Publication

<1%

72

Yuan-Ming Lee. "Molecular epidemiology of HCV genotypes among injection drug users in Taiwan: Full-length sequences of two new subtype 6w strains and a recombinant

<1%

form_2b6w", Journal of Medical Virology,
01/2010

Publication

73

L. Barrett. "Stronger hepatitis C virus-specific CD8+ T-cell responses in HIV coinfection : Anti-HCV T-cell responses in HIV coinfection", Journal of Viral Hepatitis, 04/2010

<1%

Publication

74

"Viral Infections of Humans", Springer Science and Business Media LLC, 2014

<1%

Publication

75

Frank J. Palella. "Declining Morbidity and Mortality among Patients with Advanced Human Immunodeficiency Virus Infection", New England Journal of Medicine, 03/26/1998

<1%

Publication

76

"20th International Congress of Nutrition: Granada, Spain, September 15 20, 2013", Annals of Nutrition and Metabolism, 2013

<1%

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

High rate of seronegative HCV infection in HIV-positive patients

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
