

# Inhibition of hepatitis C virus replication by chalepin and pseudane IX isolated from *Ruta angustifolia* leaves

*by* Maria Lusida

---

**Submission date:** 23-Jun-2020 09:37PM (UTC+0800)

**Submission ID:** 1348571437

**File name:** lepin\_and\_pseudane\_IX\_isolated\_from\_Ruta\_angustifolia\_leaves.pdf (1.2M)

**Word count:** 6877

**Character count:** 37333

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/267573380>

# 37 Inhibition of hepatitis C virus replication by chalepin and pseudane IX isolated from *Ruta angustifolia* leaves

Article in *Fitoterapia* · December 2014

DOI:10.1016/j.fitote.2014.10.011

CITATIONS

26

READS

168

10 authors, including:



**Tutik Sri Wahyuni**  
Airlangga University

28 PUBLICATIONS 262 CITATIONS

[SEE PROFILE](#)



**Aty Widyawaruyanti**  
Airlangga University

108 PUBLICATIONS 812 CITATIONS

[SEE PROFILE](#)



**Maria Lusida**  
Airlangga University

106 PUBLICATIONS 923 CITATIONS

[SEE PROFILE](#)



**Achmad Fuad**  
Airlangga University

15 PUBLICATIONS 105 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Representative of Airlangga University for SATREPS Project 2015-2020 [View project](#)



10 Association between quasispecies variants of hepatitis B virus, as detected by high-throughput sequencing, and progression of advanced liver disease in Indonesian patients [View project](#)



37

## Inhibition of hepatitis C virus replication by chalepin and pseudane IX isolated from *Ruta angustifolia* leaves



Tutik Sri Wahyuni<sup>a,b</sup>, Aty Widyawaruyanti<sup>b</sup>, Maria Inge Lusida<sup>c</sup>, Achmad Fuad<sup>b</sup>, Soetjipto<sup>c</sup>, Hiroyuki Fuchino<sup>d</sup>, Nobuo Kawahara<sup>d</sup>, Yoshitake Hayashi<sup>e</sup>, Chie Aoki<sup>a</sup>, Hak Hotta<sup>a,\*</sup>

<sup>a</sup> Division of Microbiology, Kobe University Graduate School of Medicine, 7-5-1 Kusunoki-cho, Chuo-ku, Kobe 650-0017, Japan

<sup>b</sup> Department of Pharmacognocny and Phytochemistry, Faculty of Pharmacy, Airlangga University, Jl. Dharmawangsa Dalam, Surabaya 60286, Indonesia

<sup>c</sup> Institute of Tropical Disease, Airlangga University, Jl. Mulyorejo, Surabaya 60115, Indonesia

<sup>d</sup> Research Center for Medicinal Plant Resources, National Institute of Biomedical Innovation, 1-2 Hachimandai, Tsukuba, Ibaraki 305-0843, Japan

<sup>e</sup> Division of Infectious Disease Pathology, Kobe University Graduate School of Medicine, 7-5-1 Kusunoki-cho, Chuo-ku, Kobe 650-0017, Japan

35

## ARTICLE INFO

## Article history:

Received 7 August 2014

Accepted in revised form 11 October 2014

Accepted 14 October 2014

Available online 22 October 2014

26

## Chemical compounds studied in this article:

Chalepin (PubChem CID 119066)

Scopoletin (PubChem CID 5280460)

Arborinine (PubChem CID 5281832)

 $\gamma$ -Fagarine (PubChem CID 107936)

Kokusaginine (PubChem CID 10227)

Pseudane IX (PubChem data not available)

## Keywords:

Hepatitis C virus

*Ruta angustifolia*

Rutaceae

Post-entry inhibition

Alkaloid

Coumarin

53

## ABSTRACT

Hepatitis C virus (HCV) infection is highly prevalent among global populations, with an estimated number of infected patients being 170 million. Approximately 70–80% of patients acutely infected with HCV will progress to chronic liver disease, such as liver cirrhosis and hepatocellular carcinoma, which is a substantial cause of morbidity and mortality worldwide. New therapies for HCV infection have been developed, however, the therapeutic efficacies still need to be improved. Medicinal plants are promising sources for antivirals against HCV. A variety of plants have been tested and proven to be beneficial as antiviral drug candidates against HCV. In this study, we examined extracts, their subfractions and isolated compounds of *Ruta angustifolia* leaves for antiviral activities against HCV in cell culture. We isolated six compounds, chalepin, scopoletin,  $\gamma$ -fagarine, arborinine, kokusaginine and pseudane IX. Among them, chalepin and pseudane IX showed strong anti-HCV activities with 50% inhibitory concentration (IC<sub>50</sub>) of  $1.7 \pm 0.5$  and  $1.4 \pm 0.2$   $\mu\text{g/ml}$ , respectively, without apparent cytotoxicity. Their anti-HCV activities were stronger than that of ribavirin ( $2.8 \pm 0.4$   $\mu\text{g/ml}$ ), which has been widely used for the treatment of HCV infection. Mode-of-action analyses revealed that chalepin and pseudane IX inhibited HCV at the post-entry step and decreased the levels of HCV RNA replication and viral protein synthesis. We also observed that arborinine, kokusaginine and  $\gamma$ -fagarine possessed moderate levels of anti-HCV activities with IC<sub>50</sub> values being  $6.4 \pm 0.7$ ,  $6.4 \pm 1.6$  and  $20.4 \pm 0.4$   $\mu\text{g/ml}$ , respectively, whereas scopoletin did not exert significant anti-HCV activities at 30  $\mu\text{g/ml}$ .

© 2014 Elsevier B.V. All rights reserved.

## 1. Background

Hepatitis C virus (HCV) is an enveloped virus that belongs to the *Hepacivirus* genus within the *Flaviviridae* family. The viral genome is a single-stranded, positive-sense RNA of 9.6 kb with highly structured 5'- and 3'-untranslated regions [1,2]. It encodes a polyprotein precursor consisting of about 3000 amino acid residues, which is cleaved by the host and viral

proteases to generate 10 mature proteins, such as core, E1, E2, a putative ion channel p7, and nonstructural proteins NS2, NS3, NS4A, NS4B, NS5A and NS5B. The core, E1 and E2 are cleaved off by the signal peptidase and signal peptide peptidase of the host cell and, together with the viral genome, form the virus particles. The E1 and E2 glycoproteins are responsible for binding to a number of different virus receptor molecules on the cell surface, such as scavenger receptor class B type I, CD81, claudin 1 and occludin. On the other hand, nonstructural proteins play crucial roles in virus replication. NS2 possesses a metalloprotease activity that mediates cleavage between NS2

22

\* Corresponding author. Tel.: +81 78 382 5500; fax: +81 78 382 5519.  
E-mail address: [hotta@kobe-u.ac.jp](mailto:hotta@kobe-u.ac.jp) (H. Hotta).

63

<http://dx.doi.org/10.1016/j.fitote.2014.10.011>

0367-326X/© 2014 Elsevier B.V. All rights reserved.

and NS3. After this cleavage takes place, NS3 exerts a serine protease activity that is responsible for the cleavage at the remaining cleavage sites of the polyprotein. NS3 also possesses a helicase activity in its C-terminal domain, which is required for viral RNA replication. NS4A stabilizes NS3 by forming a complex with it and also acts as an inducer of membrane alterations. NS4B is a hydrophobic protein and is involved in the membranous web formation, a characteristic feature of HCV-infected cells. NS5A is a phosphoprotein with versatile functions and is required for viral RNA replication and particle assembly [1,3]. NS5B possesses an RNA-dependent RNA polymerase activity. It is known that the HCV replication cycle is linked to the lipid metabolism of the host cells. It should also be noted that the HCV genome exhibits a considerable degree of sequence heterogeneity, based on which HCV is currently classified into 7 genotypes (1 to 7) and more than 70 subtypes (1a, 1b, 2a, 2b, etc.) [4].

HCV infection is highly prevalent among global populations, especially in Africa and Asia, with an estimated number of infected patients being 170 million worldwide. Approximately 3 million people are newly infected with HCV worldwide every year [3,5]. Seventy to 80% of newly infected patients progress to chronic infection. Patients with chronic HCV infection have a high risk to develop severe liver diseases such as cirrhosis and hepatocellular carcinoma, and also to develop extra hepatic manifestations, including glucose and lipid metabolic disorders [6,7]. A standard care of HCV infection using pegylated interferon (Peg-IFN)- $\alpha$  and ribavirin can achieve sustained virological response (SVR) in ca. 50% of patients infected with HCV genotype 1 or 4 [5]. Triple combination therapy using Peg-IFN- $\alpha$ , ribavirin and an NS3 protease inhibitor increased the SVR rate to 70 to 80%. Moreover, recent approval of other direct-acting antivirals (DAA), including NS5A inhibitors, can further improve the SVR rate. However, they are not equally effective for all of the seven HCV genotypes and, more importantly, serious adverse effects are observed with some patients [5,8]. This highlights the need for a new alternative and/or complementary agent(s) for treatment of HCV.

A wide variety of medicinal plants and their phytochemical constituents have been reported to inhibit HCV infection. For example, an extract of *Phyllanthus amarus* root significantly inhibited HCV NS3 protease with a 50% inhibitory concentration (IC<sub>50</sub>) of 5  $\mu$ g/ml whereas *P. amarus* leaves inhibited HCV NS5B polymerase with the same IC<sub>50</sub> value [9]. We tested ethanol extracts of Indonesia plants for their anti-HCV activities and reported that *Toona sureni* leaves, *Melicope latifolia* leaves, *Melanolepis multiglandulosa* stem and *Ficus fistulosa* leaves possessed anti-HCV activities [10]. We also reported that extracts of *Glycyrrhiza uralensis* root and isolated compounds, such as glycycomarin, glycerin, glycyrol, and liquiritigenin, and extracts of *Morinda citrifolia* leaves, an isolated compound, pheophorbide a, and its related compound, pyropheophorbide a, exhibited anti-HCV activities [11,12]. Likewise, silymarin, iridoid, epigallocatechin-3-gallate were reported to inhibit HCV infection at the entry step while diosgenin, luteolin, quercetin, 3-hydroxy caruilignan C, excoecariphenol D and apigenin at the post-entry step [13,14]. Although a number of novel antivirals against HCV are being developed, further studies are still needed to identify a safer, more effective and cheaper anti-HCV substance(s). Medicinal plants are a good target for the study.

*Ruta angustifolia* belongs to the Rutaceae family. Plants in the *Ruta* genus have been used as traditional remedy, such as antiseptics, antihelminthics and anti-inflammatory, wound-healing and pain-relief drugs, to cure malconditions during pregnancy and disorders in the gastrointestinal, respiratory, nervous, skin and musculoskeletal systems [15]. In Indonesia, *R. angustifolia* has been known as traditional medicine for liver disease and jaundice. It contains coumarin, alkaloid and flavonoid compounds. Angustifolin and four aromatic derivatives, moskachan A, B, C and D, have been identified as constituents of *R. angustifolia* [16,17]. In this study, we examined the anti-HCV activities of extracts from *R. angustifolia* and its constituents.

## 2. Materials and methods

### 2.1. Cells and viruses

Huh7.5 cells and the plasmid pFL-J6/JFH1 to produce the J6/JFH1 strain of HCV genotype 2a [18] were kindly provided by Dr. C. M. Rice, The Rockefeller University, New York, NY, USA. Huh7.5 cells were cultivated in Dulbecco's modified Eagle's medium (Wako, Osaka, Japan) supplemented with fetal bovine serum (Biowest, Nuaille, France), non-essential amino acids (Invitrogen, Carlsbad, CA), penicillin (100 IU/ml) and streptomycin (100  $\mu$ g/ml) (Invitrogen). Cells were grown at 37 °C in a 5% CO<sub>2</sub> incubator.

### 2.2. Collection, extraction, fractionation and compound isolation of *R. angustifolia* leaves

*R. angustifolia* leaves were collected at Lembang, a mountain area of the West Java region, Indonesia. The collected samples were verified by botanical researchers at the Purwadadi Botanical Garden, Purwadadi, Indonesia. Leaves of the plants were dried at room temperature, pulverized and extracted by means of two different extraction procedures; (i) 96% ethanol and (ii) *n*-hexane, dichloromethane and methanol, successively. Maceration process was repeated over 3 days. The obtained filtrates were concentrated under reduced pressure to yield ethanol, *n*-hexane, dichloromethane and methanol extracts. The dichloromethane extract was subjected to the open column chromatography with silica gel (development solvent: gradient of chloroform–methanol system). A bioactivity-positive fraction(s) was further fractionated under open column chromatography with silica gel and mobile phase gradient of hexane–ethyl acetate system. Based on thin layer chromatography (TLC) profiles, several fractions were combined and passed through an activated charcoal column and eluted by each 2 l of methanol (100%), 30% of chloroform–methanol, and chloroform (100%) [19]. Each fraction was concentrated in vacuo and further subfractionated by recycling high-performance liquid chromatography (HPLC) (solvent system: 100% methanol, column: GS-320 + GS-310, 21.5 mm ID  $\times$  1000 mm, flow rate: 5.0 ml/min, detection: UV 210 nm) and preparative HPLC (column: Waters XBridge C18 10  $\times$  250 mm, solvent system: gradient solvent of 0.1% trifluoroacetic acid (TFA) – acetonitrile, flow rate: 2.5 ml/min, column temperature: 30 °C). Preparative HPLC was run on JASCO LC-2000 plus series. Recycling preparative HPLC was performed on a Japan Analytical Industry LC-908W.



To determine the structure of the isolated compounds, liquid chromatography–mass spectrometry (LC–MS),  $^1\text{H}$ -nuclear magnetic resonance (NMR) and  $^{13}\text{C}$  NMR analyses were performed.  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra were measured with BRUKER Ascend 600 spectrometer (600 MHz). LC–MS was performed with a Thermofischer Scientific Orbitrap Elite equipped with an electrospray ionization (ESI) source [20].

### 2.3. Analysis of anti-HCV activities

*R. angustifolia* extracts and isolated compounds were dissolved in dimethyl sulfoxide (DMSO) to obtain stock solutions at a concentration of 100 mg/ml. The stock solutions were stored at  $-20^\circ\text{C}$  until used. Ribavirin was purchased (Sigma-Aldrich, Steinheim, Germany) and used as a positive control.

Huh7.5 cells were seeded in 24-well plates ( $1.9 \times 10^5$  cells/well). A fixed amount of the J6/JFH1-P47 strain of HCV genotype 2a [10,21], with a multiplication of infection (MOI) of 0.5 focus-forming units (ffu)/cell, was mixed with serial dilutions of the extracts (100, 30, 10, 1 and 0.1  $\mu\text{g}/\text{ml}$ ) and compounds (30, 10, 3, 1 and 0.1  $\mu\text{g}/\text{ml}$ ), and inoculated to the cells. After 2 h, the cells were washed with medium to remove the residual virus and further incubated in the medium containing the same concentrations of the test samples as those during virus inoculation.

Time-of-addition experiments were performed to assess the mode of action of the samples, as described previously [10–12]. In brief, three sets of experiments were done in parallel. (i) To assess the antiviral effect at the entry step, the mixture of HCV and a sample was inoculated to the cells. After virus adsorption for 2 h, the residual virus and the sample were removed, and cells were refed with fresh medium without the sample for 46 h. (ii) To assess the antiviral effect at the post-entry step, HCV was inoculated to the cells in the absence of the sample. After virus adsorption for 2 h, the residual virus was removed and cells were refed with fresh medium containing the sample for 46 h. (iii) As a positive control, HCV mixed with the sample was inoculated to the cells. After virus adsorption for 2 h, the residual virus and the sample were removed, and cells were refed with fresh medium containing the sample for 46 h. Culture supernatants were obtained at 1 and 2 days post-infection (dpi) and titrated for virus infectivity [21]. Virus and cells treated with medium containing 0.1% DMSO served as a control. The percent inhibition of virus infectivity by the samples was calculated by comparing to the control using SPSS probit analysis, and  $\text{IC}_{50}$  values were determined.

### 2.4. Real-time quantitative RT-PCR

Total RNA was extracted from the cells using a RNeasy RNA cell miniprep system (Promega, Madison, WI) according to the manufacturer's instructions. One  $\mu\text{g}$  of total RNA was reverse transcribed using a GoScript Reverse Transcription system (Promega) with random primers and subjected to quantitative real-time PCR analysis using SYBR Premix Ex Taq (TaKaRa, Kyoto, Japan) in a MicroAmp 96-well reaction plate and an ABI PRISM 7500 system (Applied Biosystems, Foster City, CA). The primers used to amplify an NS5A region of the HCV genome were 5'-AGACGTATTGAGGTCCATGC-3' (sense) and 5'-CCGACGCGGTGCTGATAG-3' (antisense). As an internal

control, human glyceraldehyde-3-phosphate dehydrogenase (GAPDH) gene expression levels were measured using primers 5'-GCCATCAATGACCCCTTCATT-3' (sense) and 5'-TCTCGCTCTGGAAGATGG-3' (antisense).

### 2.5. Immunoblotting

Cells were lysed and separated with SDS-polyacrylamide gel electrophoresis as described previously [10,11]. The samples were transferred onto a polyvinylidene difluoride membrane (Millipore, Bedford, MA), which was then incubated with the respective primary antibodies. The primary antibodies used were mouse monoclonal antibodies against HCV NS3 and GAPDH (Millipore). Horseradish peroxidase-conjugated goat anti-mouse immunoglobulin (Invitrogen) was used to visualize the respective proteins by means of an enhanced chemiluminescence detection system (ECL; GE Healthcare, Buckinghamshire, UK). The relative band intensity was quantified using densitometry analysis with ImageJ software. The NS3 protein expression levels were normalized to their respective GAPDH protein levels.

### 2.6. WST-1 assay for cytotoxicity

WST-1 assay was performed as described previously [10]. In brief, Huh7.5 cells in 96-well plates were treated with serial dilutions of the test samples or 0.1% DMSO as a control for 48 h. After the treatment, 10  $\mu\text{l}$  of WST-1 reagent (Roche, Mannheim, Germany) was added to each well and cells were cultured for 4 h. The WST-1 reagent is absorbed by the cells and converted to formazan by mitochondrial dehydrogenases. The amount of formazan, which correlates with the number of living cells, was determined by measuring the absorbance of each well using a microplate reader at 450 nm and 630 nm. Percent cell viability compared to the control was calculated for each dilution of substances and 50% cytotoxic concentration ( $\text{CC}_{50}$ ) values were determined by SPSS probit analysis.

## 3. Results

### 3.1. Bioactivity-guided fractionation and purification of extracts from *R. angustifolia* leaves and isolation of compounds

Dried and pulverized *R. angustifolia* leaves were extracted with ethanol, *n*-hexane, dichloromethane and methanol as described in the Materials and methods section, and examined for antiviral activities against the J6/JFH1-P47 strain of HCV genotype 2a [10,21]. The results revealed that the dichloromethane extract of *R. angustifolia* leaves had potent anti-HCV activity with  $\text{IC}_{50}$  of  $1.6 \pm 0.3 \mu\text{g}/\text{ml}$  (Table 1). The dichloromethane extract was further purified by open column chromatography to

**Table 1**  
Anti-HCV activity ( $\text{IC}_{50}$ ), cytotoxicity ( $\text{CC}_{50}$ ) and selectivity index (SI) of *R. angustifolia* leaves extracts.

Sample	$\text{IC}_{50}$ ( $\mu\text{g}/\text{ml}$ )	$\text{CC}_{50}$ ( $\mu\text{g}/\text{ml}$ )	SI
Ethanol extract	$3.0 \pm 1.4^a$	>100	>30.3
<i>n</i> -Hexane extract	$15.6 \pm 5.2$	>100	>6.4
Dichloromethane extract	$1.6 \pm 0.3$	$49.2 \pm 3.6$	30.8
Methanol extract	$8.1 \pm 2.0$	>100	>12.3

<sup>a</sup> Mean  $\pm$  SEM of data from two independent experiments.

obtain 6 fractions, of which fraction 4 showed potent anti-HCV activity with  $IC_{50}$  of 0.7  $\mu\text{g/ml}$  (Table 2). Fraction 4 was further fractionated under open column chromatography with silica gel and mobile phase gradient of hexane–ethyl acetate system to obtain 29 fractions. Based on the TLC profile, some of the subfractions were combined and 4 groups of subfractions were passed through an activated charcoal column, which were eluted by each 2 l of 100% methanol, 30% chloroform–methanol and 100% chloroform to give 3 fractions (a, 100% methanol; b, 30% chloroform–methanol; c, 100% chloroform) [19]. Each fraction was concentrated in vacuo. Fraction 1a was subjected to recycling HPLC to afford F1a-1 (3.1 mg) and F1a-2 (96.7 mg), which were determined as an identical compound (compound 1). Fraction 2b was separated using preparative HPLC to obtain F2b-1 (compound 2), F2b-2 and F2b-3 (compound 3). Fraction 3c was separated by filtration method with methanol to yield F3c-1 (compound 4) and F3c-2, the latter of which was subjected to preparative HPLC to obtain 3c-2A (compound 5) and 3c-2B (compound 4). Fraction 4a was subjected to preparative HPLC to obtain F4a-3 (compound 6). The structures of the isolated compounds were determined by LC–MS and NMR analyses (Supplementary Information).

Compound 1 (100.7 mg) was isolated as colorless amorphous powder with a molecular formula of  $C_{19}H_{22}O_4$  by orbitrap MS,  $m/z$  315.15719  $[M + H]^+$  and was identified as chalepin by comparison with NMR literature data [22]. Compound 2 (5.0 mg) was a yellow amorphous powder with a molecular formula of  $C_{10}H_9O_4$ ,  $m/z$  193.04833  $[M + H]^+$  and was identified as scopoletin [23]. Compound 3 (3.0 mg) was identified as  $\gamma$ -fagarine [24], an alkaloid compound, which has a molecular formula of  $C_{13}H_{12}O_3N$ ,  $m/z$  230.07985  $[M + H]^+$ . Compound 4 (28.7 mg) was identified as arborinine [23], another alkaloid compound with a molecular formula of  $C_{16}H_{16}O_4N$ ,  $m/z$  286.10952  $[M + H]^+$ . Compound 5 (6.5 mg) with a molecular formula of  $C_{14}H_{14}O_4N$  was identified as kokusaginine [23,25], an alkaloid compound. Compound 6 (3.7 mg) with a molecular formula of  $C_{18}H_{26}ON$ ,  $m/z$  272.20089 was identified as pseudane IX [26]. The structures of the compounds are shown in Fig. 1.

### 3.2. Anti-HCV activities of the isolated compounds


Anti-HCV activities of the isolated compounds were tested. Ribavirin, which has been widely used for the treatment of HCV infection, was used as a positive control. The results obtained revealed that chalepin (compound 1) and pseudane IX (compound 6) possessed strong anti-HCV activities, with  $IC_{50}$  being  $1.7 \pm 0.5$  and  $1.4 \pm 0.2$   $\mu\text{g/ml}$ , respectively (Table 3).

**Table 2**  
Anti-HCV activity ( $IC_{50}$ ), cytotoxicity ( $CC_{50}$ ) and selectivity index (SI) of fractions from a dichloromethane extract of *R. angustifolia* leaves.

Sample	$IC_{50}$ ( $\mu\text{g/ml}$ )	$CC_{50}$ ( $\mu\text{g/ml}$ )	SI
Fraction 1	>100 <sup>a</sup>	$\geq 100$	n.a. <sup>b</sup>
Fraction 2	$\geq 100$	>100	n.a. <sup>b</sup>
Fraction 3	>100	>100	n.a. <sup>b</sup>
Fraction 4	0.7	42.1	64.8
Fraction 5	>100	>100	n.a. <sup>b</sup>
Fraction 6	7.4	>100	>13.5

<sup>a</sup> Not detectable at the concentration of 100  $\mu\text{g/ml}$ .

<sup>b</sup> Not applicable.



**Fig. 1.** Molecular structures of the compounds isolated from *R. angustifolia* leaves. Chalepin (compound 1); scopoletin (compound 2);  $\gamma$ -fagarine (compound 3); arborinine (compound 4); kokusaginine (compound 5); pseudane IX (compound 6).

Their anti-HCV activities were stronger than that of ribavirin ( $2.8 \pm 0.4$   $\mu\text{g/ml}$ ). Also,  $\gamma$ -fagarine (compound 3), arborinine (compound 4) and kokusaginine (compound 5) showed weaker but significant anti-HCV activities, with  $IC_{50}$  being  $20.4 \pm 0.4$ ,  $6.4 \pm 0.7$  and  $6.4 \pm 1.6$   $\mu\text{g/ml}$ , respectively. On the other hand, scopoletin (compound 2) did not show any significant inhibitory effect at the concentration of 30  $\mu\text{g/ml}$ . Dose-dependent profiles of anti-HCV activities and cytotoxicity of those compounds are shown in Fig. 2.

### 3.3. Mode-of-action of anti-HCV activities of chalepin and pseudane IX

Time-of-addition experiments were performed to determine whether the compounds inhibit HCV at the entry or post-entry step [10]. Percent HCV inhibitions by chalepin and pseudane IX at the concentrations of 30, 10, 3, 1 and 0.1  $\mu\text{g/ml}$  were measured in the experiments where the treatment was done either during (at the entry step), after virus inoculation (at the post-entry step) or both (Fig. 3). The  $IC_{50}$  values of chalepin for treatment at the entry step, post-entry step and both were


**Table 3**  
Anti-HCV activity ( $IC_{50}$ ), cytotoxicity ( $CC_{50}$ ) and selectivity index (SI) of the isolated compounds.

Isolate code	Compound	$IC_{50}$ ( $\mu\text{g/ml}$ )	$CC_{50}$ ( $\mu\text{g/ml}$ )	SI
Compound 1	Chalepin	$1.7 \pm 0.5^a$	$14.0 \pm 2.4$	8.2
Compound 2	Scopoletin	>30 <sup>b</sup>	>30 <sup>b</sup>	n.a. <sup>c</sup>
Compound 3	$\gamma$ -Fagarine	$20.4 \pm 0.4$	>30 <sup>b</sup>	>1.5
Compound 4	Arborinine	$6.4 \pm 0.7$	$16.3 \pm 6.2$	2.5
Compound 5	Kokusaginine	$6.4 \pm 1.6$	>30 <sup>b</sup>	>4.7
Compound 6	Pseudane IX	$1.4 \pm 0.2$	$26 \pm 0.9$	18.6
Positive control	Ribavirin	$2.8 \pm 0.4$	>30 <sup>b</sup>	>10.7


<sup>a</sup> Mean  $\pm$  SEM of data from two independent experiments.

<sup>b</sup> Not detectable at the concentration of 30  $\mu\text{g/ml}$ .

<sup>c</sup> Not applicable.



86  
Fig. 2. Dose-dependent inhibition of HCV infection by isolated compounds and their cytotoxicity. Various concentrations of isolated compounds, chalepin (A), scopoletin (B), γ-fagarine (C), arborinine (D), kokusagine (E) and pseudane IX (F), were mixed with an equal amount of HCV to obtain a final concentration of 30 to 0.1 µg/ml and inoculated to Huh 7.5 cells (MOI = 0.5). After virus adsorption, the cells were cultured with the same concentrations of compounds for 46 h. The culture supernatants were harvested and titrated for virus infectivity. Percent inhibitions of HCV infectivity by each compound are shown. In parallel, cytotoxicity of the compounds was measured by WST-1 assay. Con, control. The data represent means ± SEM of data from two independent experiments.



14  
Fig. 3. Mode-of-action analysis of chalepin and pseudane IX. Huh7.5 cells infected with HCV were treated with various concentrations of chalepin (A) and pseudane IX (B) at different timings; (i) only during virus inoculation (entry), (ii) only after virus inoculation (post entry) and (iii) both during and after virus inoculation (both), with a total incubation time of 48 h. Con, control. Percent inhibitions at each concentration and IC<sub>50</sub> values are shown. The data represent mean ± SEM of data from two independent experiments.



26.7 ± 1.3, 5.2 ± 0.7 and 1.7 ± 0.5 µg/ml, respectively. Also, those for pseudane IX were 11.5 ± 0.2, 3.0 ± 0.9 and 1.4 ± 0.9 µg/ml, respectively. These results suggested that chalepin and pseudane IX inhibited HCV predominantly at the post-entry step.


102  
3.4. Inhibition of HCV RNA replication and HCV protein synthesis by chalepin and pseudane IX

To further confirm that chalepin and pseudane IX inhibit HCV infection at the post-entry step of HCV life cycle, we investigated the effect of those compounds on HCV RNA replication, viral protein synthesis and infectious virus production. Real-time quantitative RT-PCR analysis revealed that chalepin and pseudane IX at 3 and 10 µg/ml inhibited

HCV RNA replication (Fig. 4A). Consistently, immunoblotting analysis demonstrated that both compounds inhibited HCV protein synthesis (Fig. 4B). We confirmed in the same experiment that they inhibited HCV production in the culture (Fig. 4C).

4. Discussion

Medicinal plants are good resources to search a novel drug candidate(s). A wide variety of phytochemicals that inhibit virus infections have been isolated and reported. In the present study, we examined the possible anti-HCV activity of *R. angustifolia* extracts and its constituents. *R. angustifolia* is widely distributed throughout the world and has been used as folk medicine for treatment of certain diseases. Plants of the genus *Ruta* have been commonly used



24  
5  
Fig. 4. Inhibition of HCV RNA replication, HCV protein synthesis and infectious virus production by chalepin and pseudane IX. (A) Huh7.5 cells infected with HCV (MOI = 2) and treated with chalepin or pseudane IX (10 and 3 µg/ml) and the untreated control were subjected to real-time quantitative RT-PCR analysis. HCV RNA levels were normalized to GAPDH mRNA expression levels. Data represent mean ± SEM of data from two independent experiments, and the value for the untreated control at 1 dpi was arbitrarily expressed as 1. Chal, chalepin; Pseu, pseudane. \*, p < 0.001; †, p < 0.05. (B) Levels of HCV NS3 protein synthesis in the cells described in (A) were measured by immunoblotting analysis using anti-NS3 monoclonal antibody at 1 and 2 dpi. GAPDH served as an internal control to verify equal amounts of sample loading. Signal intensities of NS3 were normalized to the corresponding GAPDH signal. (C) Virus infectivity in the culture supernatants of the cells described in (A) was measured. \*, p < 0.001; †, p < 0.05.



in both ancient and modern medicine practices in the Mediterranean region to cure pulmonary diseases, rheumatic diseases and helminthic infections [15,27]. A large number of chemical constituents have been isolated from the genus *Ruta*, such as coumarins, alkaloids, benzoquinones, flavone glycosides, sterols, triterpenoids, acridone alkaloids, stigmasterol, lupeol, 5-methoxyarborinine, 5-hydroxyarborinine, ostruthin, bergapten, psoralen, xanthotoxin, limonoid obacunone, isopimpinellin, integriquinolone, kokusaginine, dictamnine, furoquinolin alkaloid, xanthyletin and xanthoxyletin [28,29]. It was also reported that coumarin compounds are the major constituents of the plants in the Rutaceae family, with about two hundred different coumarins being identified [30,31]. Specifically, four aromatic compounds, such as moskachans A, B, C and D have been identified from *R. angustifolia* [17]. Another study on a chloroform extract of aerial part of *R. angustifolia* identified angustifolin, alkaloid graveolin, scoparone and 6,7,8-trimethoxycoumarin [16].

In this study, *R. angustifolia* leaves were subjected to extraction in different polarities of solvents and antiviral activities of the extracts were examined against the J6/JFH1-P47 strain of HCV. The results revealed that a dichloromethane extract of *R. angustifolia* leaves possessed the most potent activity (Table 1), suggesting that a semi-polar compound(s) extracted by dichloromethane was involved in the anti-HCV activity. We further fractionated the dichloromethane extract to obtain 6 fractions. Anti-HCV assay showed that fraction 4 inhibited HCV with  $IC_{50}$  of 0.7  $\mu\text{g/ml}$  without apparent cytotoxicity (Table 2). As the TLC profile of the positive fractions suggested the presence of chlorophyll or its related substance(s) in this fraction (data not show), we used activated-charcoal in column chromatography to remove them. Finally, we isolated six compounds and determined their structures by combination of HPLC, LC-MS and NMR; they were chalepin, scopoletin,  $\gamma$ -fagarine, arborinine, kokusaginine, and pseudane IX. Chalepin and scopoletin are classified as coumarins while the remaining four ( $\gamma$ -fagarine, arborinine, kokusaginine and pseudane IX) are alkaloids.

Chalepin, which has been isolated from *R. chalepensis* [32], *Stauranthus perforates* [33], *Clausena anisata* [34] and *Clausena lansium* [35], belongs to furocoumarin compounds with furan ring fused to the coumarin structure. It was reported to possess antimicrobial activities against *Pseudomonas aeruginosa* and *Trichomonas* as well as anti-coagulant activities [34,35]. However, there has been no report so far regarding its antiviral activity against HCV. To the best of our knowledge, the present study is the first to demonstrate anti-HCV activities of chalepin with  $IC_{50}$  being  $1.7 \pm 0.5 \mu\text{g/ml}$  ( $5.4 \pm 0.5 \mu\text{M}$ ) (Table 3). It was reported that chalepin inhibited respiration of isolated rat liver mitochondria by 40% at the concentration of  $16 \mu\text{M}$  ( $5.0 \mu\text{g/ml}$ ) [36]. Under our experimental conditions using cultured cells, chalepin exerted only marginal cytotoxicity, if any, at  $30 \mu\text{g/ml}$  (Fig. 2).

We have reported that other coumarin compounds, such as glycycomarin, glycyrin and glycyrol, possess anti-HCV activities, with  $IC_{50}$  of 8.8, 7.2, and  $4.6 \mu\text{g/ml}$ , respectively [11]. The basic structure (1,2-benzopyron) of coumarin appears to be important for binding to HCV [37]. Fourteen compounds out of 24 coumarin derivatives were reported to inhibit HCV NS5B RNA polymerase activities with  $IC_{50}$  values between 17 and  $63 \mu\text{M}$ . The activities of those compounds were influenced by the position of methylation or hydroxylation groups in the

benzopyron ring [37]. Recently, benzimidazole derivatives of coumarins have been reported to possess increased inhibitory effect on RNA polymerase of HCV NS5B [38,39]. On the other hand, scopoletin, which is the other coumarin isolated in the present study, did not inhibit HCV at the concentration of  $30 \mu\text{g/ml}$  ( $155 \mu\text{M}$ ). Scopoletin isolated from several plants, such as *Erycibe obtusifolia* Benth, *Aster tataricus* and *Foeniculum vulgare*, and its synthetic derivatives have been extensively studied [40]. Further detailed analyses of derivatives of chalepin and scopoletin would help us understand the structural basis of anti-HCV activity of chalepin and generate a more potent anti-HCV compound(s).

$\gamma$ -Fagarine, arborinine and kokusaginine, which are alkaloid compounds, showed moderate inhibition with  $IC_{50}$  of  $20.4 \pm 0.4$ ,  $6.4 \pm 0.7$  and  $6.4 \pm 1.6 \mu\text{g/ml}$ , respectively (Table 3). These compounds have been isolated from several plants, including the Rutaceae family [41–44] and arborinine was previously reported to inhibit human rhinovirus with  $IC_{50}$  of  $3.19 \mu\text{M}$  by virtual model [45]. Pseudane IX, another alkaloid, also known as 2-nonyl-4 (1H)-quinolone, 2-nonyl-4-hydroxyquinoline (NHQ) or 4-hydroxy-2-nonylquinolone [46], showed potent anti-HCV activities with  $IC_{50}$  of  $1.4 \pm 0.2 \mu\text{g/ml}$  ( $5.1 \pm 0.2 \mu\text{M}$ ) (Table 3). A wide variety of quinolones have been used as antimicrobial, anticancer and antiallergenic agents. Quinolones are known as broad-spectrum antibacterial agents with the main structure of 1,4 dihydro-4-oxo-quinolinyl moiety. Quinolones inhibit prokaryotic type II topoisomerases through direct binding to bacterial chromosome, and likewise, it may bind to viral nucleic acids and/or nucleoprotein complexes to act as antivirals [47]. Quinolones consist of heterobicyclic aromatic compounds and the moiety of  $C_9H_{19}$  at carbon number 2 of pseudane IX may play an important role in its activities. Quinolones have been reported to act as inhibitors of HCV NS5B RNA polymerase by binding to the allosteric site II (non-nucleoside inhibitor-site 2) of this protein [48]. Twelve fluoroquinolone derivatives were reported to inhibit HCV replication and HCV NS3 helicase activity in cultured cells. Among them, feroxacin, difloxacin, ofloxacin, 8-quinolinol, enoxacin, lumifloxacin and flumiquine were reported to inhibit HCV with  $IC_{50}$  of 3.8, 2.5, 2.4, 2.2, 1.0, 1.9 and  $1.7 \mu\text{M}$ , respectively [49].

The key steps in HCV life cycle include entry into the host cells, uncoating and replication of the viral genome, translation of virus proteins, and assembly and release of the virion [8,18]. To determine the impact of active compounds on HCV life cycle, we conducted time-of-addition experiments. We observed that chalepin and pseudane IX exhibited their anti-HCV activities at the post-entry step, inhibiting HCV RNA replication and NS3 protein synthesis.

In conclusion, we have identified chalepin and pseudane IX as anti-HCV compounds. These compounds could be good candidates to develop novel anti-HCV drugs.

## 1 Acknowledgments

The authors are grateful to Dr. C. M. Rice (The Rockefeller University, New York, NY) for providing Huh-7.5 cells and pFL-J6/JFH1. This work was supported in part by Science and Technology Research Partnerships for Sustainable Development (SATREPS) supported by the Japan Science and Technology Agency, and the Japan International Cooperation Agency (55)

and the Ministry of Research and Technology (RISTEK), Republic of Indonesia. This work was also carried out as part of Japan Initiative for Global Research Network on Infectious Diseases (J-GRID), Ministry of Education, Culture, Sports, Science and Technology, Japan, and the Global Center of Excellence (G-COE) Program at Kobe University Graduate School of Medicine.

### 3 Appendix A. Supplementary data

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.fitote.2014.10.011>.

### References

- [1] Rupp D, Bartenschlager R. Targets for antiviral therapy of hepatitis C. *Semin Liver Dis* 2014;34:9–21.
- [2] Moradpour D, Penin F, Rice CM. Replication of hepatitis C virus. *Nat Rev Microbiol* 2007;5:453–63.
- [3] Lange CM, Jacobson IM, Rice CM, Zeuzem S. Emerging therapies for the treatment of hepatitis C. *EMBO Mol Med* 2014;6:4–15.
- [4] Gottwein JM, Scheel TK, Jensen TB, Lademann JB, Prentoe JC, Knudsen ML, et al. Development and characterization of hepatitis C virus genotype 1–7 cell culture systems: role of CD81 and scavenger receptor class B type I and effect of antiviral drugs. *Hepatology* 2009;49:364–77.
- [5] Ireton RC, Gale Jr M. Pushing to a cure by harnessing innate immunity against hepatitis C virus. *Antiviral Res* 2014;108:156–64.
- [6] Shoji I, Deng L, Hotta H. Molecular mechanism of hepatitis C virus-induced glucose metabolic disorders. *Front Microbiol* 2011;2:278.
- [7] Deng L, Shoji I, Ogawa W, Kaneda S, Soga T, Jiang DP, et al. Hepatitis C virus infection promotes hepatic gluconeogenesis through an NS5A-mediated, FoxO1-dependent pathway. *J Virol* 2011;85:8556–68.
- [8] Ploss A, Dubuisson J. New advances in the molecular biology of hepatitis C virus infection: towards the identification of new treatment targets. *Gut* 2012;61(Suppl. 1):i25–35.
- [9] Ravikummar YS, Ray U, Nandhitha M, Perween A, Raja Naika H, Khanna N, et al. Inhibition of hepatitis C virus replication by herbal extract: *Phyllanthus amarus* as potent natural source. *Virus Res* 2011;158:89–97.
- [10] Wahyuni TS, Tumewu L, Permanasari AA, Apriani E, Adianti M, Rahman A, et al. Antiviral activities of Indonesian medicinal plants in the East Java region against hepatitis C virus. *Virology* 2013;10:259.
- [11] Adianti M, Aoki C, Komoto M, Deng L, Shoji I, Wahyuni TS, et al. Anti-hepatitis C virus compounds obtained from *Glycyrrhiza uralensis* and other *Glycyrrhiza* species. *Microbiol Immunol* 2014;58:180–7.
- [12] Ratnoglik SL, Aoki C, Sudarmono P, Komoto M, Deng L, Shoji I, et al. Antiviral activity of extracts from *Morinda citrifolia* leaves and chlorophyll catabolites, pheophorbide a and pyropheophorbide a, against hepatitis C virus. *Microbiol Immunol* 2014;58:188–94.
- [13] Ashfaq UA, Idrees S. Medicinal plants against hepatitis C virus. *World J Gastroenterol* 2014;20:2941–7.
- [14] Calland N, Dubuisson J, Rouille Y, Seron K. Hepatitis C virus and natural compounds: a new antiviral approach? *Viruses* 2012;4:2197–217.
- [15] Pollio A, De Natale A, Appetiti E, Aliotta G, Touwaide A. Continuity and change in the Mediterranean medical tradition: *Ruta* spp. (rutaceae) in Hippocratic medicine and present practices. *J Ethnopharmacol* 2008;116:469–82.
- [16] Del Castillo JB, Luis FR, Secundino M. Angustifolin, a coumarin from *Ruta angustifolia*. *Phytochemistry* 1984;23:2095–6.
- [17] Del Castillo JB, Secundino M, Luis FR. Four aromatic derivatives from *Ruta angustifolia*. *Phytochemistry* 1986;25:2209–10.
- [18] Lindenbach BD, Evans MJ, Syder AJ, Wolk B, Tellinghuisen TL, Liu CC, et al. Complete replication of hepatitis C virus in cell culture. *Science* 2005;309:623–6.
- [19] Fuchino H, Sekita S, Mori K, Kawahara N, Satake M, Kiuchi F. A new leishmanicidal saponin from *Brunfelsia grandiflora*. *Chem Pharm Bull* 2008;56:93–6.
- [20] Fuchino H, Daikonya A, Kumagai T, Goda Y, Takahashi Y, Kawahara N. Two new labdane diterpenes from fresh leaves of *Leonurus japonicus* and their degradation during drying. *Chem Pharm Bull* 2013;61:497–503.
- [21] Bungyoku Y, Shoji I, Makine T, Adachi T, Hayashida K, Nagano-Fujii M, et al. Efficient production of infectious hepatitis C virus with adaptive mutations in cultured hepatoma cells. *J Gen Virol* 2009;90:1681–91.
- [22] Nicoletti M, Delle Monache F, Marini-Bettolo GB. Carbon-13 spectral data of prenylated coumarins. *Planta Med* 1982;45:250–1.
- [23] Wu T-S, Shi L-S, Wang J-J, Lou S-C, Chang H-C, Chen Y-P, et al. Cytotoxic and antiplatelet aggregation principles of *Ruta Graveolens*. *J Chin Chem Soc* 2003;50:171–8.
- [24] Min YD, Kwon HC, Yang MC, Lee KH, Choi SU, Lee KR. Isolation of limonoids and alkaloids from *Phellodendron amurense* and their multidrug resistance (MDR) reversal activity. *Arch Pharm Res* 2007;30:58–63.
- [25] Bodendiek SB, Mahieux C, Hänsel W, Wulff H. 4-Phenoxybutoxy-substituted heterocycles – a structure–activity relationship study of blockers of the lymphocyte potassium channel Kv1.3. *Eur J Med Chem* 2009;44:1838–52.
- [26] Somanathan R, Smith KM. Synthesis of some 2-alkyl-4-quinolone and 2-alkyl-4-methoxyquinoline alkaloids. *J Heterocycl Chem* 1981;18:1077–9.
- [27] Kannan R, Babu UV. Identity and pharmacognosy of *Ruta graveolens* Linn. *Anc Sci Life* 2012;32:16–9.
- [28] Wijeratne EMK, Bandara BMR, Gunatilaka AAL, Tezuka Y, Kikuchi T. Chemical constituents of three Rutaceae species from Sri Lanka. *J Nat Prod* 1992;55:1261–9.
- [29] Mohan PS, Ramesh M, Shanmugam P. Studies on the Indian Rutaceae. Chemical investigation of *Limonia alata* and *Evodia lunu-ankenda*. *J Nat Prod* 1985;48:501.
- [30] Gray AI, Waterman PG. Coumarins in the Rutaceae. *Phytochemistry* 1978;17:845–64.
- [31] Mileti S, Massot B, Gontier E, Bourgaud F, Guckert A. *Ruta graveolens* L.: a promising species for the production of furanocoumarins. *Plant Sci* 2001;161:189–99.
- [32] Mohr N, Budzikiewicz H, El-Tawil BAH, El-Beih FKA. Further furoquinolone alkaloids from *Ruta chalepensis*. *Phytochemistry* 1982;21:1838–9.
- [33] Anaya AL, Macías-Rubalcava M, Cruz-Ortega R, García-Santana C, Sánchez-Monterrubio PN, Hernández-Bautista BE, et al. Allelochemicals from *Stauranthus perforatus*, a Rutaceous tree of the Yucatan Peninsula, Mexico. *Phytochemistry* 2005;66:487–94.
- [34] Emerole G, Thabrew MI, Anosa V, Okorie DA. Structure–activity relationship in the toxicity of some naturally occurring coumarins-chalepin, imperatorin and oxypeucedanin. *Toxicology* 1981;20:71–80.
- [35] Adebajo AC, Iwalewa EO, Obuotor EM, Ibikunle GF, Omisore NO, Adewunmi CO, et al. Pharmacological properties of the extract and some isolated compounds of *Clauseria lamsium* stem bark: anti-trichomonal, antidiabetic, anti-inflammatory, hepatoprotective and antioxidant effects. *J Ethnopharmacol* 2009;122:10–9.
- [36] Olorunsogo OO, Emerole GO, Malomo SO, Thabrew MI. Inhibition of mitochondrial respiration by chalepin, a naturally occurring furcoumarin. *Toxicol Lett* 1983;15:323–7.
- [37] Nichols DB, Leao RA, Basu A, Chudayeu M, de Moraes Pde F, Talele TT, et al. Evaluation of coumarin and neoflavone derivatives as HCV NS5B polymerase inhibitors. *Chem Biol Drug Des* 2013;81:607–14.
- [38] Neyts J, De Clercq E, Singha R, Chang YH, Das AR, Chakraborty SK, et al. Structure–activity relationship of new anti-hepatitis C virus agents: heterobicyclic-coumarin conjugates. *J Med Chem* 2009;52:1486–90.
- [39] Peng XM, Damu GL, Zhou C. Current developments of coumarin compounds in medicinal chemistry. *Curr Pharm Des* 2013;19:3884–930.
- [40] Liu W, Hua J, Zhou J, Zhang H, Zhu H, Cheng Y, et al. Synthesis and in vitro antitumor activity of novel scopoletin derivatives. *Bioorg Med Chem Lett* 2012;22:5008–12.
- [41] Rethy B, Zupko I, Minorics R, Hohmann J, Ocsovszki I, Falkay G. Investigation of cytotoxic activity on human cancer cell lines of arborinine and furanocoumarins isolated from *Ruta graveolens*. *Planta Med* 2007;73:41–8.
- [42] Haque MM, Begum S, Sohrab MH, Ahsan M, Hasan CM, Ahmed N, et al. Secondary metabolites from the stem of *Ravenia spectabilis* Lindl. *Pharmacogn Mag* 2013;9:76–80.
- [43] Cuca Suarez LE, Patarrayo ME, Lozano JM, Delle Monache F. Biological activity of secondary metabolites from *Peltostigma guatemalense*. *Nat Prod Res* 2009;23:370–4.
- [44] Veiga TA, King-Diaz B, Marques AS, Sampaio OM, Vieira PC, da Silva MF, et al. Furoquinoline alkaloids isolated from *Balfourodendron riedelianum* as photosynthetic inhibitors in spinach chloroplasts. *J Photochem Photobiol B* 2013;120:36–43.
- [45] Röllinger JM, Schuster D, Danzl B, Schwaiger S, Markt P, Schmidtke M, et al. In silico target fishing for rationalized ligand discovery exemplified on constituents of *Ruta graveolens*. *Planta Med* 2009;75:195–204.
- [46] Heeb S, Fletcher MP, Chhabra SR, Diggle SP, Williams P, Camara M. Quinolones: from antibiotics to autoinducers. *FEMS Microbiol Rev* 2011;35:247–74.
- [47] Richter S, Parolin C, Palumbo M, Palu G. Antiviral properties of quinolone-based drugs. *Curr Drug Targets Infect Disord* 2004;4:111–6.
- [48] Kumar DV, Rai R, Brameld KA, Somoza JR, Rajagopalan R, Janc JW, et al. Quinolones as HCV NS5B polymerase inhibitors. *Bioorg Med Chem Lett* 2011;21:82–7.
- [49] Khan IA, Siddiqui S, Rehmani S, Kazmi SU, Ali SH. Fluoroquinolones inhibit HCV by targeting its helicase. *Antivir Ther* 2012;17:467–76.

# Inhibition of hepatitis C virus replication by chalepin and pseudane IX isolated from *Ruta angustifolia* leaves

---

## ORIGINALITY REPORT

---

25%

SIMILARITY INDEX

13%

INTERNET SOURCES

22%

PUBLICATIONS

0%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

- 1** Kenji Nakashima. "Inhibition of hepatitis C virus replication through adenosine monophosphate-activated protein kinase-dependent and -independent pathways : Role of AMPK in HCV replication", *Microbiology and Immunology*, 11/2011

Publication

1%
- 2** S. Noer, A. Abinawanto, A. Basith. "Molecular identification of the medicinal herb plant *Ruta angustifolia* from Lembang Bandung using chloroplast DNA markers", AIP Publishing, 2018

Publication

1%
- 3** Jardim, A.C.G., Z. Igloi, J.F. Shimizu, V.A.F.F.M. Santos, L.G. Felipe, B.F. Mazzeu, Y. Amako, M. Furlan, M. Harris, and P. Rahal. "Natural compounds isolated from Brazilian plants are potent inhibitors of hepatitis C virus replication in vitro", *Antiviral Research*, 2015.

Publication

<1%
- 4** Diggle, S.P.. "Functional Genetic Analysis Reveals a 2-Alkyl-4-Quinolone Signaling

<1%



System in the Human Pathogen *Burkholderia pseudomallei* and Related Bacteria",  
Chemistry & Biology, 200607

Publication

---

5

Yu, Jung-Sheng, Wei-Chun Chen, Chin-Kai Tseng, Chun-Kuang Lin, Yao-Chin Hsu, Yen-Hsu Chen, and Jin-Ching Lee. "Sulforaphane Suppresses Hepatitis C Virus Replication by Up-Regulating Heme Oxygenase-1 Expression through PI3K/Nrf2 Pathway", PLoS ONE, 2016.

<1 %

Publication

---

6

Hanane Elayadi, Michael Smietana, Christophe Pannecouque, Pieter Leyssen, Johan Neyts, Jean-Jacques Vasseur, Hassan B. Lazrek. "Straightforward synthesis of triazoloacyclonucleotide phosphonates as potential HCV inhibitors", Bioorganic & Medicinal Chemistry Letters, 2010

<1 %

Publication

---

7

Chung, Chueh-Yao, Ching-Hsuan Liu, Thierry Burnouf, Guey-Horng Wang, Shun-Pang Chang, Alagie Jasse, Chen-Jei Tai, Cheng-Jeng Tai, Ching-Jang Huang, Christopher D. Richardson, Ming-Hong Yen, Chun-Ching Lin, and Liang-Tzung Lin. "Activity-based and fraction-guided analysis of *Phyllanthus urinaria* identifies loliolide as a potent inhibitor of hepatitis C virus entry", Antiviral Research, 2016.

<1 %

8

[patents.justia.com](http://patents.justia.com)

Internet Source

<1 %

---

9

[www.wjgnet.com](http://www.wjgnet.com)

Internet Source

<1 %

---

10

[tessera.spandidos-publications.com](http://tessera.spandidos-publications.com)

Internet Source

<1 %

---

11

Samatchaya Boonchawalit. "Molecular Evolution of HIV-1 CRF01\_AE Env in Thai Patients", PLoS ONE, 11/02/2011

Publication

<1 %

---

12

Michael J. Sofia, Wonsuk Chang, Phillip A. Furman, Ralph T. Mosley, Bruce S. Ross. "Nucleoside, Nucleotide, and Non-Nucleoside Inhibitors of Hepatitis C Virus NS5B RNA-Dependent RNA-Polymerase", Journal of Medicinal Chemistry, 2012

Publication

<1 %

---

13

Oriana Tabarrini, Serena Massari, Dirk Daelemans, Miguel Stevens et al. "Structure-Activity Relationship Study on Anti-HIV 6-Desfluoroquinolones", Journal of Medicinal Chemistry, 2008

Publication

<1 %

---

14

Eun-Mee Park, Lam N. Nguyen, Yun-Sook Lim, Soon B. Hwang. "Farnesyl-diphosphate farnesyltransferase 1 regulates hepatitis C virus propagation", FEBS Letters, 2014

<1 %

15 Erica Silberstein, Kathleen Mihalik, Laura Uritzky, Ewan P. Plant et al. "Persistent Growth of a Human Plasma-Derived Hepatitis C Virus Genotype 1b Isolate in Cell Culture", PLoS Pathogens, 2010  
Publication

---

16 [jvi.asm.org](http://jvi.asm.org)  
Internet Source

---

17 [mdpi.com](http://mdpi.com)  
Internet Source

---

18 Daniel Rupp, Ralf Bartenschlager. "Targets for Antiviral Therapy of Hepatitis C", Seminars in Liver Disease, 2014  
Publication

---

19 [www.science.gov](http://www.science.gov)  
Internet Source

---

20 Nag, A., J. M. Robotham, and H. Tang. "Suppression of Viral RNA Binding and the Assembly of Infectious Hepatitis C Virus Particles In Vitro by Cyclophilin Inhibitors", Journal of Virology, 2012.  
Publication

---

21 Alexander Ploss, Jean Dubuisson. "New advances in the molecular biology of hepatitis C virus infection: towards the identification of new treatment targets", Gut, 2012  
Publication

---



22

www0.nih.go.jp

Internet Source

<1 %

23

Yukihiro Furusawa, Yuka Yamanouchi, Takashi Iizumi, Qing-Li Zhao et al.

"Checkpoint kinase 2 is dispensable for regulation of the p53 response but is required for G2/M arrest and cell survival in cells with p53 defects under heat stress", Apoptosis, 2017

Publication

<1 %

24

Eyre, Nicholas S, Karla J Helbig, and Michael R Beard. "Current and future targets of antiviral therapy in the hepatitis C virus life cycle", Future Virology, 2014.

Publication

<1 %

25

"Hepatitis", Wiley, 2006

Publication

<1 %

26

Bushra Sultana, Sadaf Yaqoob, Zohaib Zafar, Haq Nawaz Bhatti. "Escalation of liver malfunctioning: A step toward Herbal Awareness", Journal of Ethnopharmacology, 2018

Publication

<1 %

27

Jian-Rui Li, Wen-Jing Li, Jun-Jun Cheng, Meng-Hao Huang et al. "A Simple but Accurate Method for Evaluating Drug-Resistance in Infectious HCVcc System", BioMed Research International, 2017

Publication

<1 %

28

[aip.scitation.org](http://aip.scitation.org)

Internet Source

&lt;1%

29

Robert G. Gentles, Min Ding, John A. Bender, Carl P. Bergstrom et al. "Discovery and Preclinical Characterization of the Cyclopropylindolobenzazepine BMS-791325, A Potent Allosteric Inhibitor of the Hepatitis C Virus NS5B Polymerase", *Journal of Medicinal Chemistry*, 2014

Publication

&lt;1%

30

Julien Martinez, Ben Longdon, Simone Bauer, Yuk-Sang Chan et al. "Symbionts Commonly Provide Broad Spectrum Resistance to Viruses in Insects: A Comparative Analysis of Wolbachia Strains", *PLoS Pathogens*, 2014

Publication

&lt;1%

31

He, Changyu, Qinglai Yang, Lianjiang Tan, Bingya Liu, Zhenggang Zhu, Bing Gong, Yu-Mei Shen, and Zhifeng Shao. "Design and synthesis of redox and oxidative dual responsive block copolymer micelles for intracellular drug delivery", *European Polymer Journal*, 2016.

Publication

&lt;1%

32

Jantan, I.. "Inhibitory effect of compounds from Zingiberaceae species on human platelet aggregation", *Phytomedicine*, 20080425

Publication

&lt;1%

33	<a href="http://virologyj.biomedcentral.com">virologyj.biomedcentral.com</a> Internet Source	<1 %
34	Eugene Lin. "An artificial neural network approach to the drug efficacy of interferon treatments", <i>Pharmacogenomics</i> , 10/2006 Publication	<1 %
35	<a href="http://www.escience.cn">www.escience.cn</a> Internet Source	<1 %
36	<a href="http://tel.archives-ouvertes.fr">tel.archives-ouvertes.fr</a> Internet Source	<1 %
37	Current Topics in Microbiology and Immunology, 2013. Publication	<1 %
38	Costin-Ioan Popescu, Nathalie Callens, Dave Trinel, Philippe Roingard et al. "NS2 Protein of Hepatitis C Virus Interacts with Structural and Non-Structural Proteins towards Virus Assembly", <i>PLoS Pathogens</i> , 2011 Publication	<1 %
39	Walid Hamdy El-Tantawy, Abeer Temraz. "Natural products for the management of the hepatitis C virus: a biochemical review", <i>Archives of Physiology and Biochemistry</i> , 2018 Publication	<1 %
40	Yu Wei, Jinlong Li, Jie Qing, Mingjie Huang et al. "Discovery of Novel Hepatitis C Virus NS5B Polymerase Inhibitors by Combining Random	<1 %



## Forest, Multiple e-Pharmacophore Modeling and Docking", PLOS ONE, 2016

Publication

41

S.-J. Jiang. "Isolation and phytotoxicity of a metabolite from *Curvularia eragrostidis* and characterisation of its modes of action", *Annals of Applied Biology*, 2/2008

Publication

<1 %

42

Yamanaka, Atsushi, Yuko Tabuchi, Kris C. Mulyatno, Helen Susilowati, Eryk Hendrianto, Soegeng Soegijanto, and Eiji Konishi. "Dengue virus infection-enhancing and neutralizing antibody balance in children of the Philippines and Indonesia", *Microbes and Infection*, 2012.

Publication

<1 %

43

[www.labome.org](http://www.labome.org)

Internet Source

<1 %

44

Wang, X., K. Yamataka, M. Okamoto, S. Ikeda, and M. Baba. "Potent and Selective Inhibition of Tat-Dependent HIV-1 Replication in Chronically Infected Cells by a Novel Naphthalene Derivative JTK-101", *Antiviral Chemistry & Chemotherapy*, 2007.

Publication

<1 %

45

[heterocycles.jp](http://heterocycles.jp)

Internet Source

<1 %

46

[ip-science.thomsonreuters.com.au](http://ip-science.thomsonreuters.com.au)

Internet Source

<1 %

---

47 Kuo, Yi-Chen, I-Yin Chen, Shin C. Chang, Shun-Chi Wu, Tzu-Min Hung, Po-Huang Lee, Kunitada Shimotohno, and Ming-Fu Chang. "Hepatitis C virus NS5A protein enhances gluconeogenesis through upregulation of Akt-/JNK-PEPCK signalling pathways", Liver International, 2013. <1%

Publication

---

48 F. Chen, Y. Zhao, M. Liu, D. Li, H. Wu, H. Chen, Y. Zhu, F. Luo, J. Zhong, Y. Zhou, Z. Qi, X.-L. Zhang. "Functional Selection of Hepatitis C Virus Envelope E2-Binding Peptide Ligands by Using Ribosome Display", Antimicrobial Agents and Chemotherapy, 2010. <1%

Publication

---

49 Daisuke Akazawa, Kenichi Morikawa, Noriaki Omi, Hitoshi Takahashi et al. "Production and characterization of HCV particles from serum-free culture", Vaccine, 2011. <1%

Publication

---

50 [jbiomedsci.biomedcentral.com](http://jbiomedsci.biomedcentral.com) <1%

Internet Source

---

51 [www.freepatentsonline.com](http://www.freepatentsonline.com) <1%

Internet Source

---

52 [www.jpmsonline.com](http://www.jpmsonline.com) <1%

Internet Source

---

[www.artificial-organs.com](http://www.artificial-organs.com)

53

Internet Source

<1 %

54

Jessica Wagoner, Amina Negash, Olivia J. Kane, Laura E. Martinez et al. "Multiple effects of silymarin on the hepatitis C virus lifecycle", *Hepatology*, 2010

Publication

<1 %

55

[citeseerx.ist.psu.edu](http://citeseerx.ist.psu.edu)

Internet Source

<1 %

56

[eprints.nottingham.ac.uk](http://eprints.nottingham.ac.uk)

Internet Source

<1 %

57

[www.eijppr.com](http://www.eijppr.com)

Internet Source

<1 %

58

Oliva, Anna, Kumudini M. Meepagala, David E. Wedge, Dewayne Harries, Amber L. Hale, Giovanni Aliotta, and Stephen O. Duke. "Natural Fungicides from *Ruta graveolens* L. Leaves, Including a New Quinolone Alkaloid", *Journal of Agricultural and Food Chemistry*, 2003.

Publication

<1 %

59

Yao, W., H. Cai, X. Li, T. Li, L. Hu, and T. Peng. "Endoplasmic Reticulum Stress Links Hepatitis C Virus RNA Replication to Wild-Type PGC-1 /Liver-Specific PGC-1 Upregulation", *Journal of Virology*, 2014.

Publication

<1 %

Li, Y.P.. "Non-genotype-specific role of the



- 
- 60 hepatitis C virus 5' untranslated region in virus production and in inhibition by interferon", *Virology*, 20111220  
Publication <1%
- 
- 61 Jun Ishihara, Tetsuya Sugimoto, Akio Murai. "Asymmetric total synthesis of syringolide 1, a nonproteinaceous elicitor", *Tetrahedron*, 1997  
Publication <1%
- 
- 62 Aleksandar Včev, Jelena Jakab, Lucija Kuna, Martina Smolić. "Chapter 13 Health-Related Quality of Life in Antiviral-Treated Chronic Hepatitis C Patients", IntechOpen, 2017  
Publication <1%
- 
- 63 [koreascience.or.kr](http://koreascience.or.kr)  
Internet Source <1%
- 
- 64 T. Schaller, N. Appel, G. Koutsoudakis, S. Kallis, V. Lohmann, T. Pietschmann, R. Bartenschlager. "Analysis of Hepatitis C Virus Superinfection Exclusion by Using Novel Fluorochrome Gene-Tagged Viral Genomes", *Journal of Virology*, 2007  
Publication <1%
- 
- 65 [www.med.kobe-u.ac.jp](http://www.med.kobe-u.ac.jp)  
Internet Source <1%
- 
- 66 [www.teriin.org](http://www.teriin.org)  
Internet Source <1%
- 
- 67 [www.intmedpress.com](http://www.intmedpress.com)  
Internet Source <1%
-

68

Muhammad Faraz Anwar, Ramsha Khalid, Alina Hasanain, Sadaf Naeem, Shamshad Zarina, Syed Hani Abidi, Syed Ali. "Integrated Cheminformatics-Molecular Docking Approach to Drug Discovery Against Viruses", Infectious Disorders - Drug Targets, 2020

Publication

&lt;1 %

69

[legacy.impactjournals.com](http://legacy.impactjournals.com)

Internet Source

&lt;1 %

70

[mapsdatabase.com](http://mapsdatabase.com)

Internet Source

&lt;1 %

71

Minjung Lee, Myoungki Son, Eunhyun Ryu, Yu Su Shin, Jong Gwang Kim, Byung Woog Kang, Hyosun Cho, Hyojeung Kang.

"Quercetin-induced apoptosis prevents EBV infection", Oncotarget, 2015

Publication

&lt;1 %

72

[core.ac.uk](http://core.ac.uk)

Internet Source

&lt;1 %

73

Yong Ren, Xuan Jiang, Deng Pan, Hai-Quan Mao. "Charge Density and Molecular Weight of Polyphosphoramidate Gene Carrier Are Key Parameters Influencing Its DNA Compaction Ability and Transfection Efficiency", Biomacromolecules, 2010

Publication

&lt;1 %

74

[repository.unair.ac.id](http://repository.unair.ac.id)

Internet Source

&lt;1 %

75

Wei-Sheng Lo, Yun-Ping Lim, Chien-Chih Chen, Chih-Chien Hsu, Pavel Souček, Chul-Ho Yun, Wen Xie, Yune-Fang Ueng. "A dual function of the furanocoumarin chalepensis in inhibiting Cyp2a and inducing Cyp2b in mice: the protein stabilization and receptor-mediated activation", Archives of Toxicology, 2012

Publication

&lt;1 %

76

[www.tandfonline.com](http://www.tandfonline.com)

Internet Source

&lt;1 %

77

Gomez, C. E., B. Perdiguero, M. V. Cepeda, L. Mingorance, J. Garcia-Arriaza, A. Vandermeeren, C. O. S. Sorzano, and M. Esteban. "High, Broad, Polyfunctional, and Durable T Cell Immune Responses Induced in Mice by a Novel Hepatitis C Virus (HCV) Vaccine Candidate (MVA-HCV) Based on Modified Vaccinia Virus Ankara Expressing the Nearly Full-Length HCV Genome", Journal of Virology, 2013.

Publication

&lt;1 %

78

[d-nb.info](http://d-nb.info)

Internet Source

&lt;1 %

79

[www.doria.fi](http://www.doria.fi)

Internet Source

&lt;1 %

80

C. Staedel. "A functional selection of viral genetic elements in cultured cells to identify hepatitis C virus RNA translation inhibitors",

&lt;1 %

81

Sandra Westhaus, Dorothea Bankwitz, Stefanie Ernst, Katrin Rohrmann et al. "

Characterization of the inhibition of hepatitis C virus entry by -generated and patient-derived oxidized low-density lipoprotein ", Hepatology, 2013

Publication

<1 %

82

[www.pubfacts.com](http://www.pubfacts.com)

Internet Source

<1 %

83

Tanaka, M.. "Single-point mutations of hepatitis C virus NS3 that impair p53 interaction and anti-apoptotic activity of NS3", Biochemical and Biophysical Research Communications, 20060217

Publication

<1 %

84

[www.future-science.com](http://www.future-science.com)

Internet Source

<1 %

85

[doktori.bibl.u-szeged.hu](http://doktori.bibl.u-szeged.hu)

Internet Source

<1 %

86

Masanori Ikeda. "Different anti-HCV profiles of statins and their potential for combination therapy with interferon", Hepatology, 07/2006

Publication

<1 %

87

Qu, Di, Jing Li, Xiao-Hui Yang, Zi-Dan Zhang, Xiao-Xing Luo, Ming-Kai Li, and Xia Li. "New Biscoumarin Derivatives: Synthesis, Crystal

<1 %



Structure, Theoretical Study and Antibacterial Activity against Staphylococcus aureus", Molecules, 2014.

Publication

88

Guan-Hua Du. "Natural Small Molecule Drugs from Plants", Springer Science and Business Media LLC, 2018

Publication

<1 %

89

[www.thieme-connect.com](http://www.thieme-connect.com)

Internet Source

<1 %

90

A. Felipe, J. Bielanska, N. Comes, A. Vallejo, S. Roig, S. Ramon y Cajal, E. Condom, J. Hernandez-Losa, J.C. Ferreres. "Targeting the Voltage-Dependent K<sup>+</sup> Channels Kv1.3 and Kv1.5 as Tumor Biomarkers for Cancer Detection and Prevention", Current Medicinal Chemistry, 2012

Publication

<1 %

91

"Bioactive Compounds in Underutilized Fruits and Nuts", Springer Science and Business Media LLC, 2020

Publication

<1 %

92

[www.patents.com](http://www.patents.com)

Internet Source

<1 %

93

Manfroni, Giuseppe, Rolando Cannalire, Maria Letizia Barreca, Neerja Kaushik-Basu, Pieter LEYSEN, Johan Winqvist, Nunzio Iraci, Dinesh Manvar, Jan Paeshuyse, Rupa

<1 %

Guhamazumder, Amartya Basu, Stefano Sabatini, Oriana Tabarrini, U. Helena Danielson, Johan Neyts, and Violetta Cecchetti. "The versatile nature of the 6-aminoquinolone scaffold: identification of submicromolar hepatitis C virus NS5B inhibitors", Journal of Medicinal Chemistry, 2013.

Publication

94

[www.jbc.org](http://www.jbc.org)

Internet Source

<1 %

95

Donna Sir, Wen-ling Chen, Jinah Choi, Takaji Wakita, T.S. Benedict Yen, Jing-hsiung James Ou. "Induction of incomplete autophagic response by hepatitis C virus via the unfolded protein response", Hepatology, 2008

Publication

<1 %

96

[www.jisikworld.com](http://www.jisikworld.com)

Internet Source

<1 %

97

[scholar.sun.ac.za](http://scholar.sun.ac.za)

Internet Source

<1 %

98

C J Pachuk. "RNA interference-mediated prevention and therapy for hepatocellular carcinoma", Oncogene, 06/26/2006

Publication

<1 %

99

[www.medsci.org](http://www.medsci.org)

Internet Source

<1 %

100

[link.springer.com](http://link.springer.com)

<1 %

101

Eike Steinmann. "Antiviral effects of amantadine and iminosugar derivatives against hepatitis C virus", *Hepatology*, 08/2007

Publication

<1 %

102

Marcello, T.. "Interferons @a and @I Inhibit Hepatitis C Virus Replication With Distinct Signal Transduction and Gene Regulation Kinetics", *Gastroenterology*, 200612

Publication

<1 %

103

T. Li. "Characterization of self-assembled virus-like particles of rat hepatitis E virus generated by recombinant baculoviruses", *Journal of General Virology*, 08/24/2011

Publication

<1 %

104

Gundo Diedrich. "How does hepatitis C virus enter cells?", *FEBS Journal*, 9/2006

Publication

<1 %

105

Nestor G. Iglesias. "Dynamic RNA structures in the dengue virus genome", *RNA Biology*, 03/01/2011

Publication

<1 %

106

Ashfaq, Usman A. "Medicinal plants against hepatitis C virus", *World Journal of Gastroenterology*, 2014.

Publication

<1 %

**107** Shahid Akbar. "Handbook of 200 Medicinal Plants", Springer Science and Business Media LLC, 2020 <1 %  
Publication

---

**108** Li, Chia-Ying, Wei-Jern Tsai, Amooru Gangaiah Damu, E-Jian Lee, Tian-Shung Wu, Nguyen Xuan Dung, Tran Dinh Thang, and Le Thanh. "Isolation and Identification of Antiplatelet Aggregatory Principles from the Leaves of Piper lolot", Journal of Agricultural and Food Chemistry, 2007. <1 %  
Publication

---

**109** Bloom Jonathan, Zapf Christoph, Levin Jeremy, Flint Michael. "Recent Advances in Hepatitis C Therapies", Wiley, 2010 <1 %  
Publication

---

Exclude quotes  On

Exclude matches  Off

Exclude bibliography  On

# Inhibition of hepatitis C virus replication by chalepin and pseudane IX isolated from *Ruta angustifolia* leaves

---

## GRADEMARK REPORT

---

FINAL GRADE

**/100**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---

PAGE 9

---