

# Anti-hepatitis C virus compounds obtained from Glycyrrhiza uralensis and other Glycyrrhiza species

*by* Maria Lusida

---

**Submission date:** 24-Jun-2020 01:09AM (UTC+0800)

**Submission ID:** 1348644799

**File name:** ned\_from\_Glycyrrhiza\_uralensis\_and\_other\_Glycyrrhiza\_species.pdf (282.53K)

**Word count:** 5857

**Character count:** 32050

## ORIGINAL ARTICLE

35  
**Anti-hepatitis C virus compounds obtained from  
*Glycyrrhiza uralensis* and other *Glycyrrhiza* species**Myrna Adianti<sup>1,2</sup>, Chie Aoki<sup>1,3</sup>, Mari Komoto<sup>1</sup>, Lin Deng<sup>1</sup>, Ikuo Shoji<sup>1</sup>, Tutik Sri Wahyuni<sup>1,2</sup>,  
Maria Inge Lusida<sup>2</sup>, Soetjipto<sup>2</sup>, Hiroyuki Fuchino<sup>4</sup>, Nobuo Kawahara<sup>4</sup> and Hak Hotta<sup>1</sup>77  
<sup>1</sup>Division of Microbiology, Kobe University Graduate School of Medicine, 7-5-1 Kusunoki-cho, Chuo-ku, Kobe 650-0017, <sup>2</sup>Institute of Tropical Disease, Airlangga University, Jl. Mulyorejo, Surabaya 60115, <sup>3</sup>Japan Science and Technology/Japan International Cooperation Agency Science and Technology Research Partnership for Sustainable Development Laboratory (JST/JICA SATREPS), Faculty of Medicine, University of Indonesia, Jl. Salemba 4, Jakarta 10430, Indonesia and <sup>4</sup>Research Center for Medicinal Plant Resources, National Institute of Biomedical Innovation, 1–2, Hachimandai, Tsukuba City, Ibaraki Prefecture 305-0843, Japan81  
**ABSTRACT**

Development of complementary and/or alternative drugs for treatment of hepatitis C virus (HCV) infection is still much needed from clinical and economic points of view. Antiviral substances obtained from medicinal plants are potentially good targets to study. *Glycyrrhiza uralensis* and *G. glabra* have been commonly used in both traditional and modern medicine. In this study, extracts of *G. uralensis* roots and their components were examined for anti-HCV activity using an HCV cell culture system. It was found that a methanol extract of *G. uralensis* roots and its chloroform fraction possess anti-HCV activity with 50%-inhibitory concentrations (IC<sub>50</sub>) of 20.0 and 8.0 µg/mL, respectively. Through bioactivity-guided purification and structural analysis, glycoumarin, glycyrin, glycyrol and liquiritigenin were isolated and identified as anti-HCV compounds, their IC<sub>50</sub> being 8.8, 7.2, 4.6 and 16.4 µg/mL, respectively. However, glycyrrhizin, the major constituent of *G. uralensis*, and its monoammonium salt, showed only marginal anti-HCV activity. It was also found that licochalcone A and glabridin, known to be exclusive constituents of *G. inflata* and *G. glabra*, respectively, did have anti-HCV activity, their IC<sub>50</sub> being 2.5 and 6.2 µg/mL, respectively. Another chalcone, isoliquiritigenin, also showed anti-HCV activity, with an IC<sub>50</sub> of 3.7 µg/mL. Time-of-addition analysis revealed that all *Glycyrrhiza*-derived anti-HCV compounds tested in this study act at the post-entry step. In conclusion, the present results suggest that glycoumarin, glycyrin, glycyrol and liquiritigenin isolated from *G. uralensis*, as well as isoliquiritigenin, licochalcone A and glabridin, would be good candidates for seed compounds to develop antivirals against HCV.

**Key words** antiviral substance, coumarin, *Glycyrrhiza uralensis*, hepatitis C virus.

34  
Hepatitis C virus is a member of the genus *Hepacivirus* and the family *Flaviviridae*. Based on the heterogeneity of the viral genome, HCV is currently classified into seven genotypes (1–7) and more than 67 subtypes (1a, 1b, 2a, 2b etc.) (1, 2). The viral genome, a single-stranded, positive-sense RNA of 9.6 kb, encodes a25  
polyprotein precursor consisting of about 3000 amino acid residues that is cleaved by host and viral proteases to generate 10 mature proteins, namely core, E1, E2, a putative ion channel p7, and nonstructural proteins NS2, NS3, NS4A, NS4B, NS5A and NS5B (3). Core, E1 and E2 are components of the infectious virus particle together**Correspondence**25  
Hak Hotta, Division of Microbiology, Kobe University Graduate School of Medicine, 7-5-1 Kusunoki-cho, Chuo-ku, Kobe, 650-0017, Japan.  
Tel: +81 78 382 5500; fax: +81 78 382 5519. email: hotta@kobe-u.ac.jp

Received 4 December 2013; revised 20 December 2013; accepted 25 December 2013.

58  
**List of Abbreviations:** CC<sub>50</sub>, 50%-cytotoxic concentrations; E, envelope; Fr, fraction; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; HCV, hepatitis C virus; IC<sub>50</sub>, 50%-inhibitory concentration; ID, internal diameter; SI, selectivity index.

with the viral genome; however, the nonstructural proteins constitute the viral replication complex, where replication of the viral genome takes place. The HCV proteins also play essential roles in the pathological processes associated with HCV infection, such as carcinogenesis and glucose and lipid metabolic disorders (4, 5).

Hepatitis C virus is among the major causative agents of chronic hepatitis, hepatic cirrhosis and hepatocellular carcinoma (5–7). The global prevalence of HCV is >2.5%; thus, about 180 million people are chronically infected with this virus worldwide. A variety of standard treatment regimens using combinations of pegylated interferon, ribavirin and other direct-acting agents, such as HCV-specific inhibitors against NS3 protease and NS5A, have been adopted with considerable success. However, some clinically important issues remain unsolved, such as the emergence of drug-resistant virus and the cost of these drugs. Therefore, development of complementary and/or alternative drugs, especially those from medicinal plants, for treating HCV infection is still much needed from both clinical and economic points of view (8, 9).

*Glycyrrhiza uralensis* and *G. glabra* have been widely used as supplementary treatments in both traditional herbal medicine and modern medicine (10, 11). The radix of *Glycyrrhiza* spp. is commonly known as “gan cao” in Chinese and licorice in English. Bioactive constituents of *Glycyrrhiza* species can be classified into triterpenoids (such as glycyrrhizic acid), coumarins (such as glycyricoumarin, glycyrin and glycyrol), flavones (such as liquiritin and liquiritigenin), chalcones (such as isoliquiritigenin and licochalcone A), isoflavans (such as glabridin), stilbenoids and other miscellaneous compounds (11). Glycyrrhizic acid, also known as glycyrrhizin and considered the principal component of *Glycyrrhiza* spp., is a glycosylated triterpenoid saponin that consists of one molecule of glycyrrhetic acid and two molecules of D-glucuronic acid. Upon hydrolysis, the aglycone, 18 $\beta$ -glycyrrhetic acid (simply called glycyrrhetic acid), and two molecules of D-glucuronic acid are released. Glycyrrhizin and other compounds isolated from *Glycyrrhiza* species reportedly have antiviral activity against a variety of viruses, including HIV, herpes simplex virus, influenza virus, severe acute respiratory syndrome coronavirus, hepatitis viruses and enteroviruses (11–15). As for hepatitis viruses, glycyrrhizin has been used to treat liver diseases, including chronic hepatitis B and C (10). Although glycyrrhizin decreases serum alanine aminotransferase concentrations in HCV-infected patients, it does not significantly reduce amounts of HCV RNA (16, 17). It has been reported that a glycyrrhizin-containing preparation

reduces hepatic steatosis in transgenic mice expressing the full-length HCV polyprotein (18). Recently, anti-HCV activity of glycyrrhizin *in vitro* was reported (19, 20). However, clear evidence for it still appears to be lacking.

In this study, we used an HCV cell culture system to examine a methanol extract and a chloroform sub-fraction of *G. uralensis* and certain isolated compounds, as well as commercially available purified compounds, such as glycyrrhizin and glycyrrhetic acid, for their anti-HCV activity. We report here that glycyricoumarin, glycyrin, glycyrol and liquiritigenin isolated from *G. uralensis* showed anti-HCV activity whereas glycyrrhizin showed only a marginal anti-HCV activity. We also found that some other constituents of *G. uralensis* or of *G. inflata* and *G. glabra*, such as isoliquiritigenin, licochalcone A and glabridin, showed anti-HCV activity.

## MATERIALS AND METHODS

### Cells and viruses

Huh7.5 cells and the plasmid pFL-J6/JFH1 (21) were kindly provided by Dr. C. M. Rice (Rockefeller University, New York, NY, USA). Huh7.5 cells were cultured in Dulbecco's modified Eagle's medium supplemented with FBS (Biowest, Nuaillé, France), non-essential amino acids (Invitrogen, Carlsbad, CA, USA), penicillin (100 IU/mL) and streptomycin (100  $\mu$ g/mL) (Invitrogen) at 37 °C in a 5% CO<sub>2</sub> incubator. A cell culture-adapted strain of HCV genotype 2a (J6/JFH1-P47) was prepared as described previously (22) and used in this study at an MOI of 2.0.

### Extraction, sub-fractionation and purification of *G. uralensis* roots

*G. uralensis* roots were purchased from Tochimoto Tenkaido (Osaka, Japan). A methanol extract of *G. uralensis* roots was prepared and subjected to purification procedures, as described previously (23–26). In brief, *G. uralensis* roots were dried at room temperature and pulverized. They were then extracted with methanol at 50 °C for 6 hr. The extracts were filtered and the filtrates concentrated by using an evaporator at temperatures not exceeding 40 °C. The residues obtained were re-suspended in water and successively partitioned between chloroform and *n*-butanol. The chloroform extract was subjected to recycling preparative HPLC (solvent system, 100% methanol; column, GS-320 + GS-310, 21.5 mm ID  $\times$  1000 mm, flow rate: 5.0 mL/min; detection, UV 210 nm; Condition A) to afford 10 fractions (Fr.1 to Fr.10). Fr.7 was subjected to HPLC

separation (solvent system, acetonitrile–water; column, Imtakt Unison UK-C18C (Kyoto, Japan), 4.6 mm ID × 250 mm; flow rate, 2.0 mL/min; detection, UV 254 nm) to give 12 fractions (Fr.7–1 to 7–12) and glycyrin (2.5 mg; Fr.7–9). Fr.7–6 was purified by recycling HPLC (Condition A) to afford glycycomarin (0.7 mg). Fr.8- to Fr.10 were combined and then re-chromatographed by HPLC (solvent system, acetonitrile–water; column, Imtakt Unison UK-C18C, 4.6 mm ID × 250 mm; flow rate, 2.0 mL/min; detection, UV 254 nm) to give 15 fractions (Fr.8–1 to Fr.8–15). Fr.8–3 was subjected to recycling HPLC (Condition A) to give liquiritigenin (1.2 mg). Fr.8–9 was purified by recycling HPLC (Condition A) to afford glycyrol (1.1 mg). The <sup>1</sup>H- and <sup>13</sup>C-NMR spectra were measured with a Jeol ECA 500 spectrometer (500 MHz; Tokyo, Japan). HPLC was performed on a JASCOLC-2000 plus system (Tokyo, Japan).

### Chemicals

Glycyrrhizic acid (cat. no. 074-03481), glycyrrhizic acid mono-ammonium salt *n*-hydrate (cat. no. 075-02171), glycyrrhetic acid (cat. no. 072-02181) and glabridin (cat. no. 070-04821) were purchased from Wako Pure Chemical Industries (Osaka, Japan). Liquiritin (cat. no. L8045), liquiritigenin (cat. no. 78825) and licochalcone A (cat. no. 68783) were purchased from Sigma–Aldrich (Tokyo, Japan) and isoliquiritigenin (cat. no. I0822) from Tokyo Chemical Industry (Tokyo, Japan). Licorice-saponins G2 (cat. no. P2502) and H2 (cat. no. P2503), and glycyrrhetic acid 3-*O*-glucuronide (cat. no. NH080502) were purchased from Funakoshi (Tokyo, Japan). Glycycomarin, glycyrol, glycyrin and liquiritigenin were isolated from *G. uralensis* extracts in this study, as described above.

### Analysis of anti-HCV activity of plant extracts and purified compounds

Test samples were weighed and dissolved in DMSO to obtain stock solutions at 10 or 30 mg/mL. The stock solutions were stored at –20 °C until used. Huh7.5 cells were seeded in 24-well plates ( $1.6 \times 10^5$  cells/well). HCV was mixed with serial dilutions of the test samples (100, 30, 10, 3 and 1 µg/mL) and inoculated into the cells. After 2 hr, the cells were washed with medium to remove residual virus and further incubated in medium containing the same concentrations of the samples as those during virus inoculation. In time-of-addition experiments, treatment with the samples was performed only during or after virus inoculation in order to assess the mode of action of the samples examined. Culture supernatants were collected 1 and 2 days post-infection

and titrated for virus infectivity, as described below. Virus and cells treated with medium containing 0.1% DMSO served as controls. Percent inhibition of the virus infectivity for each dilution of the samples was calculated by comparison with mock-treated controls and IC<sub>50</sub> determined.

### Virus titration

Virus samples were diluted serially 10-fold in complete medium and inoculated onto Huh7.5 cells seeded on glass coverslips in a 24-well plate. After virus adsorption for 2 hr, the cells were washed with medium to remove residual virus and cultured for 24 hr. The virus-infected cells were stained with an indirect immunofluorescence method as reported previously (27). In brief, the virus-infected cells were washed with PBS, fixed with 4% paraformaldehyde for 15 min and permeabilized with 0.1% Triton X-100 in PBS for 15 min at room temperature. After being washed three times with PBS, the cells were incubated with HCV-infected patient's serum for 1 hr, followed by incubation with FITC-conjugated goat anti-human IgG (MBL, Nagoya, Japan). The cells were counterstained with Hoechst 33342 (Molecular Probes, Eugene, OR, USA) for 5 min and HCV-infected cells were counted under a BZ-9000 fluorescence microscope (Keyence, Osaka, Japan).

### Cytotoxicity assay

The cytotoxicity of the samples was assessed by WST-1 assay as described previously with a slight modification (27). In brief, Huh7.5 cells in 96-well plates were treated with serial dilutions of the samples or 0.1% DMSO as a control for 48 hr. At the end of the treatment, 10 µL of WST-1 reagent (Roche, Mannheim, Germany) was added to each well and the cells cultured for 1 hr. The WST-1 reagent is absorbed by the cells and converted to formazan by mitochondrial dehydrogenases. The amount of formazan, which correlates with the number of living cells, was determined by measuring the absorbance of each well using a microplate reader at 450 and 630 nm. Percent cell viability compared to the control was calculated for each dilution of the samples and CC<sub>50</sub> were determined.

### Immunoblotting

Cells were lysed with an SDS sample buffer, after which equal amounts of protein were subjected to SDS–polyacrylamide gel electrophoresis and transferred onto a polyvinylidene difluoride membrane (Millipore, Bedford, MA, USA), as described previously (28, 29). The membranes were incubated with the respective primary



antibodies. The primary antibodies used were mouse monoclonal antibodies against HCV NS3 and GAPDH (Millipore). Horseradish peroxidase-conjugated goat anti-mouse immunoglobulin (Invitrogen) was used to visualize the respective proteins by means of an enhanced chemiluminescence detection system (ECL; GE Healthcare, Buckinghamshire, UK).

### 22 Real-time quantitative RT-PCR

Real-time quantitative RT-PCR was performed as described previously (28, 29). In brief, total RNA was extracted from the cells using a ReliaPrep RNA cell miniprep system (Promega, Madison, WI, USA) according to the manufacturer's instructions. One microgram of total RNA was reverse transcribed using a GoScript Reverse Transcription system (Promega) with random primers and subjected to real-time quantitative PCR analysis using SYBR Premix Ex Taq (TaKaRa, Kyoto, Japan) in a MicroAmp 96-well reaction plate and an ABI PRISM 7500 system (Applied Biosystems, Foster City, CA, USA). The HCV-specific primers to amplify an NS5A region of the viral genome were 5'-AGACGTATTGAGGTCCATGC-3' (sense) and 5'-CCGCAGCGACGGTGCTGATAG-3' (antisense). Human GAPDH gene expression was measured using primers 5'-GCCATCAATGACCCCTTCATT-3' (sense) and 5'-TCTCGCTCCTGGAAGATGG-3', which served as an internal control.

## RESULTS

### Anti-HCV activity of a methanol extract and sub-fractions of *G. uralensis* roots

First, we examined a crude methanol extract of *G. uralensis* roots for anti-HCV activity using the HCV J6/JFH1-P47 strain and Huh7.5 cells. The IC<sub>50</sub> and CC<sub>50</sub> values of the crude methanol extract were 20.0 and 300 µg/mL, respectively, the selectivity indexes (SI: CC<sub>50</sub>/IC<sub>50</sub>) being 15 (Table 1). We further partitioned the methanol extract using different solvents and found that the anti-HCV activity was concentrated into a chloroform partition, whose IC<sub>50</sub> and CC<sub>50</sub> were 8.0 and 93 µg/mL, respectively, the SI being 11.6.

**Table 1.** Anti-HCV activity (IC<sub>50</sub>), cytotoxicity (CC<sub>50</sub>) and selectivity index (SI) of extracts of *G. uralensis* roots

Sample	IC <sub>50</sub> (µg/mL)	CC <sub>50</sub> (µg/mL)	SI
Methanol extract	20.0	300	15.0
Chloroform partition	8.0	93	11.6
<i>n</i> -butanol partition	17.5	250	14.3
Water partition	95.0	>500	>5.3

**Table 2.** Anti-HCV activity (IC<sub>50</sub>), cytotoxicity (CC<sub>50</sub>) and selectivity index (SI) of recycling HPLC fractions of chloroform partition of *G. uralensis* roots

Sample	IC <sub>50</sub> (µg/mL)	CC <sub>50</sub> (µg/mL)	SI
Fraction 1	>30	>30	na
Fraction 2	>30	>30	na
Fraction 3	>30	>30	na
Fraction 4	>30	>30	na
Fraction 5	20	>30	>1.5
Fraction 6	2.9	65	22
Fraction 7	3.6	40	11
Fraction 8	2.9	40	14
Fraction 9	2.9	>70	>24
Fraction 10	4.9	>70	>14

na, not applicable.

Next, we sub-fractionated the chloroform partition by using recycling HPLC and examined the sub-fractions for anti-HCV activity. We found significant anti-HCV activity with fractions 6–10, IC<sub>50</sub> ranging between 2.9 and 4.9 µg/mL (Table 2). We further tried to purify a major component(s) in fractions 6–10 by using recycling HPLC: we identified glycycomarin (30), glycyrin (31), glycyrol (32) and liquiritigenin (33) by NMR spectrum analysis (data not shown).

### Anti-HCV activity of *G. uralensis*-derived purified compounds and related chemical compounds obtained from other *Glycyrrhiza* species

We then examined the purified compounds obtained from the *G. uralensis* extracts (glycycomarin, glycyrin, glycyrol and liquiritigenin) and other *Glycyrrhiza*-derived chemical compounds, such as glycyrrhizin and its derivatives. We found that glycyrrhizin (glycyrrhizic acid) and its monoammonium salt (glycyrrhizic acid monoammonium), the latter being more water-soluble than the former, possessed only marginal anti-HCV activity, IC<sub>50</sub> values being 180 and 320 µg/mL, respectively (Table 3). The aglycone of glycyrrhizin, glycyrrhetic acid, showed more significant anti-HCV activity than did glycyrrhizin, its IC<sub>50</sub> being 40 µg/mL. It should be noted that their anti-HCV activities were weaker than those of the recycling HPLC fractions 6–10 of the *G. uralensis* extract. The monoglucuronide form of glycyrrhizin, glycyrrhetic acid 3-*O*-glucuronide, did not show significant anti-HCV activity at the concentration tested. These results suggest that another compound(s) in fractions 6–10 was responsible for the anti-HCV activity. Two other saponins tested in this study, licorice-saponins G2 and H2, did not show detectable anti-HCV activity. On the other hand, coumarins, such as

**Table 3.** Anti-HCV activity ( $IC_{50}$ ), cytotoxicity ( $CC_{50}$ ) and selectivity index (SI) of *G. uralensis*- and other *Glycyrrhiza* spp.-derived compounds

Sample	$IC_{50}$ ( $\mu\text{g/mL}$ )	$CC_{50}$ ( $\mu\text{g/mL}$ )	SI
<b>Triterpenoids</b>			
Glycyrrhizin	180	560	3.1
Glycyrrhizic acid monoammonium	320	690	2.2
Glycyrrhetic acid 3-O-glucuronide	>30 <sup>†</sup>	>30	na
Glycyrrhetic acid	40.0	77.4	1.9
Licorice-saponin G2	>30 <sup>†</sup>	>30	na
Licorice-saponin H2	>30 <sup>†</sup>	>30	na
<b>Coumarins</b>			
Glycoumarin (isolated)	8.8	69.0	7.8
Glycyrin (isolated)	7.2	25.0	3.5
Glycyrol (isolated)	4.6	35.5	7.7
<b>Flavonoids</b>			
<b>Flavanones</b>			
Liquiritin	75.0	570	7.6
Liquiritigenin	16.4	125	7.6
<b>Chalcones</b>			
Isoliquiritigenin	3.7	11.0	3.0
Licochalcone A†	2.5	20.0	8.0
<b>Isoflavan</b>			
Glabridin†	6.2	22.7	3.7

<sup>†</sup>Not detected in *G. uralensis* (34). <sup>††</sup>No detectable HCV inhibition at 30  $\mu\text{g/mL}$ . na, not applicable.

glycoumarin, glycyrin and glycyrol, which were isolated in this study (purity >90%) from the *G. uralensis* extracts, showed potent anti-HCV activity,  $IC_{50}$  being 4.6–8.8  $\mu\text{g/mL}$ . Liquiritigenin, which was also isolated from the *G. uralensis* extracts, showed weaker anti-HCV activity than the coumarins. Also, isoliquiritigen, licochalcone A and glabridin exhibited potent anti-HCV activity, their  $IC_{50}$  being 3.7, 2.5 and 6.2  $\mu\text{g/mL}$ , respectively. It should be noted that licochalcone A and glabridin have been reported to be absent in *G. uralensis* but present in *G. inflata* and *G. glabra* (34).

#### Mode-of-action of *G. uralensis*-derived isolated compounds and related compounds obtained from other *Glycyrrhiza* species

To determine whether the anti-HCV effects of the compounds of *Glycyrrhiza* species are exerted on the entry or post-entry stage, we performed the following time-of-addition experiments.

1. HCV was mixed with a test compound and the mixture inoculated into the cells. After virus adsorption for 2 hr, the residual virus and test sample were removed and the cells re-fed with fresh medium

**Table 4.** Mode-of-action analysis of anti-HCV activities of *G. uralensis*- and other *Glycyrrhiza* spp.-derived compounds

Compound	Conc. ( $\text{mg/mL}$ ) <sup>†</sup>	Anti-HCV activity (% inhibition)		
		During <sup>†</sup>	After <sup>†</sup>	During & after <sup>†</sup>
Chloroform partition	30	28.6	100	100
Glycoumarin	20	16.7	100	100
Glycyrin	15	18.4	98.3	99.6
Glycyrol	10	21.3	100	100
Liquiritigenin	30	15.5	90.0	87.2
Isoliquiritigenin	8	14.1	91.0	82.5
Licochalcone A	5	0	94.4	93.8
Glabridin	12	0	91.0	93.8

<sup>†</sup>The concentrations of each compound used were  $2 \times IC_{50}$ . <sup>††</sup>Treatment with the compound was administered only during, only after or both during and after virus inoculation.

without the test sample for 46 hr. This experiment examines the antiviral effect at the entry step.


2. HCV was inoculated into the cells in the absence of a test sample. After virus adsorption for 2 hr, the residual virus was removed and the cells treated with the test samples for 46 hr. This experiment examines the antiviral effect at the post-entry step.

3. As a positive control, HCV mixed with the test sample was inoculated into the cells. After virus adsorption for 2 hr, the residual virus and test sample were removed and the cells treated with the test samples for 46 hr.

We found that all the *Glycyrrhiza* species-derived compounds as well as the chloroform partition of the *G. uralensis* extract exerted their antiviral effects after virus inoculation (Table 4). These results suggest that all the anti-HCV compounds of *Glycyrrhiza* species tested in this study, namely, glycoumarin, glycyrin, glycyrol, liquiritigenin, isoliquiritigenin, licochalcone A and glabridin, as well as the chloroform extract, act primarily at the post-entry step.

#### Inhibition of HCV RNA replication and HCV protein synthesis by glycoumarin, glycyrin, and glycyrol obtained from *G. uralensis*

To further confirm that *G. uralensis*-derived compounds exert their anti-HCV activities at the post-entry step, Huh7.5 cells were inoculated with HCV for 2 hr, followed by treatment with each of compounds for 1–2 days. The results obtained clearly demonstrated that glycoumarin, glycyrin and glycyrol inhibit HCV RNA replication, resulting in decreased HCV protein



**Fig. 3.** Effects of glycy coumarin, glycyrin and glycyrol isolated from *G. uralensis* on HCV RNA replication and protein synthesis. (a) Huh 7.5 cells infected with HCV J6/JFH1 and treated with either glycy coumarin (20  $\mu$ g/mL), glycyrin (15  $\mu$ g/mL), glycyrol (10  $\mu$ g/mL) or left untreated were subjected to western blot analysis using monoclonal antibody against the HCV NS3 protein at 1 and 2 days post-infection. GAPDH served as an internal control to verify equal amounts of sample loading. Signal intensities of NS3 were normalized to the corresponding GAPDH signal. (b) Amounts of HCV RNA in the cells described in (a) were measured by real-time quantitative RT-PCR analysis. These amounts were normalized to GAPDH mRNA expression. Data represent means  $\pm$  SEM of data from two independent experiments. The value for the untreated control at 1 day post-infection is arbitrarily expressed as 1.0. \* $P < 0.001$ , compared with the control. (c) Amounts of HCV infectious particles in the supernatants of the cultures described in (a) and (b) were determined: data for glycy coumarin, glycyrin, glycyrol and the untreated control are shown. Data represent means  $\pm$  SEM of data from two independent experiments. \* $P < 0.05$ ; † $P < 0.01$ , compared with the untreated control; dpi, days post infection.

synthesis as demonstrated by both real-time quantitative RT-PCR and immunoblotting analyses (Fig. 1a, b). We also confirmed that production of HCV infectious particles is inhibited by glycy coumarin, glycyrin and glycyrol at 1 and 2 days post-infection (Fig 1c).

## DISCUSSION

*Glycyrrhiza* species possess a variety of bioactive compounds, such as: (i) triterpenoids, for example,

glycyrrhizin, glycyrrhetic acid 3-O-glucuronide, glycyrrhetic acid and various licorice-saponins; (ii) coumarins, for example, glycy coumarin, glycyrin and glycyrol; (iii) flavanones, for example, liquiritin and liquiritigenin; (iv) chalcones, for example, isoliquiritigenin and licochalcone; (v) isoflavans, for example, glabridin; (vi) stilbenoids, for example, dihydrostilbenes; and other miscellaneous compounds (11). In this connection, flavanones, chalcones and isoflavans are members of flavonoids.



Glycyrrhizin is the major constituent of *Glycyrrhiza* species, representing 1–9% of the total (14). Glycyrrhizin and other compounds isolated from *Glycyrrhiza* species reportedly possess antiviral activity against a variety of viruses (11–15). With regard to HCV, it was recently reported that glycyrrhizin inhibits HCV infection in Huh-7 cells (19, 20). On the other hand, controversial observations have been reported in clinical settings; namely, that glycyrrhizin has no significant effect on amounts of HCV RNA whereas it reduces serum alanine aminotransferase concentrations in HCV-infected patients (16, 17). These apparently contradictory findings might be explained by a membrane-stabilizing effect of glycyrrhizin; clear evidence for antiviral activity of glycyrrhizin is still lacking (10). In the present study, we observed that glycyrrhizin and glycyrrhizic acid monoammonium showed only marginal anti-HCV activity (Table 3). Glycyrrhetic acid showed significant anti-HCV activity. However, their anti-HCV activities were significantly weaker than that of the HPLC fractions obtained from *G. uralensis* extract. On the other hand, we found that certain coumarins, such as glycycomarin, glycyrin and glycyrol, and a flavanone, liquiritigenin, isolated from *G. uralensis*, as well as other compounds contained in *Glycyrrhiza* species, possess potent anti-HCV activities. Mode-of-action analysis revealed that all the *Glycyrrhiza*-derived anti-HCV compounds tested act primarily at the post-entry step. Consistent with our observations, other researchers have also reported that two flavonoids, isoliquiritigenin and glycycomarin, extracted from *Glycyrrhizae radix*, either *G. uralensis* or *G. glabra*, inhibit replication of an HCV subgenomic RNA replicon *in vitro* (35). The IC<sub>50</sub> values of isoliquiritigenin and glycycomarin against HCV subgenomic replicon were 6.2 and 15.5 µg/mL, respectively. On the other hand, to the best of our knowledge, there are no published reports so far regarding anti-HCV activities of glycyrin and glycyrol. It has been reported that glycyrin, which is known to possess peroxisome proliferator-activated receptor-γ ligand-binding activity, shows antibacterial activity against *Streptococcus pyogenes*, *Haemophilus influenzae* and *Moraxella catarrhalis* (36–38). Glycyrol reportedly exerts anti-inflammatory effects by binding to calcineurin to inhibit its activity (39, 40). However, the possible antimicrobial activity of glycyrol has not been reported. Further study is needed to elucidate the detailed mechanism of anti-HCV activities of glycyrin and glycyrol, and also possible antiviral activities against viruses other than HCV.

In conclusion, our results suggest that glycycomarin, glycyrin, glycyrol and liquiritigenin isolated from *G. uralensis*, as well as isoliquiritigenin, licochalcone A

and glabridin, would be good candidates for seed compounds to develop antivirals against HCV.

## 14 ACKNOWLEDGMENTS

The authors are grateful to Dr. C. M. Rice (Rockefeller University, New York, NY, USA) for providing Huh7.5 cells and pFL-J6/JFH1. This study was supported in part by Science and Technology Research Partnerships for Sustainable Development (SATREPS) from Japan Science and Technology Agency (JST) and Japan International Cooperation Agency (JICA). This study was also carried out as part of the Japan Initiative for Global Research Network on Infectious Diseases (J-GRID), Ministry of Education, Culture, Sports, Science and Technology, Japan.

## DISCLOSURE

51 The authors have no conflicts of interest to declare.

## REFERENCES

- Gottwein J.M., Scheel T.K., Jensen T.B., Lademann J.B., Prentoe J.C., Knudsen M.L., Hoegh A.M., Bukh J. (2009) Development and characterization of hepatitis C virus genotype 1–7 cell culture systems: role of CD81 and scavenger receptor class B type I and effect of antiviral drugs. *Hepatology* **49**: 364–77.
- Smith D.B., Bukh J., Kuiken C., Muerhoff A.S., Rice C.M., Stapleton J.T., Simmonds P. (2013) Expanded classification of hepatitis C Virus into 7 genotypes and 67 Subtypes: updated criteria and assignment web resource. *Hepatology* **59**: 318–27.
- Moradpour D., Penin F., Rice C.M. (2007) Replication of hepatitis C virus. *Nat Rev Microbiol* **5**: 453–6.
- Shoji I., Deng L., Hotta H. (2011) Molecular mechanism of hepatitis C virus-induced glucose metabolic disorders. *Front Microbiol* **2**: 278.
- Arzumanyan A., Reis H.M., Feitelson M.A. (2013) Pathogenic mechanisms in HBV- and HCV-associated hepatocellular carcinoma. *Nat Rev Cancer* **13**: 123–35.
- Shepard C.W., Finelli L., Alter M.J. (2005) Global epidemiology of hepatitis C virus infection. *Lancet Infect Dis* **5**: 558–67.
- Mohd Hanafiah K., Groeger J., Flaxman A.D., Wiersma S.T. (2013) Global epidemiology of hepatitis C virus infection: new estimates of age-specific antibody to HCV seroprevalence. *Hepatology* **57**: 1333–42.
- Calland N., Dubuisson J., Rouille Y., Seron K. (2012) Hepatitis C virus and natural compounds: a new antiviral approach? *Viruses* **4**: 2197–217.
- Ploss A., Dubuisson J. (2012) New advances in the molecular biology of hepatitis C virus infection: towards the identification of new treatment targets. *Gut* **61** (Suppl 1): i25–35.
- Stickel F., Schuppan D. (2007) Herbal medicine in the treatment of liver diseases. *Dig Liver Dis* **39**: 293–304.
- Asl M.N., Hosseinzadeh H. (2008) Review of pharmacological effects of *Glycyrrhiza* sp. and its bioactive compounds. *Phytother Res* **22**: 709–24.



12. Pompei R., Flore O., Marccialis M.A., Pani A., Loddo B. (1979) Glycyrrhizic acid inhibits virus growth and inactivates virus particles. *Nature* **281**: 689–90.
13. Cinatl J., Morgenstern B., Bauer G., Chandra P., Rabenau H., Doerr H.W. (2003) Glycyrrhizin, an active component of liquorice roots, and replication of SARS-associated coronavirus. *Lancet* **361**: 2045–6.
14. Fiore C., Eisenhut M., Krause R., Ragazzi E., Pellati D., Armanini D., Bielenberg J. (2008) Antiviral effects of *Glycyrrhiza* species. *Phytother Res* **22**: 141–8.
15. Wang J., Chen X., Wang W., Zhang Y., Yang Z., Jin Y., Ge H.M., Li E., Yang G. (2013) Glycyrrhizic acid as the antiviral component of *Glycyrrhiza uralensis* Fisch. against coxsackievirus A16 and enterovirus 71 of hand foot and mouth disease. *J Ethnopharmacol* **147**: 114–21.
16. van Rossum T.G., Vulto A.G., Hop W.C., Brouwer J.T., Niesters H.G., Schalm S.W. (1999) Intravenous glycyrrhizin for the treatment of chronic hepatitis C: a double-blind, randomized, placebo-controlled phase I/II trial. *J Gastroenterol Hepatol* **14**: 1093–9.
17. Orlent H., Hansen B.E., Willems M., Brouwer J.T., Huber R., Kullak-Ublick G.A., Gerken G., Zeuzem S., Nevens F., Tielmans W.C., Zondervan P.E., Lagging M., Westin J., Schalm S.W. (2006) Biochemical and histological effects of 26 weeks of glycyrrhizin treatment in chronic hepatitis C: a randomized phase II trial. *J Hepatol* **45**: 539–46.
18. Korenaga M., Hidaka I., Nishina S., Sakai A., Shinozaki A., Gondo T., Furutani T., Kawano H., Sakaida I., Hino K. (2011) A glycyrrhizin-containing preparation reduces hepatic steatosis induced by hepatitis C virus protein and iron in mice. *Liver Int* **31**: 552–60.
19. Ashfaq U.A., Masoud M.S., Nawaz Z., Riazuddin S. (2011) Glycyrrhizin as antiviral agent against hepatitis C virus. *J Transl Med* **9**: 112.
20. Matsumoto Y., Matsuura T., Aoyagi H., Matsuda M., Hmwe S.S., Date T., Watanabe N., Watashi K., Suzuki R., Ichinose S., Wake K., Suzuki T., Miyamura T., Wakita T., Aizaki H. (2013) Antiviral activity of glycyrrhizin against hepatitis C virus in vitro. *PLoS ONE* **8**: e68992.
21. Lindenbach B.D., Evans M.J., Syder A.J., Wolk B., Tellinghuisen T.L., Liu C.C., Maruyama T., Hynes R.O., Burton D.R., McKeating J.A., Rice C.M. (2005) Complete replication of hepatitis C virus in cell culture. *Science* **309**: 623–6.
22. Bungyoku Y., Shoji I., Makine T., Adachi T., Hayashida K., Nagano-Fujii M., Ide Y., Deng L., Hotta H. (2009) Efficient production of infectious hepatitis C virus with adaptive mutations in cultured hepatoma cells. *J Gen Virol* **90**: 1681–9.
23. Li S.Y., Fuchino H., Kawahara N., Sekita S., Satake M. (2002) New phenolic constituents from *Smilax bracteata*. *J Nat Prod* **65**: 262–6.
24. Fuchino H., Sekita S., Mori K., Kawahara N., Satake M., Kiuchi F. (2008) A new leishmanicidal saponin from *Brunfelsia grandiflora*. *Chem Pharm Bull* **56**: 93–6.
25. Fuchino H., Kawano M., Mori-Yasumoto K., Sekita S., Satake M., Ishikawa T., Kiuchi F., Kawahara N. (2010) *In vitro* leishmanicidal activity of benzophenanthridine alkaloids from *Bocconia pearcei* and related compounds. *Chem Pharm Bull* **58**: 1047–50.
26. Fuchino H., Daikonya A., Kumagai T., Goda Y., Takahashi Y., Kawahara N. (2013) Two new labdane diterpenes from fresh leaves of *Leonurus japonicus* and their degradation during drying. *Chem Pharm Bull* **61**: 497–503.
27. Deng L., Adachi T., Kitayama K., Bungyoku Y., Kitazawa S., Ishido S., Shoji I., Hotta H. (2008) Hepatitis C virus infection induces apoptosis through a Bax-triggered, mitochondrion-mediated, caspase 3-dependent pathway. *J Virol* **82**: 10375–8.
28. Deng L., Shoji I., Ogawa W., Kaneda S., Soga T., Jiang D.P., Ide Y.H., Hotta H. (2011) Hepatitis C virus infection promotes hepatic gluconeogenesis through an NS5A-mediated, FoxO1-dependent pathway. *J Virol* **85**: 8556–68.
29. Wahyuni T.S., Tumewu L., Permanasari A.A., Apriani E., Adianti M., Rahman A., Widyawaruyanti A., Lusida M.I., Fuad A., Soetjipto D., Nasronudin D., Fuchino H., Kawahara N., Shoji I., Deng L., Aoki C., Hotta H. (2013) Antiviral activities of Indonesian medicinal plants in the East Java region against hepatitis C virus. *Virol J* **10**: 259.
30. Demizu S., Kajiyama K., Takahashi K., Hiraga Y., Yamamoto S., Tamura Y., Okada K., Kinoshita T. (1988) Antioxidant and antimicrobial constituents of licorice: isolation and structure elucidation of a new benzofuran derivative. *Chem Pharm Bull* **36**: 3474–9.
31. Kinoshita T., Saitoh T., Shibata S. (1978) A new 3-aryl coumarin from licorice root. *Chem Pharm Bull* **26**: 135–40.
32. Shiozawa T., Urata S., Kinoshita T., Saitoh T. (1989) Revised structures of glycyrol and isoglycyrol, constituents of the root of *Glycyrrhiza uralensis*. *Chem Pharm Bull* **37**: 2239–40.
33. Chokchaisiri R., Suaisom C., Sriphota S., Chindaduang A., Chuprajob T., Suksamrarn A. (2009) Bioactive flavonoids of the flowers of *Butea monosperma*. *Chem Pharm Bull* **57**: 428–32.
34. Kondo K., Shiba M., Nakamura R., Morota T., Shoyama Y. (2007) Constituent properties of licorices derived from *Glycyrrhiza uralensis*, *G. glabra*, or *G. inflata* identified by genetic information. *Biol Pharm Bull* **30**: 1271–7.
35. Sekine-Osajima Y., Sakamoto N., Nakagawa M., Itsui Y., Tasaka M., Nishimura-Sakurai Y., Chen C.H., Suda G., Mishima K., Onuki Y., Yamamoto M., Maekawa S., Enomoto N., Kanai T., Tsuchiya K., Watanabe M. (2009) Two flavonoids extracts from *Glycyrrhizae radix* inhibit *in vitro* hepatitis C virus replication. *Hepatol Res* **39**: 60–9.
36. Mae T., Kishida H., Nishiyama T., Tsukagawa M., Konishi E., Kuroda M., Mimaki Y., Sashida Y., Takahashi K., Kawada T., Nakagawa K., Kitahara M. (2003) A licorice ethanolic extract with peroxisome proliferator-activated receptor- $\gamma$  ligand-binding activity affects diabetes in KK-Ay mice, abdominal obesity in diet-induced obese C57BL mice and hypertension in spontaneously hypertensive rats. *J Nutr* **133**: 3369–77.
37. Kuroda M., Mimaki Y., Sashida Y., Mae T., Kishida H., Nishiyama T., Tsukagawa M., Konishi E., Takahashi K., Kawada T., Nakagawa K., Kitahara M. (2003) Phenolics with PPAR- $\gamma$  ligand-binding activity obtained from licorice (*Glycyrrhiza uralensis* roots) and ameliorative effects of glycyrin on genetically diabetic KK-A(y) mice. *Bioorg Med Chem Lett* **13**: 4267–72.
38. Tanaka Y., Kikuzaki H., Fukuda S., Nakatani N. (2001) Antibacterial compounds of licorice against upper airway respiratory tract pathogens. *J Nutr Sci Vitaminol* **47**: 270–3.
39. Peng L., Qi Y., Wu H., Wei Q. (2011) Interaction of glycyrol with calcineurin A studied by spectroscopic methods and docking. *IUBMB Life* **63**: 14–20.
40. Shin E.M., Zhou H.Y., Guo L.Y., Kim J.A., Lee S.H., Merfort I., Kang S.S., Kim H.S., Kim S., Kim Y.S. (2008) Anti-inflammatory effects of glycyrol isolated from *Glycyrrhiza uralensis* in LPS-stimulated RAW264.7 macrophages. *Int Immunopharmacol* **8**: 1524–32.

# Anti-hepatitis C virus compounds obtained from Glycyrrhiza uralensis and other Glycyrrhiza species

## ORIGINALITY REPORT

25%

SIMILARITY INDEX

13%

INTERNET SOURCES

24%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 N. K. Mohd-Ismail, L. Deng, S. K. Sukumaran, V. C. Yu, H. Hotta, Y.-J. Tan. "The Hepatitis C Virus Core Protein Contains a BH3 Domain That Regulates Apoptosis through Specific Interaction with Human Mcl-1", *Journal of Virology*, 2009

Publication 1%
- 2 Sri Hartati, Chie Aoki, Muhammad Hanafi, Marissa Angelina, Pratiwi Soedarmono, Hak Hotta. "Antiviral effect of *Archidendron pauciflorum* leaves extract to hepatitis C virus: An *in vitro* study in JFH-1 strain", *Medical Journal of Indonesia*, 2018

Publication 1%
- 3 Yoshinori Tanaka. "Establishment of an indicator cell system for hepatitis C virus", *Microbiology and Immunology*, 04/2010

Publication 1%
- 4 Fuchino, Hiroyuki, Fumiyuki Kiuchi, Azusa Yamanaka, Akihiro Obu, Hiroshi Wada, Kanami

1%

Mori-Yasumoto, Nobuo Kawahara, Diana Flores, Olga Palacios, Setsuko Sekita, and Motoyoshi Satake. "New Leishmanicidal Stilbenes from a Peruvian Folk Medicine, *Lonchocarpus nicou*", CHEMICAL & PHARMACEUTICAL BULLETIN, 2013.

Publication

---

5

Hanane Elayadi, Michael Smietana, Christophe Pannecouque, Pieter Leyssen, Johan Neyts, Jean-Jacques Vasseur, Hassan B. Lazrek. "Straightforward synthesis of triazoloacyclonucleotide phosphonates as potential HCV inhibitors", Bioorganic & Medicinal Chemistry Letters, 2010

Publication

---

6

Mori, K.. "Mechanism of action of ribavirin in a novel hepatitis C virus replication cell system", Virus Research, 201104

Publication

---

7

Pfaender, Stephanie, Richard JP Brown, Thomas Pietschmann, and Eike Steinmann. "Natural reservoirs for homologs of hepatitis C virus", Emerging Microbes & Infections, 2014.

Publication

---

8

Ishihara, Kanako, Mieko Saito, Natsumi Shimokubo, Yasukazu Muramatsu, Shigeki Maetani, and Yutaka Tamura. "Methicillin-

1%

1%

1%

1%



resistant Staphylococcus aureus carriage among veterinary staff and dogs in private veterinary clinics in Hokkaido, Japan : MRSA in veterinary clinics of Japan", Microbiology and Immunology, 2014.

Publication

---

9

T. Li. "Characterization of self-assembled virus-like particles of rat hepatitis E virus generated by recombinant baculoviruses", Journal of General Virology, 08/24/2011

<1%

Publication

---

10

Gian Luca Daino, Aldo Frau, Cinzia Sanna, Daniela Rigano et al. "Identification of Myricetin as an Ebola Virus VP35–Double-Stranded RNA Interaction Inhibitor through a Novel Fluorescence-Based Assay", Biochemistry, 2018

<1%

Publication

---

11

Yasuhiro Sumino. "Growth mechanism of satellite cells in human urethral rhabdosphincter", Neurourology and Urodynamics, 07/2007

<1%

Publication

---

12

Erica Silberstein, Kathleen Mihalik, Laura Uritzky, Ewan P. Plant et al. "Persistent Growth of a Human Plasma-Derived Hepatitis C Virus Genotype 1b Isolate in Cell Culture", PLoS

<1%

13

Fiore, C.. "A history of the therapeutic use of liquorice in Europe", *Journal of Ethnopharmacology*, 20050714

Publication

---

<1%

14

George Koutsoudakis. "Cell Culture Replication of a Genotype 1b Hepatitis C Virus Isolate Cloned from a Patient Who Underwent Liver Transplantation", *PLoS ONE*, 08/24/2011

Publication

---

<1%

15

Elsebai, Mahmoud Fahmi, George Koutsoudakis, Verónica Saludes, Gemma Pérez-Vilaró, Ari Turpeinen, Sampo Mattila, Anna Maria Pirttilä, Fabien Fontaine-Vive, Mohamed Mehiri, Andreas Meyerhans, and Juana Diez. "Pan-genotypic hepatitis C virus inhibition by natural products derived from the wild Egyptian artichoke", *Journal of Virology*, 2015.

Publication

---

<1%

16

Yukihiro Furusawa, Yuka Yamanouchi, Takashi Iizumi, Qing-Li Zhao et al. "Checkpoint kinase 2 is dispensable for regulation of the p53 response but is required for G2/M arrest and cell survival in cells with p53 defects under heat stress", *Apoptosis*, 2017

Publication

<1%

17

A. Kaul, I. Woerz, P. Meuleman, G. Leroux-Roels, R. Bartenschlager. "Cell Culture Adaptation of Hepatitis C Virus and In Vivo Viability of an Adapted Variant", *Journal of Virology*, 2007

Publication

&lt;1%

18

Y.-H. Kou, M.-F. Chang, Y.-M. Wang, T.-M. Hung, S. C. Chang. "Differential Requirements of NS4A for Internal NS3 Cleavage and Polyprotein Processing of Hepatitis C Virus", *Journal of Virology*, 2007

Publication

&lt;1%

19

Changcai Bai, Xiuping Zhou, Lu Han, Yongjie Yu, Nan Li, Ming Zhang, Zhuo Qu, Pengfei Tu. "Two new 18, 19-seco Triterpenoids from *Ilex asprella* (Hook. et Arn.) Champ. ex Benth", *Fitoterapia*, 2018

Publication

&lt;1%

20

[ncbiupdatesg3n3.blogspot.com](http://ncbiupdatesg3n3.blogspot.com)

Internet Source

&lt;1%

21

[www.tandfonline.com](http://www.tandfonline.com)

Internet Source

&lt;1%

22

[hub.hku.hk](http://hub.hku.hk)

Internet Source

&lt;1%

23

[digitalcommons.library.tmc.edu](http://digitalcommons.library.tmc.edu)

Internet Source

&lt;1%



24 Young-Eun Cho, In-Sook Kwun. " (licorice) extracts increase cell proliferation and bone marker enzyme alkaline phosphatase activity in osteoblastic MC3T3-E1 cells ", Journal of Nutrition and Health, 2018  
Publication <1%

---

25 Hiroshi Gomyo, Manabu Shimoyama, Kentaro Minagawa, Kimikazu Yakushijin et al. "Effective Anti-viral Therapy for Hemophagocytic Syndrome Associated with B-cell Lymphoma", Leukemia & Lymphoma, 2009  
Publication <1%

---

26 [www.omicsonline.org](http://www.omicsonline.org)  
Internet Source <1%

---

27 I. Imhof, P. Simmonds. "Development of an Intergenotypic Hepatitis C Virus (HCV) Cell Culture Method To Assess Antiviral Susceptibilities and Resistance Development of HCV NS3 Protease Genes from HCV Genotypes 1 to 6", Journal of Virology, 2010  
Publication <1%

---

28 Jun-Jun Cheng, Jian-Rui Li, Meng-Hao Huang, Lin-Lin Ma et al. "CD36 is a co-receptor for hepatitis C virus E1 protein attachment", Scientific Reports, 2016  
Publication <1%

---

Jingjing Wang, Xiaoqing Chen, Wei Wang,

29 Yating Zhang, Ziying Yang, Yu Jin, Hui Ming Ge, Erguang Li, Guang Yang. "Glycyrrhizic acid as the antiviral component of Glycyrrhiza uralensis Fisch. against coxsackievirus A16 and enterovirus 71 of hand foot and mouth disease", *Journal of Ethnopharmacology*, 2013  
Publication

---

30 [www.biolreprod.org](http://www.biolreprod.org)  
Internet Source

---

31 Hu, Linzhen, Zhenzhen Wang, Jinwen Zhang, Yuanyuan Lu, Kaiping Wang, Yongbo Xue, Yu Zhang, and Yonghui Zhang. "Two New Bioactive  $\alpha$ -Pyrones from *Hypericum japonicum*", *Molecules*, 2016.  
Publication

---

32 Charles E. Carraher, Ngoc Trang Caroline Truong, Michael R. Roner, Alisa Moric, Nancy T. Trang. "Synthesis of organoarsenic, organoantimony, and organobismuth poly(ether esters) from reaction with glycyrrhetic acid and their preliminary activity against pancreatic cancer cell lines", *Journal of the Chinese Advanced Materials Society*, 2013  
Publication

---

33 *Viruses and Human Cancer*, 2014.  
Publication

---

34

Internet Source

&lt;1%

35

[nrid.nii.ac.jp](http://nrid.nii.ac.jp)

Internet Source

&lt;1%

36

[repo.flib.u-fukui.ac.jp](http://repo.flib.u-fukui.ac.jp)

Internet Source

&lt;1%

37

Hanoch Kaphzan. "Co-application of NMDA and dopamine-induced rapid translation of RSK2 in the mature hippocampus", *Journal of Neurochemistry*, 7/20/2007

Publication

&lt;1%

38

Hum Nath Jnawali, Eunjung Lee, Ki-Woong Jeong, Areum Shin, Yong-Seok Heo, Yangmee Kim. " Anti-inflammatory Activity of Rhamnetin and a Model of Its Binding to c-Jun NH - Terminal Kinase 1 and p38 MAPK ", *Journal of Natural Products*, 2014

Publication

&lt;1%

39

Wu, P.. "Supercritical fluid extraction assisted isolation of sesquiterpene lactones with antiproliferative effects from *Centipeda minima*", *Phytochemistry*, 201204

Publication

&lt;1%

40

C. Chapel. "Reduction of the infectivity of hepatitis C virus pseudoparticles by incorporation of misfolded glycoproteins induced

&lt;1%



by glucosidase inhibitors", Journal of General  
Virology, 04/01/2007

Publication

- 
- 41 [www.pubfacts.com](http://www.pubfacts.com) <1 %  
Internet Source
- 
- 42 [ar.scribd.com](http://ar.scribd.com) <1 %  
Internet Source
- 
- 43 [www.jneurosci.org](http://www.jneurosci.org) <1 %  
Internet Source
- 
- 44 Hung-Yu Sun, Chun-Chieh Lin, Pei-Ju Tsai,  
Wei-Jen Tsai et al. "Lipoprotein lipase liberates  
free fatty acids to inhibit HCV infection and  
prevent hepatic lipid accumulation", Cellular  
Microbiology, 2017 <1 %  
Publication
- 
- 45 Simeon, Rudo L., and Zhilei Chen. "A Screen  
for Genetic Suppressor Elements of Hepatitis C  
Virus Identifies a Supercharged Protein Inhibitor  
of Viral Replication", PLoS ONE, 2013. <1 %  
Publication
- 
- 46 Mingorance, L., M. Friesland, M. Coto-Llerena,  
S. Perez-del-Pulgar, L. Boix, J. M. Lopez-Oliva,  
J. Bruix, X. Forns, and P. Gastaminza. <1 %  
"SELECTIVE INHIBITION OF HEPATITIS C  
VIRUS INFECTION BY HYDROXYZINE AND  
BENZTROPIN.", Antimicrobial Agents and

## Chemotherapy, 2014.

Publication

- 
- 47 Nabil M. Abdel-Hamid, Shimaa A. Abass, Ahmed A. Mohamed, Daniah Muneam Hamid. "Herbal management of hepatocellular carcinoma through cutting the pathways of the common risk factors", Biomedicine & Pharmacotherapy, 2018  
Publication <1%
- 
- 48 [www.pnas.org](http://www.pnas.org)  
Internet Source <1%
- 
- 49 Stanley M. Lemon. "Induction and Evasion of Innate Antiviral Responses by Hepatitis C Virus", Journal of Biological Chemistry, 2010  
Publication <1%
- 
- 50 Niki Vassilaki. "The HCV ARFP/F/core+1 protein: Production and functional analysis of an unconventional viral product", IUBMB Life, 07/2009  
Publication <1%
- 
- 51 [obgyn.onlinelibrary.wiley.com](http://obgyn.onlinelibrary.wiley.com)  
Internet Source <1%
- 
- 52 [studentsrepo.um.edu.my](http://studentsrepo.um.edu.my)  
Internet Source <1%
- 
- 53 [www.nrcresearchpress.com](http://www.nrcresearchpress.com)  
Internet Source <1%
-

- 
- 54 Tutik Sri Wahyuni, Adita Ayu permatasari, Tri Widiandani, Achmad Fuad, Aty Widyawaruyanti, Chie Aoki-Utsubo, Hak Hotta. " Antiviral Activities of Genus against Hepatitis C Virus ", Natural Product Communications, 2018  
Publication <1%
- 
- 55 Yao, Qifeng, Jie Wang, Xiaoyong Zhang, Xuhua Nong, Xinya Xu, and Shuhua Qi. "Cytotoxic Polyketides from the Deep-Sea-Derived Fungus Engyodontium album DFFSCS021", Marine Drugs, 2014.  
Publication <1%
- 
- 56 obrnutafaza.hr  
Internet Source <1%
- 
- 57 Shu Zhu. "Survey of Glycyrrhizae Radix resources in Mongolia: chemical assessment of the underground part of Glycyrrhiza uralensis and comparison with Chinese Glycyrrhiza Radix", Journal of Natural Medicines, 04/2009  
Publication <1%
- 
- 58 pubmed.ncbi.nlm.nih.gov  
Internet Source <1%
- 
- 59 "Dietary Components and Immune Function", Springer Science and Business Media LLC, 2010  
Publication <1%
-



60

bioone.org

Internet Source

&lt;1%

61

Stickel, F.. "Herbal medicine in the treatment of liver diseases", Digestive and Liver Disease, 200704

Publication

&lt;1%

62

Jianping Liu. "Medicinal Herbs for Hepatitis C Virus Infection: A Cochrane Hepatobiliary Systematic Review of Randomized Trials", The American Journal of Gastroenterology, 3/2003

Publication

&lt;1%

63

Rajesh Naithani, Loredana Huma, Louis Holland, Deepak Shukla, David McCormick, Rajendra Mehta, Robert Moriarty. "Antiviral Activity of Phytochemicals: A Comprehensive Review", Mini-Reviews in Medicinal Chemistry, 2008

Publication

&lt;1%

64

S.-C. Teng. "Using Both Chemical and Biological Fingerprints for the Quality Study of Estrogenic Licorice (*Glycyrrhiza uralensis*)", Journal of Food Science, 9/2003

Publication

&lt;1%

65

WATANABE, Manabu, Naomi SUMIDA, Koji YANAI, and Takeshi MURAKAMI. "Cloning and Characterization of Saponin Hydrolases from *Aspergillus oryzae* and *Eupenicillium*

&lt;1%

brefeldianum", Bioscience Biotechnology and Biochemistry, 2005.

Publication

66

[krex.k-state.edu](http://krex.k-state.edu)

Internet Source

<1%

67

Abe, K.i.. "Serum-free cell culture system supplemented with lipid-rich albumin for hepatitis C virus (strain O of genotype 1b) replication", Virus Research, 200705

Publication

<1%

68

T. Fukai, T. Nomura. "Fortschritte der Chemie organischer Naturstoffe/Progress in the Chemistry of Organic Natural Products", Springer Science and Business Media LLC, 1998

Publication

<1%

69

Seong Eun Jin, You Kyung Son, Byung-Sun Min, Hyun Ah Jung, Jae Sue Choi. "Anti-inflammatory and antioxidant activities of constituents isolated from Pueraria lobata roots", Archives of Pharmacal Research, 2012

Publication

<1%

70

[tessera.spandidos-publications.com](http://tessera.spandidos-publications.com)

Internet Source

<1%

71

[eprints.nottingham.ac.uk](http://eprints.nottingham.ac.uk)

Internet Source

<1%

72	<a href="http://ddtjournal.com">ddtjournal.com</a> Internet Source	<1%
73	<a href="http://ruor.uottawa.ca">ruor.uottawa.ca</a> Internet Source	<1%
74	<a href="http://link.springer.com">link.springer.com</a> Internet Source	<1%
75	<a href="http://tel.archives-ouvertes.fr">tel.archives-ouvertes.fr</a> Internet Source	<1%
76	"Plant and Human Health, Volume 3", Springer Science and Business Media LLC, 2019 Publication	<1%
77	<a href="http://lppm.ipb.ac.id">lppm.ipb.ac.id</a> Internet Source	<1%
78	Kyoko Mori, Youki Ueda, Yasuo Ariumi, Hiromichi Dansako, Masanori Ikeda, Nobuyuki Kato. "Development of a drug assay system with hepatitis C virus genome derived from a patient with acute hepatitis C", Virus Genes, 2012 Publication	<1%
79	Zhuo Shang, Xiaoming Li, Li Meng, Chunshun Li, Shushan Gao, Caiguo Huang, Bingui Wang. "Chemical profile of the secondary metabolites produced by a deep-sea sediment-derived fungus <i>Penicillium commune</i> SD-118", Chinese Journal of Oceanology and Limnology, 2012 Publication	<1%

---

**80** Fujii, Shunsuke, Osamu Morinaga, Takuhiro Uto, Shuichi Nomura, and Yukihiro Shoyama. "Development of a Monoclonal Antibody-Based Immunochemical Assay for Liquiritin and Its Application to the Quality Control of Licorice Products", *Journal of Agricultural and Food Chemistry*  
Publication <1%

---

**81** Conteduca, Vincenza, Domenico Sansonno, Sabino Russi, Fabio Pavone, and Franco Dammacco. "Therapy of chronic hepatitis C virus infection in the era of direct-acting and host-targeting antiviral agents", *Journal of Infection*, 2013.  
Publication <1%

---

**82** Haagmans, B.L.. "Coronaviruses and their therapy", *Antiviral Research*, 200609  
Publication <1%

---

---

Exclude quotes  On

Exclude matches  Off

Exclude bibliography  On

# Anti-hepatitis C virus compounds obtained from Glycyrrhiza uralensis and other Glycyrrhiza species

---

GRADEMARK REPORT

---

FINAL GRADE

**/100**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---