

# Detection of human herpesvirus-8 antigen in HIV-infected patients in East Java, Indonesia

*by* Maria Lusida

---

**Submission date:** 23-Jun-2020 11:18PM (UTC+0800)

**Submission ID:** 1348604741

**File name:** s-8\_antigen\_in\_HIV-infected\_patients\_in\_East\_Java,\_Indonesia.pdf (73.62K)

**Word count:** 2815

**Character count:** 15392

DETECTION OF HUMAN HERPESVIRUS-8 ANTIGEN IN HIV-INFECTED PATIENTS IN EAST JAVA, INDONESIA

Devi Oktafiani<sup>1\*</sup>, Ni Luh Ayu Megasari<sup>1</sup>, Elsa Fitriana<sup>1</sup>, Nasronudin<sup>2</sup>, Maria Inge Lusida<sup>3,4</sup>, Soetjipto<sup>2,3,5</sup>

<sup>1</sup>Doctoral Degree Program, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia; <sup>2</sup>Universitas Airlangga Hospital, Surabaya, Indonesia; <sup>3</sup>Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia; <sup>4</sup>Department of Medical Microbiology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia; <sup>5</sup>Department of Medical Biochemistry, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia.

\*Corresponding Author's E-mail: [devi.oktafiani-2015@fk.unair.ac.id](mailto:devi.oktafiani-2015@fk.unair.ac.id)

**Article History**

Received: March. 17, 2017

Revised Received: Dec. 13, 2017

Accepted: Dec. 13, 2017

Published Online: June. 18, 2018

37

**Abstract**

**Background:** Co-infection of human herpesvirus-8 (HHV-8) in HIV-positive people might cause Kaposi's sarcoma. Early detection of HHV-8 may prevent the onset of clinical manifestations. In Indonesia, detection of HHV-8 antigen in HIV-positive patients has yet to be reported. The intention of this research was to examine the presence of HHV-8 antigen in HIV-positive patients in East Java, Indonesia.

**Material and Methods:** 103 serum samples were collected from HIV-positive patients in Surabaya and Tulungagung, East Java, Indonesia. Serums were then tested for the presence of HHV-8 antigen by using sandwich ELISA.

**Results:** Human Herpesvirus-8 antigen was detected in 15 samples (14.5%). The presence of HHV-8 infection in HIV-positive patients did not present differently in males and females and among different age groups. Human immunodeficiency virus-positive serum samples were collected from 23 homosexual men, 25 intravenous drug users (IVDUs) and 52 heterosexuals. In the male homosexual group, HHV-8 antigen was detected in 21.7% (5/23) of the samples, while in the intravenous drug user group (IVDUs), 16% (4/25) of the samples were found to have HHV-8 antigen.

**Conclusion:** This research found the presence of HHV-8 antigen in 14.5% of patients in East Java, Indonesia. It is recommended that patients with a positive result should receive further examination to detect any clinical manifestations related to HHV-8 infection, especially in the form of Kaposi's sarcoma lesions, so that the illness can be appropriately managed.

**Key Words:** HHV-8 antigen, HIV-positive, East Java, Indonesia.

**Introduction**

The human immunodeficiency virus (HIV) primarily attacks T helper (CD4+) lymphocytes. The human immunodeficiency virus (HIV) infection causes the immune system to weaken, so that the host becomes more susceptible to various infections and malignancies (Grant et al., 2005). One common co-infection found in people living with HIV/AIDS is a herpesvirus infection, including herpes simplex virus types 1 and 2 (HSV-1, HSV-2), varicella zoster virus, cytomegalovirus (CMV), Epstein-Barr virus (EBV), herpesviruses 6 and 7, and human herpesvirus 8 (HHV-8 or Kaposi sarcoma-associated herpesvirus [KSHV]) (Brooks et al., 2013). HHV-8 infection in HIV/AIDS patients may cause Kaposi's Sarcoma (Dourminishev et al., 2003). Kaposi's sarcoma is a tumor developed from cells on lymph nodes or in blood vessels, and can also develop in other parts of the body (Dourminishev et al., 2003).

HHV-8 or KSHV, the only known human herpesvirus (rhadinovirus), is the most recently discovered tumor virus (Viejo-Borbolla et al., 2003). HHV-8 has a diameter of 140 nm and a genome of between approximately 125 kb (Burbelo et al., 2015), 165 kb (Brooks et al., 2013) and 170 kb (Russo et al., 1996). The virus has a double stranded linear DNA and icosahedral capsid envelope (Viejo-Borbolla et al., 2003). It is covered by a tegument containing protein, and closed during

budding of the cell. The membrane is derived from outer envelope of lipid membrane from various host and specific virus glycoprotein (Swaminathan, 2003).

Kaposi's sarcoma is more prevalent in immunosuppressed patients than healthy people. It often occurs in people with Jewish, Mediterranean, African, and Middle Eastern origins. Sero-epidemiological evidence of HHV-8 infection can be used to reflect the epidemiology of Kaposi's sarcoma (Grulich et al., 2001). Ablashi and colleagues' research reports that in 1999, seropositive prevalence of HHV-8 infection in general population of USA, Northern Europe, and Asia was only around zero to five percent; this was also the case in a seroepidemiological study in Malaysia, Hong Kong, and Sri Lanka where the seroprevalence of HHV-8 was found to be around 3,7%. However, a higher prevalence was found in Southern Europe, around five to ten percent. (Ablashi et al., 1999). There has been no data regarding viremia detection using ELISA method and no epidemiological reporting on the prevalence of HHV-8 infection in Indonesia. The aim of this study was to correct this omission with an examination of the presence of HHV-8 antigen in HIV-positive patients in East Java, Indonesia.

19

## Materials and Methods

This study was approved by the Ethics Committee of Universitas Airlangga Hospital. A hundred and three serum samples were collected from HIV-positive patients in Universitas Airlangga Hospital, Surabaya and Dokter Iskak Tulungagung General Hospital, East Java, Indonesia. The inclusions criteria are positive HIV infected with naïve or treated with antiretroviral, all age (in this case 7-5 years old, and the mean age is 37 years old). Exclusion criteria in this study were patients who did not agree to follow the study. Informed consent was obtained from each patient prior to sample collection. Patients were also asked to complete a questionnaire to provide general information and the possible route of transmission. The patients were then subjected to an HHV-8 antigen test using sandwich ELISA (Abbexa, Cambridge, UK), according to manufacturer instructions. Data obtained were analyzed using statistical software (SPSS version 21.0).

## Results

HIV-positive patients involved in this study comprised of 64 males and 39 females, with a mean age of 37 years (range 7–64 years). This research showed that 15 of 103 samples (14.56%) were found to be HHV-8 antigen positive. In regards to patients' gender, ten men (15.6%) and five women (12.8%) tested positive. The presence of HHV-8 antigen did not show any significant differences in age ( $p=0.05$ ) and gender ( $p=0.78$ ). HHV-8 infection in homosexual men was 21.7%, while 16% intravenous drug users (IVDUs) were positive.

18

40

**Table 1.** Determinants of HHV-8 infection in HIV positive patients

Characteristic	N	HHV-8 positive N (%)	P value p<0,05, S (Significant), NS (Not Significant)
<b>Age group (years)</b>			
A: ≤ 30	31	4 (12,9%)	p= 0,5 (NS)
B: 31-40	35	7 (20%)	
C: >40	37	4 (10,8%)	
<b>Sex</b>			
Male	64	10 (15,6%)	p= 0,78 (NS)
Female	39	5 (12,8%)	
<b>Sexual orientation</b>			
Homo/bisexual	25	4 (16%)	p= 0,5 (NS)
Heterosexual	78	11 (14,1%)	
<b>Marital status</b>			
Married	68	10 (15,6%)	p=0,6 (NS)
Not married	35	5 (12,8%)	

## Discussion

The results of the study conducted in East Java, Indonesia are an interesting comparison with other studies around the world. It must be reiterated here that the participants in the study were not screened for Kaposi's sarcoma and the ELISA method was not used to collect the data regarding HHV-8 viremia. While the presence of HHV-8 antigen was detected in 14.56% of HIV-positive patients in East Java, Indonesia, previous studies in Xinjiang, China (Zhang et al., 2012), Nigeria (Ogoina et al., 2011) and Cameroon (Mbondji-Wonje et al., 2013) showed a greater prevalence, that is

31.2%, 62% and 70% respectively. Human herpesvirus 8 antigen was detected in 21.7% HIV-positive homosexual men. A study in Thailand reported the presence of HHV-8 lytic antigen in 12% HIV-positive homosexual men, 16% heterosexual men, and 9% IVDUs (Ayuthaya et al., 2002). HHV-8 without KS in serum samples are detected 4% and AIDS-KS are 42.8% in Brazil (Machado et al., 2016). Detection of herpes virus DNA in lymphocytes could possibly represent latent infection, while detectable DNA in serum or plasma is usually associated with disease (Polstra et al., 2003). In HIV-positive patients without Kaposi Sarcoma, 4% were positive using PCR method. These patients certainly present a higher risk of developing AIDS-KS (Machado et al., 2016).

Kaposi's sarcoma was the most common neoplasm found in patients infected with HIV. Kaposi's sarcoma is a tumor of blood vessels, which appears as lesions on the skin, and then spreads to other organs. It is noted here that there has been a decrease in the incidence of Kaposi's with other opportunistic illnesses, since the introduction of highly active antiretroviral therapy (HAART) (Jacobson et al., 1999). This downward trend is most likely due to secondary effects of antiretroviral therapy on the immune system (Cattelan et al., 2001 dan De Milito et al., 1999) rather than direct effect to HHV-8 (Kedes dan Ganem, 1997).. A study utilizing polymerase chain reaction (PCR) demonstrated that HHV-8 was detected in more than 90% of Kaposi's sarcoma lesions (Mendez et al., 1998). Prospective studies indicated that the detection of HHV-8 could predict the development of Kaposi's sarcoma in asymptomatic subjects (Whitby et al., 1995 dan Moore et al., 1996). Several studies using different serological tests found that almost 100% patients with different clinical forms of Kaposi's sarcoma had a high titer of the HHV-8 antibody, which had been detected before the appearance of clinical lesions of Kaposi's sarcoma (Whitby et al., 1995; Gao et al., 1996 dan Moore et al., 1996). Current evidence suggests that HHV-8 infection is a prerequisite in the pathogenesis of Kaposi's sarcoma, but factors relating to the host immuno-deficiency may greatly increase the incidence of Kaposi's sarcoma among subjects infected with HHV-8.

The presence of HHV-8 infection in HIV-positive patients is possibly related to the risk of developing opportunistic diseases, including Kaposi's sarcoma (classic, AIDS-related KS, endemic and iatrogenic) and other proliferative diseases, such as primary effusion lymphoma and multicentric Castleman disease. Among immuno-competent people, only a small proportion of HHV-8 infections develop into Kaposi sarcoma. Various tests have detected HHV-8 antibodies in the majority of asymptomatic individuals with increased tumor risk, including HIV-positive homosexual men (Martin et al., 1998), adults from Italy and various regions of sub-Saharan Africa (Sitas et al., 1999; Calabro et al., 1998), and some low risk people, such as blood donors from the UK, USA, and Jamaica (Simpson et al., 1996; Engels et al., 1999).

## Conclusion

The presence of HHV-8 antigen was found in 14.5% of the HIV-positive patients who participated in this study which is the first of its kind in Indonesia. Based on these results found in patients in East Java, further investigations should be conducted on patients living with HIV. This would assist with early detection and subsequent management of any clinical manifestations related to HHV-8 infection, especially in the form of Kaposi's sarcoma lesion.

**Conflict of Interest:** The authors declare no conflict of interest.

**Acknowledgements:** Thanks to LSM Mahameru, KDS Couple Community, and all the colleagues who have helped in this study.

## References

1. Ablashi, D., Chatlynne, L., Cooper, H., Thomas, D., Yadav, M., Norhanom, A.W., Chandana, A.K., Churdboonchart, V., Kulpradist, S.A., Patnaik, M., Liegmann, K., Masood, R., Reitz, M., Cleghorn, F., Manns, A., Levine, P.H., Rabkin, C., Biggar, R., Jensen, F., Gill, P., Jack, N., Edwards, J., Whitman, J., Boshoff, C (1999). Seroprevalence of human herpesvirus-8 (HHV-8) in countries of Southeast Asia compared to the USA, the Caribbean and Africa. *British Journal of Cancer*. 81: 893–897.
2. Ayuthaya, P.I.N., Katano, H., Inagi, R., Auwanit, W., Sata, T., Kurata, T., and Yamanishi, K. (2002). The seroprevalence of human herpesvirus 8 infection in the Thai population. *Southeast Asian Journal Tropical Medicine Public Health*, 33 (2): 297-305
3. Brooks, G.F., Carol, K.C., Butel, J.S., Morse, S.A., Mietzner, T.A. (2013). *Jawetz, Melnick, & Adelberg's Medical Microbiology: Twenty-six Edition* ed. USA: McGraw Hill, p.467-488
4. Calabro, M.L., Sheldon, J., Favero, A., Simpson, G.R., Fiore, R., Gomes, E., Angarano, G., Chieco-Bianchi, L. and Schulz, T.F. (1998). Seroprevalence of Kaposi's sarcoma-associated herpesvirus/human herpesvirus 8 in several regions of Italy. *Journal of Human Virology*, 1: 207–213

5. Cattelan, A., Calabro, M., Gasperini, P., Aversa, S., Zanchetta, M., Meneghetti, F., De Rossi, A., Chieco-Bianchi, L. (2001). Acquired immunodeficiency syndrome-related Kaposi's sarcoma regression after highly active antiretroviral therapy: biologic correlates of clinical outcome. *Journal of the National Cancer Institute Monographs*. 28:44–49.
6. De Milito, A., Catucci, M., Venturi, G., Romano, L., Incandela, L., Valensin, P.E., Zazzi, M. (1999). Antiretroviral therapy with protease inhibitors in human immunodeficiency virus type 1- and human herpesvirus 8-coinfected patients. *Journal of Medical Virology*. 1999;57:140–144.
7. Engels, EA, Eastman, H., Ablashi, DV, Wilks, RJ, Braham, J, and Manns, A. (1999). Risk of transfusion-associated transmission of human herpesvirus 8. *Journal of the National Cancer Institute*. 91, 1773–1775
8. Gao, SJ, Kingsley, L., Li, M., Zheng, W., Parravicini, C., Ziegler, J., Newton, R., Rinaldo, C.R., Saah, A., Phair J, Detels, R, Chang, Y, Moore, P.S. (1996). KSHV antibodies among Americans, Italians and Ugandans with and without Kaposi's sarcoma. *Nature Medicine* 2:925–928
9. Grulich, A., Rawlinson, W.D., Cunningham, P. (2001). Human herpesvirus 8: a newly described sexually transmissible infection. *Venerology* 14, 174-180
10. Jacobson, L.P., Yamashita, T.E., Detels, R., Margolick, J.B., Chmiel, J.S., Kingsley, L.A., Melnick, S., Mulnoz, A. (1999). Impact of potent antiretroviral therapy on the incidence of Kaposi's sarcoma and non-Hodgkin's lymphomas among HIV-1 infected individuals. Multicenter AIDS Cohort Study. *Journal of Acquire Immune Deficiency Syndrome*;21(Suppl 1):S34–41.
11. Kedes, D.H. and Ganem, D. (1997). Sensitivity of Kaposi's sarcoma-associated herpesvirus replication to antiviral drugs. Implications for potential therapy. *The Journal of Clinical Investigation*; 99:2082–2086.
12. Machado, I., Farias, K., Pereira, M., Freitas, P., Fonseca, B. (2016). Human herpesvirus 8 (HHV-8) detected by nested polymerase chain reaction (PCR) in HIV patients with or without Kaposi's sarcoma. An analytic cross-sectional study. *Sao Paulo Medical Journal*. 2016; 134(3):187-192
13. Martin, J.N., Ganem, D.E., Osmond, D.H., Page-Shafer, KA, Macrae, D. and Kedes, D.H. (1998). Sexual transmission and the natural history of human herpesvirus 8 infection. *The New England Journal of Medicine*; 338, 948–954
14. Mbondji-Wonje, C., Ragupathy, V., Lee, S., Wood, O., Awazi, B., and Hewlett, I.K. (2013). Seroprevalence of Human Herpesvirus-8 in HIV-1 Infected and Uninfected Individuals in Cameroon. *Viruses* 2013, 5, 2253-2259
15. Mendez, J.C., Procop, G.W., Espy, M.J., Paya, C.V., Smith, T.F. (1998). Detection and semiquantitative analysis of human herpesvirus 8 DNA in specimens from patients with Kaposi's sarcoma. *Journal of Clinical Microbiology*; 36:2220–2222
16. Moore, P.S., Kingsley, L.A., Holmberg, S.D., Spira, T., Gupta, P., Hoover, D.R., Parry, J.P., Conley, L.J., Jaffe, H.W., Chang, Y. (1996). Kaposi's sarcoma-associated herpesvirus infection prior to onset of Kaposi's sarcoma. *AIDS* 10:175–180
17. Ogoina, D., Onyemelukwe, G., Musa, B.O., Babadoko, A. (2011). Seroprevalence and determinants of human herpes virus 8 infection in adult Nigerians with and without HIV-1 infection. *African Health Sciences* 2011; 11(2): 158 – 162
18. Polstra, A.M., Van Den Burg R., Goudsmit, J., Cornelissen, M. (2003). Human herpesvirus 8 load in matched serum and plasma samples of patients with AIDS-associated Kaposi's sarcoma. *Journal of Clinical Microbiology*. 2003;41(12):5488-5491.
19. Russo, J.J., Bohenzky, R.A., Chien, M-C., Chen, J., Yan, M., Maddalena, D., Parry, J.P., Peruzzi, D., Edelman, I.S., Chang, Y., and Moore, P.S. (1996). Nucleotide sequence of the Kaposi sarcoma-associated herpesvirus (HHV8). *Proceeding of the National Academy of Sciences of the United States of America*. 93 (25), 14862–14867.
20. Simpson, G.R., Schulz, T.F., Whitby, D., Cook, P.M., Boshoff, C., Rainbow, L., Howard, M.R., Gao, Shou-Jiang., Bohenzky, R.A., Simmonds, P., Lee, C., Ruitter, A., Hatzakis, A., Tedder, R.S., Weller, I., Weiss, R.A., Moore, S. (1996). Prevalence of Kaposi's sarcoma associated herpesvirus infection measured by antibodies to recombinant capsid protein and latent immunofluorescence antigen. *Lancet*, 349, 1133–1138
21. Sitas, F., Carrara, H., Beral, V., Newton, R., Reeves, G., Bull, D., Jentsch, U., Pacella-Norman, R., Bourbouliia, D., Whitby, D., Boshoff, C. and Weiss, R. (1999). Antibodies against human herpesvirus 8 in black South African patients with cancer. *The New England Journal of Medicine*; 340, 1863–1871
22. Swaminathan, S. (2003). Molecular biology of Epstein-Barr virus and Kaposi's sarcoma-associated herpesvirus. *Seminars in Hematology*; 40, 107–115.
23. Viejo-Borbolla, A., Ottinger, M., Schulz, T.F. (2003). Human herpesvirus 8: Biology and role in the pathogenesis of Kaposi's sarcoma and other AIDS-related malignancies. *Current HIV/AIDS Reports*. 5, 169–175.
24. Whitby, D., Howard, M.R., Tenant-Flowers, M., Brink, N.S., Copas, A., Boshoff, C., Hatzioannou, T., Suggett, F.E., Aldam, D.M., Denton, A.S. (1995). Detection of Kaposi sarcoma associated herpesvirus in peripheral blood of HIV-infected individuals and progression to Kaposi's sarcoma. *Lancet* 346:799–802
25. Zhang, T., Shao, X., Chen, Y., Minhas, V., Wood, C., and He, N. (2012). Human Herpesvirus 8 Seroprevalence, China, *Virology Papers*. Paper 238.

# Detection of human herpesvirus-8 antigen in HIV-infected patients in East Java, Indonesia

---

## ORIGINALITY REPORT

---

25%

SIMILARITY INDEX

14%

INTERNET SOURCES

24%

PUBLICATIONS

1%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

- 1** Y. Sergerie, Y. Abed, J. Roy, G. Boivin. "Comparative Evaluation of Three Serological Methods for Detection of Human Herpesvirus 8-Specific Antibodies in Canadian Allogeneic Stem Cell Transplant Recipients", *Journal of Clinical Microbiology*, 2004 1%

Publication

---
- 2** Harold W. Jaffe. "Human Herpesvirus 8 and Kaposi's Sarcoma — Some Answers, More Questions", *New England Journal of Medicine*, 06/17/1999 1%

Publication

---
- 3** Zahra Mozaheb. "Chapter 8 Epidemiology of Lymphoid Malignancy in Asia", IntechOpen, 2012 1%

Publication

---
- 4** Ronaldo B. FREITAS, Maria Rute FREITAS, Alexandre C. LINHARES. "Prevalence of human herpesvirus 8 antibodies in the population of Belém, Pará, Brazil", *Revista do Instituto de* 1%

5

N W S Davies. "Factors influencing PCR detection of viruses in cerebrospinal fluid of patients with suspected CNS infections", Journal of Neurology, Neurosurgery & Psychiatry, 2005

Publication

---

1%

6

M Amin, A Bakhtiar, M Subarjo, EB Aksono, P Widiyanti, K Shimizu, Y Mori. "Screening for Middle East respiratory syndrome coronavirus among febrile Indonesian Hajj pilgrims: A study on 28,197 returning pilgrims", Journal of Infection Prevention, 2018

Publication

---

1%

7

"OS08: Critical Care Medicine 1 : Oral Sessions", Respirology, 2013.

Publication

---

1%

8

Maduja V.M. Divarathna, Rukshan A.M. Rafeek, Faseeha Noordeen. "A review on epidemiology and impact of human metapneumovirus infections in children using TIAB search strategy on PubMed and PubMed Central articles", Reviews in Medical Virology, 2019

Publication

---

1%

9

R Newton. "Review of the distribution of Kaposi's sarcoma-associated herpesvirus (KSHV) in Africa in relation to the incidence of

1%

Kaposi's sarcoma", British Journal of Cancer,  
01/13/2003

Publication

---

10

Gerhard Sissolak. "AIDS-related Kaposi's sarcoma: epidemiological, diagnostic, treatment and control aspects in sub-Saharan Africa", Tropical Medicine and International Health, 10/2005

Publication

---

11

Monica Gandhi, Ruth M Greenblatt. "Human herpesvirus 8, Kaposi's sarcoma, and associated conditions", Clinics in Laboratory Medicine, 2002

Publication

---

12

[iedea.org](http://iedea.org)  
Internet Source

1%

---

13

[www.phaaustralia.com](http://www.phaaustralia.com)  
Internet Source

1%

---

14

[epdf.tips](http://epdf.tips)  
Internet Source

1%

---

15

Souza, V.A.U.F., L.M. Sumita, W. Freire, H.K. Sato, J.L. Grandi, L.C. Pierrotti, M.C. Nascimento, and C.S. Pannuti. "Prevalence of antibodies to human herpesvirus-8 in populations with and without risk for infection in São Paulo State", Brazilian Journal of Medical

1%



## and Biological Research, 2004.

Publication

16

Abel Viejo-Borbolla. "Kaposi's Sarcoma-Associated Herpesvirus (Human Herpesvirus 8)", Principles and Practice of Clinical Virology, 04/23/2004

Publication

1%

17

[www.jove.com](http://www.jove.com)

Internet Source

1%

18

[www.infectagentscancer.com](http://www.infectagentscancer.com)

Internet Source

1%

19

Lígia Camera Pierrotti, Laura Masami Sumita, Wilton Santos Freire, Hélio Hehl Caiaffa Filho, Vanda Akico Ueda Fick de Souza. "Detection of human herpesvirus 8 DNA and antibodies to latent nuclear and lytic-phase antigens in serial samples from aids patients with Kaposi's sarcoma", Journal of Clinical Virology, 2000

Publication

1%

20

[www.frontiersin.org](http://www.frontiersin.org)

Internet Source

1%

21

[www.aidsreviews.com](http://www.aidsreviews.com)

Internet Source

1%

22

[smjournals.com](http://smjournals.com)

Internet Source

1%

23	<a href="http://content.nejm.org">content.nejm.org</a> Internet Source	<1%
24	<a href="http://www.iarc.fr">www.iarc.fr</a> Internet Source	<1%
25	K. E. Foreman. "Propagation of a Human Herpesvirus from AIDS-Associated Kaposi's Sarcoma", <i>New England Journal of Medicine</i> , 1/16/1997 Publication	<1%
26	Kai U. Chow, Paris S. Mitrou, Kerstin Geduldig, Eilke B. Helm, Dieter Hoelzer, H. Reinhard Brodt. "Changing Incidence and Survival in Patients with AIDS-Related Non-Hodgkin's Lymphomas in the Era of Highly Active Antiretroviral Therapy (HAART)", <i>Leukemia &amp; Lymphoma</i> , 2009 Publication	<1%
27	Gutierrez, F.. "Occult lymphadenopathic Kaposi's sarcoma associated with severe pulmonary hypertension: A clinical hint about the potential role of HHV-8 in HIV-related pulmonary hypertension?", <i>Journal of Clinical Virology</i> , 200610 Publication	<1%
28	<a href="http://mdpi.com">mdpi.com</a> Internet Source	<1%

29 Alicia C. McDonald. "Human herpesvirus 8 seroprevalence among Tobago women and the role of sexual lifestyle behavior", Journal of Medical Virology, 02/2009 <1%

Publication

---

30 [medcraveonline.com](http://medcraveonline.com) <1%

Internet Source

---

31 [www.nature.com](http://www.nature.com) <1%

Internet Source

---

32 Weinstein, Jessica E., and Carol L. Karp. "Ocular surface neoplasias and human immunodeficiency virus infection :", Current Opinion in Infectious Diseases, 2013. <1%

Publication

---

33 [preview-infectagentscancer.biomedcentral.com](http://preview-infectagentscancer.biomedcentral.com) <1%

Internet Source

---

34 [worldwidescience.org](http://worldwidescience.org) <1%

Internet Source

---

35 Gisele Zandman-Goddard, Yehuda Shoenfeld, Gisele Zandman-Goddard, Yehuda Shoenfeld. "Infections and SLE", Autoimmunity, 2009 <1%

Publication

---

36 [intl-vir.sgmjournals.org](http://intl-vir.sgmjournals.org) <1%

Internet Source

---

37 [www.oalib.com](http://www.oalib.com)

<1%

38

[theses.gla.ac.uk](https://theses.gla.ac.uk)

Internet Source

<1%

39

Michael J. Cannon. "Blood-Borne and Sexual Transmission of Human Herpesvirus 8 in Women with or at Risk for Human Immunodeficiency Virus Infection", *New England Journal of Medicine*, 03/2001

Publication

<1%

40

Mario Luppi. "HHV-8 Infection in the Transplantation Setting: A Concern Only for Solid Organ Transplant Patients?", *Leukemia & Lymphoma*, 1/1/2002

Publication

<1%

41

Adina Cohen, Dana G. Wolf, Emma Guttman-Yassky, Ronit Sarid. "KAPOSI'S SARCOMA—ASSOCIATED HERPESVIRUS: Clinical, Diagnostic, and Epidemiological Aspects", *Critical Reviews in Clinical Laboratory Sciences*, 2008

Publication

<1%

42

K. K. Uldall. "Adherence in people living with HIV/AIDS, mental illness, and chemical dependency: a review of the literature", *AIDS Care*, 1/1/2004

Publication

<1%

43

S TALBOT. "Kaposi's Sarcoma and Human Herpesvirus 8", HIV and the New Viruses, 1999

Publication

<1%

44

Vanda Akico Ueda Fick de Souza. "Seroreactivity to Kaposi's sarcoma-associated herpesvirus (human herpesvirus 8) latent nuclear antigen in AIDS-associated Kaposi's sarcoma patients depends on CD4+ T-cell count", Journal of Medical Virology, 10/2007

Publication

<1%

45

"Viral Infections of Humans", Springer Science and Business Media LLC, 2014

Publication

<1%

46

"DNA Tumor Viruses", Springer Science and Business Media LLC, 2009

Publication

<1%

47

Frank J. Palella. "Declining Morbidity and Mortality among Patients with Advanced Human Immunodeficiency Virus Infection", New England Journal of Medicine, 03/26/1998

Publication

<1%

48

S. V. Nuvor, H. Katano, W. K. Ampofo, J. S. Barnor, T. Sata. "Higher Prevalence of Antibodies to Human Herpesvirus 8 in HIV-Infected Individuals than in the General Population in Ghana, West Africa", European Journal of Clinical Microbiology and Infections

<1%

# Diseases, 2001

Publication

---

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On

# Detection of human herpesvirus-8 antigen in HIV-infected patients in East Java, Indonesia

---

## GRADEMARK REPORT

---

FINAL GRADE

**/100**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---