

Analysis of genetic and serology of hepatitis A virus infection during and after outbreak in two junior high schools in Surabaya, Indonesia

by Juniastuti Juniastuti

Submission date: 26-Feb-2021 03:57PM (UTC+0800)

Submission ID: 1518677303

File name: Analysis_of_genetic.pdf (3.8M)

Word count: 5703

Character count: 31882

Analysis of genetic and serology of hepatitis A virus infection during and after outbreak in two junior high schools in Surabaya, Indonesia

Juniastuti^{1,2,3} | Dedy Wahyuddin² | Nihayatussa'adah³ | Mochamad Amin³ |
Laura Navika Yamani^{3,4} | Takako Utsumi^{3,5} | Florentina Sustini⁶ | Maria Inge Lusida^{1,3}

¹Department of Microbiology, School of Medicine, Universitas Airlangga, Surabaya, East Java, Indonesia

²Master Program of Tropical Medicine, School of Medicine, Universitas Airlangga, Surabaya, East Java, Indonesia

³Institute of Tropical Disease, Universitas Airlangga, Campus C Universitas Airlangga Jalan Mulyorejo, Surabaya, East Java, Indonesia

⁴Department of Epidemiology, Faculty of Public Health, Universitas Airlangga, Campus C Universitas Airlangga Jalan Mulyorejo, Surabaya, East Java, Indonesia

⁵Center for Infectious Diseases, Kobe University Graduate School of Medicine, Kobe, Hyogo, Japan

⁶Department of Public Health, School of Medicine, Universitas Airlangga, Surabaya, East Java, Indonesia

Correspondence

Juniastuti, Department of Microbiology, School of Medicine, Universitas Airlangga, Jalan Mayjen Prof Dr Moestopo 47, Surabaya 60131, East Java, Indonesia.
Email: junsf@fk.unair.ac.id

Abstract

Outbreaks of hepatitis A have occurred in some cities in Indonesia. In Surabaya, the capital city of East Java province, Indonesia, hepatitis A outbreaks have been reported since 2013, with a marked increase in the number of cases in 2015. The aim of the present study was to analyze the genetic and serology of acute symptomatic cases (early infection) during a hepatitis A outbreak and asymptomatic cases after the outbreak in two junior high schools in Surabaya in 2015 to 2016. Students with acute symptomatic hepatitis A during the outbreak and other students who were asymptomatic 3 to 4 months after the outbreak were enrolled. Asymptomatic students had no symptoms from the outbreak until they were enrolled. Sera were collected to identify anti-hepatitis A virus (HAV) IgM (by enzyme-linked immunosorbent assay) and HAV genetic variations/genotypes (using polymerase chain reaction [PCR]-sequencing and phylogenetic analysis). A total of 33 (97.1%) out of 34 sera of students with acute symptoms were positive for anti-HAV IgM and 18% of them were positive by PCR, identified as HAV subgenotype IA. No prominent amino acid variations were observed from reported HAV sequences from Indonesia. Among 38 sera of asymptomatic students, most (55.3%) were positive for anti-HAV IgM, while none were positive by PCR. In conclusion, HAV-IA was the only subgenotype identified in acute symptomatic cases during the outbreak. The percentage of HAV-specific IgM-positive cases was very high among acute symptomatic students, but that was also high among asymptomatic students, which might contribute as the important source of infection during the outbreak.

KEYWORDS

genetic variation, hepatitis A virus, immunoglobulin

1 | INTRODUCTION

Hepatitis A virus (HAV) infection has long been and remains a major public health issue in many countries.¹ In developing countries, HAV

infection is endemic, with most individuals in these countries being exposed to HAV during early childhood.² Due to improvements in sanitation conditions, the pattern of HAV endemicity in several developing countries has changed from high to intermediary in recent decades.³⁻⁶ HAV infection is closely associated with unsafe water or food, inadequate sanitation, and poor personal hygiene. The

Juniastuti and Dedy Wahyuddin contributed equally to this work.

18 virus is primarily spread via the fecal-oral route, either by person-to-person contact or through the ingestion of food or water that has been contaminated by the fecal matter of infected individuals.^{7,8}

Hepatitis A is the most common form of acute viral hepatitis worldwide. A typical symptomatic presentation includes nonspecific prodromal symptoms with various combinations of fever, malaise, weakness, anorexia, nausea, vomiting, arthralgia, and myalgias. Prodromal symptoms tend to decrease with the onset of jaundice, which lasts for several weeks and is followed by a convalescent period. Peak infectivity occurs during the 2 weeks before the onset of jaundice. When jaundice appears, stool viral concentrations decrease and most patients are noninfectious after 1 week.^{1,9} Viremia occurs within 1 to 2 weeks after HAV exposure and presents for approximately 3 to 4 weeks before the onset of jaundice,¹⁰ while IgM antibodies to HAV are detectable at or before the onset of clinical illness and decrease within 3 to 6 months.¹¹ Infected children under the age of 6 years do not generally exhibit noticeable symptoms, and only 10% develop jaundice. Starting from 6 years of age to adulthood, infection typically causes more severe symptoms, with jaundice occurring in more than 70% of cases.⁸ It is important to note that in most cases, asymptomatic cases are not recognized as hepatitis A, whereas they can shed HAV in feces. The asymptomatic and nonjaundiced HAV-infected individuals, particularly children, are frequently the source of infection for others.¹² Other previous findings have also suggested that HAV is mainly transmitted by children with either asymptomatic or subclinical infection.^{13,14}

HAV is a nonenveloped RNA virus of 27 to 32 nm diameter in size, with an icosahedral symmetry, which belongs to the genus *Hepatovirus* of the *Picornaviridae* family, and has a positive-polarity, single-stranded 7.5-kb genome.¹⁵ The amplification and sequencing of variable regions within capsid proteins, that is, the VP3/VP1 and VP1/P2A junctions of wild-type HAV isolates from different regions of the world, revealed significant nucleotide sequence heterogeneity, but limited amino acid heterogeneity.^{10,16,17} Based on nucleotide sequence heterogeneity, HAV isolates may be differentiated into seven unique genotypes.¹⁸ Viruses from four of the genotypes (I, II, III, and VII) were detected in human cases of hepatitis A, whereas viruses from the other three genotypes (IV, V, and VI) were isolated from animal species only.¹⁹⁻²²

Outbreaks of hepatitis A in Indonesia have been consecutively reported in the following cities: Bogor (West Java province) in 1998; Jember and Bondowoso (East Java province) in 2006; Tangerang (West Java province) in 2007; Yogyakarta (Special Region of Yogyakarta province) in 2008; and Ngawi (East Java province) in 2009, with icteric cases in each area accounting for 74, 50, 17, 1160, and 146 individuals, respectively (Sub-directorate of Surveillance and Outbreak Response, Directorate General of Disease Control and Environmental Health, Ministry of Health, personal communication). Hepatitis A outbreaks occurred in 6 provinces with 495 cases in 2013, with East Java province being the most prevalent, and in 3 provinces with 460 cases in 2014.^{23,24} The subgenotype Indonesian IA sublineage HAV was identified in 34 sera during outbreaks of hepatitis A in some regions between 2003 and 2007 in Indonesia.²⁵

In Surabaya (the capital of East Java province), the number of hepatitis A cases increased from 20 in 2013 to 40 in 2014 (Public Health Office of Surabaya, 2013 and 2014, unpublished). In 2015, the number of hepatitis A cases markedly increased to 59, and occurred in 2 junior high schools in Surabaya (Primary Health Care of PacarKeling, 2015, unpublished). This hepatitis A outbreak was defined by the initial detection of three students who tested positive for anti-HAV IgM. The present study reported the results of genetic and serological examinations of hepatitis A in symptomatic cases (early infection) during the outbreak and in asymptomatic cases a few months after the outbreak.

33 2 | MATERIALS AND METHODS

2.1 | Samples

The outbreak of hepatitis A in two junior high schools (a private school and a public school) in Surabaya occurred between October 2015 and January 2016. Students with acute symptoms including those with acute symptoms during the sample collection and those with a history of acute symptoms for the past 1 month were enrolled in January 2016. An acute symptomatic case was defined by the onset of acute illness with a minimum of two of the following clinical symptoms: fever, malaise, flatulence, anorexia, nausea, vomiting, dark urine, and jaundice, which are consistent with hepatitis A during the outbreak and serological/genetic confirmation of acute hepatitis A (serum IgM antibody to HAV and/or HAV RNA being positive). A few months after the outbreak (between March and April 2016), students without acute hepatitis A symptoms (from the outbreak until the sampling period) were enrolled. Sera were collected from symptomatic and asymptomatic students.

Ethical clearance for this study was obtained from the Ethics Committee of the School of Medicine, Universitas Airlangga, Surabaya. Informed consent for participation in this study was obtained from the parents of each individual.

2.2 | Detection of anti-HAV IgM

An anti-HAV IgM enzyme immunoassay was performed using the BIOS HAV-IgM Enzyme Immunoassay Test Kit (Chemux BioScience, Inc, San Francisco, CA) according to the manufacturer's instructions.

2.3 | RNA extraction and reverse transcription polymerase chain reaction

Hepatitis A virus RNA was extracted from 140 μ l plasma using a commercially available kit (QIAamp Viral RNA kit; Qiagen, Tokyo, Japan).

5 Extracted RNA was reverse-transcribed and amplified using SuperScript One-Step RT-PCR (Invitrogen, Tokyo, Japan) and a set of primers. Polymerase chain reaction (PCR) amplification was subsequently performed to detect the regions of the VP1-2A and VP3-VP1 junctions. First- and second-round PCRs using primers

representing the VP1-P2A junction were performed with 35 cycles at 95°C for 2 minutes, 55°C for 1 minute, 72°C for 1 minute, and a final extension step at 72°C for 10 minutes. The first round of PCR using primers representing the VP3-VP1 junction was performed at 94°C for 5 minutes, 40 cycles at 94°C for 45 seconds, 57°C for 1 minute, 72°C for 1 minute, and a final extension step at 72°C for 10 minutes, whereas the second round was performed at 94°C for 5 minutes, 40 cycles at 94°C for 45 seconds, 55°C for 1 minute, 72°C for 1 minute, and a final extension step at 72°C for 10 minutes (Table 1).^{10,16,17,26}

PCR products were electrophoresed on a 2% agarose gel containing ethidium bromide and visualized under UV illumination. Amplified cDNA fragments were sequenced by a direct sequencing method with the BigDye Terminator v1.1 Cycle Sequencing Kit and an ABI Prism 310 sequencer (Applied Biosystems, Foster City, CA).

2.4 | Sequence analysis

Nucleotide sequences were aligned with reference strains by the program Molecular Evolutionary Genetic Analysis (MEGA) X (<http://www.megasoftware.net>). Genotypes have more than 15% nucleotide variations and subgenotypes have 7% to 7.5% nucleotide variations.²² Phylogenetic trees were constructed by the Neighbor-Joining method and bootstrap resampling was performed 1000 times.

3 | RESULTS

The outbreak occurred in two junior high schools between October 2015 and January 2016. The location of the two schools was very close. Among 1219 students during the outbreak, 59 (4.8%) hepatitis A cases were reported in the 2 schools. The definition of a case of hepatitis A was an individual exhibiting acute hepatitis A symptoms. A total of 34 students with acute symptoms from the 2 schools in January 2016 and 38 asymptomatic students between March and April 2016 (3-4 months after the outbreak) were enrolled. All 34

students with acute symptoms had jaundice accompanied by other symptoms. All students (with and without acute symptoms) were aged between 12 and 14 years old and were not vaccinated. The workflow and results obtained are shown in Figure 1.

In consideration of possible sources of HAV transmission, investigations focused on canteens, water supply systems, toilets, and the locations of septic tanks. Washbasins were not adequate in canteens (2-3 canteens in the 2 schools), with only a bucket/sink filled with water being available for dish washing, but not for hand washing. Water supplies in both schools were from three to seven wells, respectively, and also from tap water. Although the isolation of HAV was not performed from the water supplies, a bacterial analysis revealed that water from the seven wells and tap water were contaminated with *Escherichia coli* (unpublished data from the Center for Health Laboratory, Surabaya). Many toilets were available in the 2 schools: 6 toilets in the private school and 19 in the public school. Although toilets were separated from other rooms, the majority of the septic tanks were located close to the wells (less than 10 m).

3.1 | Detection of anti-HAV IgM

Positivity for anti-HAV IgM was 97.1% among 34 students with acute symptoms (33/34), and 55.3% among 38 asymptomatic students (21/38).

3.2 | Detection of the HAV RNA genome in serum

Among 33 samples from students with acute symptoms and testing positive for anti-HAV IgM, 6 (18%) were positive for HAV RNA, whereas 1 sample from a student with acute symptoms, but negative for anti-HAV IgM was negative for HAV RNA. Among 38 samples from asymptomatic students who tested positive for anti-HAV IgM ($n = 21$) and negative for anti-HAV IgM ($n = 17$), none were positive for HAV RNA. The VP1/P2A junction region was detected in 4 sera, while the VP3/VP1 junction region was detected in 2.

TABLE 1 Primers used for HAV RNA amplification

| Region | Name of primers | Sequence | Nucleotide number | Product size (bp) | |
|---------|-----------------|--------------------|---------------------------|-------------------|-----|
| VP1/P2A | 1st PCR | BR-5 ^a | TTGTCTGTCACAGAACAAATCAG | 2950-2972 | 361 |
| | | BR-9 ^a | AGTCACACCTCTCCAGGAAAACCTT | 3310-3286 | |
| | 2nd PCR | RJ-3 ^b | TCCAGAGCTCCATTGAA | 2984-3002 | 234 |
| | | BR-6 ^a | AGGAGGTGGAAGCACTTCATTGA | 3217-3193 | |
| VP3/VP1 | 1st PCR | HAV-1 ^c | GCTCCTCTTTATCATGCTATGGAT | 2172-2196 | 244 |
| | | HAV-2 ^c | CAGGAAATGTCTCAGGTACTTTCT | 2415-2391 | |
| | 2nd PCR | HAV-3 ^c | ATGTTACTACACAAGTTGGAGAT | 2195-2218 | 186 |
| | | HAV-4 ^c | GATCCTCAATTGTTGTGATAGCT | 2380-2357 | |

Abbreviation: PCR, polymerase chain reaction.

^a (17).

^b (16).

^c (10).




FIGURE 1 Workflow with the obtained results. HAV, hepatitis A virus; PCR, polymerase chain reaction

3.3 | Genetic analysis of HAV isolates

The nucleotide sequence identities of the VP1/P2A junction region with the HAV subgenotype IA strain (AB917146) from Surabaya, Indonesia ranged between 99% and 100%. The nucleotide sequence identities of the VP3/VP1 junction region with HAV subgenotype IA strain (AB839697) from Tangerang, Indonesia were 98%. No prominent amino acid variations were observed from those of other reported sequences from Indonesia (Figures 2 and 3).

Phylogenetic analyses of sequences revealed that all strains ($n = 6$) belonged to subgenotype IA (Figures 4 and 5).

4 | DISCUSSION

Anti-HAV IgM has been used diagnostically as the primary marker of acute infection,²⁷ and mainly comprises antibodies against capsid proteins. IgM anti-HAV enzyme immunoassays are available commercially.²⁸ The sensitivity, specificity, and positive predictive value of IgM anti-HAV measurements for acute hepatitis were previously reported to be 100%, 99%, and 88%, respectively.²⁹ IgM antibodies to HAV are detectable at or before the onset of clinical illness, and decrease within approximately 3 to 6 months.¹¹ A previous study reported that IgM levels were often elevated in acute hepatitis A infection (85%).³⁰ Anti-HAV IgM was shown to be negative 4 months after the onset of infection in 70% of patients and by 7 months in 86%.¹¹ Furthermore, lower concentrations were 4 to 6 months after the onset of infection.³¹

In the two junior high schools, high percentages of anti-HAV IgM were observed in symptomatic (97.1%, 33/34) and asymptomatic students (55.3%, 21/38). Of note, the percentage of anti-HAV IgM in asymptomatic students was higher than expected. These asymptomatic students had no symptoms from the time of outbreak until the time of this study (3–4 months after the outbreak), but the presence of antibodies (anti-HAV IgM) after the outbreak can confirm that they were in fact related to the outbreak. The result indicates that hepatitis A represented a great threat in the two schools, with

asymptomatic students potentially facilitating the transmission of HAV during the outbreak. CDC surveillance data³² showed that children aged 5 to 14 years historically have the highest incidence of hepatitis A, although the incidence of HAV infection may be the highest among those <5 years old.³³ Among older children (older than 6 years old) and adults, HAV infection is generally symptomatic with jaundice occurring in >70% of patients.⁸ The ratio of symptomatic to asymptomatic students in an outbreak appears to reflect the age and immunity of individuals. In a previous outbreak of hepatitis A in a religious community, a household serosurvey detected anti-HAV IgM in 15 individuals younger than 20 years, 2 (13%) of whom developed jaundice.³⁴ In this present study, a higher percentage of students with jaundice (33/54, 61%) was noted among those who tested positive for anti-HAV IgM, while the remainder (39%) did not have jaundice, whereas the age of all students ranged between 12 and 14 years. The unidentified source shedding HAV may be important for the spread of HAV to other individuals. Students with less scrupulous hygiene may serve as a major source of transmission. The lack of proper hygiene facilitates HAV transmission from these children to their relatives and friends.^{13,14}

Previous hepatitis A outbreaks in schools were reported in 2001 in Malaysian boarding schools and in a large private school in Manila.³⁵ Environmental issues, such as the chlorination of water and improvements in the personal hygiene standards of food handlers working in the schools were addressed. A number of recommendations made as a result of the investigation into this outbreak included continued surveillance for the disease, the provision of health education, and better sanitation practices in the canteen, as well as encouraging students to bring their own food and drink to school. Investigations on the two schools in Surabaya found that poor hygiene and sanitation in canteens (no available washbasins), water supplies (contaminated with *E. coli*, a known marker of feces contamination), and the close proximity of septic tanks to wells may have contributed to the spread of HAV. In the present study, the water supply was the likely transmission source because it may have been contaminated by the septic tanks. Some recommendations for these two schools may be similar to those for previous outbreaks of hepatitis A in other schools.

| | 770 | 780 | 790 | 800 | 810 | 820 |
|------------------------------------|------------|--------------|----------------|----------|-----------|--------|
| AB917146 (IA) Surabaya, Indonesia | SRIAAGDLES | SVDDPRSEEDRR | FESHIECRKPYKEL | RLRLEVGR | QRLRYAQEE | LSNEVL |
| S3K | | | | | | |
| S5K | | | | | | |
| S7K | | | | | | |
| S8K | | | | | | |
| KM261585 (IA) Netherland | | | | | | |
| AB939714 (IA) Jember, Indonesia | | | | | | |
| AB295166 (IA) Bali, Indonesia | | | | | | |
| AB939709 (IA) Bali, Indonesia | | | | | | |
| AB939718 (IA) Mataram, Indonesia | | | | | | |
| AB939695 (IA) Mataram, Indonesia | | | | | | |
| AB939694 (IA) Makassar, Indonesia | | | | | | |
| AB939724 (IA) Solo, Indonesia | | | | | | |
| AB939726 (IA) Solo, Indonesia | | | | | | |
| AB939697 (IA) Tangerang, Indonesia | | | | | | |
| AB939731 (IA) Tangerang, Indonesia | | | | | | |
| AB623053 (IA) Japan | | | | | X | |
| EU073748 (IA) South Korea | | | | | S | |
| LC049340 (IA) Mongolia | | | | | | |
| AF357222 (IA) Ghana | | | | D | | |
| EF207320 (IA) Thailand | | | S | | | |
| X93302 (IA) Italy | | | | | | |
| EU251188 (IA) Russia | | | | | | |
| K02990 (IA) USA | | | | | | |
| KC182559 (IA) Mexico | | | | | | |
| EUS26059 (IA) Uruguay | | | | | K | |
| EUS26038 (IA) Uruguay | | | | | | |
| HMT69724 (IA) Argentina | | | | | | |
| M14707 (IB) Australia | | | | | | K |
| KF724018 (IB) Spain | | | | | | K |
| KF724021 (IB) Spain | | | | | | K |
| AY741663 (IB) Egypt | | | | | | K |
| M20273 (IB) North Africa | | | | | | K |
| FJ829494 (IIA) France | | | | | S | |
| GUS90568 (IIA) France | | | | | S | |
| DQ141214 (IIA) France | | | | | S | |
| GUS90576 (IIA) France | | | | | S | |
| AY644670 (IIB) Sierra Leone | | | | | S | |
| AC968417 (IIIA) United Kingdom | D | L | | K | | K |
| EU011791 (IIIA) India | D | L | | K | | K |
| DQ010414 (IIIA) India | D | L | | K | | R |
| FJ360735 (IIIA) India | D | L | | K | | K |
| AB973882 (IIIA) Japan | D | L | | K | | K |
| AB256339 (IIIB) Japan | D | L | | K | | K |
| AB300205 (IIIB) Saga, Japan | D | L | | K | | K |
| AB245929 (IIIB) Nagoya, Japan | D | L | | K | | K |
| AB425359 (IIIB) Fukuoka, Japan | D | L | | K | | K |

FIGURE 2 Alignments of amino acid sequences in the VP1/P2A junction region of 4 HAV isolates from Surabaya (code: S, shown in bold) and 43 reported sequences of HAV genotypes/subgenotypes. HAV, hepatitis A virus

| | 500 | 510 | 520 | 530 | 540 |
|------------------------------------|----------|----------|----------|----------|----------|
| AB109057 (IA) Tangerang, Indonesia | F | S | T | T | V |
| S15K | F | S | T | T | V |
| S28K | F | S | T | T | V |
| AB109054 (IA) Makassar, Indonesia | | | | | |
| AB939714 (IA) Mataram, Indonesia | | | | | |
| LC049340 (IA) Mongolia | | | | | |
| AB012262 (IA) Japan | | | | | |
| EF181000 (IA) Uruguay | | | | | |
| MF741663 (IB) Spain | | | | | |
| MF740011 (IB) Spain | | | | | |
| DQ141214 (IIA) France | | | | | |
| FJ360735 (IIIA) France | | | | | |
| DQ141214 (IIA) France | | | | | |
| AY644670 (IIB) Sierra Leone | | | | | |
| EU011791 (IIIA) India | | | | | |
| FJ360735 (IIIA) India | | | | | |
| AB111111 (IIIB) Saga, Japan | | | | | |
| AB425359 (IIIB) Fukuoka, Japan | | | | | |

FIGURE 3 Alignments of amino acid sequences in the VP3/VP1 junction region of 2 HAV isolates from Surabaya (code: S, shown in bold) and 16 reported sequences of HAV genotypes/subgenotypes. HAV, hepatitis A virus




FIGURE 4 Neighbor-joining phylogenetic tree of nucleotide sequences in the VP1/P2A junction region of 4 HAV isolates from Surabaya (code: S*, shown in bold) and 43 reference strains. The genotypes/subgenotypes are indicated on the branches. Bootstrap values are indicated for each branch. The nucleotide sequences reported in this study were deposited in the DDBJ database under accession numbers LC432511-LC432514. HAV, hepatitis A virus

A study in China also found that an increased risk of HAV was associated with no hand washing before food preparation and cooking, before eating, and after defecation,³⁶ and a study in Brazil reported that a higher number of water taps in the home, which may promote

the use of water for washing, was associated with a decreased risk of HAV.³⁷ The prevalence of hepatitis A in Indonesia is generally very high, and mass hepatitis A vaccinations have not been performed for the prevention and control of HAV infection.³⁸ The hepatitis A




FIGURE 5 Neighbor-joining phylogenetic tree of nucleotide sequences in the VP3/VP1 junction region of 2 HAV isolates from Surabaya (code: S*, shown in bold) and 16 reference strains. The genotypes/subgenotypes are indicated on the branches. Bootstrap values are indicated for each branch. The nucleotide sequences reported in this study were deposited in the DDBJ database under accession numbers LC432515-LC432516. HAV, hepatitis A virus

vaccination currently has few indications in areas of the world in which infection is highly endemic and where most of the population is already immune¹; however, as an outbreak response, site-specific vaccinations have been recommended to control community-wide outbreaks in small communities, such as in schools.⁸

A total of 6 (18%) out of the 33 symptomatic students who tested positive for anti-HAV IgM and none of the asymptomatic students (positive or negative for anti-HAV IgM) were positive for HAV RNA. Viremia occurs within 1 to 2 weeks of HAV exposure, presents for approximately 3 to 4 weeks before the onset of jaundice, and persists until the icteric phase in immunocompetent individuals.³⁹ The duration of jaundice is quite variable, the mean length was 7 days, with a range of 4 to >22 days.²⁷ Blood collected before the onset of symptoms appears to have the highest potential risk of transmitting HAV infection because of high viral concentrations and essentially no antibodies. Serum HAV RNA was generally detected before IgM anti-HAV. However, Bower et al¹⁰ showed that viremia may persist long after the onset of symptoms (average, 79 days) and may present for a markedly longer period during the convalescent phase of hepatitis A. It should be noted that they used the PCR method, which detects HAV RNA at a level of ~4 HAV particles/mL and appears to have been more sensitive than assays used in other studies.¹⁰ In this present study, a low frequency of positivity for HAV RNA was obtained in the sera of symptomatic students who tested positive for IgM anti-HAV. The sampling time should be considered as a possible reason for this finding. This may have been due to a low HAV titer in blood at the time of specimen collection or the viremia period of HAV infection, which generally only occurred for 3 to 4 weeks, having passed.³⁹ In this study, not all blood samples were collected at the time of jaundice, some of them were collected within 1 month of

jaundice onset. Time of collecting samples is critical, however, outbreaks always represent challenges in the investigation. The low frequency of positivity for HAV RNA was consistent with a previous finding. Mulyanto et al reported that only 21.52% of 158 serum samples from patients with acute hepatitis during hepatitis A outbreaks were positive for HAV RNA.²⁵

Genetic analyses have played a crucial role in epidemiological investigations. To study the degree of variability among HAV isolates during the outbreak, a genetic analysis of selected genome regions (VP1/P2A and VP3/VP1 amino acids) was performed. The VP1/P2A junction region was detected in four sera, while the VP3/VP1 junction region was detected in two. Diverse primer pairs that amplify different regions of the HAV genome may affect their sensitivity. Based on phylogenetic analyses, all six isolated HAV strains were clustered with subgenotype IA strains (Figures 4 and 5), whereas there were no unique amino acid sequence changes from other reported isolates in other areas of Indonesia (Figures 2 and 3). Subgenotype IA is the most abundant type worldwide. Previous studies showed that subgenotype IA was the only genotype of isolated HAV strains in some areas of Indonesia, including an imported case from Surabaya.^{25,40} The results of this outbreak in Surabaya confirmed the findings of previous studies.

In conclusion, most students with acute symptoms (97.1%) during the outbreak and even those without symptoms (55.3%) after the outbreak were positive for anti-HAV IgM. Among the 6 (18%) sera from symptomatic students with anti-HAV IgM, all were identified as HAV subgenotype IA. In addition to symptomatic cases, asymptomatic cases may have played a role as a crucial source during the outbreak. Control measures need to be adapted, according to the further epidemiology investigation.

ACKNOWLEDGMENTS

The authors are grateful to all participants who donated their sera to this study. The authors would also like to thank the heads and teachers of the two schools and also the parents of the enrolled students for their kindness in supporting the collection of data and specimens.

CONFLICTS OF INTEREST

The authors declare that they have no conflicts of interest.

ORCID

Juniastuti  <https://orcid.org/0009-0001-2212-1117>

REFERENCES

- Wasley A, Fiore A, Bell BP. Hepatitis A in the era of vaccination. *Epidemiol Rev.* 2006;28:101-111.
- Jacobsen KH, Koopman JS. Declining hepatitis A seroprevalence: a global review and analysis. *Epidemiol Infect.* 2004;132:1005-1022.

3. Kunasol P, Cooksley G, Chan VF, et al. Hepatitis A virus: declining seroprevalence in children and adolescents in Southeast Asia. *Southeast Asian J Trop Med Public Health*. 1998;29:255-262.
4. Ruttiman R, Tapia-Conyer R, Urdaneta E, et al. Hepatitis A in Latin America. A changing epidemiologic pattern. *Am J Trop Med Hyg*. 1999;61:825-829.
5. Barzaga NG. Hepatitis A shifting epidemiology in South-East Asia and China. *Vaccine*. 2000;18:561-564.
6. Tanaka J. Hepatitis A shifting epidemiology in Latin America. *Vaccine*. 2000;18:557-560.
7. Pereira FEL, Gonçalves CS. Hepatitis A. *Rev Soc Bras Med Trop*. 2003;36:387-400.
8. World Health Organization (WHO). 2017. Hepatitis A. <http://www.who.int/mediacentre/factsheets/fs328/en/>
9. Koff RS. Hepatitis A. *Lancet*. 1998;341:1643-1649.
10. Bower WA, Nainan OV, Han X, Margolis HS. Duration of viremia in hepatitis A virus infection. *J Infect Dis*. 2000;182:12-17.
11. Kao HW, Ashcavai M, Redeker AG. The persistence of hepatitis A IgM antibody after acute clinical hepatitis A. *Hepatology*. 1984;4:933-936.
12. Staes CJ, Schlenker TL, Risk I, et al. Sources of infection among persons with acute hepatitis A and no identified risk factors during a sustained community-wide outbreak. *Pediatrics*. 2000;106:E54.
13. Castelli F, Matteelli A, Signorini L, et al. Pediatric migration and hepatitis A risk in host population. *J Travel Med*. 1999;6:204-206.
14. Poovorawan Y, Theamboonlers A, Chongsrisawat V, Jantaradsamee P, Chutsirimongkol S, Tangkijvanich P. Clinical features and molecular characterization of hepatitis A virus outbreak in a childcare center in Thailand. *J Clin Virol*. 2005;32:24-28.
15. Nainan OV, Xia G, Vaughan G, Margolis HS. Diagnosis of hepatitis A virus infection: A molecular approach. *Clin Microbiol Rev*. 2006;19:63-79.
16. Jansen RW, Siegl G, Lemon SM. Molecular epidemiology of human hepatitis A virus defined by an antigen-capture polymerase chain reaction method. *Proc Natl Acad Sci USA*. 1990;87:2867-2871.
17. Robertson BH, Khanna B, Nainan OV, Margolis HS. Epidemiologic patterns of wild-type hepatitis A virus determined by genetic variation. *J Infect Dis*. 1991;163:286-292.
18. Taylor MB. Molecular epidemiology of South African strains of hepatitis A virus: 1982-1996. *J Med Virol*. 1997;51:273-279.
19. Nainan OV, Margolis HS, Robertson BH, Balayan M, Brinton MA. Sequence analysis of a new hepatitis A virus naturally infecting cynomolgus macaques (*Macaca fascicularis*). *J Gen Virol*. 1991;72:1685-1689.
20. Tsarev SA, Emerson SU, Balayan MS, Ticchurst J, Purcell RH. Simian hepatitis A virus (HAV) strain AGM27: comparison of genome structure and growth in cell culture with other HAV strains. *J Gen Virol*. 1991;72:1677-1683.
21. Byun KS, Kim JH, Song KJ, et al. Molecular epidemiology of hepatitis A virus in Korea. *J Gastroenterol Hepatol*. 2001;16:519-524.
22. Robertson BH, Jansen RW, Khanna B, et al. Genetic relatedness of hepatitis A virus strains recovered from different geographical regions. *J Gen Virol*. 1992;73:1365-1377.
23. National Institute of Health Research and Development, Ministry of Health, Republic of Indonesia (2013). *Report on Result of National Basic Health Research (RISKESDAS)*.
24. National Institute of Health Research and Development, Ministry of Health, Republic of Indonesia (2014). *Report on Result of National Basic Health Research (RISKESDAS)*.
25. Mulyanto, Wibawa IDN, Suparyatno JB, et al. The complete genomes of subgenotype IA hepatitis A virus strains from four different islands in Indonesia form a phylogenetic cluster. *Arch Virol*. 2014;159:935-945.
26. Hussain Z, Das B, Husain S, et al. Hepatitis A viral genotypes and clinical relevance: clinical and molecular characterization of hepatitis A virus isolates from northern India. *Hepatol Res*. 2005;32:16-24.
27. Cuthbert JA. Hepatitis A: old and new. *Clin Microbiol Rev*. 2001;14:38-58.
28. Poddar U, Thapa BR, Prasad A, Singh K. Changing spectrum of sporadic acute viral hepatitis in Indian children. *J Trop Pediatr*. 2002;2002(48):210-213.
29. Hirata R, Hoshino Y, Sakai H, Marumo F, Sato C. Patients with hepatitis A with negative IgM-HA antibody at early stages. *Am J Gastroenterol*. 1995;90:1168-1169.
30. Mosley JW, Visoná KA, Villarejos VM. Immunoglobulin M level in the diagnosis of type A hepatitis. *Am J Clin Pathol*. 1981;75:86-87.
31. Stapleton JT. Host immune response to hepatitis A virus. *J Infect Dis*. 1995;171:S9-S14.
32. Centers for Disease Control and Prevention (CDC) (2000). *Hepatitis surveillance* (Report No. 57). Atlanta: Centers for Disease Control and Prevention.
33. Armstrong GL, Bell BP. Hepatitis A virus infections in the United States: model-based estimates and implications for childhood immunization. *Pediatrics*. 2002;109:839-845.
34. Pavia AT, Nielsen L, Armington L, Thurman DJ, Tierney E, Nichols CR. A community-wide outbreak of hepatitis A in a religious community: impact of mass administration of immune globulin. *Am J Epidemiol*. 1990;131:1085-1093.
35. David AM. Steering Committee for Prevention and Control of Infectious Diseases. Hepatitis A outbreaks-methods of intervention in South-East Asian countries. *Int J Infect Dis*. 2004;8:201-209.
36. Mausezahl D, Cheng F, Zhang SQ, Tanner M. Hepatitis A in a Chinese urban population: the spectrum of social and behavioural risk factors. *Int J Epidemiol*. 1996;25:1271-1279.
37. De Almeida LM, Amaku M, Azevedo RS, Cairncross S, Massad E. The intensity of transmission of hepatitis A and heterogeneities in social-environmental risk factors in Rio de Janeiro, Brazil. *Trans R Soc Trop Med Hyg*. 2002;96:605-610.
38. Mulyanto. Viral hepatitis in Indonesia: past, present, and future. *Euroasian J Hepato-Gastroenterol*. 2016;6:65-69.
39. Rosenblum LS, Villarino ME, Nainan OV, et al. Hepatitis A outbreak in a neonatal intensive care unit: risk factors for transmission and evidence of prolonged viral excretion among preterm infants. *J Infect Dis*. 1991;164:476-482.
40. Utsumi T, Yano Y, Amin M, et al. Acute hepatitis due to hepatitis A virus subgenotype IA as an imported infectious disease from Indonesia. *Kobe J Med Sci*. 2014;60:E43-E47.

How to cite this article: Juniastuti, Wahyuddin D, Nihayatussa'adah, et al. Analysis of genetic and serology of hepatitis A virus infection during and after outbreak in two junior high schools in Surabaya, Indonesia. *J Med Virol*. 2019;91:1048-1055.

Analysis of genetic and serology of hepatitis A virus infection during and after outbreak in two junior high schools in Surabaya, Indonesia

ORIGINALITY REPORT

23%

SIMILARITY INDEX

17%

INTERNET SOURCES

17%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Haesun Yun. "Genetic analysis of HAV strains isolated from patients with acute hepatitis in Korea, 2005–2006", *Journal of Medical Virology*, 05/2008
Publication 4%
- 2 Franco, Elisabetta. "Hepatitis A: Epidemiology and prevention in developing countries", *World Journal of Hepatology*, 2012.
Publication 2%
- 3 sites.kowsarpub.com
Internet Source 1%
- 4 www.ejohg.com
Internet Source 1%
- 5 repository.unair.ac.id
Internet Source 1%
- 6 K. H. JACOBSEN, J. S. KOOPMAN. "Declining hepatitis A seroprevalence: a global review and analysis", *Epidemiology and Infection*, 2004 1%

| | | |
|----|---|-----|
| 7 | s3.amazonaws.com Internet Source | 1% |
| 8 | jcm.asm.org Internet Source | 1% |
| 9 | docplayer.net Internet Source | 1% |
| 10 | www.cbbsweb.org Internet Source | 1% |
| 11 | aem.asm.org Internet Source | 1% |
| 12 | www.freepatentsonline.com Internet Source | 1% |
| 13 | gamalesmat.com Internet Source | <1% |
| 14 | www.scielo.br Internet Source | <1% |
| 15 | www.budilukmanto.org Internet Source | <1% |
| 16 | J. A. Cuthbert. "Hepatitis A: Old and New", Clinical Microbiology Reviews, 2001 Publication | <1% |
| 17 | journal.fkm.ui.ac.id Internet Source | <1% |

18

www.vindy.com

Internet Source

<1%

19

S.-H. Park. "Molecular characterization of hepatitis A virus isolated from acute gastroenteritis patients in the Seoul region of Korea", *European Journal of Clinical Microbiology & Infectious Diseases*, 05/26/2009

Publication

<1%

20

whqlibdoc.who.int

Internet Source

<1%

21

www.ajtmh.org

Internet Source

<1%

22

www.google.de

Internet Source

<1%

23

I Gde Rurus Suryawan, Andrianto Andrianto, Arifita Devi Anggaraeni, Arisya Agita, Ricardo Adrian Nugraha. "The Role of Human Platelet-Rich Plasma to Enhance the Differentiation from Adipose derived Mesenchymal Stem Cells into Cardiomyocyte: An Experimental Study", Cold Spring Harbor Laboratory, 2021

Publication

<1%

24

link.springer.com

Internet Source

<1%

25

Haesun Yun. "Hepatitis A Virus genotype and its correlation with the clinical outcome of acute

<1%

hepatitis A in Korea: 2006-2008", Journal of Medical Virology, 12/2011

Publication

26

icmr.gov.in

Internet Source

<1%

27

www.futuremedicine.com

Internet Source

<1%

28

pure.uva.nl

Internet Source

<1%

29

Ahmad Nejati, Manochehr Makvandi, Alireza Samarbafzadeh, Niloofar Neisi, Hamid Moradzadegan. "Molecular epidemiology of hepatitis A virus in patients in the ahwaz region of Iran", Journal of Medical Virology, 2012

Publication

<1%

30

David, A.M.. "Hepatitis A outbreaks-methods of intervention in South-East Asian countries", International Journal of Infectious Diseases, 200407

Publication

<1%

31

Hussain, Z.. "Hepatitis A viral genotypes and clinical relevance: Clinical and molecular characterization of hepatitis A virus isolates from northern India", Hepatology Research, 200505

Publication

<1%

32

arca.icict.fiocruz.br

Internet Source

<1%

33

Endo, K.. "Full-length sequences of subgenotype IIIA and IIIB hepatitis A virus isolates: Characterization of genotype III HAV genomes", Virus Research, 200706

Publication

<1%

34

repository.tudelft.nl

Internet Source

<1%

35

www.nejm.org

Internet Source

<1%

36

www.tdx.cat

Internet Source

<1%

37

www.turkjbiochem.com

Internet Source

<1%

38

synapse.koreamed.org

Internet Source

<1%

39

virologyj.biomedcentral.com

Internet Source

<1%

40

www.cdc.gov

Internet Source

<1%

41

aprenderly.com

Internet Source

<1%

42

jurnal.permi.or.id

Internet Source

<1%

43

pediatrics.aappublications.org

Internet Source

<1%

44

Mike G Catton. "Epidemiology", Viral Hepatitis, 07/07/2005

Publication

<1%

45

eddymercado.files.wordpress.com

Internet Source

<1%

46

G. L. Armstrong, B. P. Bell. "Hepatitis A Virus Infections in the United States: Model-Based Estimates and Implications for Childhood Immunization", PEDIATRICS, 2002

Publication

<1%

47

Daniel Shouval. "Chapter 17 Hepatitis A Virus", Springer Science and Business Media LLC, 2014

Publication

<1%

48

cmbdata.gsfc.nasa.gov

Internet Source

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Analysis of genetic and serology of hepatitis A virus infection during and after outbreak in two junior high schools in Surabaya, Indonesia

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8
