

Post – vaccinated asymptomatic rotavirus infections : A community profile study of children in Surabaya, Indonesia

by Maria Lusida

Submission date: 24-Jun-2020 12:18AM (UTC+0800)

Submission ID: 1348626540

File name: A_community_profile_study_of_children_in_Surabaya,_Indonesia.pdf (793.09K)

Word count: 3887

Character count: 22721



Post-vaccinated asymptomatic rotavirus infections: A community profile study of children in Surabaya, Indonesia



Emily Gunawan^{a,c}, Takako Utsumi^{b,d}, Rury M. Wahyuni^b, Zayyin Dinana^b,
Subijanto M. Sudarmo^{a,e}, Ikuo Shoji^d, Soetjipto^{b,c,a}, Maria I. Lusida^{b,c,a,*}

^a Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

^b Indonesia-Japan Collaborative Research Center for Emerging and Re-emerging Infectious Diseases, Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia

^c Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia

^d Center for Infectious Diseases, Kobe University Graduate School of Medicine, Kobe, Japan

^e Department of Child Health, Dr. Soetomo General Hospital, Universitas Airlangga, Surabaya, Indonesia

14

ARTICLE INFO

Article history:

Received 8 August 2018

Received in revised form 22 January 2019

Accepted 10 February 2019

Keywords:

Rotavirus

Asymptomatic

Post-vaccination

Genotyping

ABSTRACT

Background: Rotavirus gastroenteritis accounts for significant childhood morbidity and mortality worldwide. Vaccination using Rotarix™ (GSK) and RotaTeq® (Merck) was introduced due to the tremendous disease burden. The possibility of asymptomatic infections following vaccinations was poorly understood. This study examined rotavirus cases in post-vaccinated children, their clinical manifestations and the genotypes of isolated strains.

Methods: Stool samples of healthy, vaccinated children under 5 years of age in Surabaya were collected monthly for 1 year between January 2016 and February 2017. Episodes of gastroenteritis were reported, and samples were collected. Rotavirus was identified using multiplex reverse transcription Polymerase Chain Reaction (QIAGEN, Inc., Valencia, CA). Clinical manifestations were measured using the Vesikari score. The genotype was analyzed by Applied Biosystems (Foster, CA).

Results: A total of 109 stool samples were collected from 30 subjects, of which 22 received Rotarix; 8 RotaTeq. Nine out of 109 samples were collected during diarrhea episodes of 8 subjects. Two asymptomatic rotavirus infections were identified by RT-PCR. The genotypes isolated were G1P[8] and G3P[8].

Conclusions: Asymptomatic rotavirus infections can occur in post-vaccinated children. Strains identified were homologous to serotypes eliciting gastroenteritis in unvaccinated children of the same community.

© 2019 The Authors. Published by Elsevier Limited on behalf of King Saud Bin Abdulaziz University for Health Sciences. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Rotavirus infection accounts for significant morbidity and mortality in infants and young children worldwide. The World Health Organization (WHO) estimated 215,000 rotavirus-associated deaths among children under 5 years of age in 2013 [1]. The latest study in Indonesia reported rotavirus as the leading cause of severe acute gastroenteritis in children under 5 years of age, being responsible for 60% of hospitalizations and 41% of clinical visits presenting with diarrhea [2]. So far, 12 G- (G1-G6, G8-G12, G20) and 14 P- (P[1], P[3]-P[11], P[14], P[19], P[25], and P[28]) genotype combinations have been identified in humans, with the global epidemiological emphasis on G1-G4 and G9 [3].

Rotavirus enters humans through the oral route and destroys epithelial cells of mature villi in the small intestine, causing malabsorption and therefore osmotic diarrhea. Animal models also demonstrate the role of the enteric nervous system and NSP4 enterotoxin in inducing osmotic diarrhea. The vomiting center is simultaneously activated although its mechanism remains unclear [4]. Mortality arises when severe dehydration is treated inadequately.

Rotavirus vaccine was introduced in 2006 considering the tremendous global disease burden. Currently, the two licensed vaccines are RotaTeq® (Merck & Co., Inc., West Point, PA, USA) and Rotarix® (GSK Biologicals, Belgium) [5,6].

Many studies have examined the efficacy of rotavirus vaccination in preventing severe gastroenteritis, ranging from 40 to 64% in developing countries [7–10] and >90% in developed countries [11–15]. However, the relationship among rotavirus vaccination, the possible subsequent rotavirus infections in children

* Corresponding author at: Institute of Tropical Disease, Kampus C, Jl. Unair, Mulyorejo, Gubeng, Surabaya, East Java, 60286, Indonesia.
E-mail address: ingelusida@itd.unair.ac.id (M.I. Lusida).

<https://doi.org/10.1016/j.jiph.2019.02.015>

1876-0341/© 2019 The Authors. Published by Elsevier Limited on behalf of King Saud Bin Abdulaziz University for Health Sciences. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



Fig. 1. Map of the participants' residences in Surabaya (Figure adapted from Fig. 1 of [17]).

and changes in local circulating strains are poorly understood. In this study, rotavirus infection status post-vaccination was examined in vaccinated children in Surabaya, Indonesia.

Materials & methods

This study was approved by the Ethics Committee at the Institute of Research and Innovation (*Lembaga Penelitian dan Inovasi Universitas Airlangga*) and complied with the National Health and Medical Research Council National Statement for the Ethical Conduct in Research Involving Humans (1999). This was a prospective cohort study of representative vaccinated children under 5 years of age in Surabaya between January 2016 and February 2017. The primary outcome was to identify the presence of rotavirus in the stools of these children and to examine their clinical manifestations.

Community setting and participant recruitment

In Indonesia, rotavirus vaccination is included in the universal immunization schedule under the optional category without government funding [16]. There are no precise data on rotavirus vaccine coverage in Indonesia. Until the end of the study period, these vaccines were relatively new and remained exclusively for those who could afford it.

A total of 30 immunocompetent children under 5 years of age who had completed either RotaTaq® (Merck & Co., Inc., West Point, PA, USA) or Rotarix® (GSK Biologicals, Belgium) vaccinations were recruited from a young mothers' community in Surabaya. Surabaya is the capital city of East Java Province, Indonesia (Fig. 1), and has a warm, tropical climate. The period of recruitment was January to March 2016. Immunocompetence was defined as being HIV negative and not undergoing immunosuppressive or other immunomodulation therapies. The residences of these participants are scattered throughout the city to ensure representative results (Fig. 1).

Stool sample collection

Stool samples from each participant were collected monthly at home in clean containers provided over the one-year study period.

An initial stool sample from each study participant was obtained within 2 days after the first meeting with his/her parents. The specific date of each follow-up stool sample was dependent on the day that the parents were willing to provide a sample, ending between December 2016 and February 2017. The majority of parents were unable to provide samples in July and December 2016 due to *Idul Fitri* and the Christmas holidays.

Participants who had diarrhea between monthly collections were also instructed to provide the diarrhea sample. Diarrhea was defined as at least 3 uncommonly loose or watery stools within 24 h [18]. At the time of recruitment, two participants had diarrhea.

Stool specimens were transported promptly in refrigerated boxes to the Viral Diarrhea Laboratory, Institute of Tropical Disease Universitas Airlangga, Surabaya, where they were frozen at -80°C until tested for rotavirus.

Subject characteristics and clinical information

Informed consent was obtained from the parents of the participants upon recruitment. Subject characteristics and clinical information were attained directly from the parents of the participants using a questionnaire developed for the study. For diarrhea episodes, clinical severity was assessed using the Vesikari scoring system [19].

Identification of rotavirus

All stool specimens were subsequently thawed and aliquoted. Viral RNA was extracted using a QIAamp Viral RNA Extraction Kit® (QIAGEN, Inc., Valencia, CA) as described in the manufacturer's protocol. Identification of rotavirus was confirmed by a two-step reverse transcription-polymerase chain reaction (RT-PCR) assay against VP7 protein using the corresponding oligonucleotide primer sets (Table 1) according to the WHO manual [20]. Positive controls were also used in this study. The RT-PCR is favored due to its much lower detection threshold, whereby it can identify more asymptomatic infections with significantly lower viral loads [21,22].

Table 1
Primer sets for rotavirus identification [20].

Primer	Sequence (5'–3')	Position	Amplicon size (bp)
First amplification			
Beg9	GGCTTTAAAGAGAGAATTCCGTCTGG	nt 1-29	
End9	GGTACACATCATACAATTCTAATCTAAG	nt 1062-1036	1062
Second amplification			
RVG9	GCTACACATCATACAATTCT	nt 1062-1044	
aAT8	GTCACACCAATTGTAAATTCG	nt 178-198	885
aBT1	CAAGTACTCAAATCAATGATGG	nt 314-335	749
aCT2	CAATGATATTAACACATTTCTGTG	nt 411-435	652
aDT4	CGTTCTGTGAGGAGATTG	nt 480-498	583
aET3	CGTTGAAGAAGTTGCAACAG	nt 689-709	374
aFT9	CTAGATGTAACATACTACTAC	nt 757-776	306

35 Further genotyping was performed through nucleic acid sequencing and phylogenetic analysis of the VP7 gene. The PCR product was directly sequenced using a BigDye terminator cycle sequencing kit and Applied Biosystems 3500XL Genetic Analyzer (Applied Biosystems, Foster, CA). The sequencing results were then aligned using CLUSTAL X software. The phylogenetic analysis was conducted using Molecular Evolutionary Genetic Analysis (MEGA4) software (<http://www.megasoftware.net>).

Results

Vaccine types

9 Eight of 30 (26.7%) study participants received RotaTaq[®] (Merck & Co., Inc., West Point, PA, USA) while the remaining 22 (73.3%) received Rotarix[®] (GSK Biologicals, Belgium).

Detection of rotavirus

28 A total of 109 stool specimens were collected from the 30 study participants. The number of samples obtained from each participant ranged from 1 to 12 over the study course. Rotavirus was detected by RT-PCR in 2 stool specimens collected from 2 different asymptomatic participants. These samples were collected on 21 and 23 February 2016, respectively. On their corresponding collection days, no symptoms of gastroenteritis were reported by these participants. Stool passage frequency and consistency remained normal.

During the course of the study, eight participants experienced diarrhea at least once. One participant reported diarrhea twice, in April and June 2016. Diarrhea episodes ranged from three to ten days. The mean Vesikari score was 7.1. These children were promptly presented to their individual pediatricians and received adequate treatments. This prevented aggravating conditions as reflected by the result of clinical severity scoring. All nine diarrhea specimens were negative for rotavirus.

Characteristics of children with asymptomatic rotavirus infection

Both asymptomatic participants whose stools tested rotavirus-positive were vaccinated with Rotarix[®] (GSK Biologicals, Belgium). On specimen collection, these participants were 10 and 12 months old. The first asymptomatic participant was male, overweight, received breastmilk until 9 and a half months of age and started formula milk since 2 months of age. He was vaccinated in June (2 months old) and August (4 months old) 2015. The second asymptomatic participant was female with a normal nutritional status, breastfed until the end of the study period (23 months old), and vaccinated in April 2015 (2 months old) and 15 June 2015 (4 months old). Both also had access to safe and clean water, good sanitation and parents with adequate education.

Rotavirus genotypes

On gel electrophoresis, the migration pattern of the first positive sample corresponds to the G1 genotype. However, the second positive sample had shorter RNA migration patterns in both the first and second amplification products, which did not correspond to any G-type. Therefore, further genotyping was conducted.

Sequencing of the VP7 gene confirmed that the rotavirus genotypes were G1P[8] and G3P[8]. Phylogenetic analysis revealed that isolated strains were closely related to genotypes reported in Japan and Spain, respectively (Fig. 2).

Discussion

Rotavirus remains the global leading cause of childhood hospitalizations and deaths from severe gastroenteritis. Approximately 37% of diarrheal deaths in children under 5 years old is attributed to this infection [1]. As of 2016, 81 countries have included rotavirus vaccines in their national immunization schedules [23]. Indonesia has not been registered in the list due to the optionality of rotavirus vaccine, available only through the private market.

The analysis of stool specimens from post-vaccinated children in the community enabled us to evaluate subsequent rotavirus infections in individuals with rotavirus-specific protection, which has not been performed before. Indonesia is a suitable location to conduct this analysis because given the uneven vaccine coverage setting, the vaccinated population is more likely to be exposed to rotavirus.

In this study, asymptomatic rotavirus infections occurred in two vaccinated children. There are no available comparison data on rotavirus asymptomatic infection in vaccinated children. However, previous studies showed that the prevalence of asymptomatic rotavirus infections in the general population varied from 3% to 30% [24–27] according to community setting and age composition.

Contrastingly, stool samples from diarrhea episodes in this study tested negative for rotavirus. These findings are not consistent with previous studies which demonstrated that severe rotavirus gastroenteritis could occur in 3.7% [7] and 1.9% [10] of post-vaccinated children. A possible explanation for this might be the differences in number of samples used.

Recruitment of study participants was done before knowing their rotavirus infection status, whereas the duration of rotavirus shedding in asymptomatic infection can range from 8 to 25 days [28]. Hence, some children who were potentially infected during the period between specimen collections may not have been detected.

The genotypes of rotavirus identified in this study suggest that these asymptomatic infections were prevalent infections. G1P[8] is the most common dominant genotype worldwide with a prevalence of 52% from 1994 to 2003 [29]. A multi-center study




Fig. 2. Neighbor-joining Maximum Composite Likelihood (MCL) Phylogenetic Tree of Human Rotavirus VP7 sequences and other G representative genotypes recognized thus far (partial and/or complete). Numbers on the nodes represent the bootstrap support from 1000 replications. Strains from asymptomatic infection in the present study are denoted by an asterisk.

conducted in Indonesia reported 11% G1P[8] distribution in 2006 [2]. Similarly, G1P[8] accounted for 11.4% of cases in the 2013 one-year-surveillance in Surabaya [30].

In accordance with previous studies, we found a uniquely short RNA migration pattern which, upon further analysis, revealed genotype G3P[8] that is closely related to Spanish equine-like G3P[8] [31]. This finding supports a recent study by *Utsumi et al.* on the predominance of G3 strains with an equine-like VP7 gene and a DS-1-like genetic backbone in Indonesia between 2015–2016 [32]. However, the relationship between these vaccinated children with asymptomatic infection and unvaccinated children with intestinal infectious disease in the aforementioned study is a complex causality dilemma. Nonetheless, it is likely that asymptomatic infections

are responsible for the continuing circulation of strains within a community.

These findings suggest that rotavirus infections could still happen in post-vaccinated children. While the number of samples in this study is too small to analyze, it is presumable that both rotavirus vaccines are beneficial in preventing severe gastroenteritis. Former studies showed that the efficacy for both RotaTeq® (Merck & Co., Inc., West Point, PA, USA) and Rotarix® (GSK Biologicals, Belgium) is similar in developing countries [7–10]. Moreover, Rotarix® apparently provides cross-protection for other rotavirus serotypes, despite its monovalent G1P[8] content [33].

In asymptomatic infections, rotavirus could still be excreted in formed stools, although in lower quantities [28], thereby facilitat-

ing its spread within local communities or to other communities when the host travels. With uneven vaccine coverage, unvaccinated children would become more susceptible to severe rotavirus-associated diseases. This study result emphasizes the importance of rotavirus vaccination for children across the globe.

Conclusion

Our study reports asymptomatic rotavirus infections in vaccinated children whose serotypes are homologous to the prevalent serotypes eliciting severe clinical manifestations in unvaccinated children. Complementary to previous studies, our findings emphasize the importance of rotavirus vaccination for all children.

13

Funding

No funding Sources.

Competing interests

None declared.

Ethical approval

Not required.

1

Acknowledgements

This research was supported by the Japan Initiative for Global Research Network on Infectious Diseases (J-GRID) from Ministry of Education, Culture, Sport, Sciences & Technology in Japan, and Japan Agency for Medical Research and Development (AMED: 18fm0108004h0004). The authors acknowledge Dato' Sri Prof. Tahir for supporting this research through Tahir Professorship, Indonesia.

References

- [1] Tate JE, Burton AH, Boschi-Pinto C, Parashar UD, Agoos M, Serhan F, et al. Global, regional, and national estimates of rotavirus mortality in children <5 years of age, 2000–2013. *Clin Infect Dis* 2016;62(Suppl. 2):S96–105.
- [2] Soenarto Y, Aman AT, Bakri A, Waluya H, Firmansyah A, Kadim M, et al. Burden of severe rotavirus diarrhoea in Indonesia. *J Infect Dis* 2009;200(s1):S188–94. Available from: <https://academic.oup.com/jid/lookup/doi/10.1086/6055338>.
- [3] Afrad MH, Matthijnsens J, Moni S, Kabir F, Ashrafi A, Rahman MZ, et al. Genetic characterization of a rare bovine-like human VP4 mono-reassortant G6P[8] rotavirus strain detected from an infant in Bangladesh. *Infect Genet Evol* 2013;19(July):120–6.
- [4] Hagbom M. Rotavirus disease mechanisms diarrhea, vomiting and inflammation – how and why? 2015:1–57, 1463.
- [5] Vesikari T, Matson DO, Dennehy P, Van Damme P, Santosham M, Rodriguez Z, et al. Safety and efficacy of a pentavalent human-bovine (WC3) reassortant rotavirus vaccine. *N Engl J Med* 2006;354(1):23–33. Available from: <https://doi.org/10.1056/NEJMoa052664>.
- [6] Ruiz-Palacios GM, Pérez-Schael I, Velázquez FR, Abate H, Breuer T, Clemens SC, et al. Safety and efficacy of an attenuated vaccine against severe rotavirus gastroenteritis. *N Engl J Med* 2006;354(1):11–22. Available from: <https://doi.org/10.1056/NEJMoa052434>.
- [7] Zaman K, Ansh DD, Victor JC, Shin S, Yunus M, Dallas MJ, et al. Efficacy of pentavalent rotavirus vaccine against severe rotavirus gastroenteritis in infants in developing countries in Asia: a randomised, double-blind, placebo-controlled trial. *Lancet* 2010;376(9741):615–23.
- [8] Armah GE, Sow SO, Breiman RF, Dallas MJ, Tapia MD, Feikin DR, et al. Efficacy of pentavalent rotavirus vaccine against severe rotavirus gastroenteritis in infants in developing countries in sub-Saharan Africa: a randomised, double-blind, placebo-controlled trial. *Lancet* 2010;376(9741):606–14. Available from: [https://doi.org/10.1016/S0140-6736\(10\)60889-6](https://doi.org/10.1016/S0140-6736(10)60889-6).
- [9] Tissera MS, Cowley D, Bogdanovic-Sakran N, Hutton ML, Lyras D, Kirkwood CD, et al. Options for improving effectiveness of rotavirus vaccines in developing countries. *Hum Vaccin Immunother* 2017;13(4):921–7. Available from: <https://doi.org/10.1080/21645515.2016>.
- [10] Madhi SA, Cunliffe NA, Steele D, Witte D, Kirsten M, Louw C, et al. Effect of human rotavirus vaccine on severe diarrhea in African infants. *N Engl J Med* 2010;362(4):289–98.
- [11] Vesikari T, Matson DO, Dennehy P, Van Damme P, Santosham M, Rodriguez Z, et al. Safety and efficacy of a pentavalent human-bovine (WC3) reassortant rotavirus vaccine. *N Engl J Med* 2006;354(1):23–33. Available from: <http://www.nejm.org/doi/abs/10.1056/NEJMoa052664>.
- [12] Block SL, Vesikari T, Goveia MG, Rivers SB, Adeyi BA, Dallas MJ, et al. Efficacy, immunogenicity, and safety of a pentavalent human-bovine (WC3) reassortant rotavirus vaccine at the end of shelf life. *Pediatrics* 2007;119(1):11–8. Available from: <http://pediatrics.aappublications.org/cgi/doi/10.1542/peds.2006-2058>.
- [13] Goveia MG, Dinubile MJ, Dallas MJ, Heaton PM, Kuter BJ. Efficacy of pentavalent human-bovine (WC3) reassortant rotavirus vaccine based on breastfeeding frequency. *Pediatr Infect Dis J* 2008;27(7):656–8.
- [14] Ruiz-Palacios GM, Pérez-Schael I, Velázquez FR, Abate H, Breuer T, Clemens SC, et al. Safety and efficacy of an attenuated vaccine against severe rotavirus gastroenteritis. *N Engl J Med* 2006;354(1):11–22.
- [15] Vesikari T, Karvonen A, Pymula R, Schuster V, Tejedor JC, Cohen R, et al. Efficacy of human rotavirus vaccine against rotavirus gastroenteritis during the first 2 years of life in European infants: randomised, double-blind controlled study. *Lancet* 2007;370(9601):1757–63.
- [16] Menteri Kesehatan RI. Peraturan Menteri Kesehatan Republik Indonesia Nomor 12 Tahun 2017 tentang Penyelenggaraan Imunisasi. Kementerian Kesehatan 2017:1–162. Available from: http://hukor.kemkes.go.id/uploads/produk_hukum/PMK.No.12.tg.PenyelenggaraanImunisasi.pdf.
- [17] Nagao Y, Rachmie EM, Ochi S, Padmidewi MM. Geographical association between socioeconomic and age of dengue haemorrhagic fever patients in Surabaya, Indonesia. *Dengue Bull* 2011;35(January):116–30.
- [18] WHO. Essential concepts concerning diarrhoea. In: The treatment of diarrhoea: a manual for physicians and other senior health workers; 2005. p. 1–44. Available from: <http://apps.who.int/iris/bitstream/handle/10665/43209/9241593180.pdf?sequence=1>.
- [19] Ruuska T, Vesikari T. Rotavirus disease in Finnish children: use of numerical scores for clinical severity of diarrhoeal episodes. *Scand J Infect Dis* 1990;22(3):259–67. Available from: <https://doi.org/10.1093/003655490009027046>.
- [20] WHO. Rotavirus characterisation: molecular methods. Manual of rotavirus detection and characterization methods 2009, 143 p.
- [21] McLauchlin J, Amar CFL, East CL, Gray J, Iturriza-Gomara M, Maclure EA. Detection by PCR of eight groups of enteric pathogens in 4,627 faecal samples: re-examination of the English case-control Infectious Intestinal Disease Study (1993–1996). *Eur J Clin Microbiol Infect Dis* 2007;26(5):311–23. Available from: <https://doi.org/10.1007/s10096-007-0290-8>.
- [22] Kang G, Iturriza-Gomara M, Wheeler JG, Crystal P, Monica B, Ramani S, et al. Quantitation of group A rotavirus by real-time reverse-transcription-polymerase chain reaction: correlation with clinical severity in children in South India. *J Med Virol* 2004;73(1):118–22. Available from: <https://doi.org/10.1002/jmv.20053>.
- [23] PATH. Country national immunization program (NIP) introductions of rotavirus vaccine 2016:1–2. Available from: <http://www.vaccineresources.org/files/PATH-Country-Introduction-Table-EN-2016.05.01.pdf>.
- [24] Barron-Romero BL, Barreda-Gonzalez J, Doval-Ugalde R, Zermeno-Eguia Liz J, Huerta-Pena M. Asymptomatic rotavirus infections in day care centers. *J Clin Microbiol* 1985;22(1):116–8.
- [25] Abiodun PO, Ihongbe JC, Ogbimi A. Asymptomatic rotavirus infection in Nigerian day-care centres. *Ann Trop Paediatr* 1985;5(3):163–5. Available from: <https://doi.org/10.1080/02724936.1985.11748385>.
- [26] Endara P, Trueba G, Solberg OD, Bates SJ, Ponce K, Cevallos W, et al. Symptomatic and subclinical infection with rotavirus P[8]G9, rural Ecuador. *Emerg Infect Dis* 2007;13(4):574–80.
- [27] Phillips G, Lopman B, Rodrigues LC, Tam CC. Asymptomatic rotavirus infections in England: prevalence, characteristics, and risk factors. *Am J Epidemiol* 2010;171(9):1023–30.
- [28] Mukhopadhyay I, Sarkar R, Menon VK, Babji S, Paul A, Rajendran P, et al. Rotavirus shedding in symptomatic and asymptomatic children using reverse transcription-quantitative PCR. *J Med Virol* 2013;85(9):1661–8.
- [29] Santos N, Hoshino Y. Global distribution of rotavirus serotypes/genotypes and its implication for the development and implementation of an effective rotavirus vaccine. *Rev Med Virol* 2005;15:29–56. Available from: <https://doi.org/10.1002/rmv.448>.
- [30] Sudarmo SM, Shigemura K, Athiyah AF, Osawa K, Wardana OP, Darma A, et al. Genotyping and clinical factors in pediatric diarrhea caused by rotaviruses: one-year surveillance in Surabaya, Indonesia. *Gut Pathog* 2015;7(1):1–10.
- [31] Arana A, Miriam A, Montes M, Jere KC, Iturriza-Gomara M, Cilla G. Emergence and spread of G3P[8] rotaviruses possessing an equine-like VP7 and a DS-1-like genetic backbone in the Basque Country (North of Spain), 2015. *Infect Genet Evol* 2016;44:137–44. Available from: <https://doi.org/10.1016/j.meegid.2016.06.048>.
- [32] Utsumi T, Wahyuni RM, Doan YH, Dinana Z, Soegijanto S, Fujii Y, et al. Equine-like G3 rotavirus strains as predominant strains among children in Indonesia in 2015–2016. *Infect Genet Evol* 2018;61:224–8. Available from: <https://doi.org/10.1016/j.meegid.2018.03.027>.
- [33] Steele AD, Neuzil KM, Cunliffe NA, Madhi SA, Bos P, Ngwira B, et al. Human rotavirus vaccine Rotarix™ provides protection against diverse circulating rotavirus strains in African infants: a randomized controlled trial. *BMC Infect Dis* 2012;12(1):213. Available from: *BMC Infectious Diseases*.

Post – vaccinated asymptomatic rotavirus infections : A community profile study of children in Surabaya, Indonesia

ORIGINALITY REPORT

25%

SIMILARITY INDEX

18%

INTERNET SOURCES

24%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Takako Utsumi, Maria Inge Lusida, Zayyin Dinana, Rury Mega Wahyuni et al. "Occurrence of norovirus infection in an asymptomatic population in Indonesia", *Infection, Genetics and Evolution*, 2017
Publication 2%
- 2 safelir.tbzmed.ac.ir
Internet Source 1%
- 3 e-journal.unair.ac.id
Internet Source 1%
- 4 Mokibul Hassan Afrad, Jelle Matthijnsens, Sayra Moni, Farzana Kabir et al. "Genetic characterization of a rare bovine-like human VP4 mono-reassortant G6P[8] rotavirus strain detected from an infant in Bangladesh", *Infection, Genetics and Evolution*, 2013
Publication 1%
- 5 Tonietti, P. O., A. S. Hora, F. D. F. Silva, V. L. A. Ruiz, and F. Gregori. "Phylogenetic Analyses 1%

of the VP4 and VP7 Genes of Porcine Group A Rotaviruses in Sao Paulo State, Brazil: First identification of G5P[23] in Piglets", Journal of Clinical Microbiology, 2013.

Publication

6	uclouvain.be Internet Source	1%
7	academic.oup.com Internet Source	1%
8	Takako Utsumi, Rury Mega Wahyuni, Yen Hai Doan, Zayyin Dinana et al. "Equine-like G3 rotavirus strains as predominant strains among children in Indonesia in 2015–2016", Infection, Genetics and Evolution, 2018 Publication	1%
9	www.who.int Internet Source	1%
10	dhr.gov.in Internet Source	1%
11	"Acknowledgement to Reviewers", Journal of Infection and Public Health, 2019 Publication	1%
12	Fan, Z.C.. "Generation and characterization of an N ^p r ^o -disrupted marker bovine viral diarrhea virus derived from a BAC cDNA", Journal of Virological Methods, 200808	1%

13	s3-eu-west-1.amazonaws.com Internet Source	1%
14	hal.archives-ouvertes.fr Internet Source	1%
15	Ainara Arana, Milagrosa Montes, Khuzwayo C. Jere, Miriam Alkorta, Miren Iturriza-Gómara, Gustavo Cilla. "Emergence and spread of G3P[8] rotaviruses possessing an equine-like VP7 and a DS-1-like genetic backbone in the Basque Country (North of Spain), 2015", <i>Infection, Genetics and Evolution</i> , 2016 Publication	1%
16	staging.mja.com.au Internet Source	1%
17	www.portalgaruda.org Internet Source	1%
18	bmcpediatr.biomedcentral.com Internet Source	<1%
19	fieldresearch.msf.org Internet Source	<1%
20	Devi Oktafiani, Ni Luh Ayu Megasari, Elsa Fitri Ana, . Nasronudin, Maria Inge Lusida, . Soetjipto. "	<1%

First Report on HHV-6 Infection Among HIV-Infected Individuals Residing in Surabaya, Indonesia

", HIV/AIDS - Research and Palliative Care, 2020

Publication

21

cubeplayland.blogspot.com

Internet Source

<1%

22

bmcinfectdis.biomedcentral.com

Internet Source

<1%

23

Elmira Flem, Kirsti Vainio, Henrik Døllner, Cathrine Midgaard et al. "Rotavirus gastroenteritis in Norway: Analysis of prospective surveillance and hospital registry data", *Scandinavian Journal of Infectious Diseases*, 2009

Publication

<1%

24

Ching-Min Wang, Shou-Chien Chen, Kow-Tong Chen. "Current status of rotavirus vaccines", *World Journal of Pediatrics*, 2015

Publication

<1%

25

Mustafa Altindis. "Molecular characterization of rotaviruses in mid-western Turkey, 2006–2007", *Central European Journal of Medicine*, 12/07/2009

Publication

<1%

26	onlinelibrary.wiley.com Internet Source	<1%
27	rotacouncil.org Internet Source	<1%
28	www.gutpathogens.com Internet Source	<1%
29	www.nature.com Internet Source	<1%
30	www.mdpi.com Internet Source	<1%
31	pediatric-infectious-disease.imedpub.com Internet Source	<1%
32	Nischay Mishra, Lina Reslan, Mohsen El-Husseini, Hawraa Raouf et al. "Full genome characterization of human G3P[6] and G3P[9] rotavirus strains in Lebanon", Infection, Genetics and Evolution, 2020 Publication	<1%
33	Ratana Tacharoenmuang, Satoshi Komoto, Ratigorn Guntapong, Sompong Upachai et al. "High prevalence of equine-like G3P[8] rotavirus in children and adults with acute gastroenteritis in Thailand", Journal of Medical Virology, 2019 Publication	<1%
34	Yano, Yoshihiko. "Hepatitis B virus infection in	

Indonesia", World Journal of Gastroenterology, 2015.

Publication

<1%

35

Shyamal Kumar Paul. "Phylogenetic analysis of rotaviruses with genotypes G1, G2, G9 and G12 in Bangladesh: evidence for a close relationship between rotaviruses from children and adults", Archives of Virology, 11/2008

Publication

<1%

36

Pérez-Vilar, Silvia, Javier Díez-Domingo, Joan Puig-Barberà, Ruth Gil-Prieto, and Silvana Romio. "Intussusception following rotavirus vaccination in the Valencia Region, Spain", Human Vaccines & Immunotherapeutics, 2015.

Publication

<1%

37

Putri Sari Wulandari, Juniastuti, Rury Mega Wahyuni, Mochamad Amin et al. "Predominance of Norovirus GI.4 From Children With Acute Gastroenteritis In Jambi, Indonesia, 2019", Journal of Medical Virology, 2020

Publication

<1%

38

Marlen Martinez-Gutierrez, Victor Arcila-Quiceno, Juanita Trejos-Suarez, Julian Ruiz-Saenz. "Prevalence and molecular typing of rotavirus in children with acute diarrhoea in Northeastern Colombia", Revista do Instituto de Medicina Tropical de São Paulo, 2019

Publication

<1%

39

kce.fgov.be

Internet Source

<1%

40

tampub.uta.fi

Internet Source

<1%

41

Yamanaka, Atsushi, Yuko Tabuchi, Kris C. Mulyatno, Helen Susilowati, Eryk Hendrianto, Soegeng Soegijanto, and Eiji Konishi. "Dengue virus infection-enhancing and neutralizing antibody balance in children of the Philippines and Indonesia", *Microbes and Infection*, 2012.

Publication

<1%

42

Shahina Tabassum. "Genomic characterization of rotavirus strains obtained from hospitalized children with diarrhoea in Bangladesh", *Journal of Medical Virology*, 05/1994

Publication

<1%

43

Subijanto Sudarmo, Katsumi Shigemura, Alpha Athiyyah, Kayo Osawa et al. "Genotyping and clinical factors in pediatric diarrhea caused by rotaviruses: one-year surveillance in Surabaya, Indonesia", *Gut Pathogens*, 2015

Publication

<1%

44

Jelle Matthijnssens. "Rotavirus disease and vaccination: impact on genotype diversity", *Future Microbiology*, 12/2009

Publication

<1%

45

Ruggeri, F. M., and L. Fiore. "Advances in understanding of rotaviruses as food- and waterborne pathogens and progress with vaccine development", *Viruses in food and water*, 2013.

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Post – vaccinated asymptomatic rotavirus infections : A community profile study of children in Surabaya, Indonesia

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
