

**DAFTAR PUSTAKA**

- Apriastuti DA. (2012). *Kejadian Hiperbilirubinemia Akibat Inkompatibilitas ABO di RSUD Pandan Arang boyolali*. Skripsi. Universitas Sebelas Maret
- Auliasari, NA. (2019) *Faktor Risiko Kejadian Ikterus Neonatorum di RSUD DR. Soetomo*. Skripsi. Universitas Airlangga.
- Behrman, Kliegman. Arvin. (2014). *Nelson Ilmu Kesehatan Anak Edisi: 15 Volume 2*. Jakarta : EGC. 854 – 856.
- Bethesda, Dean L. 2005. National Center for Biotechnology Information (US); diakses di <https://www.ncbi.nlm.nih.gov/books/NBK2266/>. Diakses pada tanggal: 25 Juli 2020.
- Biade, *et al.* (2016). Faktor Risiko Hiperbilirubinemia pada Bayi Lahir dari Ibu Diabetes Mellitus . *Sari Pediatri*, **18**(1), pp 6.
- Bozkurt, Ö., Yücesoy, E., Oğuz, B., Akinel, Ö., Palali, M. F., & Ataş, N. (2020). Severe neonatal hyperbilirubinemia in the southeast region of turkey. *Turkish Journal of Medical Sciences*, 50(1), 103–109. <https://doi.org/10.3906/sag-1906-22>
- Brits, Hanneke., Adendorff, Jeanie.,Huisamen, Dyanti., *et al.* 2018. The prevalence of neonatal jaundice and risk factors in healthy term neonates at National District Hospital in Bloemfontein. *African Journal of Primary Health Care & Family Medicine*, **10**(1), 1-6, <https://doi.org/10.4102/phcfm.v10i1.1582>.
- Bulbul A, Cayonu N, Sanli ME, Uslu S. Evaluation of risk factors for development of severe hyperbilirubinemia in term and near term infants in Turkey. *Pak J Med Sci*. 2014;30:1113–8.

- Campbell Wagemann, S. and P. Mena Nannig (2019). Severe hyperbilirubinemia in newborns, risk factors and neurological outcomes. *Rev Child Pediatr.* 90(3); 267-274
- Chand, N. & Sanyal, A.J.(2007). Sepsis-Induced Cholestasis. *Hepatology.* 45(1).
- Cohen, et al (2010). Understanding Neonatal Jaundice: A prespective on causation. *Taiwan pediatric assosiation.* 51(3), pp 143-148
- Cohudhury, Panna., Bagga, A., Chugh, K., Ramji, S., Gupta, P. (2011). Principels of Pediatrics & Neonatal Emergencies Third Edition. *The Indian Academy of Pediatrics: India.*
- Devi DS, Vijaykumar B (2017). Risk Factors For Neonatal Hyperbilirubinemia: A Case Control Study. *Journal of Reproduction, Contraception, Obstetrics and Gynecology,* 6, 198-202
- El-Kabbany, et al. (2017). Does anaesthesia in mothers during delivery affect bilirubin levels in their neonates?. *Korean J Pediatr,* 60(12), pp 385–389.
- England, Carole (2009). Bayi dengan Berat Badan Lahir Rendah yang sehat dalam Diane M. Frasel dan Margareth MA. *Buku Ajar Bidan Myles edisi 14.* Jakarta : ECG
- Garosi E, Mohammadi F, Ranjkesh. (2016). The Relationship Between Neonatal Jaundice and Maternal and Neonatal Factors. *Iranian Journal of Neonatology,* 7(1), pp 38-39.
- Harrison, VC. (2008). *The Newborn Babyi 5<sup>th</sup> Edition.* Cape Town : Juta & Company, Ltd.

- Hue Chen, Su., *et al* (2012). Association of ABO Incompatibility With Red Blood Cell Indices of Cord Blood Unit. *Taiwan Pediatric Association*. 53, pp 138-143
- Hwang, N. R., & Kim, J. K. (2018). Relationship between asymptomatic rotavirus infection and jaundice in neonates: A retrospective study. *BMC Pediatrics*, **18**(1). <https://doi.org/10.1186/s12887-018-1352-z>
- Ifalahma, D. (2015). Karakteristik Bayi Baru Lahir dengan Ikterus Patologi Di Rumah Sakit Umum Daerah Wonogiri. Akademi Kebidanan Citra Medika Surakarta. <http://ejurnal.akbidcm.ac.id>.
- Ikatan Dokter Anak Indonesia. 2013. Indikasi Terapi Sinar pada Bayi Menyusui yang Kuning. Public Article.
- Imron R, Metti D. (2015). Peningkatan Angka Kejadian Hiperbilirubinemia Pada Bayi Baru Lahir Dikarenakan Berat Badan Lahir Yang Rendah (BBLR). *Jurnal Keperawatan*, **9**(1), pp 47-49.
- Isa, H. M., Mohamed, M. S., Mohamed, A. M., Abdulla, A., & Abdulla, F. (2017). Neonatal indirect hyperbilirubinemia and glucose-6-phosphate dehydrogenase deficiency. *Korean Journal of Pediatrics*, **60**(4), 106–111. <https://doi.org/10.3345/kjp.2017.60.4.106>
- Ives, N. K. (2015). Management of Neonatal Jaundice. *Pediatric and Child Health*. **25**(6), pp. 276-281
- Kanoko, M.(2012). Buku saku : Pelayanan Kesehatan Neonatal Esensial. Jakarta: Kementerian Kesehatan.
- Kasemy, Z. A., Bahbah, W. A., El Hefnawy, S. M., & Alkalash, S. H. (2020). Prevalence of and mothers' knowledge, attitude and practice towards

glucose-6-phosphate dehydrogenase deficiency among neonates with jaundice: A cross-sectional study. *BMJ Open*, 10(2).  
<https://doi.org/10.1136/bmjopen-2019-034079>

Kaur., S *et al* (2019). Risk Factors for Low Birth Weight among Rural and Urban Malaysian Women. *BMC Public Health*. 19.

Kemenkes RI. (2019). Profil Kesehatan Indonesia 2018. Jakarta : Pusdatin Kemenkes RI.

Kirk, Jean M.(2008). Neonatal jaundice: a critical review of the role and practice of bilirubin analysis. *Annals of Clinical Biochemistry*,**45**, pp. 452–462.

Kurniawati, D.O. (2015). Profil Ibu Hamil Risiko Tinggi Berdasarkan Umur dan Paritas. Terdapat dalam [www.jurnal.akbidgriyahusada.ac.id](http://www.jurnal.akbidgriyahusada.ac.id).

Laroia, Nirupama. 2015. Birth Trauma. <http://emedicine.medscape.com/article/980112-overview#4>

Mansjoer A.(2013). *Kapita Selekta Kedokteran Jilid 2. Edisi III*. Jakarta: Media Aesculapius Fakultas Kedokteran Universitas Indonesia.

Maulida, Mentariasih. (2018). *Hubungan Kejadian Hiperbilirubinemia Dengan Inkompatibilitas ABO Pada Bayi Baru Lahir Di Rsud Abdul Moeloek Provinsi Lampung Tahun 2014 – 2015*. Skripsi. Universitas Lampung.

Marlina, L (2017). *Gambaran Karakteristik Ibu dengan Kejadian Ikterus Pada Neonatus di RSUD Wates Kulon Progo*. Skripsi. Universitas Aisyiyah Yogyakarta

Martiza, L., *et al*. (2012).*Buku Ajar Gastroenterologi-Hepatologi*. Jakarta : IDAI.

Mathindas S, Wilar R dan Wahani A. 2013. Hiperbilirubinemia Pada Neonatus. *Jurnal Biomedik*. **5**(1). Pp 4-10

- Mudmainna. (2015). Faktor Resiko Kejadian Ikterus Neonatorum Pada Neonatus Di Ruang Perinatologi Teratai Rumah sakit Umum Daerah Kabupaten Muna Tahun 2014 - 2015.
- Moeslichan *et al.* (2004). *Tatalaksana Ikterus Neonatorum di Indonesia*. Jakarta : HTA Indonesia
- Mojtahedi SY, Izadi A, Seirafi G, Khedmat L, Tavakolizadeh R. Risk Factors Associated with Neonatal Jaundice: A Cross-Sectional Study from Iran. 2018. Open Access Maced J Med Sci.;6(8):1387-1393. Published 2018 Aug 11. doi:10.3889/oamjms.2018.319
- Nartono, K. (2013). *Inflamasi Plasenta Sebagai Faktor Ikterogenik Neonatal pada Inkompatibilitas Golongan Darah ABO ibu-janin*. Jakarta: Universitas Indonesia
- Nkhoma, Ella., Poole, Charles., et al (2009). The global prevalence of glucose-6-phosphate dehydrogenase deficiency: A systematic review and meta-analysis. *Blood Cells Molecules and Diseases* 42(3):267-78. DOI: [10.1016/j.bcmd.2008.12.005](https://doi.org/10.1016/j.bcmd.2008.12.005)
- Norman, M., Åberg, K., Holmsten, K., Weibel, V., & Ekéus, C. (2015). Predicting nonhemolytic neonatal hyperbilirubinemia. *Pediatrics*, 136(6), 1087–1094. <https://doi.org/10.1542/peds.2015-2001>
- Olusanya, B.O. Osibonjo, F.B. Slusher, T.N. (2015). *Risk Factors for Severe Neonatal Hyperbilirubinemia in Low and Middle Income Countries: A Systematic Review and MetaAnalysis. PLOS ONE* | DOI: 10.1371/journal.pone.0117229.

- Omar, R.A., Algur, N., Megged, O., Hammerman, C., & Kaplan, M. (2015). Glucose-6-phosphate dehydrogenase screening in Israel-Arab and Palestinian-Arab neonates. *Journal of Pediatrics*, 167(1), 169–172. <https://doi.org/10.1016/j.jpeds.2015.04.039>
- Ozcan, M., *et al.* (2017). Hyperbilirubinemia Due to Minor Blood Group Incompability in Newborn: A Case Report. *Turkish Pediatric Association*.
- Percival, Patricia. (2009). *Ikterus dan Infeksi*. Dalam: Fraser DM & Margareth MA. Myles Buku Ajar Bidan Edisi 14. Jakarta : ECG
- Puspita, Ndaru (2018). Pengaruh Berat Badan Lahir Rendah Terhadap Kejadian Ikterus Neonatorum Di Sidoarjo. *Jurnal Berkala Epidemiologi*. 6(2), Pp : 174 - 181
- Ramachandran, Arun (2015). Neonatal hyperbilirubinaemia. *Pediatrics and Child Health*, 26(4), pp 162-168.
- Raines, Deborah A dan Jain, Sameer. (2019). Cephalhematoma. New York : StatPearls Publishing LLC. Diperoleh dari : <https://www.ncbi.nlm.nih.gov/books/NBK470192/> [3 Januari 2020]
- Rini, Kartika. (2016) *Analisis Faktor Risiko yang Berhubungan Dengan Kejadian Ikterus Neonatorum Fisiologis Di Ruang Cendrawasih RSUD Dr. Sutomo*. Skripsi. Universitas Airlangga
- Rochjati, P. (2013). *Skrining Antenatal Pada Ibu Hamil*. Surabaya: FK UNAIR
- Shah, P.
- Rougée, Luc Ra et al. “Obstetric Obesity is Associated with Neonatal Hyperbilirubinemia with High Prevalence in Native Hawaiians and Pacific

- Island Women.” *Hawai'i journal of medicine & public health : a journal of Asia Pacific Medicine & Public Health*, 75,12 (2016): 373-378.
- Roy R, dan Mulla, S.(2017). Clinical Guideline for Non- Invasive Bilirubin measurements using a Transcutaneous Bilirubinometer. *Norfolk and Norwich University Hospital*.
- Sakinah, Bulqis Inas (2019). *Faktor Risiko Kelahiran Prematur di RSUD Dr. M. Soewandhi Surabaya pada Tahun 2017*. Skripsi. Universitas Airlangga
- Scrafford CG, Mullany LC, Katz J, Khattry SK, LeClerq SC, Darmstadt GL, et al. (2013) Incidence of and risk factors for neonatal jaundice among newborns in southern Nepal. *Trop Med Int Health*, **18**(11), pp 1317–1328. Diperoleh dari : <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5055829/> [16 September 2019]
- Setareh Akhavan, *et al.* (2016). Oxytocin and Neonatal Hyperbilirubinemia: A Cohort Study. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, **7**(4), pp 20198
- Shan, K. H., Wang, T. M., & Lin, M. C. (2019). Association between rooming-in policy and neonatal hyperbilirubinemia. *Pediatrics and Neonatology*, **60**(2), 186–191. <https://doi.org/10.1016/j.pedneo.2018.06.002>
- Suhartati. (2010). Mewaspada Defisiensi Glukosa-6 Fosfat Dehidrogenasi (G6PD) dalam Upaya Mewujudkan Indonesia Sehat. *Fakultas Kedokteran Universitas Airlangga*.
- Sukadi, A. (2010). *Hiperbilirubinemia*. Dalam Kosim *et al.* (2014). *Buku Ajar Neonatologi*. Edisi I cetakan ke-4 . Jakarta : IDAI

- Susilowati, L (2017). Faktor – faktor yang Berhubungan dengan Terjadinya Hiperbilirubin pada Neonatus di RSB Permata Hati, *2*(2), pp 42-53.
- Sumangala Devi D, Vijaykumar B. (2016). Risk factors for neonatal hyperbilirubinemia: a case control study. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology*, *6*(1), pp 198. Diperoleh dari : <http://dx.doi.org/10.18203/2320-1770.ijrcog20164657> [14 September 2019]
- Suradi, Rulina dan Debby Letupeirissa. 2013. Buku Bedah ASI IDAI. Diperoleh dari: <https://www.idai.or.id/artikel/klinik/asi/air-susu-ibu-dan-ikterus>. [Diakses tanggal : 28 Juli 2020]
- Tavakolizadeh, R. *et al.* (2018) Maternal risk factors for neonatal jaundice: a hospital-based crosssectional study in Tehran. *European Journal of Translational Myology*, *28*(3), p.1. Diperoleh dari <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6176394/>
- Theis, S.R and Hashmi, M.F. (2019). *Coomb Test*. From : <https://www.ncbi.nlm.nih.gov/books/NBK547707> [13 Oktober 2019]
- Thielemans, Laurence., Trip-Hoving, Margreet., Bancone, Germana., et al. 2017. Neonatal Hyperbilirubinemia in a Marginalized Population on the ThaiMyanmar Border: a study protocol. *BMC Pediatrics*, *17*(1), pp 32. DOI 10.1186/s12887-017-0798-8
- Tresnatias, YD. (2017) *Analisis Faktor Risiko Ikterus Neonatorum di RS Haji Surabaya*. Skripsi. Universitas Airlangga
- UNICEF (2018). *Breastfeeding in the 21<sup>st</sup> Century*. Diperoleh dari : [www.unicef.org/breastfeeding](http://www.unicef.org/breastfeeding)



- Urrah et al.(2016). Hyperbilirubinemia in Neonates: Types, Causes, Clinical Examinations, Preventive Measures and Treatments: A Narrative Review Article. *Iran J Public Health*, **45**(5), pp 558–568. Diperoleh dari : <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4935699/> [18 Desember 2019]
- Victora CG, Bahl R, Barros AJ, et al( 2016). Breastfeeding in the 21st century: epidemiology, mechanisms, and lifelong effect. *Lancet*.;387(10017):475-490. doi:10.1016/S0140-6736(15)01024-7
- Wan ASL, Mat Daud S, Teh SH, et al.(2016). Management of Neonatal Jaundice in Primary Care Malays Fam Physician. **11**(2 & 3), Pp 16-19. Diperoleh dari: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5408871/>. [21 Oktober 2019]
- Welsh, Andrew (ed.). (2010) *Neonatal Jaundice*. [online] London : Royal College of Obstetricians and Gynaecologists, <https://www.nice.org.uk/guidance/cg98/evidence/full-guideline245411821>. (12 September 2019)
- Weng, Y. H., Chiu, Y. W. & Cheng, S. W. 2012.Breast milk jaundice and maternal diet with chinese herbal medicines. *Evid. Based Complement. Alternat. Med.* 2012, 150120
- Weng, Y., Chiu, Y., Cheng, S. et al. 2016. Risk assessment of gene variants for neonatal hyperbilirubinemia in Taiwan. *BMC Pediatr* 16, 144 . <https://doi.org/10.1186/s12887-016-0685-8>
- Weng, Y. H., Chiu, Y. W. & Cheng, S. W. 2018. Risk assessment of prolonged jaundice in infants at one month of age: A prospective cohort study. [www.nature.com/scientificreports](http://www.nature.com/scientificreports). 8(1), pp 6-11

- WHO. 2019. *Infant, newborn*. [Online] who.int. diperoleh dari : <https://www.who.int/infant-newborn/en/>. (13 September 2019)
- Wibowo, Satrio. (2007). Perbandingan Kadar Bilirubin Neonatus Dengan Dan Tanpa Defisiensi Glucose-6-Phosphate Dehydrogenase, Infeksi Dan Tidak Infeksi. Thesis. *Program Pasca Sarjana Magister Ilmu Biomedik Dan Program Pendidikan Dokter Spesialis -I Ilmu Kesehatan Anak Universitas Diponegoro*.
- Widagdo. (2012). *Tatalaksana Masalah Penyakit Anak dengan Ikterus*. Jakarta: CV Sagung Seto.
- Wijaya, F.A dan Suryawan, I.W.B (2019). Faktor risiko kejadian hiperbilirubinemia pada neonatus di ruang perinatologi RSUD Wangaya Kota Denpasar. *Medicina 2019*. 50(02), pp : 357 - 364
- Winkjosastro H. (2012). *Ilmu Kebidanan*. Jakarta.: Yayasan Bina Pustaka Sarwono Prawirohardjo.
- Wisra, Dewi. (2016). *Hubungan Metode Persalinan dengan Ikterus Neonatorum di RSUD Wates Yogyakarta Tahun 2016*. Skripsi. Program Studi Ilmu Keperawatan STIKES Jenderal Achmad Yani Yogyakarta.
- Yaestin, NA. (2017). *Hubungan Berat Badan Lahir Rendah Dengan Kejadian Ikterus Di RS PKU Muhammadiyah Yogyakarta*. Skripsi. Universitas Aisyiyah Yogyakarta.
- Yang, Hui., Wang, Qian., Dan Zheng, Lei, Et Al, 2016. Clinical Significance Of Ugt1a1 Genetic Analysis In Chinese Neonates With Severe Hyperbilirubinemia. *Pediatrics And Neonatology*. 57(4), Pp 307-310

Zhou, Jinzu., Yang, Changyi., Zhu, Wenbi, *et al.* 2018. Identification of Genetic Risk Factors for Neonatal Hyperbilirubinemia in Fujian Province, Southeastern China: A Case-Control Study. *BioMed Research International*. Pp 1-8