



saya

Kepada ijvs@vetmedmosul.org & 1 lainnya

3 Jun 2018 jam 13.01

 1 lampiran

To,

The Editors

Iraqi Journal of Veterinary Sciences (Iraqi J. Vet. Sci.)

Dear Sir/Ma'am

We are submitting a manuscript entitled

The Expression of IL-1 β , IL-6 And TNF- α After the Administration of Pisang Ambon (Musa Paradisiaca L.) Stem Extract in Rats' Enteritis for consideration for publication in the Iraqi Journal of Veterinary Sciences (Iraqi J. Vet. Sci.).

I, Rochmah Kurnijasanti (and Miranti Candrarisna) certify that:

- * The manuscript is original work of all authors.
- * All authors made a significant contribution to this study.
- * This manuscript is not under the consideration, submitted for publication, or has been published in any other journal.
- * All authors have read and approved the final version of the manuscript.

Thank you

Sincerely,

Rochmah Kurnijasanti

Laboratory of Pharmacology, Faculty of Veterinary Medicine, Universitas Airlangga



Bintangikan



Hapus



Pindahkan ke



Teruskan



Balas



Lainnya

Publishing article



Taha Muneer

Kepada santisam19@yahoo.co.id


6 Jun 2018 jam 02.46

2 lampiran


Dear Dr,

I receive the article which you like to publish in the Iraqi J vet Sci. you need to follow the instruction of Of Iraqi J vet sci, you can see this instructions online however I attached the instruction and i mention some instruction in your article please follow it and re sending the article with my best wishes.

Prof Dr Muneer S. Al-badrany
Secretary of Iraqi j vet sci



Stem Extr... .docx



تعليمات... .docx



Bintangkan



Hapus



Pindahkan ke



Teruskan



Balas



Lainnya

Re: Publishing article



saya

Kepada [Taha Muneer](#)

11 Jun 2018 jam 15.28

1 lampiran

Dear **Prof. Dr. Muneer S. Al-badrany,**

Thank you for the instructions. I have revised the references according to the guide. Thus, herewith I attached the revised file. Please kindly check the attached.

Sincerely,
Rochmah Kumijasanti (Dr., M.Si., drh.)

> Tampilkan pesan asli



The Expre....docx
4.5MB



Bentangkan



Hapus



Pindahkan ke



Teruskan



Balas



Lainnya

Revision article



Taha Muneer

Kepada santisam19@yahoo.co.id

23 Jul 2018 jam 16.41

Dear Dr,
We inform you that your manuscript needs improvement and you can see it in the attachment I sent. Please repair it and send it to us soon.

Prof Dr Muneer S. Al-badrany
Secretary of Iraqi j vet sci



8-144(2).pdf
54.6kB

-
- undown
 - Hapus
 - Pindahkan ke
 - Teruskan
 - Balas
 - Lainnya

The effect of pisang ambon (*Musa paradisiaca L.*) stem extract on the regulation of IL-1 β , IL-6 and TNF- α in rats' enteritis

Commented [ws1]: Please change to italic font

Abstract

Considering that NSAIDs are widely used in medication for pain, inflammatory, and fever, its side effect, mucosal erosion, needs to be minimized. The purpose of this paper is to study the potential use of *Musa paradisiaca L.* stem extract in lowering TNF α , IL-1 β and IL-6 proinflammatory cytokines in rats' enteritis. Thirty male wistar strain rats (*Rattus Norvegicus*) aged 2-3 months were divided into 5 groups with 6 repetitions. The positive and negative control groups were given a solvent of *Musa paradisiaca L.* stem extract. Groups P1, P2, and P3 were treated with the extract. The groups were treated for 14 days. On the 8th day, all of the rats were administered with Indomethacin as enteritis inducer, except for the rats in the negative control group which were only administered with indomethacin solvent. On the 15th day, the rats were necropsied, and their jejunum were taken in order to make the immunohistochemistry (IHC) preparations. The IHC test results were statistically analyzed. The results showed that the granting of *Musa paradisiaca L.* stem extract has anti-inflammatory activity against the inflamed jejunum tissue. The anti-inflammatory activity against the damage of jejunum tissue was indicated by an attenuated expression of IL-1 β , IL-6 and TNF- α proinflammatory cytokines in the jejunum. The increasing dose showed more significant effects.

Commented [ws2]: What statistical tests were used to analyze the data ?

Keywords: Enteritis, IL-1 β , IL-6, TNF- α

Commented [ws3]: Please added *Musa paradisiaca L.*

Introduction

The Non-Steroidal Anti-inflammatory Drugs (NSAIDs) are often used as medication to pain, inflammatory, and fever. The use of these medications is very effective in overcoming pain, fever, and inflammation, but it is often accompanied by side effects on mucosal erosion i.e. that can form ulcers, perforations and bleeding to cause death (1). This happens because NSAIDs have wide enterohepatic circulation. One of NSAIDs that has the most toxic side effects on intestinal mucosa is indomethacin. Granting a systemic NSAIDs also cause damage to the small intestine due to biliary excretion (2). The NSAIDs drugs showed toxic effects on the small intestine through a constitutive enzyme inhibition of cyclooxygenase-1 (COX1) which subsequently led to a decrease in the synthesis of protective prostaglandins. The NSAIDs are also thought to have local effect that damage the mucosa of the small intestine. On the cellular level, Indomethacin also induces mitochondrial damage of villenterositis which results a villus shortening (3). This situation causes a lot of increased permeability of the small intestine (4). The side effects of NSAIDs can be inhibited or reduced with the use of drugs that can increase the synthesis of protective prostaglandins.

Treatment efforts can be done by utilizing plants as medicines, mainly from sewage plants that have not been utilized optimally. Pisang Ambon (*Musa paradisiaca* L.) is one of very important plants that is economically important because all parts of the plant can be utilized. In Indonesian, pisang means banana. For example, banana flowers can be cooked, banana stems can be used as a traditional medicine, organic fertilizer, forage, and others (5). The sap of the banana tree can also be used as a wound healer (6). Methanol extract of *Musa paradisiaca* L. can be used as wound healer (7).

Pisang Ambon (*Musa paradisiaca* L.), particularly the stem, have polyphenolic compounds as antioxidant activity which serves to inhibit the concentration of fat, has a capacity of electron donor and the reduction of metal ions, as well as serve as a chelating agent to the metal ion. The methanol extract of *Musa paradisiaca* var. *Sapientum* L have the anti-inflammatory effect against enteritis in rats which were induced with Indomethacin (8).

The inflammation process in the small intestine which is induced by indomethacin involves oxidant compounds that cause oxidative damage on the mucosa. The granting of brown seaweed in rats, which have experimental enteritis by indomethacin, may reduce the inflammatory process that is allegedly happening on intestinal mucosa via the attenuation of the negative effects of oxidants. Enteritis

induced by indomethacin is a form of inflammation that has many hispathological similarities to inflammatory bowel disease. The reactive oxygen compounds play a significant role in the process of inflammation on enteritis induced by indomethacin. An increase in reactive oxygen compounds in the tissues of the small intestine causes oxidative damage that worsen the inflammation.

It can be studied by observing the expression of proinflammatory cytokines including 'tumor necrosis factor- α ' (TNF- α) and interleukin 1 β (IL-1 β), which are the major factors in the mucosa damage of the small intestine. The explanation of the effect of the *Musa paradisiaca* L. stem extract granting against inflammation can be used as a theoretical foundation for further research about the prevention and treatment of enteritis induced by NSAIDs particularly indomethacin.

The results of the research are expected to explain the influence of the *Musa paradisiaca* L. stem extract granting against inflammation which can be observed on the expression of TNF- α , IL-1 β and IL-6 proinflammatory cytokines. The explanation of the effect of the *Musa paradisiaca* L. stem extract against inflammation can be used as a theoretical foundation for further research about the prevention and treatment of enteritis induced by NSAIDs particularly indomethacin. Aim of present work was to study the potential use of *Musa paradisiaca* L. stem extract in attenuating the expression of TNF α , IL-1 β and IL-6 proinflammatory cytokines in rats' enteritis.

Materials and methods

Ethical approval

The experiment was carried out with the ethical clearance by the Commission of Health Research Ethics, Faculty of Dentistry, Universitas Airlangga

Animals

Thirty male wistar strain rats (*Rattus Norvegicus*) aged 2-3 months and weight 150-200 g. The animals were grouped into 5 groups.

Materials

Indomethacin, a banana (*Musa paradisiaca* L.) stem, 10% formalin in 0.15 M PBS (pH 7.4), formic acid, 30%/0.28 M sodium citrate, ethanol, xylene, paraffin, water for injection, mayer hematoxylyn, eosin, chloroform, glacial acetic acid, 10% NaCl, ketamine, Methyl Green 1%, rat's monoclonal antibodies of TNF α , IL-1 β and IL-6 antibodies, Goat anti mouse IgG biotin labeled, SA-HRP

Commented [ws4]: Please added in rats's enteritis

Commented [ws5]: Please give a number of ethical clearance

(Streptavidin-Horse Peroxidase), DAB (3,3 diamino benzidine tetrahydrochloride).

Instrumentations and tools

Glass objects, glass cover, electron microscope, surgical tools, test tubes, medicine pots, rotary evaporator, pH meter, balance, camera, and ANNOVA software.

Preparation of the extract

Banana stem was cleaned, cut into small pieces, and dried. Once dry, the cleaned banana stem was mashed, resulting 900 g banana powder. The powder was extracted using maceration method with ethanol (3000 ml). The obtained macerate was collected and evaporated using rotary evaporator under vacuum, resulting thick extract of 45 g. The ethanol extract of the banana stems was put in air tight containers and stored in refrigerator till time to use.

Animal treatments

The first group (negative control, K-) consists of 6 male Wistar rats were given only a solvent of banana extract and indomethacin solvent. The second group (positive control, K+) consists of 6 male Wistar rats were given an indomethacin 15 mg/kg body weight orally to trigger enteritis without giving extracts of banana stems. The third group (P1) is composed of six male Wistar rats which were administered with the banana stem extract 5 mg/kg (9) for 7 days. The fourth group (P2) is composed of six male Wistar rats which were administered with the banana stem extract 10 mg/kg (9) for 7 days. The fifth group (P3) is composed of six male Wistar rats which were administered with the banana stem extract 20 mg/kg (9) for 7 days.

On the 8th day, the rats were administered with indomethacin 15 mg/kg (single dose) of body weight orally using oral gavage as enteritis inductions (10). The administration of the banana stems extract was continued until 7 days after the first administration. All rats were dissected in order to obtain the jejunum tissue. The TNF α , IL-1 β and IL-6 immunohistochemical examination was performed in each group.

Histopathology and immunohistochemistry

The manufacture of deparaffinized paraffin blocks and the TNF α , IL-1 β and IL-6 immunohistochemical expression examination were performed in accordance with Boenisch methods (11). The paraffin blocks were cut using rotary microtome. Blocking with 3% H₂O₂ in PBS pH 7.4 and in blocking with 1% BSA in PBS for 60 minutes. Labelled the slides with primary antibodies which are anti-TNF α , IL-1 β , and IL-6 in 4 degrees Celsius 1% BSA overnight. Washed the slides with PBS pH 7.4 three times for 5 minutes. Incubated the slides with SA-HRP (Horseradish Peroxidase-Streptavidin) 1:500 for 40 minutes at room temperature. Washed the slides with PBS pH 7.4

three times for 5 minutes. Dropped the slides with the DAB (3,3 benzene diamino tetra hydrochloride) chromogen substrate for 20 minutes. Washed with PBS pH 7.4 three times for 5 minutes. Wash with H₂O three times for 5 minutes each. Counter stained with Methyl Green 1% at room temperature. The slides were soaked with tap water for 5 minutes. Preparations were made and dried overnight at room temperature & the slides were ready to use.

Statistical analysis

A statistical analysis was performed to TNF α , IL-1 β and IL-6 expression variables on the 6 small intestine tissue by scoring the value based on Knodel scoring methods (12). To find out if there is a meaningful difference in the expression of TNF α , IL-1 β and IL-6 on the jejunum tissue between the groups, ANOVA test was performed and then followed by double Duncan's Distance test with significance level of 5%.

Results

Indomethacin, the enteritis inducer, was administered with dose 10 mg/kg, 15 mg/kg, and 20 mg/kg to each in 6 animals. It was performed in an effort to obtain the optimal dose of enteritis induction without causing the death. The results showed that the most proper dose of enteritis induction was 15 mg/kg. The results of the exploration can be seen in Figure 1.




Figure 1: Rats' feces, before induction (black arrow) after induction (red arrow).

Figure 1 shows that the Indomethacin with a dose of 15 mg/kg was successfully used as enteritis induction. The indication can be seen in the rat's feces (Figure 1). The feces signify the presence of diarrhea in the rat where the feces were more rodents and there was bleeding in comparison previous feces before induction.

Commented [ws6]: How much percent of ethanol is used for maceration ?

Commented [ws7]: When the mice were examined TNF α , IL-1 β and IL-6 ?

Commented [ws8]: How many micro meter jejunum cutting with a microtome ?

Histopathological observation of rats' jejunum which has been administered with indomethacin and administered with banana (*Musa paradisiaca* L.) extract can be seen in the Figure 2.

Immunohistochemical examination was performed by using monoclonal antibody against IL-1 β , IL-6 and TNF- α proteins. Positive results are characterized by the presence of brown spots on the cytoplasm. The results were examined using a microscope with 10x40 magnification.

The immunohistochemical examination results with monoclonal antibodies against the expression of IL-1 β , to see the effect of the banana stem extract (*Musa paradisiaca* L.), can be seen in Figure 3 and Table 1.

The immunohistochemical examination results with monoclonal antibodies against the expression of IL-6, to see the effect of the banana stem extract (*Musa paradisiaca* L.), can be seen in Figure 4 and Table 2.

The immunohistochemical examination results with monoclonal antibodies against the expression of TNF α , to see the effect of the banana stem extract (*Musa paradisiaca* L.), can be seen in Figure 5 and Table 3.

Table 1: The interleukin 1 β (IL-1 β) expression on IHC examination

Groups	The average of IL-1 β expression \pm SD (cell/viewing field)
Negative Control (K-)	32.00a \pm 2.36
Positive Control (K+)	64.67e \pm 4.13
5 mg/kg banana stem extract (P1)	53.33d \pm 3.07
10 mg/kg banana stem extract (P2)	45.50c \pm 5.64
20 mg/kg banana stem extract (P3)	39.83b \pm 3.25

Table 2: The interleukin 6 (IL-6) expression on IHC examination

Groups	The average of IL-6 expression \pm SD (cell/viewing field)
Negative Control (K-)	28.33a \pm 2.16
Positive Control (K+)	60.00d \pm 2.00
5 mg/kg banana stem extract (P1)	48.83c \pm 2.61
10 mg/kg banana stem extract (P2)	42.00b \pm 8.40
40 mg/kg banana stem extract (P3)	28.17 a \pm 3.82

Commented [ws9]: Please added of rats' jejunum

Commented [ws10]: Please added of rats' jejunum

Commented [ws11]: Please correct the dosage for this group




Figure 2: Histomicrograph of epithelium, H.E color with 100x magnification using Optilab @ viewer. Explanation: (K-) there is no damage to the epithelium, (K+) many erosion (e) and ulcer (u) can be observed, (P1) there are many erosion (e), (P3) there is small desquamation (d)

Commented [ws12]: Please added of mucosa jejunum




Figure 3: The interleukin 1 β (IL-1 β) expression on the IHC examination using a microscope with amagnification of 400 times.

Commented [ws13]: Please added from rat's jejunum




Figure 4: The interleukin 6 (IL-6) expression on the IHC examination using a microscope with amagnification of 400 times.

Commented [ws14]: Please added from rat's jejunum




Figure 5: The interleukin TNF α expression on the IHC examination.

Table 3: The interleukin TNF α expression of rats' jejunum on IHC examination

Groups	The average of TNF α expression \pm SD (cell/viewing field)
Negative Control (K-)	43.66a \pm 3.20
Positive Control (K+)	79.67d \pm 3.88
5 mg/kg banana stem extract (P1)	59.33c \pm 7.31
10 mg/kg banana stem extract (P2)	53.83b \pm 3.25
20 mg/kg banana stem extract (P3)	44.83a \pm 3.25

Discussion

The results showed that the administration of banana stem extract (*Musa paradisiaca* L.) has an anti-inflammatory activity against inflamed jejunal. Anti-inflammatory activity against tissue damage in the small intestine was indicated by a decreased expression of IL-1 β , IL-6, and TNF- α proinflammatory cytokines in jejunum. In increasing doses, the results showed a more significant decrement of IL-1 β , IL-6 and TNF- α proinflammatory cytokines expressions in jejunum. This is possible because the higher the dose, the more potent anti-inflammatory

active ingredient, which was expressed by the decrement of IL-1 β , IL-6 and TNF- α proinflammatory cytokines in the jejunum. The expression of IL-1 β , IL-6 and TNF- α proinflammatory cytokines in jejunum is a systematic acute phase response caused by the combined work of IL-1, TNF- α , and IL-6. Interleukin-1, IL-6, and TNF alpha work synergistically to increase T cell activation and acute inflammatory responses. In 12-24 hours from the beginning of the acute phase response, increased IL-1, TNF- α , and IL-6 induce the production of acute phase proteins by hepatocytes. The ability of IL-1, TNF- α , and IL-6 to induce the production of acute phase proteins by liver cells occurs through the induction of the same transcription factor, NF-IL-6, after each cytokine binds to its receptor. This macrophage is an active cell that has phagocytosis activity that releases more inflammatory mediators and cytokines. The active macrophages tissue secretes IL-1, IL-6, and TNF- α cytokines, which induce many local and systemic changes in acute inflammatory responses. All three cytokines work locally by inducing coagulation and increasing vascular permeability. TNF- α and IL-1 induce increased expression of adhesion molecules in vascular endothelial cells. TNF- α stimulates the expression of Selectin E that bind to adhesion molecules on neutrophils,

Commented [ws15]: Please added from rat's jejunum

whereas IL-1 induces increased expression of ICAM-1 and VCAM-1 which bound to integrins in lymphocytes and monocytes. Neutrophils, monocytes, and lymphocytes in the circulation are attached to the blood vessel wall by recognizing the adhesion molecule which then moves through the blood vessel wall and into the tissue. IL-1 and TNF- α also work on macrophages and endothelial cells to induce the chemokines that play a role in neutrophilic influx through enhancement of their adhesion to vascular endothelium and act as strong chemotactic factors. IFN- γ and TNF- α activate macrophages and neutrophils, increase phagocytic activity, and increase release of lytic enzymes into the tissues (13).

Local acute inflammatory responses can occur without much involvement of the immune system. Cytokines, that are sometimes released at the site of inflammation, facilitate the attachment of immune cells to vascular endothelial cells and the migration of these cells through blood vessel walls into the tissues. This leads to lymphocyte, neutrophil, monocyte, eosinophil, basophil, and mast cell infestation to the site of damaged tissue, where they play a role in clearance of antigens and tissue healing (13). The timing and intensity of the local acute inflammatory response should be carefully regulated to control tissue damage and facilitate the tissue repair mechanism which is needed for healing.

Conclusion

To conclude, the provision of banana stem extract can decrease TNF α , IL-1 β and IL-6. TNF α , IL-1 β and IL-6 expression were decreased even more along with the increased dose of banana stem extract.

Commented [ws16]: Please added expression

Acknowledgement

The authors would like to thank DRPM of Directorate of Higher Education (DIKT) for providing the fund to conduct this research.


References

1. Peesa J, Yalavarthi P, Rasheed A, Mandava V. A perspective review on role of novel NSAID prodrugs in the management of acute inflammation. *J Acute Dis.* 2016;5(5).
2. Niu X, de Graaf I, van der Bij H, Groothuis G. Precision cut intestinal slices are an appropriate ex vivo model to study NSAID-induced intestinal toxicity in rats. *Toxicol Vitro.* 2014;28(7):1296-1305.
3. Basivireddy J, Vasudevan A, Jacob M, Balasubramanian K. Indomethacin-induced mitochondrial dysfunction and oxidative stress in villus enterocytes. *Biochem Pharmacol.* 2002;64(2):339-349.
4. Silver K, Littlejohn A, Thomas L, Bawa B, Lillich J. Suppression of calpain expression by NSAIDs is associated with inhibition of cell migration in rat duodenum. *Toxicol.* 2017;383:1-12.
5. Suhartanto M, Harti H. *Teknologi Sehat Budidaya Pisang: Dari Benih Sampai Pasca Panen.* 1st ed. [ebook] Bogor: Pusat Kajian Hortikultura Tropika Lembaga Penelitian Dan Pengabdian Kepada Masyarakat Institut Pertanian Bogor. Available at: <http://pkht.ipb.ac.id/wp-content/uploads/2016/02/buku-ajar-teknologi-sehat-pisang.pdf> [Accessed 24 Dec. 2017].
6. Pangestika R. Efektivitas Getah Batang Semu Pisang Ambon (*Musa acuminata*) Dan Getah Batang Semu Pisang Kepok (Musa balbisiana) Pada Penyembuhan Luka Bakar Mencit (*Mus musculus*). Undergraduate. Universitas Sanata Dharma. 2017. 69p. <http://repository.usd.ac.id/id/eprint/9424>
7. Amutha K, Selvakumari U. Wound healing activity of methanolic stem extract of *Musa paradisiaca* Linn. (Banana) in Wistar albino rats. *Int Wound J.* 2014;13(5):763-767.
8. Akinlolu A. Musa Sapientum Improv. Total Antioxidant and Lipid Profile Status of Adult Wistar Rats in Indomethacin – Induced Gastric Ulceration. *Dergipark.ulakbim.gov.tr.* Retrieved 16 November 2017, from <http://dergipark.ulakbim.gov.tr/sducmfir/article/view/1018003883/0>.
9. Salau B. Methanolic Extract of Musa Sapientum Surker Proffers Antioxidant Effects against Indomethacin-Induced Ulceration Oxidative stress in Wistar Rats. *Int J Pharm Biol Sci.* 2013;8(1):07-12. <http://dx.doi.org/10.9790/3008-0810712>
10. Odabasoglu F, Cakir A, Suleyman H, Aslan A, Bayir Y, Halici M, Kazaz C. Gastroprotective and antioxidant effects of usnic acid on indomethacin-induced gastric ulcer in rats. *Ethnopharmacol.* 2006;103(1), 59-65. 10.1016/j.jep.2005.06.043.
11. Li W, Hao D, Fan T, Huang H, Yao H, Niu X. Protective effect of chelerythrine against ethanol-induced gastric ulcer in mice. *Chemico- Biological Interactions.* 2014;208:18-27. 10.1016/j.cbi.2013.11.011
12. Rüschoff J, Dietel M, Baretton G, Arbogast S, Walch A, Monges G. HER2 diagnostics in gastric cancer—guideline validation and development of standardized immunohistochemical testing. *Virchows Archiv.* 2010;457(3):299-307. 10.1007/s00428-010-0952
13. Kindt T, Goldsby R, Osborne B, Kuby J. *Kuby immunology.* New York: W.H. Freeman; 2007. 574 p. ISBN: 9778071678

Commented [ws17]: Please added page number

← Back ↶ ↷ → Archive Move Delete Spam ...


Article Revision Yahoo/Inbox ☆



 Santi Kurnia <santisam19@yahoo.co.id> Thurs, August 2 at 7:42 PM ☆
To: Taha Muneer

Dear Prof. Dr. Taha Muneer

I hereby send a revision of my article entitled "The effect of pisang ambon (*Musa paradisiaca* L.) stem extract on the regulation of IL-1 β , IL-6, and TNF- α in rats' enteritis" and I attach it to this email. Thank you very much for your attention and cooperation.

Sincerely
Rochmah Kumijasanti
Faculty of Veterinary Medicine
Airlangga University, Surabaya, Indonesia


Revision.docx



← Back ↶ ↷ → Archive Move Delete Spam ...

Acceptance Letter Yahoo/Inbox ☆

 **Iraqi Journal of Veterinary Sciences** <vetmedmosul@gmail.com>  Fri, August 28 at 6:32 PM ☆

To: Rochmah Kurnijasanti

Dear Dr. Rochmah Kurnijasanti

Thanks for sharing the approved final proof of your article. Your article will be published soon.

Sincerely
Prof. Taha Muneer
Iraqi Journal of Veterinary Science