

Volume 31 (2020): Issue 5 (Sep 2020)

in Journal of Basic and Clinical Physiology and Pharmacology



REVIEW

Phytochemical, ethanomedicinal and pharmacological applications of escin from *Aesculus hippocastanum* L. towards future medicine

Sahar Idris, Anuradha Mishra, and Mohd Khushtar

Article Category: Review Article | Article Number: 20190115 | Published online: 10 Jul 2020

ABSTRACT

Medicinal plants are used from ancient times for treatment of various ailments. *Aesculus hippocastanum* (Horse chestnut), is the popular and most v

... Show More

PDF 



Relationship between trough level of tyrosine kinase inhibitor (imatinib and nilotinib) and BCR-ABL ratios in an Indonesian chronic-phase chronic myeloid leukemia (CML) population

Budi Suprapti, Mareta Rindang Andarsari, Pharmasinta Putri Hapsari, Junaidi Khotib, Suharjono, and Siprianus Ugroseno Yudho Bintoro

Article Category: Research Article | Article Number: 20190315 | Published online: 04 Aug 2020

ABSTRACT

Objectives

Among Chronic Myeloid Leukemia (CML) patients treated with Tyrosine Kinase Inhi

... Show More

Dietary supplementation of *Pleurotus tuber regium* in rat feed ameliorates metabolic and hematotoxicity induced by carbon tetrachloride

Kenneth Obinna Okolo, Orish Ebere Orisakwe, and Iyeopu Minakiri Siminialayi

Article Category: Research Article | Article Number: 20190188 | Published online: 11 Jan 2020

ABSTRACT

Pleurotus tuber regium, a wild edible mushroom can reduce free radical-mediated injury and oxidative stress induced by carbon tetrachloride (CCl₄)

... Show More

Protective effects and chemical composition of *Corchorus olitorius* leaf fractions against isoproterenol-induced myocardial injury through p65NF_κB-dependent anti-apoptotic pathway in rats

Babatunde Alabi, Temidayo Omobowale, Joseph Badejo, Adeolu Adedapo, Oluwole Fagbemi, and Olugbenga Iwalewa

Article Category: Research Article | Article Number: 20190108 | Published online: 21 Apr 2020

ABSTRACT

Background

The fractions of *Corchorus olitorius* leaf (COLF) were evaluated against oxidative stress, inflammation and apoptosis in isoproterenol (I

... Show More

Natural limonoids protect mice from alcohol-induced liver injury

Abacuc Valansa, Borris Rosnay Tietcheu Galani, Pascal Dieudonne Djamen Chuisseu, Armelle Tontsa Tsamo, Vincent Brice Ayissi Owona, and Nicolas Yanou Njintang

Article Category: Research Article | Article Number: 20190271 | Published online: 07 Apr 2020

ABSTRACT

Background

Alcoholic liver disease (ALD) is regarded as a global health problem with limited therapeutic options.

Previous studies highlighted some

... Show More

Production of the secondary metabolite catechin by *in vitro* cultures of *Camellia sinensis* L

Sutini, Widiwurjani, Chrismawan Ardianto, Junaidi Khotib, Djoko Agus Purwanto, and Wirdhatul Muslihatin

Article Category: Research Article | Article Number: 20190357 | Published online: 06 May 2020

ABSTRACT

Background

Catechin is one of the secondary metabolites in *Camellia sinensis* L. that is alternatively produced through *in vitro* cultures. The *in vit*

... Show More

Modulatory properties of cardiac and quercetin glycosides from *Dacryodes edulis* seeds during L-NAME-induced vascular perturbation

Peter Uchenna Amadi, Emmanuel Nnabugwu Agomuo, and Chiamaka Winifred Adumekwe

Article Category: Review Article | Article Number: 20190116 | Published online: 11 Jul 2020

ABSTRACT

Background

Numerous food wastes have been identified to possess potent bioactive compound

... Show More

CASE REPORT



Enoxaparin induced reactive thrombocytosis: a rare adverse drug reaction

Saleel Salman Meenpidiyil, Shihas Azeez, Vaisakh Prasanna Kumar, Dhanush Suresh, Sareena

Kalathathoduill, Safa Thandupara, and Muhammed Hashik Puthukudi

Article Category: Case Report | Article Number: 20190312 | Published online: 24 Jun 2020

ABSTRACT

Objectives

Enoxaparin is a low molecular weight heparin (LMWH). which belongs to the class

Budi Suprapti*, Mareta Rindang Andarsari, Pharmasinta Putri Hapsari, Junaidi Khotib, Suharjono and Siprianus Ugroseno Yudho Bintoro

Relationship between trough level of tyrosine kinase inhibitor (imatinib and nilotinib) and BCR-ABL ratios in an Indonesian chronic-phase chronic myeloid leukemia (CML) population

<https://doi.org/10.1515/jbcpp-2019-0315>

Received October 23, 2019; accepted April 7, 2020; published online August 7, 2020

Abstract

Objectives: Among Chronic Myeloid Leukemia (CML) patients treated with Tyrosine Kinase Inhibitor (TKI-imatinib-nilotinib), some showed a suboptimal response. Based on pharmacokinetic studies, TKI trough level (C_{min}^{∞}) is associated with clinical outcomes, reflected by the BCR-ABL ratio. However, the interindividual pharmacokinetic variability of imatinib and nilotinib is found to be moderate–high. This study aims to analyze the relationship between TKI C_{min}^{∞} and BCL-ABL ratio in chronic-phase CML patients.

Methods: Cross-sectional study to CML chronic-phase patients treated with imatinib 400 mg daily or nilotinib 400 or 800 mg daily for ≥ 12 months. The exclusion criteria were therapy discontinuation within 29 days (imatinib) or 8 days (nilotinib) before the sampling day. Blood samples were drawn 1 h before the next dose. Imatinib-nilotinib C_{min}^{∞} and BCR-ABL ratio were measured using HPLC and RT-qPCR. The relationship was analyzed using bivariate correlation Spearman's rho test.

Results: Twenty-three imatinib and 11 nilotinib patients met the inclusion criteria. The mean imatinib and nilotinib C_{min}^{∞} were $1,065.46 \pm 765.71$ and $1,445 \pm 1,010.35$ ng/mL respectively. There were large interindividual variations in both groups (71.87% vs. 69.88%). Half of the patients in each group were found to reach C_{min}^{∞} target ($\geq 1,000$ ng/mL,

imatinib; ≥ 800 ng/mL nilotinib), but only 12 (35,29%) of them result in BCR-ABL ratio $\leq 0.1\%$. C_{min}^{∞} imatinib was found to be significantly associated with BCR-ABL ratio. But, not with the nilotinib group.

Conclusions: There were high interindividual variations of imatinib and nilotinib correlated with BCR-ABL ratio, but no correlation in nilotinib.

Keywords: BCR-ABL ratio; C_{min}^{∞} ; CML; imatinib; nilotinib.

Introduction

The development of tyrosine kinase inhibitors (TKI) has changed the management of Chronic Myeloid Leukemia (CML) therapy from lethal cancer to controlled chronic disease [1, 2]. Imatinib and nilotinib selectively inhibit tyrosine kinase activity by occupying the Adenosine Triphosphate (ATP) binding domain in ABL so as to prevent substrate phosphorylation. The inhibition of phosphorylation of the substrate will inactivate the nucleus and cytoplasmic signal transduction pathways, including RAS, phosphatidylinositol-3 kinase – alpha serine/threonine kinase (PI3K-Akt) and Janus kinase – signal transducers and activators of transcription (JAK-STAT) which cause a decreased proliferation and increased cell apoptosis [3].

Imatinib mesylate is the first generation of TKI, which can induce therapeutic responses including hematological response (leucocyte), cytogenetic response (presence/absence of cells containing the Philadelphia chromosome), and molecular response (BCR-ABL ratio) in more than 80% of CML patients [4, 5]. Although imatinib is known to produce a high cytogenetic response, some patients do not respond to imatinib therapy or relapse after a primary response. Nilotinib is one of the second generations of TKI that is approved by the FDA to be used as therapy in CML patients who are resistant or intolerant to imatinib [6, 7].

The BCR-ABL ratio is a parameter of the TKI molecular response, expressed the ratio of BCR-ABL transcription

*Corresponding author: Budi Suprapti, Department of Clinical Pharmacy, Airlangga University, Campus C Jalan Mulyorejo, Surabaya, East Java, 60115, Indonesia, E-mail: budi-s@ff.unair.ac.id

Mareta Rindang Andarsari, Pharmasinta Putri Hapsari, Junaidi Khotib and Suharjono: Faculty of Pharmacy, Department of Clinical Pharmacy, Airlangga University, Surabaya, East Java, Indonesia
Siprianus Ugroseno Yudho Bintoro: Hematology Oncology Division, Department of Internal Medicine, Dr Soetomo General Hospital, Surabaya, East Java, Indonesia

level to ABL with a target of $\leq 1\%$ after 12 months of therapy. CML patients treated with imatinib or nilotinib showed that decreasing of BCR-ABL transcript levels occurred most rapidly in 6 months on TKI and reached the plateau at 12–15 months after therapy [8]. Data in Dr. Soetomo General Hospital, Surabaya, showed that there was an increase in CML cases from 58 patients in 2006 to 160 patients in 2014 [9] and molecular response attained in 40% patients only.

Based on pharmacokinetic studies, there is a relationship between of TKI C_{min}^{∞} and therapeutic response [10]. Additionally, there was high variability in TKI C_{min}^{∞} . The coefficient of interindividual variation in C_{min}^{∞} of imatinib and nilotinib is 45–50% [11]. One of the factors that influence the suboptimal response to TKI therapy is the variation of pharmacokinetic factors and/or drug interactions that affect the pharmacokinetics of TKI [12]. This study was conducted to determine the relationship between the TKI C_{min}^{∞} (imatinib and nilotinib) and BCR-ABL ratio in patients with chronic phase CML who had used TKI, either imatinib or nilotinib, for ≥ 12 months therapy.

Materials and methods

Study design and population

This observational study was conducted on adult patients (age ≥ 18 years) with a diagnosis of chronic phase CML who had been treated with TKI, both imatinib 400 mg/day or nilotinib 400 or 800 mg/day for ≥ 12 months. The exclusion criteria were the discontinuation of therapy within 29 days (imatinib) or 8 days (nilotinib) before the sampling day. All subjects gave informed consent before the study. The study was approved by Dr. Soetomo General Hospital Ethical Review Committee and conducted in accordance with the criteria set by the declaration of Helsinki.

Study protocol

About 8 mL of blood was drawn from each patient within ± 1 h before the next dose. Blood samples were collected in two separate vacutainers for determination of C_{min}^{∞} of TKI (5 mL) and BCR-ABL ratio (3 mL). Imatinib and nilotinib levels were measured using High-Performance Liquid Chromatography (HPLC) method, and the BCR-ABL ratio is measured using RT-qPCR.

Measurements of C_{min}^{∞} of imatinib and nilotinib

Trough levels of imatinib and nilotinib in plasma were measured by the HPLC method. About 500 μ L of plasma added, then 500 μ L acetonitrile and 50 μ L of the internal standard solution with a concentration of 50 μ g/mL in methanol (the internal standard used for imatinib assay was nilotinib, and for nilotinib assay was imatinib), then centrifuged 6,000 \times g for 10 min. The supernatant obtained was

evaporated to dryness with N_2 gas, then resuspended in a 500 μ L mobile phase, and then injected into the HPLC column. The instrument used was the DAD-Agilent 1100 series HPLC machine with RP-C18 columns. The mobile phase used consisted of 40% solvent A (72.5% water, 25% methanol, and 2.5% triethylamine), 20% methanol, and 40% acetonitrile, which was flowed at a rate of 0.9 mL/min at 35 $^{\circ}$ C and isocratic conditions. Elution was observed at wavelengths of 267 nm [13].

Measurements of BCR-ABL ratios

Three mL of blood sample collected in EDTA vacutainer was stored at a temperature of 2–8 $^{\circ}$ C and processed within 48 h of the initial sampling to avoid degradation of RNA. Measurements of the BCR-ABL ratio in whole blood CML patients were carried out by the RT-qPCR method with the GeneXpert tools. The cartridge preparation process must be carried out within 15 min.

Statistical analysis

The relationship between the imatinib/nilotinib C_{min}^{∞} and BCR-ABL ratio was analyzed by the bivariate correlation Spearman rho test.

Results

Twenty-three imatinib patients and 11 nilotinib patients met the inclusion criteria and signed informed consent. All subjects in the nilotinib group had a history of being treated with imatinib before being treated with nilotinib. Other patient characteristics data are presented in Table 1.

There were 10 (43.48%) patients who used imatinib could reach the C_{min}^{∞} target ($>1,000$ ng/mL). In the nilotinib group, there were 7 (63.64%) patients could achieve the C_{min}^{∞} target (>800 ng/mL). The distribution of C_{min}^{∞} from the two groups is presented in Figure 1.

Table 2 shows that the average C_{min}^{∞} is $1,065.46 \pm 765.71$ ng/mL for imatinib and $1445 \pm 1,010.35$ ng/mL for nilotinib. The data showed there were large interindividual variations in both groups (71.87% vs. 69.88%).

The distribution of the BCR-ABL ratio values of the two groups is shown in Figure 2.

Ten (43.48%) patients who received imatinib achieved BCR-ABL ratio of $\leq 0.1\%$, while in nilotinib group achieved in 7 (63.64%) patients.

Within imatinib group, there was a correlation between C_{min}^{∞} and BCR-ABL ratios ($p=0.043$) and the mean of C_{min}^{∞} in the group of patients who had a BCR-ABL ratio $\leq 0.1\%$ tended to be greater than the group of patients who had a BCR-ABL ratio of $>0.1\%$ (1,381.95 vs. 822.01 ng/mL) (Table 3). In addition, the group of patients who had C_{min}^{∞} imatinib $\geq 1,000$ ng/mL had a 2.564 times greater chance of

Table 1: Basic characteristic of chronic phase-CML patients treated with imatinib or nilotinib.

| Patient's characteristic | | Imatinib | | | Nilotinib | | |
|--------------------------|-------------------|----------------------|------|------------------|----------------------|------|------------------|
| | | Total patient (n=23) | | | Total patient (n=11) | | |
| | | n | % | $\bar{x} \pm SD$ | n | % | $\bar{x} \pm SD$ |
| Sex | Man | 11 | 47.8 | – | 5 | 45.5 | – |
| | Woman | 12 | 52.2 | – | 6 | 54.5 | – |
| Age (when CML diagnosed) | 18–20 y.o | 1 | 4.3 | 46.70 ± 12.60 | – | – | 45.45 ± 13.69 |
| | 21–30 y.o | 3 | 13.1 | Median: 45 y.o | 1 | 9.1 | Median: 46 y.o |
| | 31–40 y.o | 8 | 34.8 | | 4 | 36.4 | |
| | 41–50 y.o | 6 | 26.1 | | 2 | 18.2 | |
| | 51–60 y.o | 4 | 17.4 | | 2 | 18.2 | |
| | 61–70 y.o | 1 | 4.3 | | 2 | 18.2 | |
| Weight | 31–40 kg | 2 | 8.7 | 62.30 ± 14.61 kg | – | – | 59.45 ± 7.61 kg |
| | 41–50 kg | 1 | 4.3 | | – | – | |
| | 51–60 kg | 11 | 47.8 | | 7 | 63.6 | |
| | 61–70 kg | 3 | 13.1 | | 2 | 18.2 | |
| | 71–80 kg | 3 | 13.1 | | 2 | 18.2 | |
| | 81–90 kg | 2 | 8.7 | | – | – | |
| | 91–100 kg | 0 | – | | – | – | |
| | 101–110 kg | 1 | 4.3 | | – | – | |
| | Imatinib duration | 1–5 years | 14 | 60.9 | 4.3 ± 2.08 years | 11 | 100 |
| | 6–10 years | 9 | 39.1 | | – | – | |
| (+ Allopurinol) | Yes | 16 | 69.6 | – | 5 | 45.5 | – |
| | No | 7 | 30.4 | – | 6 | 54.5 | – |




Figure 1: Distribution of imatinib and nilotinib C_{min}^{∞} .

Table 2: Imatinib and nilotinib average C_{min}^{∞} .

| | Patient (n) | Average C_{min}^{∞} (ng/mL) | CV (%) |
|-----------|-------------|------------------------------------|--------|
| Imatinib | 23 | 1,065.46 ± 765.71 | 71.87 |
| Nilotinib | 11 | 1445 ± 1,010.35 | 69.88 |

achieving a decrease in BCR-ABL transcripts compared to the group of patients who had C_{min}^{∞} imatinib <1,000 ng/mL (Figure 3). In contrast, there is no correlation between C_{min}^{∞} and BCR-ABL ratio in the nilotinib group.





Figure 2: BCR-ABL ratio distribution from imatinib and nilotinib group.

Discussion

CML is one of the myeloproliferative disorders in hematopoietic stem cells characterized by leukocytosis (WBC level >100 × 10⁹/L) and splenomegaly. About 90% of CML patients have genetic abnormalities characterized by the presence of the Philadelphia chromosome (Ph) with the fusion BCR-ABL gene. This fusion gene increases tyrosine kinase activity, which contributes to cell transformation

Table 3: Patient's clinical response with BCR-ABL $\leq 0.1\%$ and BCR-ABL $> 0.1\%$.

| | BCR-ABL ratio $\leq 0.1\%$ (MMR) | BCR-ABL ratio $> 0.1\%$ (non-MMR) |
|--|-------------------------------------|--------------------------------------|
| Imatinib | | |
| Average Imatinib C_{min}^{∞} (ng/mL) | 1,381.95 | 822.01 |
| Number of patients with $C_{min}^{\infty} \geq 1,000$ ng/mL | 7 (30.43%) | 3 (13.04%) |
| Number of patients with $C_{min}^{\infty} < 1,000$ ng/mL | 3 (13.04%) | 10 (43.48%) |
| Nilotinib | | |
| Average Nilotinib C_{min}^{∞} (ng/mL) | 1,360.47 | 1,595.29 |
| Number of patients with $C_{min}^{\infty} \geq 800$ ng/mL | 5 (45.45%) | 2 (18.18%) |
| Number of patients with $C_{min}^{\infty} < 800$ ng/mL | 2 (18.18%) | 2 (18.18%) |

**Figure 3:** Imatinib C_{min}^{∞} between patients with BCR-ABL ratio $> 0.1\%$ and patients with BCR-ABL $\leq 0.1\%$.

and causes uncontrolled leukemic cell growth in hematopoietic cells [14].

Imatinib and nilotinib are oral targeted-therapy of TKI groups that selectively occupy the ATP domain in ABL, which inhibits the activity of Bcr-Abl tyrosine kinase protein. Inhibition to substrate phosphorylation will inhibit the signaling pathway, which causes leukemogenesis [15].

Imatinib mesylate dosage of 400 mg/day is a standard therapy in the management of chronic phase of CML. Nilotinib has also been approved as a first-line therapy in chronic phase CML patients, clinically used in CML patients who resistant or tolerant to imatinib. By administering imatinib therapy at a dose of 400 mg/day, CML patients can reach C_{min}^{∞} target (1,000 ng/mL), a C_{min}^{∞} which can inhibit tyrosine kinase activity and therefore induce therapeutic responses. The parameter of molecular

response was the ratio of BCR-ABL [16, 17]. Meanwhile, the recommended target of nilotinib C_{min}^{∞} for TDM practice, in general, is ≥ 800 ng/mL [18, 19].

The results of this study showed that there was high interindividual variability both in imatinib C_{min}^{∞} and nilotinib C_{min}^{∞} . This high variability was also reported in various previous studies. Data from the phase III IRIS study by Larson et al. showed a wide interindividual variability of imatinib C_{min}^{∞} that measured in steady-state, 1 h before the morning dose on 29th day in 351 patients who received a dose of imatinib 400 mg/day. In another study, wide interindividual variability of imatinib C_{min}^{∞} also obtained with the coefficient of variation of 40–60% [10, 20, 21].

Some of the factors that cause this interindividual variability include body size, age, sex, liver and renal function, interaction with other drugs which are taken together, adherence, and enzyme or transporter polymorphisms associated with PK/PD drugs [22]. In this study, there was no therapy other than imatinib/or nilotinib with/without allopurinol, which was administered together by the patient, and there were no drug interactions between them. Compliance has been monitored by interviewing before sampling to ensure there is no missing dose at least 28 days before the day of sampling.

It has been reported in various studies that imatinib has a proportional dose–response relationship. The results of correlation analysis between C_{min}^{∞} and molecular response (BCR-ABL ratio) showed that there was a correlation in the imatinib group [20].

The mean of imatinib C_{min}^{∞} in patients who achieved the BCR-ABL ratio target has been found higher than the mean of imatinib C_{min}^{∞} in patients that did not reach the target BCR-ABL ratio (1,065.46 ng/mL vs. 822.01 ng/mL). The $C_{min}^{\infty} \geq 1,000$ ng/mL was associated with an increased chance of achieving a BCR-ABL ratio of $\leq 0.1\%$ (major molecular response or MMR). These results support the results of previous studies, which reported that an increase in imatinib C_{min}^{∞} correlated with a decrease in BCR-ABL transcripts characterized by a BCR-ABL ratio of $\leq 0.1\%$ [20, 22]. BCR-ABL ratio $\leq 0.1\%$ (MMR) can indicate a decrease in the progression toward an acceleration or blastic phase. Unlike imatinib, in the nilotinib group, there is no correlation between C_{min}^{∞} and BCR-ABL ratio even though higher C_{min}^{∞} tends to have a higher MMR level. The absence of a correlation between nilotinib exposure and the response was thought to be due to the wide interindividual variability of the nilotinib pharmacokinetic parameters. Larson also revealed that the relationship between nilotinib exposure and clinical response was not as clear as that observed in patients with imatinib. This condition may be due to nilotinib being more potent than imatinib [11]. In a

retrospective analysis of the Evaluating Nilotinib Efficacy and Safety in clinical Trials-newly diagnosed (ENESTnd) study, it was found that there was a faster decrease of BCR-ABL ratio in the nilotinib group. The 6th month of the median BCR-ABL ratio in the nilotinib group was found to be similar to the 18th month of BCR-ABL ratio in the imatinib group. There were reported to be more patients with nilotinib doses of 300 and 400 mg, which achieved a BCR-ABL ratio of <10% in the 3rd month compared to imatinib (74, 78, and 61%) [23]. The results of the Exploring Nilotinib BCR-ABL Effects (ENABL) study in the US on patients with suboptimal response to imatinib therapy showed that a decrease in the BCR-ABL ratio occurred fastest in the first 3–9 months after nilotinib use. After that, it was tended to be a plateau [24]. In the present study, the BCR-ABL ratio was observed after the use of nilotinib for more than 12 months, while the majority of patients had used nilotinib for more than 24 months and 7 (63.63%) patients had achieved a BCR-ABL ratio of $\leq 0.1\%$. Allegedly in the period of the implementation of the study, the BCR-ABL ratio has entered the plateau period so that it may cause no correlation between C_{min}^{∞} and the BCR-ABL ratio.

The distribution profile of the BCR-ABL/ABL ratio achieved by patients in the nilotinib group (Figure 2) showed that some patients experienced suboptimal responses. According to the guidelines of the European LeukemiaNet (ELN), all responses under the conditions of a major molecular response (BCR-ABL $\leq 0.1\%$ IS) at the 18th month after the start of TKI therapy are considered suboptimal [25]. Reflecting on previous studies, nilotinib $C_{min}^{\infty} \geq 800$ ng/mL was associated with the achievement of the BCR-ABL ratio $\leq 0.1\%$ IS in the 12th month, which indicates that nilotinib can inhibit tyrosine kinase activity well [19, 26]. This may explain that patients with nilotinib $C_{min}^{\infty} \geq 800$ ng/mL can achieve a BCR-ABL ratio of $\leq 0.1\%$. The phenomenon of having a patient with high nilotinib C_{min}^{∞} , but still unable to achieve the BCR-ABL ratio $\leq 0.1\%$ was thought to be due to TKI resistance.

Leukemia cell resistance to TKI can occur due to genetic mutations in the ABL tyrosine kinase domain and changes in expression of influx–efflux transporters [27]. Patients can experience multiple mutations in the BCR-ABL gene at the same time and result in heavier resistance. More than 90 different mutations have been found in different kinase domains [28]. *In-vitro* studies result showed that nilotinib has activity against almost all cell cultures that are resistant to imatinib due to BCR-ABL mutations, except for T315I mutants [29]. In contrast to imatinib, which requires the human organic cation transporter-1 (h-OCT1) to be uptaken into leukemia cells, nilotinib activity was found to be independent from h-OCT1 [30]. In contrast, various

studies have shown the role of ATP Binding Cassette Subfamily B member 1 (ABCB1) and ATP Binding Cassette Subfamily G member 2 (ABCG2) in the efflux mechanism of nilotinib from leukemia cells. In one study, it was found that the inhibition capacity of nilotinib tyrosine kinase decreased with the increasing levels of ABCB1. Therefore, ABCB1 overexpression can play a role in causing resistance to nilotinib [31–33]. Further research is needed to see mutations in the BCR-ABL gene and the expression of ABCB1 transporters in this group of patients to determine the possibility of nilotinib resistance.

Conclusion

There were high interindividual variations of imatinib and nilotinib C_{min}^{∞} in CML patients. Imatinib C_{min}^{∞} correlated with BCR-ABL ratios, but no correlation in nilotinib.

Acknowledgment: We would like to thank Dr. Soetomo General Hospital for granting permission to use the data provided in this study and the Tahir Foundation.

Research funding: This study was funded by the Minister of Research Technology and Higher Education of Indonesia and the Tahir Foundation.

Author contributions: All authors have accepted responsibility for the entire content of this manuscript and approved its submission.


Conflict of interest: The authors state that they have no conflict of interest.

Informed consent: Informed consent was obtained from all individuals included in this study.

References

1. Iqbal N, Iqbal N. Imatinib: a breakthrough of targeted therapy in cancer. *Chemother Res Pract* 2014;1–9. <https://doi.org/10.1155/2014/357027>.
2. Hamad A, Sahli Z, Sabban ME, Mouterik M, Nasr R. Emerging therapeutic strategies for targeting chronic myeloid leukemia stem cells. *Stem Cells Inter* 2013;2013:1–12.
3. Quintas-Cardama A, Cortes J, Kantarjian H, O'Brien S. Chronic myelogenous leukemia. In: Provan D, Gribben J, editors. *Molecular hematology*. UK: Wiley-Blackwell; 2010:76–87.
4. Savage DG, Antman KH. Imatinib mesylate-A new oral targeted therapy. *N Engl J Med* 2002;346:683–93.
5. Biswal S. Novel agents in CML therapy: tyrosine kinase inhibitors and beyond. *WebMedCentral* 2012;3:1–12.
6. Kantarjian HM, Talpaz M, Giles F, O'Brien S, Cortes J. New insights into the pathophysiology of chronic myeloid leukemia and imatinib resistance. *Ann Intern Med* 2006;145:913–23.
7. Deiniger MW. Nilotinib CCR drug updates. *Clin Cancer Res* 2008; 14:4027–31.
8. Hardling M, Wei Y, Palmqvist L, Swolin B, Stockelberg D, Gustavsson B, et al. Serial monitoring of BCR-ABL transcripts in

- chronic myelogenous leukemia (CML) treated with imatinib mesylate. *Med Oncol* 2004;21:349–58.
9. Bintoro SU. Sejarah CML, Karakteristik Klinik CML, Resistensi terhadap imatinib. Dalam: Su B, editor. *Chronic myelogenous leukemia: patogenesis, diagnosis dan terapi*. Surabaya: Global Persada Press; 2014:1-26.
 10. Larson RA, Druker BJ, Guilhot F, O'Brien SG, Riviere GJ, Krahnke T, et al. Imatinib pharmacokinetics and its correlation with response and safety in chronic-phase chronic myeloid leukemia: a subanalysis of the IRIS study. *Am Soc Hematol* 2008;111:4022–8.
 11. Larson RA, Yin OQP, Hochhaus A, Saglio G, Clark RE, Nakamae H, et al. Population pharmacokinetic and exposure-response analysis of nilotinib in patients with newly diagnosed Ph+ chronic myeloid leukemia in chronic phase. *Eur J Clin Pharmacol* 2012;68:723–33.
 12. Takahashi N, Miura M, Scott SA, Kagaya H, Kameoka Y, Tagawa H, et al. Influence of CYP3A5 and drug transporters polymorphisms on imatinib trough concentration and clinical response among patients with chronic phase chronic myeloid leukemia. *J Hum Genet* 2010;55:731–7.
 13. Pirro E, Francia SD, Martino FD, Fava C, Ulisciani S, Cambrin GR, et al. A new HPLC-UV validated method for therapeutic drug monitoring of tyrosine kinase inhibitors in leukemic patients. *J Chromatogr Sci* 2011;49:753–7.
 14. Goldman JM, Melo JV. Chronic myeloid leukemia – advances in biology and new approaches to treatment. *NEJM* 2003;349:1451–61.
 15. Patel D, Suthar MP, Patel V, Singh R. BCR ABL kinase inhibitors for cancer therapy. *Int J Pharm Sci Drug Res* 2010;2:80–90.
 16. Ishikawa Y, Kiyoi H, Watanabe K, Miyamura K, Nakano Y, Kitamura K, et al. Trough plasma concentration of imatinib reflects BCR-ABL kinase inhibitory activity and clinical response in chronic-phase chronic myeloid leukemia: a report from the BINGO study. *Cancer Sci* 2010;101:2186–92.
 17. Pavon V, Gomez R, Jaime JC, Hernandez P, Arencibia A, Espinosa-Martinez E. Introduction of imatinib as first-line therapy for chronic myeloid leukemia in Cuba. *MEDICC Rev* 2011;13:35–9.
 18. Gao B, Yeap S, Clements A, Balakrishnar B, Wong M, Gurney H. Evidence for therapeutic drug monitoring of targeted anticancer therapies. *J Clin Oncol* 2012;30:4017–22.
 19. Miura M. Therapeutic drug monitoring of imatinib, nilotinib and dasatinib for patients with chronic myeloid leukemia. *Biol Pharm Bull* 2015;38:645–54.
 20. Picard S, Titier K, Etienne G, Teilhet E, Ducint D, Bernard M, et al. Trough imatinib plasma levels are associated with both cytogenetic and molecular responses to standard-dose imatinib in chronic myeloid leukemia. *Blood* 2007;109:3496–8.
 21. Peng B, Hayes M, Resta D, Racine-Poon A, Druker BJ, Talpaz M, et al. Pharmacokinetics and pharmacodynamics of imatinib in a phase I trial with chronic myeloid leukemia patients. *J Clin Oncol* 2004;22:935–42.
 22. Takahashi N, Wakita H, Miura M, Scott SA, Nishii K, Masuko M, et al. Correlation between imatinib pharmacokinetics and clinical response in Japanese patients with chronic –phase chronic myeloid leukemia. *Clin Pharmacol Ther* 2010;88: 809–13.
 23. Savona MR, Saglio G. Identifying the time to change BCR-ABL inhibitor therapy in patients with chronic myeloid leukemia. *Acta Haematol* 2013;130:268–78.
 24. Ailawadhy S, Akard LP, Miller CB, Jillella A, DeAngelo DJ, Ericson SG. Exploratory study on the impact of switching to nilotinib in 18 patients with chronic myeloid leukemia in chronic phase with suboptimal response to imatinib. *Ther Adv Hematol* 2017;8:3–12.
 25. Baccarani M, Deininger MW, Rosti G, Hochhaus A, Soverini S, Apperley JF. European leukemia net recommendations for the management of chronic myeloid leukemia. *Blood* 2013;122: 872–80.
 26. Takahashi N, Miura M, Kuroki J, Mitani K, Kitabayashi A, Sasaki O, et al. Multicenter phase II clinical trial of nilotinib for patients with imatinib-resistant or –intolerant chronic myeloid leukemia from the east Japan CML Study Group evaluation of molecular response and the efficacy and safety of nilotinib. *Biomark Res* 2014;2:1–9.
 27. Galinsky I, Buchanan S. Guide to interpreting disease responses in chronic myeloid leukemia. *J Adv Pract Oncol* 2012;3:225–36.
 28. Jabbour E, Parikh SA, Kantarjian H, Cortes J. Chronic myeloid leukemia – mechanism of resistance and treatment. *Hematol Oncol Clin North Am* 2015;25:981.
 29. Jabbour E, Cortes J, Kantarjian H. Nilotinib for the treatment of chronic myeloid leukemia: an evidence-based review. *Core Evid* 2009;4:207–13.
 30. van Erp NP, Gelderblom H, Guchelaar HJ. Clinical pharmacokinetics of tyrosine kinase inhibitors. *Cancer Treat Rev* 2009;35:692–706.
 31. Dohse M, Scharenberg C, Shukla S, Robey RW, Volkmann T, Deeken JF, et al. Comparison of ATP-binding cassette transporter interactions with the tyrosine kinase inhibitors imatinib, nilotinib and dasatinib. *Drug Metab Dispos* 2010;38:1371–80.
 32. Eadie LN, Saunders VA, Hughes TP, White DL. Degree of kinase inhibition achieved in vitro by imatinib and nilotinib is decreased by high levels of ABCB1 but not ABCG2. *Leuk Lymphoma* 2013;54: 569–78.
 33. Eadie LN, Hughes TP, White DL. Interaction of the efflux transporter ABCB1 and ABCG2 with imatinib, nilotinib and dasatinib. *Clin Pharmacol Ther* 2014;95:294–306.



Journal of Basic and Clinical Physiology and Pharmacology

Editor In Chief: Michal Horowitz

Editorial Board: Karen Avraham, Kusal K. Das, Yoram Epstein, Elliot S. Gershon MD, Einat Kodesh, Ron Kohen, David Lichtstein, Alina Maloyan, Raphael Mechoulam, Joachim Roth, Suzanne Schneider, Esther Shohami, Haim Sohmer, Toshikazu Yoshikawa, and Joseph Tam



Editor-in-Chief

Michal Horowitz

m.horowitz@mail.huji.ac.il

Editorial Board

Karen Avraham, Tel Aviv University, Tel Aviv, Israel

Kusal Das, BLDE University, Bijapur, India

Yoram Epstein, Tel Aviv, Tel Aviv University, Israel

Elliot S. Gershon, The University of Chicago Medicine, Chicago, IL

Einat Kodesh, University of Haifa, Haifa, Israel

Ron Kohen, Jerusalem, Hebrew University of Jerusalem, Israel

David Lichtstein, Hebrew University Jerusalem, Jerusalem, Israel

Alina Maloyan, University of Texas Health Science Center, San Antonio, TX

Raphael Mechoulam, The Hebrew University Jerusalem, Jerusalem, Israel

Joachim Roth, Justus-Liebig-University Giessen, Giessen, Germany

Suzanne Schneider, University of New Mexico, Albuquerque, NM

Esther Shohami, Hebrew University of Jerusalem, Jerusalem, Israel

Haim Sohmer, Hebrew University of Jerusalem, Jerusalem, Israel

Joseph Tam, Hebrew University of Jerusalem, Jerusalem, Israel

Editorial Office

Katharina Appelt

Walter de Gruyter GmbH

Genthiner Str. 13

10785 Berlin

Germany

Tel. +49-30-26005-325

E-mail: jbcpp.editorial@degruyter.com

€99.00

Online

ADD TO CART

INSTITUTIONAL PRICING 

SEARCH

Search within Journal...



Volume

Issue

Page



DETAILS



Online ISSN: 2191-0286

Type: Journal

Language: English

Publisher: De Gruyter

First published: 01 Dec 1986

Publication frequency: 6 issues per year

Audience: researchers and health professionals in the field of clinical physiology and pharmacology

De Gruyter: Your publication is now available online

1 message

noreply.degruyter@degruyter.com <noreply.degruyter@degruyter.com>
To: budi-s@ff.unair.ac.id

Sat, Aug 8, 2020 at 11:13 AM



Dear author,

We are pleased to inform you that your following publication is now published on <https://www.degruyter.com> and available free of charge to you as author.



Relationship between trough level of tyrosine kinase inhibitor (imatinib and nilotinib) and BCR-ABL ratios in an Indonesian chronic-phase chronic myeloid leukemia (CML) population

Journal of Basic and Clinical Physiology and Pharmacology

ISSN: 2191-0286

FREE ONLINE ACCESS TO YOUR PUBLICATION**Step 1:**

New to our website?

- Please [register](#) for free.
- **Important:** Use the email address this notification was sent to.

Already have a De Gruyter account with this email address?

- Please [log in](#).

Step 2:

After logging in, you can view your publications under <https://www.degruyter.com/profile/myworks>. Note that the PDF versions are intended for your personal use only.

HELP

Problems? Questions? Please look at our [help page](#). There you can also find the contact details of our customer service.

Thank you for choosing us as a publisher. We look forward to working with you again in the future.

Kind regards,
De Gruyter

To help ensure continued delivery of important notices from De Gruyter, please add noreply.degruyter@degruyter.com to your address book or approved sender list.



Customer service
Unsubscribe here
Privacy statement
Legal information
degruyter.com

jovis