

Effect of Atorvastatin Treatment on Vascular Aterogenic Factors (Lipid Profiles and VCAM-1) in Patient Diabetes with Dyslipidemia

by Budi Suprapti

Submission date: 18-Mar-2020 11:54AM (UTC+0800)

Submission ID: 1277385669

File name: id_Profiles_and_VCAM-1_in_Patient_Diabetes_with_Dyslipidemia.pdf (863.75K)

Word count: 2325

Character count: 12615

Effect of Atorvastatin Treatment on Vascular Atherogenic Factors (Lipid Profiles and VCAM-1) in Patient Diabetes with Dyslipidemia

Asri Wido Mukti¹, Budi Suprapti¹, and Sony Wibisono²

¹ Magister of Clinical Pharmacy, Universitas Airlangga, Surabaya, Indonesia

² Division Endocrinology and Metabolism - Department of Internal Medicine, Dr. Soetomo Teaching Hospital School of Medicine Airlangga University

Info Article

Submitted: 23-04-2018

Revised: 18-03-2019

Accepted: 09-04-2019

*Corresponding author
Asri Wido Mukti

Email
ninoasri@gmail.com

ABSTRACT

To analyze effectiveness of atorvastatin 20mg on lipid profiles and adhesion molecule VCAM-1 in patient with diabetes dyslipidemia. An observational prospective cohort study was conducted from November 2016 to March 2017. Patients who fulfilled the inclusion criteria were taken twice for their lipid profiles and VCAM-1 measurements (before initiation of study and after 6 weeks treatment of atorvastatin 20mg). There were 13 patients who met the inclusion criteria. The results of 13 patients showed that after 6 weeks of atorvastatin therapy, there was a 28% decrease in total cholesterol ($t_0=223.77\pm49.69$, $t_1=160.92\pm24.69$), 39% LDL decrease ($t_0=152.59\pm44.25$, $t_1=93\pm21.44$), a decrease in TG 38.6% ($t_0=200.85\pm101.53$, $t_1=123.30\pm62.77$) and a statistically significant decrease in VCAM-1 7.47% ($t_0=729.59\pm208.06$, $t_1=675.06\pm182.88$). The results of the correlation test between total cholesterol and VCAM-1 ($p=0.185$, $r=0.268$), LDL and VCAM-1 ($p=0.127$, $r=0.307$), TG and VCAM-1 ($p=0.198$, $r=0.261$) showed no correlation. Based on the results of the study, it can be concluded that atorvastatin therapy can provide improvements in atherogenic factors such as decreased lipid profile and VCAM-1, and there was no correlation between lipid profile and VCAM-1 in type 2 DM patients with dyslipidemia.

Keywords : Atorvastatin, Adhesion molecules, Diabetes mellitus, , dyslipidemia, VCAM-1.

8

INTRODUCTION

Diabetes mellitus (DM) is a metabolic disorder disease characterized by hyperglycemia (Triplitt *et al.*, 2014). The most common lipid pattern in people with type 2 diabetes consists of hypertriglyceridemia (hyper-TG), low high-density lipoprotein cholesterol (HDL-C), and relatively normal plasma concentrations of low-density lipoprotein cholesterol (LDL-C) (Hegele *et al.*, 2013). The condition of hyperglycemia and increased free fatty acids can cause endothelial damage, endothelial become more permeable, presence of reactive oxygen species (ROS), and increase adhesion molecules (George and Lyon, 2010). One of the adhesion molecules which has increased is VCAM-1. VCAM-1 is a member of the protein of immunoglobulin superfamily which is the framework for the migration of leukocytes to subendothelial space. If this leukocyte migration

not prevented will cause monocytes to mature into macrophages and form foam cell and fatty streak that will cause atherosclerosis (Wang and Huo, 2010).

Levels of soluble adhesion molecules have been shown to correlate with various cardiovascular risk factors including hypercholesterolemia and or hypertriglyceride, decreased HDL, hypertension, DM, and smoking. In two prospective studies of patients with type 2 diabetes, VCAM-1 became a strong predictor of cardiovascular mortality. VCAM-1 levels are also significantly associated with intima-media thickness, an index of early atherosclerotic (Wang and Huo, 2010).

Atorvastatin is a class of lipid-lowering agent, HMG CoA reductase inhibitor. HMG CoA is a precursor of mevalonate formation which is a synthesis of cholesterol (Malloy and Kane, 2012).

Atorvastatin also has a pleiotropic effect that can inhibit inflammation by inhibition of leukocytes by decreasing the expression of adhesion molecules, therefore the aim of this study was to analyze the effect of atorvastatin on lipid profile and pleiotropic effects as inhibition of inflammation.

MATERIAL AND METHODS

This study conducted a prospective observational design from November 2016 to March 2017 was approved by Ethics Committee of the faculty of Medicine Airlangga University Surabaya. We included patients aged 35-79 years who were diagnosed with diabetic dyslipidemia, HbA1c 8-13%, lipid profiles during initiation of LDL therapy >100mg/dL and / or TG>150 mg/dL, free or at least 2-3 weeks after acute inflammation, and have not received atorvastatin therapy previously. Patients with elevated liver function tests (>3x normal values), patients with severe renal failure, patients with nephrotic syndrome with dyslipidemia, and patients taking progestin hormones, corticosteroids, and antiretroviral drugs (protease inhibitor class) were excluded from the study. Blood measurements were performed on lipid profiles (total cholesterol, LDL, and TG) and VCAM-1 before therapy of atorvastatin 20mg and after 6 weeks therapy. VCAM-1 were measured with ELISA method.

Analysis data started with normality test with Saphiro Wilk test. A comparative hypothesis test was performed to determine the differences of lipid profiles and VCAM-1 before therapy (t_0) and after therapy (t_1) with paired sample T-test and Wilcoxon test. In addition, a correlation test was performed to determine the correlation between lipid profile and VCAM-1 by using spearman correlation test.

RESULTS AND DISCUSSION

The number of male and female patients is almost equal with the most age range in the middle age range (46-59 years) (Table I). Based on research conducted by Mihardja et al in Indonesia also shows that the prevalence of DM increases with age with the highest age range 45-55 years (Mihardja et al, 2014). HbA1c levels indicate uncontrolled sugar ranges (8-13%) and most patients are overweight (BMI 25-18kg/m²). Overweight and obesity are the most potent risk factors for DM (WHO, 2016). Three patients had comorbidities of hypertension and two patients with CHD. In a study conducted by Lastra (et al.) also found that hypertension was found in more

than 50% of patients with DM and contributed significantly to micro and macrovascular complications in DM. The risk factor for cardiovascular disease is 4 times higher in DM patients with hypertension (Lastra et al., 2014).

The NCEP-ATP III guidelines divided LDL and TG levels within the range that can be used as indicative references when lipid-lowering therapy may be given. When TG levels are within a borderline high range, LDL-C is the main therapeutic target. Similarly, when TG levels are in the high range, LDL-C is still a major therapeutic target. Add lipid-lowering drugs such as fibrates to help reduce them. Individuals with very high TG levels (>500mg/dL) are at risk for acute pancreatitis, therefore the primary target is the reduction of TG first with a low-fat diet, weight management and physical activity, and the use of fibrates or nicotinic acid. If TG levels reach <500 mg/dl then it can be replaced again with LDL-lowering therapy (NCEPATP III, 2002).

Most patients had total cholesterol data in the range of borderline high (200-239mg/dL) are 62% with an average of 223.77±49.69mg/dL, LDL in the borderline high range (130-159mg/dL) and high (160-189mg/dL) are 38% with an average of 152.59±44.25mg/dL, triglycerides in the optimal range (<150mg/dl) are 54% with mean 200.85±101.53mg/dL, the patient's HDL level was mostly at the level of >40mg/dL (69%) with an average of 44.21±9.97mg/dL. Diabetes with dyslipemia usually characterized by three conditions: high triglyceride concentrations, low HDL concentrations, and high small dense LDL concentrations (Chehade et al., 2013). The baseline HDL values of patients were different than in those where only 31% of patients had low HDL values (<40mg/dL) (Table II).

It was found that after 6 weeks of treatment, there was a significant decrease in lipid profile (cholesterol, LDL, and TG) and VCAM-1 levels ($p < 0,05$). These results are consistent with previous studies conducted by Hogue et al, suggesting that the administration of atorvastatin 20mg for 6 weeks potentially altering lipid profile and also reduce inflammation (lowering CRP levels), oxidation (lowering levels of ox-LDL), and monocyte adhesion (Lowering levels of sICAM-1, sVCAM-1, sEselectin) in DM patients with hypertriglyceride (Hogue et al., 2008) (Table III).

Correlation test is conducted to know the relationship between lipid profile level and VCAM-1. VCAM-1 increase in DM patients has been reported in previous studies, which is related to the

Table I. Baseline characteristic of patients

Patient Characteristics	Total Patients (N=13)		$\bar{x} \pm SD$ (range)	
	n	%		
12 Gender	Male	6	46%	-
	Female	7	54%	-
Age	36-45 years	1	8%	54.84±6.70
	46-59 years	10	77%	
	60-74 years	2	15%	
HbA1c	8-10 %	7	54%	10.75±1.68
	11-13 %	6	46%	
6 BMI	18.5-24.9 kg/m ²	5	38%	26.79±5.11
	25.0-29.0 kg/m ²	6	46%	
	>30 kg/m ²	2	16%	
Comorbidities	Hypertension	3	23%	-
	CHD	2	15%	-

Table II. Baseline profil lipid pasien (t₀)

Levels of Lipid Profile	t ₀		t ₁		
	n	%	n	%	
Total Cholesterol	Desirable (<200mg/dL)	3	23%	12	92%
	Borderline High (200-239mg/dL)	8	62%	1	8%
	High (≥240mg/dL) 3	2	15%	-	-
	Optimal (<100mg/dL)	1	8%	9	69%
LDL	Near optimal (100-129 mg/dL)	1	8%	3	23%
	Borderline high (130-159 mg/dL)	5	38%	1	8%
	High (160-189mg/dL)	5	38%	-	-
	Very high (≥190mg/dL)	1	8%	-	-
TG	Optimal (<150mg/dL)	7	54%	10	77%
	Near optimal (150-129mg/dL)	-	-	-	-
	Borderline high (200-499mg/dL)	6	46%	3	23%
HDL	High (≥500mg/dL)	-	-	-	-
	<40mg/dL	4	31%	3	23%
	>40mg/dL	9	69%	10	77%

Table III. Average changes in lipid profile and VCAM-1

Lipid Profiles	Mean ± SD		P
	t ₀	t ₁	
Total Cholesterol	223.77±49.69 (165-369)	160.92±24.69 (126-205)	0.002
LDL	152.59±44.25 (79-266)	93±21.44 (60-136)	0.001
TG	200.85±101.53 (103-377)	123.30±62.77 (68-246)	0.016
VCAM-1	729.59±208.06	675.06±182.88	0.019

severity of atherosclerotic disease (Fasching *et al.*, 1996). The result of correlation analysis between serum lipid profile level with serum VCAM-1 level (Figure 1, 2 and 3) showed no significant correlation (p>0.05). Another study from Hubel *et*

al found that VCAM-1 correlated with LDL (r = 0.50, P<0.03), but did not show any association with other lipid components (Hubel *et al.*, 1998). Wh4 in the study in Indonesia showed only correlation in patients with type 2 diabetes showed a tendency of




Figure 1. Correlation between total cholesterol and VCAM-1




Figure 2. Correlation Between LDL and VCAM-1




Figure 3. Correlation Between Triglycerida and VCAM-1

elevated plasma VCAM-1 levels along with ¹¹ increase of albuminuria (Wibisono *et al.*, 2012). The limitation of this study is the relatively small sample size (13 patients), and the variation of lipid profiles among patients is very large because it is influenced by many factor. This can cause statistical results that are not meaningful.

CONCLUSION

Atorvastatin therapy is able to improved atherogenic factors such as decreased lipid profile (total cholesterol, LDL, TG) and VCAM-1

inflammatory markers in DM patients with dyslipidemia, and no correlation between lipid profile (total cholesterol, LDL, TG) and VCAM -1.

ACKNOWLEDGEMENT

All of subjects are acknowledged for participating in this study. Sony Wibisono, DR. dr., SpPD, KEMD., FINASIM is acknowledgment for constructing comments in manuscript. The authors also thank you to members of Magister of Clinical Pharmacy, Airlangga University, Surabaya and private clinic of endocrine consultant in Surabaya.

REFERENCES

- Cehade JM., Gladysz M. and Mooradian AD., 2013. Dyslipidemia in Type 2 Diabetes : Prevalence, Pathophysiology, and Management. *Drugs* 73:327-339.
- Fasching P, Veitl M, Rohac M, *et al*, 1996. Elevated concentrations of circulating adhesion molecules and their association with microvascular complications in insulin-dependent diabetes mellitus. *Journal of Clinical Endocrinology and Metabolism* ;81:4313-17.
- George SJ., and Lyon C., 2010. Pathogenesis of Atherosclerosis, In : *Atherosclerosis Molecular and Cellular Mechanisms*. Weinheim : WILEY-VCH Verlag GmbH & Co. KGaA, pp-3-14.
- John MGB, Hegele RA., and Leiter LA., 2013. Dyslipidemia. *Canadian Journal of Diabetes* 37 : S110eS116.
- Hogue JC., Lamarche B., Tremblay AJ., Bergeron J., Gagné C. and Couture P. 2008. Differential effect of atorvastatin and fenofibrate on plasma oxidized low-density lipoprotein, inflammation markers, and cell adhesion molecules in patients with type 2 diabetes mellitus. *Metabolism Clinical and Experimental* 57. 380 – 386.
- Hubel CA., Fiona L., Lisa W., Robin E. Gandley; and James M. Roberts, 1998. Small Low-Density Lipoproteins and Vascular Cell Adhesion Molecule-1 Are Increased in Association With Hyperlipidemia in Preeclampsia. *Metabolism*. 47, No 10(October),1998:pp 1281-1288.
- Lastra G., Syed S., Romaine LK., MD., and Camila M., James RS., 2014. Type 2 diabetes mellitus and hypertension: An update. *Endocrinology and Metabolism Clinics of North America* 2014 March ; 43(1): 103-122.
- Malloy MJ. and John PK., 2012. Agents Used in Hyperlipidemia In : *Basic & Clinical Pharmacology* 10th edition. USA : McGraw-Hill. pp. 625
- Laurentia M., Soetrisno U. and Sidartawan Soegondo, 2014. Prevalence and clinical profile of diabetes mellitus in productive aged urban Indonesians. *Journal of Diabetes Investigation* 2014; 5: 507-512.
- NCEP-expert-panel, third report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report, *Circulation* 106 (2002) 3143-3421 Ohtawa, M., Masuda, N., Akasaka, I., et al., Cellular uptake of fluvastatin, an inhibitor of HMG-CoA reductase, by rat cultured hepatocytes and human aortic endothelial cells. *British Journal of Clinical Pharmacology*, 1999, 47:383-389.
- Triplitt CL., Repas T. and Alvarez C., 2014. Diabetes Mellitus, In : *Pharmacotherapy A Pathophysiologic Approach* 9th edition . USA : McGraw Hill. pp 643-647
- Huang W. and Yuqing ., 2010. Adhesion Molecules and Atherosclerosis, In : *Atherosclerosis Molecular and Cellular Mechanisms*. Weinheim : WILEY-VCH Verlag GmbH & Co. KGaA, pp 43-55
- Wibisono C., Fajar A., Tjokroprawiro A., Soetjahjo A., Pranoto A., Adi S. and Murtiwi S., 2012. Profil Albuminuria dan Kadar sVCAM-1 Pada Penderita Diabetes Melitus Tipe 2. *Jurnal Penyakit Dalam* Vol 13 Nomor 1 Januari. Hal 28-35.

Effect of Atorvastatin Treatment on Vascular Aterogenic Factors (Lipid Profiles and VCAM-1) in Patient Diabetes with Dyslipidemia

ORIGINALITY REPORT

14%

SIMILARITY INDEX

10%

INTERNET SOURCES

11%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	issuu.com Internet Source	2%
2	Systems Biology of Free Radicals and Antioxidants, 2014. Publication	2%
3	Fernando Fonseca, Marzia Macedo, Marisa da Cunha Gonzaga, Carlos D'Aparecida Machado, Anna Batista. "Study of lipid profile in adult women with acne", Clinical, Cosmetic and Investigational Dermatology, 2015 Publication	1%
4	www.acpm.org Internet Source	1%
5	run.unl.pt Internet Source	1%
6	www.uspreventiveservicestaskforce.org Internet Source	1%
7	llnutrition.com Internet Source	

1%

8 ijcep.com
Internet Source

1%

9 www.effcohealth.org
Internet Source

1%

10 www.faqs.org
Internet Source

1%

11 Márta Kasza, J. Meleg, J. Vardai, B. Nagy, E. Szalai, J. Damjanovich, A. Csutak, B. Ujhelyi, V. Nagy. "Plasma E-selectin levels can play a role in the development of diabetic retinopathy", Graefe's Archive for Clinical and Experimental Ophthalmology, 2016
Publication

1%

12 creativecommons.org
Internet Source

1%

13 www.dovepress.com
Internet Source

1%

14 cdn.intechopen.com
Internet Source

1%

15 Mariusz Nowak. "Blood serum levels of vascular cell adhesion molecule (sVCAM-1), intercellular adhesion molecule (sICAM-1) and endothelial leucocyte adhesion molecule-1 (ELAM-1) in

<1%

diabetic retinopathy", Clinical and Experimental
Medicine, 09/2008

Publication

16

link.springer.com

Internet Source

<1%

17

Gelaye, Bizu, Michelle A. Williams, Seblewengel
Lemma, Yemane Berhane, Jesse R. Fann, Ann
Vander Stoep, and Xiao-Hua Andrew Zhou.

"Major depressive disorder and cardiometabolic
disease risk among sub-Saharan African
adults", Diabetes & Metabolic Syndrome Clinical
Research & Reviews, 2015.

Publication

<1%

18

Renier, G.. "Monocyte adhesion in diabetic
angiopathy", Journal of Diabetes and Its
Complications, 200303/04

Publication

<1%

19

P. K. Tingsgaard, T. Bock, P. L. La. "Topical
Budesonide Treatment Reduces Endothelial
Expression of Intercellular Adhesion Molecules
(Vascular Cell Adhesion Molecule-1 and P-
selectin) and Eosinophil Infiltration in Nasal
Polyps", Acta Oto-Laryngologica, 2009

Publication

<1%

20

"Systems Biology of Free Radicals and
Antioxidants", Springer Science and Business
Media LLC, 2014

<1%

21

Breuer, Hans-Willi M.. "Hypertriglyceridemia: A Review of Clinical Relevance and Treatment Options: Focus on Cerivastatin", Current Medical Research and Opinion, 2001.

<1%

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Effect of Atorvastatin Treatment on Vascular Aterogenic Factors (Lipid Profiles and VCAM-1) in Patient Diabetes with Dyslipidemia

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
