



Pharmacognosy Research

Publication of Pharmacognosy Network Worldwide

www.phcogres.com




Medknow

 Wolters Kluwer

Phcog.Net - Bringing Medicinal Plant Researchers Together

Caspar, CNKI (China National Knowledge Infrastructure), DOAJ, EBSCO Publishing's Electronic Databases, Excerpta Medica / EMBASE, Genamics JournalSeek, Google Scholar, Health & Wellness Research Center, Health Reference Center Academic, Hinari, Index Copernicus, National Science Library, OpenJGate, PrimoCentral, ProQuest, PubMed, Pubmed Central, Scimago Journal Ranking, SCOLAR, SCOPUS, SIIC databases, Summon by Serial Solutions and Ulrich's International Periodical Directory.



Users Online: 1964

e-19-4

Search Article

Search

Advanced search

[Submit Articles On-line](#) [E-mail Alerts](#) [Access Statistics](#) [Join Us](#)

Editorial Board

Editor—in-Chief

[Dr. Srisailam Keshetti, Ph.D](#)

University College of Pharmaceutical Sciences,
Satavahana University, Karimnagar, INDIA

Associate Editors

[Sushil Kumar Middha](#) Ph. D.

DBT-BIF facility
MLACW, Bangalore

[Dr. T. S. Mohamed Saleem](#), Ph.D.

Head, Department of Pharmacology & Pharmacy Practice
Annamacharya College of Pharmacy
New Boyanapalli
Rajampet-516126, A.P. India

[Dr. Tapas Sar](#)

Head, Department of Veterinary Pharmacology and Toxicology
West Bengal University of Animal and Fishery Sciences
37, K. B. Sarani, Kolkata - 700037

[Ho Yen Fong](#)

Kuala Lumpur, Malaysia.

Asst. Editor

Ms. Madhuri P

Phcog.Net
17, II Floor, Buddha Vihar Road,
Cox Town, Bengaluru,
Karnataka, INDIA

Editorial Board

[Dr. Ciddi Veeresham](#) Ph.D.

Professor of Pharmacy,
University College of Pharmaceutical Sciences,
Kakatiya University, Warangal,
Telangana 506009 India,

[Dr. R.M. Dharmadasa](#) Ph.D.

Research Fellow, Herbal Technology Section,
Industrial Technology Institute,
363, Baudhaloka Mawatha, Colombo 7,
Sri Lanka

[Dr. Mohd Sajjad Ahmad Khan](#), Ph.D.

Department of Biology, College of Medicine,
University of Dammam,
Dammam 31452, Saudi Arabia.

[Dr. A. Rajasekaran](#), Ph.D.

Principal and Professor,
KMCH College of Pharmacy,
Kovai Estate, Kalapatti Road,
Coimbatore- 641 048, Tamil Nadu, India

[Dr. Mohamed A. Farag](#) Ph.D.

Alexander von Humboldt fellow,
Professor, Chemistry Department
American University in Cairo, Cairo, Egypt

Feedback

Subscribe

About the journal

[Dr. Anindya Bose](#) Ph.D.

Associate Professor,
Department of Pharmaceutical Analysis,
School of Pharmaceutical Sciences,
Siksha O Anusandhan (Deemed to be University),
Bhubaneswar, Odisha, India

[Mesfin Yimam](#), (DVM, MS)

Director, Pre-Clinical Development, Unigen Pharmaceuticals
2121 South State Street
Suite 400
Tacoma WA-98405, USA

[Dr. Sami A. Gabr](#) Ph.D.

Rehabilitation Research Chair,
College of Applied Medical Sciences,
King Saud University P.O. Box 10219
Riyadh 11433, KSA

[Prof. Mohammad Azadbakht](#)

Department of Pharmacognosy,
Faculty of Pharmacy,
Mazandaran University of Medical Sciences, Sari, Iran

[Lai, Choon Sheen](#)

Centre for Drug Research,
Universiti Sains Malaysia,
11800 USM, Penang, Malaysia.

Prof. (Dr.) Vimal Kumar, Ph.D.

Principal, ITM School of Pharmacy,
ITM Universe, Dhanora Tank Road,
Off Halol Highway, Near Jarod, Paldi,
Vadodara, Gujarat 391510

[Dr. Venkata Ramireddy Narala](#), M.Sc., Ph.D.

Department of Animal Sciences (Zoology)
Yogi Vemana University
KADAPA-516003, INDIA

[Dr. Qazi Mohammad Sajid Jamal](#), Ph.D.

Assistant Professor and Co-ordinator,
Bioinformatics Unit,
Department of Health Information Management,
College of Applied Medical Sciences,
Buraydah Colleges, Al Qassim-Buraydah King Abdul Aziz Road,
East Qassim University, Kingdom of Saudi Arabia

[Dinithi Peiris](#), Ph.D.

Professor in Zoology,
Faculty of Applied Sciences,
University of Sri Jayewardenepura,
Gangodawilla,
Nugegoda, Sri Lanka

Dr. Komal Kumar Javarappa, MSc, PhD, FBSS

Postdoctoral Scientist,
Institute for Molecular Medicine,
Finland, University of Helsinki Fin-00710,
Helsinki Finland

[Dr. Gyan Singh Shekhawat](#)

Centre for advanced studies in Botany,
Jai Narain Vyas University, New Campus
Jodhpur- 342001, Rajasthan, India.

[Newman Osafo](#), Ph.D.

Department of Pharmacology,
K.N.U.S.T. Kumasi,
Ghana.

[Dr. Sandeep Ramchandra Pai](#) Ph.D.

Assistant Professor,
AMITY Institute of Biotechnology (AIB),
Amity University, Mumbai - Pune
Maharashtra, India

Dr. Christian Agyare

Faculty of Pharmacy and Pharmaceutical Sciences
College of Health Sciences
Kwame Nkrumah University of Science and Technology
Kumasi
Ghana

Dr. Gabriel Kigen

Department of Pharmacology and Toxicology,
Moi University School of Medicine
Eldoret, Kenya

Prof. Dejun Zhang

Vice Dean of Ecology and Environment Engineering college,
Qinghai University,
China

José Carlos Tavares Carvalho

Universidade Federal do Amapá,
Colegiado de Ciências Farmaceuticas,
Laboratório de Pesquisa em Farmacos,
Rodovia Juscelino Kubitschek - de 1670/1671 ao fim
Universidade
68903419 - M acapá, AP - Brasil

Fausto Pierdoná Guzen.

Laboratory of Experimental Neurology,
Department of Biomedical Sciences,
Health Science Center,
State University of Rio Grande do Norte,
Mossoró-RN, Brazil

Patrícia Rijo, PhD

CBIOS-Scientific Director for Communication and External Relations
Food Sciences and Phytochemistry (FSP) Group Coordinator - CBIOS
Escola de Ciências e Tecnologias da Saúde
Universidade Lusófona
Campo Grande, 376
1749-024 Lisboa - Portugal

Consolacion Ragasa, Ph.D

De La Salle University | DLSU
Department of Chemistry
Manila, Philippines

Dr. Kumarappan PhD.,

Professor (Assistant)
Department of Pharmacology & Toxicology,
School of Pharmacy
King Khalid University,
Abha 62529, Saudi Arabia

To join editorial board, you may send your online (ORCID ID, Researchgate, Loop, SCOPUS ID, Researcher ID) profile along with list of Publications to EB Selection Committee : ebscom@phcog.net

Sitemap | What's New | Feedback | Disclaimer Phcog Mag. | Phcog.Net
© Pharmacognosy Research | Published by Wolters Kluwer - Medknow
Online since 1st January, 2010

Editorial and Ethics Policies

  [Open Access](#)  [View mobile site](#)

ISSN: Print -0976-4836, Online - 0974-8490

Supports



Table of Contents



2017

Volume 9 | Issue 5 (Supplement)
Page Nos. 1-109

Online since Tuesday, December 12, 2017

Accessed 12,092 times.

PDF access policy

Full text access is free in HTML pages; however the journal allows PDF access only to subscribers.

EPub access policy

Full text in EPub is free except for the current issue. Access to the latest issue is reserved only for the paid subscribers.

 [View issue as eBook](#)

 [Author Institution Mapping](#)

 [Issue statistics](#)

 [RSS](#)

[Feedback](#)

[Subscribe](#)

[Next Issue](#)

[Previous Issue](#)


[Show all abstracts](#)

[Show selected abstracts](#)

[Export selected to](#) ▼

[Add to my list](#)

ORIGINAL ARTICLES




Antinociceptive and anti-inflammatory effects of triterpenes from *Pluchea quitoc* DC. aerial parts p. 1

Francisco Alcione Nobre da Silva, Sônia Maria de Farias Freire, Marilene Oliveira da Rocha Borges, Francisco Erivaldo Vidal Barros, Maria da Glória Teixeira de Sousa, Maria Nilce de Sousa Ribeiro, Giselle Maria Skelding Pinheiro Guilhon, Adolfo Henrique Müller, Antonio Carlos Romão Borges

DOI:10.4103/pr.pr_51_17 PMID:29333034

[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[PubMed\]](#) [\[Sword Plugin for Repository\]](#)^{Beta}




The activity of immunoglobulin Y anti-*Mycobacterium tuberculosis* on proliferation and cytokine expression of rat peripheral blood mononuclear cells p. 5

Sri Agus Sudjarwo, Koerniasari Eraiko, Giftania Wardani Sudjarwo, Koerniasari

DOI:10.4103/pr.pr_66_17 PMID:29333035

[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[PubMed\]](#) [\[Sword Plugin for Repository\]](#)^{Beta}



Autonomic receptors and nitric-oxide involvements in mediating vasorelaxation effect induced by *Syzygium polyanthum* leaves extract p. 9

Azlini Ismail, Wan Amir Nizam Wan Ahmad

DOI:10.4103/pr.pr_69_17 PMID:29333036

[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[PubMed\]](#) [\[Sword Plugin for Repository\]](#)^{Beta}

Curcumin and natural derivatives inhibit Ebola viral proteins: An *In silico* approach p. 15

Shruti Baikerkar

DOI:10.4103/pr.pr_30_17 PMID:29333037

[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[PubMed\]](#) [\[Sword Plugin for Repository\]](#)^{Beta}




Antihyperglycemic potential of saponin-enriched fraction from *Pithecellobium dulce* Benth. seed extract p. 23

Mahesh Kumar, Jeyabalan Govindarajan, Narendra Kumar Nyola

DOI:10.4103/pr.pr_18_17 PMID:29333038

[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[PubMed\]](#) [\[Sword Plugin for Repository\]](#)^{Beta}



Pharmacognostic assesment of the endemic and vulnerable medicinal climber-*Cayratia pedata* (Lam.) Gagnep. var. *glabra* Gamble and its antibacterial activity p. 27


S Sharmila, K Kalaichelvi, SM Dhivya, P Premamalini, P Abirami, G Jayanthi

DOI:10.4103/pr.pr_25_17 PMID:29333039

[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[PubMed\]](#) [\[Sword Plugin for Repository\]](#)^{Beta}




Pharmacological screening of *Trachyspermum ammi* for antihyperlipidemic activity in Triton X-100 induced hyperlipidemia rat model p. 34



Uzma Saleem, Saba Riaz, Bashir Ahmad, Saleem Mohammad
DOI:10.4103/pr.pr_37_17 PMID:29333040

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [PubMed] [Sword Plugin for Repository]^{Beta}

p. 41




Pharmacological evaluation of hepatoprotective activity of AHPL/AYTAB/0613 tablet in carbon tetrachloride-, ethanol-, and paracetamol-induced hepatotoxicity models in Wistar albino rats

Sanjay U Nipanikar, Sohan S Chitlange, Dheeraaj Nagore
DOI:10.4103/pr.pr_44_17 PMID:29333041

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [PubMed] [Sword Plugin for Repository]^{Beta}

p. 48



Gas chromatography mass spectrometry profiling in methanolic and ethyl-acetate root and stem extract of *Corbichonia decumbens* (Forssk.) exell from Thar Desert of Rajasthan, India

Sunita Arora, Manju Saini
DOI:10.4103/pr.pr_62_17 PMID:29333042

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [PubMed] [Sword Plugin for Repository]^{Beta}




Picroside I and picroside II from tissue cultures of *Picrorhiza kurroa*

Yamjala Ganeshkumar, Ajmera Ramarao, Ciddi Veeresham
DOI:10.4103/pr.pr_89_17 PMID:29333043

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [PubMed] [Sword Plugin for Repository]^{Beta}

p. 53



Antiplasmodial activity of isolated polyphenols from *Alectryon serratus* leaves against 3D7 *Plasmodium falciparum*

Uswatun Khasanah, Aty WidyaWaruyanti, Achmad Fuad Hafid, Mulyadi Tanjung
DOI:10.4103/pr.pr_39_17 PMID:29333044

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [PubMed] [Sword Plugin for Repository]^{Beta}

p. 57




In vitro antiproliferative effect of earthworm coelomic fluid of *Eudrilus eugeniae*, *Eisenia foetida*, and *Perionyx excavatus* on squamous cell carcinoma-9 cell line: A pilot study

Dominic Augustine, Roopa S Rao, Jayaraman Anbu, KN Chidambara Murthy
DOI:10.4103/pr.pr_52_17 PMID:29333045

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [PubMed] [Sword Plugin for Repository]^{Beta}

p. 61




Development and validation of high-performance thin-layer chromatography method for simultaneous determination of polyphenolic compounds in medicinal plants

CV Jayachandran Nair, Sayeed Ahamad, Washim Khan, Varisha Anjum, Rajani Mathur
DOI:10.4103/pr.pr_122_16 PMID:29333046

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [PubMed] [Sword Plugin for Repository]^{Beta}

p. 67




Assessment of nutritional quality and global antioxidant response of banana (*Musa* sp. CV. Nanjangud Rasa Bale) pseudostem and flower

Ramith Ramu, Prithvi S Shirahatti, KR Anilakumar, Shivasharanappa Nayakavadi, Farhan Zameer, BL Dhananjaya, MN Nagendra Prasad
DOI:10.4103/pr.pr_67_17 PMID:29333047

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [PubMed] [Sword Plugin for Repository]^{Beta}

p. 74




Astaxanthin ameliorates hepatic damage and oxidative stress in carbon tetrachloride-administered rats

Md. Ariful Islam, Md. Abdullah Al Mamun, Md. Faruk, Md. Tauhid Ul Islam, Md. Mizanur Rahman, Mohammad Nazmul Alam, A. F. M. Towheedur Rahman, Hasan Mahmud Reza, Md. Ashrafal Alam
DOI:10.4103/pr.pr_26_17 PMID:29333048

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [PubMed] [Sword Plugin for Repository]^{Beta}

p. 84




Marine-derived fungi extracts enhance the cytotoxic activity of doxorubicin in nonsmall cell lung cancer cells A459

Bruno Castro-Carvalho, Alice A Ramos, Maria Prata-Sena, Fernanda Malhão, Márcia Moreira, Daniela Gargiulo, Tida Dethoup, Suradet Buttachon, Anake Kijjoa, Eduardo Rocha
DOI:10.4103/pr.pr_57_17 PMID:29333049

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [PubMed] [Sword Plugin for Repository]^{Beta}

p. 92



Improvement of insulin secretion and pancreatic β -cell function in streptozotocin-induced diabetic rats treated with *Aloe vera* extract

Ayesha Noor, S Gunasekaran, MA Vijayalakshmi
DOI:10.4103/pr.pr_75_17 PMID:29333050

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [PubMed] [Sword Plugin for Repository]^{Beta}


p. 99

Therapeutic significance of *Loligo vulgaris* (Lamarck, 1798) ink extract: A biomedical approach

Sri Kumaran Nadarajah, Radha Vijayaraj, Jayaprakashvel Mani

DOI:10.4103/pr.pr_81_17 PMID:29333051

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [PubMed] [Sword Plugin for Repository]^{Beta}



[Sitemap](#) | [What's New](#) | [Feedback](#) | [Disclaimer](#)

© Pharmacognosy Research | Published by Wolters Kluwer - Medknow

Online since 1st January, 2010

[Editorial and Ethics Policies](#)

Open Access [View mobile site](#)

ISSN: Print -0976-4836, Online - 0974-8490

Supports





Avoca Quality Consortium

Collaborate with pharmaceutical industry leaders from over 130 companies.

theavocagroup.com

OPEN

International Journal of Pharmaceutical Research

Country [India](#) -  SCIMAGO INSTITUTIONS RANKINGS

Subject Area and Category [Pharmacology, Toxicology and Pharmaceutics](#)
[Pharmaceutical Science](#)
[Pharmacology, Toxicology and Pharmaceutics \(miscellaneous\)](#)

Publisher [Advanced Scientific Research](#)

Publication type [Journals](#)

ISSN [09752366](#)

Coverage [2010-2020](#)

Scope International Journal of Pharmaceutical Research (IJPR) is an intentional Journal which is published quarterly in English. Journal publishes papers, review articles, and short communications dealing with drug controlled release systems, pharmacodynamics, pharmacokinetics, pharmacogenomics, biopharmaceutics, drug and prodrug design, pharmaceutical analysis, drug stability, quality control, pharmaceutical engineering and materials science. Pharmaceutical Chemistry, Pharmaceutical Technology, pharmacognosy, natural product research, pharmaceutics, novel drug delivery, pharmaceutical & medicinal chemistry, computational chemistry and molecular drug design, pharmacology, pharmaceutical analysis, pharmacy practice, clinical and hospital pharmacy etc. IJPR would take much care in making your article published without much delay with your kind cooperation. IJPR hopes that Researchers, Research scholars, Academician, Industrialists etc. would make use of this research publications for the development of pharmaceutical science and technology.

 [Homepage](#)

[How to publish in this journal](#)

[Contact](#)

 [Join the conversation about this journal](#)

13

H Index

Impact factor 7.97, low cost


research journal, Call for Paper, paper Publication, Research I Review Paper

Impact factor 7.97, low cost


research journal, Call for Paper, paper Publication, Research I Review Paper

jetir journal


Quartiles




SJR




Citations per document




Total Cites Self-Cites



External Cites per Doc Cites per Doc



% International Collaboration



Citable documents Non-citable documents

Cited documents Uncited documents



Journal of International
Pharmaceutical Research



← Show this widget in
your own website

Just copy the code below
and paste within your html
code:

```
<a href="https://www.scimaç
```

Doodle Video Creation Software

Professional whiteboard, blackboard doodle videos in seconds

Doodly

Metrics based on Scopus® data as of April 2020

M **Mykhailo Anishchenko** 2 weeks ago

Dear editorial board! Please give information on how you can get an article from your journal in pdf format?

Best regards, Mykhailo Anishchenko

reply



Melanie Ortiz 2 weeks ago

SCImago Team

Dear Mykhailo,
thank you for contacting us.

We are sorry to tell you that SCImago Journal & Country Rank is not a journal. SJR is a portal with scientometric indicators of journals indexed in Elsevier/Scopus.

Unfortunately, we cannot help you with your request, we suggest you contact the journal's editorial staff, so they could inform you more deeply.

Best Regards, SCImago Team

N **Nada** 2 weeks ago

Antiplasmodial Activity of Isolated Polyphenols from *Alectryon serratus* Leaves Against 3D7 *Plasmodium falciparum*

Uswatun Khasanah^{1,2}, Aty WidyaWaruyanti^{3,4}, Achmad Fuad Hafid^{3,4}, Mulyadi Tanjung⁵

¹Post Graduate Program, Faculty of Pharmacy, Universitas Airlangga, Surabaya, ²Department of Pharmacy, Faculty of Medicine, Brawijaya University, Malang, ³Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Universitas Airlangga, ⁴Institute of Tropical Disease, Universitas Airlangga, ⁵Departement of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Surabaya, Indonesia

ABSTRACT


Background: *Alectryon serratus* was selected from a screening program devoted to search naturally occurring antimalarial compound from plants in Alas Purwo National Park, Banyuwangi, East Java, Indonesia. The previous studies showed that ethanol extract of *A. serratus* leaves contains some polyphenol compounds. **Objective:** This study was designed to isolate and investigate antiplasmodial activity of polyphenol compounds. **Material and Methods:** The ethanol extract of *A. serratus* leaves was fractionated using liquid-liquid fractionation and column chromatography. Isolated compounds were identified using High-performance liquid chromatography, ultraviolet-visible, nuclear magnetic resonance, and compared with references. The isolates were tested *in vitro* for antiplasmodial activity against chloroquine-sensitive 3D7 strain of *Plasmodium falciparum*. Thin blood smears were used to assess the levels of parasitemia and growth inhibition of the isolates. **Result:** Half maximal inhibitory concentration of Gallic acid (1), methyl gallate (2), and kempferol-3-O-rhamnoside (3) were 0.0722 μ M, 0.0128 μ M, and 3.4595 μ M, respectively. **Conclusion:** The results suggest that gallic acid, methyl gallate, and kempferol-3-O-rhamnoside isolated from *A. serratus* leaves have antiplasmodial activity and are potential to be developed as antimalarial drugs.

Key words: *Alectryon serratus*, antiplasmodial, polyphenol

SUMMARY

- The ethanol extract of *Alectryon serratus* leaves was successively fractionated in CH_2Cl_2 , EtOAc, and n-butanol. EtOAc fraction was fractionated using column chromatography and purified using preparative thin-layer chromatography (TLC). Isolates were studied for their antiplasmodial activity on parasites culture of chloroquine-sensitive 3D7 strain of *Plasmodium falciparum*. Parasitemia percentages, growth percentages, and inhibition

percentages of each group were calculated. The half maximal inhibitory concentration (IC_{50}) values that represent the concentration required to inhibit 50% of *Plasmodium* growth were calculated from a calibration curve using linear regression. The results suggest that isolates have antiplasmodial activity and are responsible in the antimalarial activity of *Alectryon serratus* leaves.



Abbreviations Used: S.F: Subfraction, EGCG: Epigallocatechingallate, EGC: Epigallocatechin

Correspondence:

Dr. Aty WidyaWaruyanti,
Department of Pharmacognosy and
Phytochemistry, Faculty of Pharmacy,
Universitas Airlangga, Surabaya,
East Java, Indonesia.
E-mail: aty_ww@yahoo.com
DOI: 10.4103/pr.pr_39_17

Access this article online

Website: www.phcogres.com

Quick Response Code:



INTRODUCTION

Malaria is a major parasitic infectious disease in many tropical and subtropical regions. Malaria incidence has been increasing since the emergence of drug-resistant *Plasmodium falciparum*. According to the WHO, as many as 207 million people suffer from malaria, with up to 627,000 deaths each year.^[1] Drug-resistant *P. falciparum* malaria is a major killer and becomes one of the most difficult obstacles to combat. Therefore, a development of a new class drug is an urgent matter. Screening plant extract for antiplasmodial activity is a useful way for discovering new leads.^[2] As a part of our study for novel antimalarial agents from plants, a screening program was undertaken on plants of Alas Purwo National Park, Banyuwangi, East Java, Indonesia. In our screening, leaves of *Alectryon serratus* were selected.^[3] This plant is widely distributed throughout the tropical region of Southeast Asia, and no traditional uses are reported.^[4] Ethyl acetate fraction of the leaves was found to exhibit an antimalarial activity (IC_{50} 9.45 μ g/mL) on chloroquine-sensitive 3D7 strain of *P. falciparum*.^[5] TLC profile of ethyl acetate fraction showed polyphenol compounds.^[5] In this paper, we report the isolation, structure elucidation, and antiplasmodial activity of polyphenol compounds.

MATERIALS AND METHODS

Collection of plant material

Leaves of *A. serratus* were collected from Alas Purwo National Park, Banyuwangi, East Java, Indonesia in August 2014. A voucher specimen was identified at Purwodadi Botanical Garden, Pasuruan, East Java, Indonesia, and a voucher specimen was deposited at the herbarium.

Extraction and fractionation

One kilogram of dried powder of *A. serratus* leaves was macerated using sonification method in ethanol 80% as the solvent. The extract was

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Cite this article as: Khasanah U, WidyaWaruyanti A, Hafid AF, Tanjung M. Antiplasmodial activity of isolated polyphenols from *Alectryon serratus* leaves against 3D7 *Plasmodium falciparum*. Phcog Res 2017;9:S57-60.

then removed, and the residue was pressed and dried. The residue was re-macerated twice with ethanol 80%. The dried extract was suspended in distilled water and fractionated. Liquid fractionation was done successively in CH_2Cl_2 , EtOAc, and n-butanol.

Isolation of polyphenol compounds

The ethyl acetate fraction (2 g) was separated using a column chromatography method using RP-18 F254 as stationary phase and $\text{CH}_3\text{CN}-\text{MeOH}-\text{H}_2\text{O}$ (2:1:4 v/v) as the mobile phase to yield 12 subfractions (SF.1-SF.12). White powder was obtained as a precipitate from S.F.2. The precipitate was filtered using Kiriya to yield compound 1 (12 mg). S.F.4 (30 mg) was purified using TLC preparative with $\text{CHCl}_3-\text{MeOH}$ (9:1 v/v) as mobile phase and silica gel F₂₅₄ as stationary phase to yield S.F.4.1 (4 mg) and S. F.4.2 (8 mg). S.F.4.2 was identified as compound 2. S.F.8 (40 mg) was purified using preparative TLC with $\text{CHCl}_3-\text{MeOH}$ (8:2 v/v) as mobile phase and silica gel F₂₅₄ as stationary phase to yield S.F.8.1 (5 mg), S.F.8.2 (7 mg), and S.F.8.3 (2 mg). S.F.8.3 was identified as compound 3.

Characterization of isolated polyphenols

TLC profiles of S.F.2 was done using RP-18 F₂₅₄ as the stationary phase and $\text{CH}_3\text{CN}-\text{MeOH}-\text{H}_2\text{O}$ (2:1:4 v/v) as the mobile phase. S.F.4.2 and S.F.8.3 were identified using TLC with silica gel F₂₅₄ as stationary phase and $\text{CHCl}_3-\text{MeOH}$ (9:1 v/v) as mobile phase. The spots were detected under ultraviolet (UV) (254 and 366 nm) before spraying using 10% H_2SO_4 in EtOH, and followed by heating the plate at 120°C and then detected under UV 366 nm and visible light. High-performance liquid chromatography (HPLC) chromatogram profile was performed using Shimadzu LC-06, with RP Shim-pack column 4.6 mm × 250 mm as stationary phase, and $\text{CH}_3\text{CN}-\text{MeOH}$ (7:3 v/v) as mobile phase with a flow rate of 0.7 ml/min, and was detected using UV detector.

Nuclear magnetic resonance (NMR) spectra were performed using JEOL 400 spectrometer, with tetramethylsilane as internal standard for ¹H, ¹³C NMR, heteronuclear multiple bond correlation (HMBC), and heteronuclear multiple quantum coherence. Compound 1 and 2 were measured using D6 methanol solvent; compound 3 was measured using D6 acetone solvent. The structural determination of compound 1, 2, and 3 was confirmed by comparing the results of NMR and UV data reported in the literature.¹⁶⁻¹¹

Plasmodium falciparum (3D7 strain) culture and maintenance

P. falciparum 3D7 strain (chloroquine sensitive) was obtained from the Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia. *Plasmodium* parasites were grown and maintained in culture using Trager and Jensen method with some modifications.¹² The parasites were bred *in vitro* in human type O-positive red blood cells in a complete medium (RPMI 1640 supplemented with 5.96 g HEPES, 0.05 g hypoxanthine, 2.1 g NaHCO_3 , 50 µg/ml gentamycin, and 10% human O + serum) in Petri dish with modified candle-jar method. The incubation was done at 37°C. The media was routinely changed daily, and the parasite growth was monitored through Giemsa staining in thin blood smears. The culture to be used for the experiment should be dominated by ring forms. Stock parasite cultures were further diluted with uninfected type O + human erythrocytes and complete culture medium to achieve 1% parasitemia and 50% hematocrit. These final parasite cultures were immediately used for the antimalarial assay.^{12,13}

In vitro antimalarial assay

A stock solution of 10 µg/ml was prepared from the isolated compounds. Four-fold serial dilution was prepared from each stock solution and yielded five serial concentrations (10 µg/ml, 1 µg/ml, 0.1 µg/ml, 0.01 µg/ml, and 0.001 µg/ml). To each microplate well, 500 µL diluted extract solution was added into 500 µL of final parasite culture. The plates were then incubated for 48 h at 37°C.

Thin blood smears were prepared on labeled slides and air-dried and fixed with methanol. The dried slides were stained using Giemsa and observed under light microscope at 1,000 times magnification, and parasitemia percentage was calculated.

Statistical analysis of data

Growth percentage was calculated using formula as follows:

$$\text{Growth percentage (\%growth)} = \frac{\text{Parasitemia} - D_0}{D_0} \times 100\%$$

Inhibition percentage was calculated as follows:

$$\text{Inhibition percentage (\% inhibition)} = 100\% - \left[\frac{Xu}{Xk} \times 100\% \right]$$

Note:

- D_0 = Parasitemia percentage of infected red blood cell on day 0
- Xu = Growth percentage of each isolate
- Xk = Growth percentage of negative control
- IC_{50} values that represent the concentration required to inhibit 50% of *Plasmodium* growth were calculated from a calibration curve by linear analysis using SPSS. IC_{50} values were expressed as mean value (±standard deviation).

RESULTS AND DISCUSSION

TLC profile of S.F.2 and S.F.4.2 showed the presence of dark fluorescent spots under UV 254 nm and UV 366 nm with Rf 0.75 and Rf 0.6. Dark spots were also detected on S.F.2 and S.F.4.2 after being sprayed and observed under UV 366 nm. S.F.8.3 also showed dark spot under UV 254 and UV 366 nm with Rf 0.48, and yellowish spots after the plate was being sprayed using 10% H_2SO_4 and was detected under UV 366 nm and visible light. Isolation of phenolic compounds was done for S.F.2 and S.F.4.

HPLC chromatogram of S.F.2 showed a single peak with Rt 4.3 min with a purity index of 0.98 and UV/Vis λ_{max} (MeOH) at 266 nm. S.F.4.2 also showed a single peak with Rt 4.5 min, purity index 0.97 and UV/Vis λ_{max} (MeOH) at 271 nm.

Results of NMR spectra of each isolate are described as follows:

S.F.2 (1)

White powder

¹H NMR (400 MHz, Methanol-*d*₄): 7.02 (s).

¹³C NMR (100 MHz, Methanol-*d*₄): 115.6 (C), 103.9 (CH), 140.1 (C), 133.3 (C), 140.1 (C), 164.1 (COOH).

S.F.4.2 (2)

Pale yellow powder

¹H NMR (400 MHz, Methanol-*d*₄): 7.02 (s), 3.75 (3H, s).

¹³C NMR (100 MHz, Methanol-*d*₄): 116.0 (C), 104.0 (CH), 140.4 (C), 133.0 (C), 140.4 (C), 164.4 (COOH), 46.2 (OCH₃).

S.F.8.3 (3)

Yellow powder

¹H NMR (400 MHz, Acetone-*d*₆): 7.83 (2H, dd, J = 8,4), 6.99 (2H, dd, J = 8,4), 6.45 (d, J = 2,4 Hz, 1H), 6.24 (d, J = 2,4 Hz, 1H), 12.69 (1H, s), 5.51 (1H, d, J = 1,6), 3.30–3.60 (3H, m), 4.18 (1H, m), 1.16 (3H, s), CH₂.

¹³C NMR (100 MHz, Acetone-*d*₆): 178.5 (C = O), 115.4 (CH), 130.1 (CH), 98.7 (CH), 93.5 (CH), 170.3, 164.4, 159.7, 163.8,

156.6 (oxyaryl group), 71.3, 70.6, 70.5, 70.2 (C-2", C-3", C-4", C-5"), 16.9 (CH₃).

Antiplasmodial activity of each isolate concentration was shown in Figure 1. Maximum inhibition percentage of 96.7%, 80.9%, and 62.8% were obtained for S.F.2, S.F.4.2, and S.F.8.3 at 10 µg/mL concentration, respectively. Concentration below 10 µg/mL exhibited lower inhibition percentage. The IC₅₀ values of each isolates were calculated and found to be 0.013, 0.0025, and 1.495 µg/mL [Table 1].

S. F.2 was identified as Gallic acid (1) that was isolated as white powder. TLC profile showed that S. F.2 was a phenolic compound. UV spectrum (MeOH) λ_{max} 266 nm showed that S. F.2 had a chromophore group. ¹H NMR spectrum showed signal typical to aromatic proton (δ_H 7.02). ¹³C NMR spectrum indicated the presence of 7 carbon atom signals including carboxylic carbon at δ_C 164.1. Further support for the structure (1) was also obtained from the comparison of NMR data with those reported for gallic acid.^[6-8]

S. F.4.2 was identified as methyl gallate (2) that was isolated as yellowish-pale powder. UV spectrum (MeOH) λ_{max} 271 nm showed that S. F.4.2 had a chromophore group. ¹H NMR spectrum showed signal typical to aromatic proton (δ_H 7.07) and methoxy proton (δ_H 3.75). ¹³C NMR spectrum indicated the presence of 8-carbon atoms including carboxylic carbon at δ_C 161.4 and methoxy carbon at δ_C 46.2. Further support for the structure (2) was also obtained from the comparison of NMR data with those reported for methyl gallate.^[6-8]

S. F.8.3 was identified as Kempferol-3-O-rhamnoside (3) that was isolated as yellow powder. ¹H NMR spectrum indicated the presence of 2 aromatic hydrogen signals with "meta-coupling" at δ 6.24 (1H, *d*, *J* = 1.6 Hz) and 6.45 (1H, *d*, *J* = 2.4 Hz), which were predicted by hydrogen in C-6 and C-8 of the A ring of the flavone skeleton. The signal at δ 12.69 was predicted as hydroxyl group at C-5. Accordingly, this compound was suggested to have a hydroxyl group at C-5 and C-7. Furthermore, ¹H NMR spectrum revealed two signals with "ortho-coupling" at δ 6.99 (2H, *d*, *J* = 8.4 Hz) and 7.83 (2H, *d*, *J* = 8.4 Hz), the signals were predicted as the hydrogen at C-2', C-3', C-5' and C-6' of the B ring. The absence of a specific signal for an olefinic hydrogen at C-3 and the presence of an anomeric hydrogen signal at δ 5.51 suggested that the compound was a flavonol glycoside. The appearance of an anomeric carbon signal at δ 101.8 in ¹³C NMR spectrum indicated the presence of sugar moiety. Due to a correlation between the anomeric hydrogen signal (δ 5.51) and C-3

carbon signal (δ 141.0) revealed by HMBC spectral data analysis, the position of sugar moiety was assigned to C-3 hydroxyl group. The methyl signal observed at δ 1.16 (3H, *s*) in ¹H NMR spectrum and at δ 16.9 in ¹³C NMR spectrum indicated that the sugar moiety was rhamnose. Based on the accumulated data above and data comparison, compound (3) was identified as kaempferol-3-O-rhamnoside.^[9-11]

The structure of gallic acid, methyl gallate, and kaempferol-3-O-rhamnoside were shown in Figure 2.

Antiplasmodial activity of compound 1 and 2 appeared and was associated with gallate group.^[9] The previous study showed that methyl gallate isolated from *Swintonia foxworthyi* had antiplasmodial activity against chloroquine-sensitive 3D7 strain of *P. falciparum* with IC₅₀ 3.5 µg/mL.^[14] The result of this study showed that methyl gallate exhibited antiplasmodial activity with IC₅₀ 0.0025 µg/mL against chloroquine-sensitive 3D7 strain of *P. falciparum*. Kaempferol-3-O-rhamnoside isolated from *Schima wallichii* also showed antiplasmodial properties with IC₅₀ 106 µM.^[15] In this study, Kaempferol-3-O-rhamnoside showed antimalarial activity with IC₅₀ 3.4595 µM [Table 1]. The difference of IC₅₀ value might be caused by the type of parasite strain. Antiplasmodial activity of some gallic acid derived from green tea was also tested by Sannela *et al.*, and the result showed that both epigallocatechingallate and epigallocatechin have potential antiplasmodial activity.^[16]

Oxidative stress through the generation of reactive oxygen species plays important role in the pathogenesis of malaria infection that causes hemoglobin degradation.^[17] As phenolic compounds, gallic acid, methyl gallate, and kaempferol-3-O-rhamnoside have antioxidant properties that may be responsible for antiplasmodial activity. According to Fidock *et al.*, a compound has potential antimalarial activity if IC₅₀ < 1–5 µM.^[18] It can be concluded from the data that *A. serratus* leaves contain gallic acid, methyl gallate, and kaempferol-3-O-rhamnoside that have antiplasmodial activity against chloroquine-sensitive 3D7 strain of *P. falciparum*.

Acknowledgement

The authors thank NPMRD (Natural Product Medicine Research and Development), ITD (Institute of Tropical Disease Center) Universitas Airlangga, Surabaya.

Table 1: Antimalarial activity of isolate compounds

Compounds	IC ₅₀	
	µg/ml±SD	µM±SD
Gallic acid	0.0130±0.0014	0.0722±0.0078
Methyl gallate	0.0025±0.0007	0.0128±0.0036
Kaempferol-3-O-rhamnoside	1.495±0.007	3.4595±0.0180

SD: Standard deviation




Figure 1: Various concentration of compounds determining the IC₅₀ value against *Plasmodium falciparum* after 48 h of incubation. Data are the mean value of double-independent experiments




Figure 2: Chemical structure of the antiplasmodial polyphenols from *Alectryon serratus* leaves. (1) Gallic acid; (2) Methyl gallate; (3) Kaempferol-3-O-rhamnoside

Financial support and sponsorship

This research was supported by Indonesian Directorate General of Higher Education DIPA BOPTN 2014, contract no 965/UN3/2014

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- World Health Organization. Guidelines for the Treatment of Malaria. 2nd ed. Geneva, Swiss: World Health Organization; 2013.
- Nguyen-Pouplin J, Tran H, Phan TA, Dolecek C, Farrar J, et al. Antimalarial and cytotoxic activities of ethnopharmacologically selected medicinal plants from South Vietnam. *J Ethnopharmacol* 2007;109:417-27.
- Widyawaruyanti A, Devi AP, Fatria N, Tumewu L, Tantular IS, Hafid AF. *In vitro* antimalarial activity screening of several Indonesian plants using hrp2 assay. *Int J Pharm Pharm Sci* 2014;6:6-9.
- Adema F, Leenbouts FP, Van Welzen PC. *Flora Malesiana. Series I-Spermatophyta*. Den Haag: Foundation Flora Malesiana; 1994.
- Widyawaruyanti A, Khasanah U, Tumewu L, Ilmi H, Hafid AF, Tantular IS. Antimalarial activity and cytotoxicity study of ethanol extract and fraction from *Alectryon serratus* leaves. *Int J Pharm Pharm Sci* 2015;7:250-3.
- Angela IC, Luz IR, Eduardo OB, Reto B, Mireya DCA, Mahabir PG. Evaluation of larvicidal and *in vitro* antiparasitic activities of plants in a biodiversity plot in the Altos de Campana National Park, Panama. *Pharmaceutical Biology* 2002;4:487-98.
- Zhiyong H, Wenshui X, Jie C. Isolation and structure elucidation of phenolic compounds in Chinese olive (*Canarium album* L.) fruit. *Eur Food Technol* 2008;226:1191-6.
- Ndjonka D, Bergmann B, Agyare C, Zimbres FM, Lüersen K, Hensel A, et al. *In vitro* activity of extracts and isolated polyphenols from west african medicinal plants against *Plasmodium falciparum*. *Parasitol Res* 2012;111:827-34.
- Song N, Xu W, Guan H, Liu X, Wang Y, Nie X. Several flavonoids from *Capsella bursa-pastoris* (L) Medic. *Asian J Tradit Med* 2007;2:218-22.
- Furusawa M, Tanaka T, Ito T, Nakaya K, Iliya I, Ohyama M, et al. Flavonol glycosides in leaves of two diospyros species. *Chem Pharm Bull (Tokyo)* 2005;53:591-3.
- Diantini A, Subarnas A, Lestari K, Halimah E, Susilawati Y, Supriyatna, et al. Kaempferol-3-O-rhamnoside isolated from the leaves of *Schima wallichii* Korth. Inhibits MCF-7 breast cancer cell proliferation through activation of the caspase cascade pathway. *Oncol Lett* 2012;3:1069-72.
- Trager W, Jensen JB. Human malaria parasites in continuous culture. *Science* 1976;193:673-5.
- Srivastava K, Singh S, Singh P, Puri SK. *In vitro* cultivation of plasmodium falciparum: Studies with modified medium supplemented with ALBUMAX II and various animal sera. *Exp Parasitol* 2007;116:171-4.
- Horgen FD, Madulid DA, Angerhofer CK, Pezzuto JM, Soejarto DD, Farnsworth NR, et al. Isolation of gallic acid esters as antiplasmodial constituents of *Swintonia foxyworthii* (Anacardiaceae). *Phytomedicine* 1997;4:353-6.
- Barliana MI, Suradji EW, Abdulah R, Diantini A, Hatabu T, Nakajima-Shimada J, et al. Antiplasmodial properties of kaempferol-3-O-rhamnoside isolated from the leaves of *Schima wallichii* against chloroquine-resistant *Plasmodium falciparum*. *Biomed Rep* 2014;2:579-83.
- Sannella AR, Messori L, Casini A, Francesco Vincieri F, Bilia AR, Majori G, et al. Antimalarial properties of green tea. *Biochem Biophys Res Commun* 2007;353:177-81.
- Becker K, Tilley L, Vennerstrom JL, Roberts D, Rogerson S, Ginsburg H, et al. Oxidative stress in malaria parasite-infected erythrocytes: Host-parasite interactions. *Int J Parasitol* 2004;34:163-89.
- Fidock DA, Rosenthal PJ, Croft SL, Brun R, Nwaka S. Antimalarial drug discovery: Efficacy models for compound screening. *Nat Rev Drug Discov* 2004;3:509-20.