


SCAM



# Evidence-Based Complementary and Alternative Medicine

—

---

# Natural Products as Sources of Antimalarial Drugs

Lead Guest Editor: Philip F. Uzor

Guest Editors: Vivitri D. Prasasty and Chukwuma O. Agubata

## **Natural Products as Sources of Antimalarial Drugs**

Lead Guest Editor: Philip F. Uzor

Guest Editors: Vivitri D. Prasasty and Chukwuma  
O. Agubata



---

Copyright © 2020 Hindawi Limited. All rights reserved.

This is a special issue published in "Evidence-Based Complementary and Alternative Medicine." All articles are open access articles distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Editorial Board

Mona Abdel-Tawab, Germany  
Rosaria Acquaviva, Italy  
Gabriel A. Agbor, Cameroon  
Ulysses Paulino Albuquerque, Brazil  
Samir Lutf Aleryani, USA  
Mohammed S. Ali-Shtayeh, Palestinian Authority  
Gianni Allais, Italy  
Terje Alraek, Norway  
Adolfo Andrade-Cetto, Mexico  
Isabel Andújar, Spain  
Letizia Angiolella, Italy  
Makoto Arai, Japan  
Hyunsu Bae, Republic of Korea  
Onesmo B. Balemba, USA  
Winfried Banzer, Germany  
Samra Bashir, Pakistan  
Jairo Kenupp Bastos, Brazil  
Arpita Basu, USA  
Daniela Beghelli, Italy  
Juana Benedí, Spain  
Bettina Berger, Germany  
Maria Camilla Bergonzi, Italy  
Andresa A. Berretta, Brazil  
Anna Rita Bilia, Italy  
Monica Borgatti, Italy  
Francesca Borrelli, Italy  
Gioacchino Calapai, Italy  
Giuseppe Caminiti, Italy  
Raffaele Capasso, Italy  
Francesco Cardini, Italy  
Pierre Champy, France  
Shun-Wan Chan, Hong Kong  
Kevin Chen, USA  
Evan P. Cherniack, USA  
Salvatore Chirumbolo, Italy  
Jae Youl Cho, Republic of Korea  
Kathrine Bisgaard Christensen, Denmark  
Shuang-En Chuang, Taiwan  
Yuri Clement, Trinidad And Tobago  
Ian Cock, Australia  
Marisa Colone, Italy

Lisa A. Conboy, USA  
Kieran Cooley, Canada  
Edwin L. Cooper, USA  
Maria T. Cruz, Portugal  
Roberto K. N. Cuman, Brazil  
Ademar A. Da Silva Filho, Brazil  
Giuseppe D'Antona, Italy  
Vincenzo De Feo, Italy  
Rocío De la Puerta, Spain  
Laura De Martino, Italy  
Antonio C. P. de Oliveira, Brazil  
Arthur De Sá Ferreira, Brazil  
Nunziatina De Tommasi, Italy  
Alexandra Deters, Germany  
Farzad Deyhim, USA  
Claudia Di Giacomo, Italy  
Antonella Di Sotto, Italy  
Luciana Dini, Italy  
Caigan Du, Canada  
Jeng-Ren Duann, USA  
Nativ Dudai, Israel  
Thomas Efferth, Germany  
Abir El-Alfy, USA  
Giuseppe Esposito, Italy  
Keturah R. Faurot, USA  
Yibin Feng, Hong Kong  
Nianping Feng, China  
Antonella Fioravanti, Italy  
Johannes Fleckenstein, Germany  
Filippo Fratini, Italy  
Brett Froeliger, USA  
Jian-Li Gao, China  
Dolores García Giménez, Spain  
Gabino Garrido, Chile  
Ipek Goktepe, Qatar  
Yuewen Gong, Canada  
Susana Gorzalczany, Argentina  
Sebastian Granica, Poland  
Settimio Grimaldi, Italy  
Maruti Ram Gudavalli, USA  
Narcís Gusi, Spain  
Svein Haavik, Norway

Solomon Habtemariam, United Kingdom  
Michael G. Hammes, Germany  
Kuzhuvelil B. Harikumar, India  
Ken Haruma, Japan  
Thierry Hennebelle, France  
Markus Horneber, Germany  
Ching-Liang Hsieh, Taiwan  
Benny T. K. Huat, Singapore  
Ciara Hughes, Ireland  
Attila Hunyadi, Hungary  
H. Stephen Injeyan, Canada  
Chie Ishikawa, Japan  
Angelo A. Izzo, Italy  
G. K. Jayaprakasha, USA  
Leopold Jirovetz, Austria  
Takahide Kagawa, Japan  
Atsushi Kameyama, Japan  
Wenyi Kang, China  
Shao-Hsuan Kao, Taiwan  
Juntra Karbwang, Japan  
Teh Ley Kek, Malaysia  
Deborah A. Kennedy, Canada  
Cheorl-Ho Kim, Republic of Korea  
Youn-Chul Kim, Republic of Korea  
Yoshiyuki Kimura, Japan  
Toshiaki Kogure, Japan  
Jian Kong, USA  
Tetsuya Konishi, Japan  
Karin Kraft, Germany  
Omer Kucuk, USA  
Victor Kuete, Cameroon  
Yiu-Wa Kwan, Hong Kong  
Kuang C. Lai, Taiwan  
Ilaria Lampronti, Italy  
Lixing Lao, Hong Kong  
Mario Ledda, Italy  
Christian Lehmann, Canada  
George B. Lenon, Australia  
Marco Leonti, Italy  
Lawrence Leung, Canada  
Min Li, China  
XiuMin Li, Armenia  
Chun-Guang Li, Australia  
Giovanni Li Volti, Italy  
Ho Lin, Taiwan  
Bi-Fong Lin, Taiwan  
Kuo-Tong Liou, Taiwan


Christopher G. Lis, USA  
Gerhard Litscher, Austria  
I-Min Liu, Taiwan  
Monica Loizzo, Italy  
V́ctor Ĺpez, Spain  
Anderson Luiz-Ferreira, Brazil  
Thomas Lundeberg, Sweden  
Michel Mansur Machado, Brazil  
Filippo Maggi, Italy  
Valentina Maggini, Italy  
Jamal A. Mahajna, Israel  
Juraj Majtan, Slovakia  
Toshiaki Makino, Japan  
Nicola Malafronte, Italy  
Francesca Mancianti, Italy  
Carmen Mannucci, Italy  
Arroyo-Morales Manuel, Spain  
Fatima Martel, Portugal  
Simona Martinotti, Italy  
Carlos H. G. Martins, Brazil  
Stefania Marzocco, Italy  
Andrea Maxia, Italy  
James H. McAuley, Australia  
Kristine McGrath, Australia  
James S. McLay, United Kingdom  
Lewis Mehl-Madrona, USA  
Ayikoé Guy Mensah-Nyagan, France  
Oliver Micke, Germany  
Maria G. Miguel, Portugal  
Luigi Milella, Italy  
Roberto Miniero, Italy  
Letteria Minutoli, Italy  
Albert Moraska, USA  
Giuseppe Morgia, Italy  
Mark Moss, United Kingdom  
Yoshiharu Motoo, Japan  
Kamal D. Moudgil, USA  
Yoshiki Mukudai, Japan  
Sakthivel Muniyan, USA  
Massimo Nabissi, Italy  
Hajime Nakae, Japan  
Takao Namiki, Japan  
Srinivas Nammi, Australia  
Krishnadas Nandakumar, India  
Vitaly Napadow, USA  
Michele Navarra, Italy  
Isabella Neri, Italy

Pratibha V. Nerurkar, USA  
Marcello Nicoletti, Italy  
Cristina Nogueira, Brazil  
Martin Offenbaecher, Germany  
Yoshiji Ohta, Japan  
Olumayokun A. Olajide, United Kingdom  
Ester Pagano, Italy  
Sokcheon Pak, Australia  
Siyaram Pandey, Canada  
Visweswara Rao Pasupuleti, Malaysia  
Bhushan Patwardhan, India  
Claudia Helena Pellizzon, Brazil  
Raffaele Pezzani, Italy  
Florian Pfab, Germany  
Sonia Piacente, Italy  
Andrea Pieroni, Italy  
Richard Pietras, USA  
Andrew Pipingas, Australia  
Haifa Qiao, USA  
Xianqin Qu, Australia  
Roja Rahimi, Iran  
Khalid Rahman, United Kingdom  
Elia Ranzato, Italy  
Ke Ren, USA  
Man Hee Rhee, Republic of Korea  
Daniela Rigano, Italy  
José L. Rios, Spain  
Barbara Romano, Italy  
Mariangela Rondanelli, Italy  
Omar Said, Israel  
Avni Sali, Australia  
Mohd. Zaki Salleh, Malaysia  
Andreas Sandner-Kiesling, Austria  
Manel Santafe, Spain  
Tadaaki Satou, Japan  
Michael A. Savka, USA  
Roland Schoop, Switzerland  
Sven Schröder, Germany  
Veronique Seidel, United Kingdom  
Senthamil R. Selvan PhD, USA  
Hongcai Shang, China  
Ronald Sherman, USA  
Karen J. Sherman, USA  
Yukihiro Shoyama, Japan  
Morry Silberstein, Australia  
Kuttulebbai N. S. Sirajudeen, Malaysia  
Francisco Solano, Spain


Chang G. Son, Republic of Korea  
Annarita Stringaro, Italy  
Shan-Yu Su, Taiwan  
Orazio Taglialatela-Scafati, Italy  
Takashi Takeda, Japan  
Ghee T. Tan, USA  
Norman Temple, Canada  
Mencherini Teresa, Italy  
Mayank Thakur, Germany  
Menaka C. Thounaojam, USA  
Evelin Tiralongo, Australia  
Michał Tomczyk, Poland  
Loren Toussaint, USA  
Luigia Trabace, Italy  
Yew-Min Tzeng, Taiwan  
Dawn M. Upchurch, USA  
Konrad Urech, Switzerland  
Takuhiko Uto, Japan  
Patricia Valentao, Portugal  
Sandy van Vuuren, South Africa  
Luca Vanella, Italy  
Alfredo Vannacci, Italy  
Antonio Vassallo, Italy  
Miguel Vilas-Boas, Portugal  
Aristo Vojdani, USA  
Almir Gonçalves Wanderley, Brazil  
Chong-Zhi Wang, USA  
Shu-Ming Wang, USA  
Jonathan L. Wardle, Australia  
Kenji Watanabe, Japan  
Jintanaporn Wattanathorn, Thailand  
Silvia Wein, Germany  
Janelle Wheat, Australia  
Jenny M. Wilkinson, Australia  
Christopher Worsnop, Australia  
Haruki Yamada, Japan  
Nobuo Yamaguchi, Japan  
Junqing Yang, China  
Ling Yang, China  
Albert S. Yeung, USA  
Armando Zarrelli, Italy  
Chris Zaslowski, Australia  
Suzanna M. Zick, USA

# Contents


## **Natural Products as Sources of Antimalarial Drugs**

Philip F. Uzor , Vivitri D. Prasasty, and Chukwuma O. Agubata  
Editorial (2 pages), Article ID 9385125, Volume 2020 (2020)








## **Effect of Formulated *Artocarpus champeden* Extract on Parasite Growth and Immune Response of *Plasmodium berghei*-Infected Mice**

A. Widyawaruyanti , N. Harwiningtias, L. Tumewu, A.F. Hafid, and Soetjipto  
Research Article (7 pages), Article ID 4678634, Volume 2020 (2020)



## **Alkaloids from Plants with Antimalarial Activity: A Review of Recent Studies**

Philip F. Uzor   
Review Article (17 pages), Article ID 8749083, Volume 2020 (2020)



## **In Vitro Antiplasmodium and Chloroquine Resistance Reversal Effects of Andrographolide**

Zaid O. Ibraheem , Roslaini Abd Majid , Hasidah Mohd Sidek , Sabariah Md Noor , Mun Fei Yam , Mohammad Faruq Abd Rachman Isnadi , and Rusliza Basir   
Research Article (16 pages), Article ID 7967980, Volume 2019 (2019)

## **Various Parts of *Helianthus annuus* Plants as New Sources of Antimalarial Drugs**

Wiwied Ekasari , Dwi Widya Pratiwi, Zelmira Amanda, Suciati, Aty Widyawaruyanti , and Heny Arwati  
Research Article (7 pages), Article ID 7390385, Volume 2019 (2019)


## **Evaluation of the Bioactivities of *Rumex crispus* L. Leaves and Root Extracts Using Toxicity, Antimicrobial, and Antiparasitic Assays**

Oladayo Amed Idris , Olubunmi Abosede Wintola , and Anthony Jide Afolayan  
Research Article (13 pages), Article ID 6825297, Volume 2019 (2019)





## **Exploring Antimalarial Herbal Plants across Communities in Uganda Based on Electronic Data**

Denis Okello  and Youngmin Kang   
Review Article (27 pages), Article ID 3057180, Volume 2019 (2019)

## **Phytochemical Constituents, Antioxidant, Cytotoxicity, Antimicrobial, Antitrypanosomal, and Antimalarial Potentials of the Crude Extracts of *Callistemon citrinus***

Rotimi Larayetan , Zacchaeus S. Ololade, Oluranti O. Ogunmola, and Ayodele Ladokun  
Research Article (14 pages), Article ID 5410923, Volume 2019 (2019)

## **Antimalarial Activity of *Cordia africana* (Lam.) (Boraginaceae) Leaf Extracts and Solvent Fractions in *Plasmodium berghei*-Infected Mice**

Dawit Zewdu Wondafrash , Dayananda Bhoumik , Birhanetensay Masresha Altaye , Helen Bitew Tareke , and Brhane Teklebrhan Assefa  
Research Article (14 pages), Article ID 8324596, Volume 2019 (2019)



## Research Article

# Various Parts of *Helianthus annuus* Plants as New Sources of Antimalarial Drugs

Wiwied Ekasari <sup>1</sup>, Dwi Widya Pratiwi,<sup>1</sup> Zelmira Amanda,<sup>1</sup> Suciati,<sup>1</sup>  
Aty Widyawaruyanti <sup>1</sup> and Heny Arwati<sup>2</sup>

<sup>1</sup>Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Universitas Airlangga, Surabaya 60115, Indonesia

<sup>2</sup>Department of Medical Parasitology, Faculty of Medicine, Universitas Airlangga, Surabaya 60115, Indonesia

Correspondence should be addressed to Wiwied Ekasari; [wiwied-e@ff.unair.ac.id](mailto:wiwied-e@ff.unair.ac.id)

Received 27 June 2019; Revised 16 October 2019; Accepted 11 November 2019; Published 26 November 2019

Guest Editor: Philip F. Uzor

Copyright © 2019 Wiwied Ekasari et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Background.** Each part of *H. annuus* plants is traditionally used as medicinal remedies for several diseases, including malaria. Antimalarial activity of the leaf and the seed has already been observed; however, there is no report about antimalarial activity of the other parts of *H. annuus* plants. In this study, we assess *in vitro* and *in vivo* antimalarial activity of each part of the plants and its mechanism as antimalarial agent against inhibition of heme detoxification. **Objective.** To investigate the antimalarial activity of various parts of *H. annuus*. **Methods.** Various parts of the *H. annuus* plant were tested for *in vitro* antimalarial activity against *Plasmodium falciparum* 3D7 strain (chloroquine-sensitive), *in vivo* antimalarial activity against *P. berghei* using Peters' 4-day suppressive test in BALB/c mice, curative and prophylaxis assay, and inhibition of heme detoxification by evaluating  $\beta$ -hematin level. **Results.** Ethanol extract of the roots showed the highest antimalarial activity, followed by ethanol extract of leaves, with  $IC_{50}$  values of  $2.3 \pm 1.4$  and  $4.3 \pm 2.2$   $\mu$ g/mL, respectively and the percentage inhibition of *P. berghei* of  $63.6 \pm 8.0$  and  $59.3 \pm 13.2$  at a dose of 100 mg/kg, respectively. Ethanol extract of roots produced an  $ED_{50}$  value of  $10.6 \pm 0.2$  mg/kg in the curative test and showed an inhibition of 79.2% at a dose of 400 mg/kg in the prophylactic assay. In inhibition of heme detoxification assay, root and leaf ethanol extracts yielded a lower  $IC_{50}$  value than positive (chloroquine) control with a value of  $0.4 \pm 0.0$  and  $0.5 \pm 0.0$  mg/mL, respectively. **Conclusion.** There were promising results of the ethanol extracts of root of *H. annuus* as a new source for the development of a new plant-based antimalarial agent.

## 1. Introduction

Most developing countries, including Indonesia, still depend on using plants in traditional medicine, including antimalarial agents. One of these plants is *Helianthus annuus* (Asteraceae), known as sunflower or “bunga matahari” in local language. Traditionally, each part of *H. annuus* plants is widely used in several healthcare applications, including the treatment of malaria. Sunflower tea is reported to be useful in the treatment of malarial and lung disease [1], and the leaves have long been used in infusion and decoction as traditional remedy for malaria by Ambalabe village community, Madagascar [2].

Muti'ah et al. [3] report that 80% ethanol extract of *H. annuus* leaf shows good antimalarial activity with an  $ED_{50}$

value of 4.64 mg/kg. Moreover, the combination of 80% ethanol extract of *H. annuus* leaf and *Acalypha indica* Linn. leads to synergetic effect in *in vivo* antimalarial assay with a CI value of 0.46 ( $CI < 1$ ) and an  $ED_{50}$  value of 1.23 mg/kg [4]. Intisar et al. [5] also report that methanol and petroleum ether extract from *H. annuus* seeds showed good inhibition activity against *P. falciparum* strain KI with an  $EC_{50}$  of 0.19 and 0.34  $\mu$ g/mL, respectively.

Based on the data above, it is recognized that *H. annuus* plants are potential source of antimalarial drugs. However, antimalarial activity of the other parts of *H. annuus* plants has not been reported to date. Thus, in this study, we assess *in vitro* and *in vivo* antimalarial activity of each part of the plant and its mechanism as antimalarial agent against the inhibition of heme detoxification. The inhibition of heme

detoxification leads to growth disorder and malaria parasites' mortality as a result of membrane lysis and activity disorder of some enzymes [6–9]. Hence, a substance with high heme detoxification inhibition activity is considered a potential antimalarial agent. Inhibition of heme detoxification is one of the potential biochemical targets, which is possessed by chloroquine and artemisinin [10, 11]. The highest antimalarial activity of *H. annuus* plant parts and its ability to inhibit heme detoxification are expected to be obtained through this study as it can be a new source of antimalarial drugs.

## 2. Materials and Methods

**2.1. Plant Materials.** *H. annuus* plants were obtained from Oro-oro Ombo, Malang, East Java, Indonesia, in November 2016, and determined by Materia Medica Batu, Malang, East Java. Each part of the plant, including roots, stems, seeds, flowers, and leaves, were air-dried separately. Samples were then ground to powder, once they are completely dried.

**2.2. Animal Model.** Adult male BALB/c mice with 20–30 g of bodyweight were obtained from Faculty of Veterinary Medicine, Universitas Airlangga. The mice were fed with mice pellet diet and given free access to drink clean water. The animals were allowed to acclimatize for two weeks before being treated. The permission and approval for animal studies were obtained from Faculty of Veterinary Medicine, Universitas Airlangga.

**2.3. Preparation of 96% Ethanol Extract of *H. annuus*.** Powders from each part of *H. annuus* plants (50 gram) were macerated by 96% ethanol and then evaporated on a rotary evaporator.

**2.4. In Vitro Antimalarial Assay.** In this research, cultures of *Plasmodium falciparum* strain 3D7 (chloroquine-sensitive) were used and cultivated using the Trager and Jensen method [12]. Cultures were cultivated in human O<sup>+</sup> red blood cells with 5% hematocrit in RPMI 1640 (Gibco BRL, USA) added with 22.3 mM HEPES (Sigma), hypoxanthine, sodium bicarbonate, and 10% human O<sup>+</sup> plasma. Chloroquine diphosphate was used as the positive control.

Antimalarial assay was done using a 24-well microplate with 1% initial and experimental parasitaemia (1 mL/well of suspension), followed by incubation in a multigas incubator at 37°C for 48 hours. The incubated materials were then collected and used to make a thin smear in a glass slide, fixated in methanol, and stained with Giemsa. The number of parasites was then observed under a microscope and compared with the negative control to determine the percentage of parasite growth inhibition. IC<sub>50</sub> value was calculated using Probit Analysis.

**2.5. In Vivo Antimalarial Activity Assay.** *Plasmodium berghei* strain ANKA was obtained from Eijkman Institute of Molecular Biology, Jakarta, Indonesia. About 20% of

parasitaemia-infected mice were dissected and then the intracardial blood was taken and diluted with the PBS. Each mouse used in this research was inoculated priorly intraperitoneally with 0.2 mL of solution containing  $1 \times 10^7$  *P. berghei*-infected erythrocytes [13].

*In vivo* antimalarial assay of the plant extract was done using Peter's method [14], which had been modified with 100 mg/kg/day as the selected dose. Thirty-six adult male BALB/c mice (25 g ± 3) were inoculated with parasites and randomly grouped into six groups (five treatment groups and one control group), with six mice in each group. Negative control group was treated with 0.5% sodium carboxymethyl cellulose (Na CMC) suspension. Treatment groups were treated with extract suspension of each part of the plant with the selected dose. Suspension was given orally once a day for 4 days. On the fourth day, a thin blood smear was made from the tail of each mouse, fixated in methanol, and then stained with Giemsa. The thin blood smears were then observed under a microscope to calculate the percentage of parasitaemia and parasite inhibition.

**2.6. In Vivo Curative Assay of 96% Ethanol Extract of *H. annuus* Roots.** Thirty adult male BALB/c mice (8 weeks old) were randomly divided into five groups (six mice per group). Each mouse was infected intraperitoneally with 0.2 mL of solution containing  $1 \times 10^7$  *P. berghei*-infected erythrocytes. Negative control (group 1) was treated with 0.5% Na CMC suspension. Groups 2, 3, 4, and 5 were treated with 96% ethanol extract suspension of *H. annuus* root with 1, 10, 100, and 250 mg/kg dose per mouse. Suspension was given orally once a day for 4 days. Percentage of parasitaemia was calculated by making a thin blood smear on the fourth day, which is then fixated in methanol and stained with Giemsa. The calculation was then compared with the control group. ED<sub>50</sub> value was obtained by using Probit Analysis.

**2.7. In Vivo Prophylaxis Assay of 96% Ethanol Extract of *H. annuus* Roots.** Thirty adult male BALB/c mice (8 weeks old) were randomly grouped into six groups (five mice per group). Groups 1, 2, 3, and 4 were treated with 96% ethanol extract suspension of *H. annuus* root with 100, 200, 400, and 800 mg/kg dose per mouse. Group 5 as the negative control group was treated with 0.5% Na CMC suspension, while the positive control (group 6) was given 13 mg/kg of doxycycline suspension [15]. Each suspension was given orally once a day for 4 days. On the fourth day, each mouse was inoculated with 0.2 mL of solution containing  $1 \times 10^7$  *P. berghei*-infected erythrocytes by intraperitoneal injection. Percentage of parasitaemia was determined on the third day by observing the thin blood smear and compared with the control group. Mean survival time of mice in each group was determined during 14 days (D<sub>0</sub>–D<sub>13</sub>). Mean survival time was calculated using the following equation:

$$\text{mean survival time (MST)} = \frac{\text{sum of survival days}}{\text{total observation days (14)}} \times 100. \quad (1)$$

**2.8. Heme Detoxification Inhibition Assay.** Inhibition of heme detoxification was assessed using the Basilico method [16] with modification. The extract and positive control (chloroquine diphosphate) were diluted with dimethyl sulfoxide (DMSO) a concentration range of 2–0.01 mg/mL. The amount of hematin present in each sample was calculated using a standard curve of hematin dissolved in 0.2 M NaOH. The absorbance was read using the ELISA reader at a wavelength of 405 nm. The effect of each sample against the production of  $\beta$ -hematin was calculated and compared with the negative control [17, 18].

**2.9. Statistical Analysis.** Data are expressed as mean  $\pm$  standard deviation (SD).  $IC_{50}$  values were calculated using Probit Analysis. Statistical significance was determined by Kruskal–Wallis nonparametric independent  $t$ -test and ANOVA (*one-way*).

### 3. Results and Discussion

*H. annuus* has been planted widely in Indonesia and long been used traditionally as anti-inflammatory, antimalarial, antiasthma, antioxidant, antitumor, and antimicrobial agent. It has been reported that several chemical compounds were already isolated and identified from various parts of this plant, such as heliangolides (sesquiterpene lactones),  $\alpha$ -pinene, sabinene (monoterpene), helikauranoides (diterpene), and alkaloid and phenolic group in the leaf part; helianthoside (triterpene) and saponin group in the flower part; fatty acid, tocopherol, tannin, and polyphenol group in the seed part; alkaloid and phenolic group in the stem and root parts [1]. Extraction yield from each part of *H. annuus* plants macerated with 96% ethanol is shown in Table 1.

*In vitro* screening to observe *P. falciparum* inhibition activity of the plant extract with  $IC_{50}$  value less than 10  $\mu$ g/mL is an important primary step in the development of new antimalarial drugs. Plant extracts showing  $IC_{50}$  values  $\leq 10$   $\mu$ g/mL can be considered to be active,  $IC_{50}$  values in the range  $10 < IC_{50} \leq 25$   $\mu$ g/mL can be considered to be moderately active, and those exhibiting  $IC_{50}$  values  $> 25$   $\mu$ g/mL can be considered to be inactive [19].  $IC_{50}$  values of 96% ethanol extract of various *H. annuus* plant parts are shown in Table 2. Root extract exhibits the highest *in vitro* antimalarial activity, followed by leaf and flower extracts with an  $IC_{50}$  value of  $2.3 \pm 1.4$ ,  $4.3 \pm 2.2$ , and  $4.8 \pm 0.0$   $\mu$ g/mL, respectively.

*In vivo* antimalarial activity was also determined for each extract from various parts of *H. annuus* plants, and the results are shown in Table 3. Ethanol extract of root exhibits the highest percentage inhibition against *P. berghei*, followed by leaf extract with the percentage inhibition of  $63.6 \pm 8.0$  and  $59.3 \pm 13.2$ , respectively.

Method and criteria are varied among the treatment groups examining antimalarial potency of plants by using the rodent animal model. *P. berghei*-infected mice given orally 50–250 mg/kg/day of extract exhibiting inhibition percentage  $> 60\%$  are considered to be active or very active, and those exhibiting inhibition percentage  $> 30\%$  are considered to be moderately active [20–24]. In this research, oral

TABLE 1: Extraction yield from various parts of *H. annuus* plants.

<i>H. annuus</i> plant parts	Extraction yield (%)
Roots	2.1
Stems	5.0
Seeds	2.9
Flowers	6.7
Leaves	4.4

administration at a dose of 100 mg/kg/day of ethanol *H. annuus* extract is used as an initial experiment and observation of parasite growth up to day 4 using mice model.

Based on the first screening of *in vivo* assay, it is shown that *H. annuus* root extract has the highest activity. Thus, the experiment is continued to assess curative and prophylaxis antimalarial activity of the ethanol root extract of *H. annuus*.

In *in vivo* curative assay, *P. berghei*-infected mice are treated orally with 96% ethanol root extract at a dose of 1, 10, 100, and 250 mg/kg. The results of the 4-day suppressive test show that the ethanol extract of *H. annuus* root has an  $ED_{50}$  value of  $10.6 \pm 0.2$  mg/kg (Table 4). According to Munoz et al. [25], antimalarial activity can be categorized based on the  $ED_{50}$  value, where  $ED_{50} < 100$  mg/kg is considered to have very good activity.

In prophylactic effect evaluation of 96% ethanol *H. annuus* root extract, mice are given a sample orally for 4 days before inoculating with the parasite. The results (see Figure 1) show that the percentage of inhibition parasitaemia in mice treated with the extract at a dose of 400 mg/kg is higher than that in those treated with 800 mg/kg. Despite that, this result is still lower than that of doxycycline, a standard drug for prophylactic antimalarial in Indonesia.

The 400 mg/kg dose group shows the highest percentage of prophylactic inhibition (79.2%) at day 3, and it declines to 62.5% at a dose of 800 mg/kg. Calabrese and Baldwin [26] report that the upside-down U-curve in the drug activity assessment at different doses may indicate the occurrence of the hormetic effect, which points out to the toxicity of the drug. Aside from the declining percentage of inhibition, the low mean survival time at a dose of 800 mg/kg, compared to 200 and 400 mg/kg doses, indicates that the *H. annuus* root extract at high dose (800 mg/kg) tends to have toxicity.

The results of Kruskal–Wallis nonparametric independent  $t$ -test and *one-way* ANOVA analysis show that there are significant differences ( $p$  value  $< 0.05$ ) in inhibition percentage of the positive control group compared with all the sample groups and the negative control group. There is no significant difference between 100, 200, 400, and 800 mg/kg doses.

Figure 1 shows that, 96 hours or the fourth day after parasite inoculation, there is an increase in the parasitic growth in the negative dose and control group, whereas in the doxycycline group the parasite growth is relatively stable, which is below 3%. This is thought to be related to the half-life of doxycycline which ranges from 12 to 22 hours [15, 27, 28] allowing doxycycline to last longer in the body and inhibit the growth of parasites. It is in contrast to extracts or traditional medicines that tend to have a short half-life, such as artemisinin and its derivatives [29, 30]. Based on

TABLE 2: IC<sub>50</sub> values of 96% ethanol extract of various parts of *H. annuus* plants against *P. falciparum* strain 3D7.

<i>H. annuus</i> plant parts	% inhibition at each concentration ( $\mu\text{g/mL}$ )					IC <sub>50</sub> ( $\mu\text{g/mL}$ )
	100	10	1	0.1	0.01	
Roots	86.6 $\pm$ 1.7	73.3 $\pm$ 6.9	32.0 $\pm$ 3.7	22.3 $\pm$ 0.0	7.3 $\pm$ 7.4	2.3 $\pm$ 1.4
Stems	64.4 $\pm$ 4.6	56.4 $\pm$ 1.1	30.7 $\pm$ 7.7	18.8 $\pm$ 4.9	0	10.2 $\pm$ 5.0
Seeds	84.4 $\pm$ 1.0	22.2 $\pm$ 6.9	14.2 $\pm$ 3.9	6.2 $\pm$ 10.3	0	19.3 $\pm$ 5.5
Flowers	85.7 $\pm$ 8.7	48.4 $\pm$ 2.6	32.0 $\pm$ 2.9	19.3 $\pm$ 8.7	0	4.8 $\pm$ 0.0
Leaves	96.3 $\pm$ 1.8	47.1 $\pm$ 4.7	24.3 $\pm$ 11.7	16.5 $\pm$ 6.2	7.0 $\pm$ 1.0	4.3 $\pm$ 2.2

TABLE 3: Mean percentage of parasitaemia, growth, and inhibition in *P. berghei*-infected mice treated with 96% ethanol *H. annuus* extract at a dose of 100 mg/kg orally once a day for 4 days.

<i>H. annuus</i> plant parts	Mean % parasitaemia		Mean % growth	Mean % inhibition
	Day 0	Day 4		
Roots	0.9 $\pm$ 0.2	4.6 $\pm$ 1.0	3.7 $\pm$ 0.8	63.6 $\pm$ 8.0
Stems	1.2 $\pm$ 0.2	7.3 $\pm$ 1.0	6.1 $\pm$ 0.8	40.3 $\pm$ 8.2
Seeds	1.0 $\pm$ 0.2	7.1 $\pm$ 0.5	6.0 $\pm$ 0.4	41.5 $\pm$ 4.4
Flowers	1.2 $\pm$ 0.5	7.8 $\pm$ 0.4	6.6 $\pm$ 0.5	35.9 $\pm$ 4.6
Leaves	0.9 $\pm$ 0.2	5.1 $\pm$ 1.5	4.2 $\pm$ 1.4	59.3 $\pm$ 13.2
Na CMC	1.6 $\pm$ 0.3	11.8 $\pm$ 0.9	10.3 $\pm$ 0.9	—

TABLE 4: Mean percentage of parasitaemia, growth, and inhibition against *P. berghei* of 96% ethanol *H. annuus* root extract compared with control.

Sample	Dose (mg/kg)	Mean % parasitaemia		Mean % growth	Mean % inhibition	ED <sub>50</sub> (mg/kg)
		D <sub>0</sub>	D <sub>4</sub>			
		96% ethanol extract of <i>H. annuus</i> root	1			
	10	1.0 $\pm$ 0.2	21.2 $\pm$ 0.9	5.1 $\pm$ 0.2	48.3 $\pm$ 2.8	
	100	1.0 $\pm$ 0.1	15.4 $\pm$ 1.7	3.6 $\pm$ 0.4	64.0 $\pm$ 5.3	
	250	1.2 $\pm$ 0.4	12.0 $\pm$ 1.8	2.7 $\pm$ 0.4	72.3 $\pm$ 5.3	
Na CMC	—	1.2 $\pm$ 0.3	40.3 $\pm$ 6.1	9.8 $\pm$ 1.5	—	—


FIGURE 1: Percentage of parasite growth during seven days after inoculation with parasites on *in vivo* prophylaxis assay of 96% ethanol extract of *H. annuus* roots. Average values and standard deviations from five mice per drug/dose are also shown.  $p$  value  $< 0.05$ . EEHA = ethanolic extract of *H. annuus*.

TABLE 5: Results of heme detoxification inhibition assay of the 96% ethanol extract of each part of the *H. annuus* plant.

Sample	Concentration (mg/mL)	Mean value of hematin (mM)	Mean % inhibition	IC <sub>50</sub> (mg/mL)
Negative control	—	121.0 ± 2.6	—	—
	2.00	34.2 ± 1.2	71.7 ± 1.0	
	1.00	44.6 ± 1.2	63.1 ± 1.0	
	0.50	58.3 ± 1.1	51.8 ± 0.9	0.4 ± 0.0
	0.25	67.6 ± 1.3	44.1 ± 1.1	
	0.10	81.8 ± 1.3	32.3 ± 1.1	
96% ethanol extract of root	0.01	99.8 ± 2.2	17.5 ± 1.8	
	2.00	61.6 ± 1.4	49.1 ± 1.1	
	1.00	71.0 ± 1.2	41.3 ± 1.0	
	0.50	84.9 ± 3.3	29.8 ± 2.8	2.4 ± 0.2
	0.25	97.3 ± 2.3	19.6 ± 1.9	
	0.10	105.5 ± 1.3	12.8 ± 1.1	
96% ethanol extract of seed	0.01	114.1 ± 1.8	5.7 ± 1.5	
	2.00	46.8 ± 1.1	61.4 ± 0.9	
	1.00	55.7 ± 1.3	53.9 ± 1.1	
	0.50	68.4 ± 0.9	43.4 ± 0.7	0.9 ± 0.1
	0.25	81.8 ± 0.9	32.4 ± 0.7	
	0.10	89.8 ± 0.9	25.8 ± 0.8	
96% ethanol extract of flower	0.01	99.3 ± 1.2	17.9 ± 1.0	
	2.00	38.6 ± 0.9	68.1 ± 0.7	
	1.00	48.9 ± 0.8	59.6 ± 0.7	
	0.50	62.0 ± 1.1	48.7 ± 0.9	0.5 ± 0.0
	0.25	75.5 ± 1.4	37.6 ± 1.1	
	0.10	83.5 ± 0.6	31.0 ± 0.5	
96% ethanol extract of leaf	0.01	98.1 ± 0.9	18.9 ± 0.7	
	2.00	140.1 ± 9.2	53.7 ± 3.1	
	1.00	179.8 ± 5.6	40.6 ± 1.8	
	0.50	206.3 ± 8.3	31.8 ± 2.8	1.9 ± 0.5
	0.25	241.8 ± 3.6	20.1 ± 1.2	
	0.10	264.7 ± 3.6	12.5 ± 1.2	
96% ethanol extract of stem	0.01	288.0 ± 5.5	4.8 ± 1.8	
	2.00	40.4 ± 2.1	66.6 ± 1.7	
	1.00	50.9 ± 0.6	57.9 ± 0.5	
	0.50	64.3 ± 0.9	46.8 ± 0.7	0.6 ± 0.0
	0.25	77.0 ± 0.8	36.3 ± 0.7	
	0.10	85.3 ± 1.2	29.4 ± 1.0	
Chloroquine diphosphate	0.01	94.6 ± 0.7	21.8 ± 0.6	

the antimalarial test results of *H. annuus* root *in vivo*, it can be concluded that the 96% ethanol extract of *H. annuus* root has more potential activities as curative antimalarial than prophylactic.

One of the potential biochemical targets for the development of new antimalarial drugs is heme detoxification. The results of heme detoxification inhibition assay (see Table 5) show that the 96% ethanol extract of *H. annuus* root (IC<sub>50</sub> = 0.4 ± 0.0 mg/mL) and leaf (IC<sub>50</sub> = 0.5 ± 0.0 mg/mL) has higher activity than chloroquine diphosphate, a standard antimalarial drug (IC<sub>50</sub> = 0.6 ± 0.0 mg/mL).

It was reported that free heme is toxic to parasites as it causes lysis of the parasite vacuolar membrane [31]. Thus, malaria parasites develop heme detoxification mechanism to change the toxic heme into a harmless one. The inhibition of heme detoxification leads to growth disorder and malaria parasites mortality as a result of membrane lysis and activity disorder of some enzymes [6–9].

Based on the entire assay undertaken in this research, it is established that the ethanol extract of *H. annuus* root shows the highest antimalarial activity among the other parts of plants, either in *in vitro*, *in vivo*, or in heme detoxification inhibition assay. Therefore, the ethanol extract of *H. annuus* root is a potential source of new antimalarial drugs from natural resources.

#### 4. Conclusion

Based on the results of the study, they show that 96% ethanol extract of *H. annuus* root has the highest antimalarial activity compared to stems, seeds, flowers, and leaves of *H. annuus*. 96% ethanol extract of *H. annuus* root has more potential activity as a curative antimalarial drug than a prophylactic one. Similarly, for the antimalarial mechanism in inhibiting heme detoxification, 96% ethanol extract of *H. annuus* root has the greatest inhibitory activity compared

to the other parts of *H. annuus*; even the inhibition is greater than chloroquine, a standard drug. So, it can be concluded that the 96% ethanol extract of *H. annuus* root contains a potential antimalarial substance which is a new source for the development of a new plant-based antimalarial agent.

### Data Availability

All data are included within the manuscript.

### Conflicts of Interest

The authors declare that they have no conflicts of interest.

### Acknowledgments

The authors would like to thank “Riset Mandat” of Universitas Airlangga Scheme for financial support.

### References

- [1] S. Saini and S. Sharma, “*Helianthus annuus* (Asteracea): a review,” *International Journal of Pharma Professional’s Research*, vol. 2, no. 4, pp. 381–386, 2011.
- [2] N. H. Rakotoarivelo, F. Rakotoarivony, A. V. Ramarosandratana et al., “Medicinal plants used to treat the most frequent diseases encountered in Ambalabe rural community, Eastern Madagascar,” *Journal of Ethnobiology and Ethnomedicine*, vol. 11, no. 1, p. 68, 2015.
- [3] R. Muti’ah, E. K. Hayati, and Khairunnisak, “The effectivity of ethanolic extract sunflower leaves (*Helianthus annuus*) as anti-malarial agent against *Plasmodium berghei*,” in *Proceedings of the International Conference, The 4th Green Technology Faculty of Science and Technology*, Islamic of University State Maulana Malik Ibrahim Malang, Malang, Indonesia, November 2013.
- [4] K. Ngibad, “The effectiveness of the combination of ethanol extract of sunflower leaves (*Helianthus annuus*) and anting-anting (*Acalypha indica* Linn) as an antimalarial in vivo,” *Journal of Galician Pharmacy*, vol. 5, no. 1, pp. 12–19, 2019.
- [5] E. M. Intisar, E. K. Hassan, A. M. Salah et al., “Anti-malarial activity of some medicinal Sudanese plants,” *Journal of Forest Products and Industries*, vol. 3, no. 6, pp. 236–240, 2014.
- [6] I. W. Sherman, *Malaria, Parasite Biology, Pathogenesis and Protection*, ASM Press, Washington, DC, USA, 1998.
- [7] H. L. Ziegler, D. Staerk, J. Christensen, L. Hviid, H. Hagerstrand, and J. W. Jaroszewski, “In vitro *Plasmodium falciparum* drug sensitivity assay: inhibition of parasite growth by incorporation of stomatocytogenic amphiphiles into erythrocyte membrane,” *Antimicrobial Agents and Chemotherapy*, vol. 46, no. 5, pp. 1441–1446, 2002.
- [8] S. Auparakkitanon, W. Noonpakdee, R. K. Ralph, W. A. Denny, and P. Wilairat, “Antimalarial 9-anilinoacridine compound directed at hemozoin,” *Antimicrobial Agents and Chemotherapy*, vol. 47, no. 12, pp. 3708–3712, 2003.
- [9] B. Tekwani and L. Walker, “Targeting the hemozoin synthesis pathway for the new antimalarial drug discovery: technologies for in vitro  $\beta$ -hemozoin formation assay,” *Combinatorial Chemistry & High Throughput Screening*, vol. 8, no. 1, pp. 63–79, 2005.
- [10] A. V. Pandey, H. Bisht, V. K. Babbarwal, J. Srivastava, K. C. Pandey, and V. S. Chauhan, “Mechanism of malarial haem detoxification inhibition by chloroquine,” *Biochemical Journal*, vol. 355, no. 2, pp. 333–338, 2001.
- [11] A. V. Pandey, B. L. Tekwani, R. L. Singh, and V. S. Chauhan, “Artemisinin, and endoperoxide antimalarial, disrupts the hemoglobin catabolism and heme detoxification systems in malarial parasite,” *Journal of Biological Chemistry*, vol. 274, no. 27, pp. 19383–19388, 1999.
- [12] W. Trager and J. Jensen, “Human malaria parasites in continuous culture,” *Science*, vol. 193, no. 4254, pp. 673–675, 1976.
- [13] W. Ekasari, T. S. Wahyuni, H. Arwati, and N. T. Putri, “Determination of effective dose of antimalarial from *Cassia spectabilis* leaf ethanol extract in *Plasmodium berghei*-infected mice,” *African Journal of Infectious Disease*, vol. 12, no. suppl. 1, pp. 110–115, 2018.
- [14] W. Peters, J. H. Portus, and B. L. Robinson, “The chemotherapy of rodent malaria. XXII. The value of drug-resistant strains of *P. berghei* in screening for blood schizontocidal activity,” *Annals of Tropical Medicine & Parasitology*, vol. 69, no. 2, pp. 155–171, 1975.
- [15] K. R. Tan, A. J. Magill, P. M. Arguin, and M. E. Parise, “Doxycycline for malaria chemoprophylaxis and treatment: report from the CDC expert meeting on malaria chemoprophylaxis,” *The American Journal of Tropical Medicine and Hygiene*, vol. 84, no. 4, pp. 517–531, 2011.
- [16] N. Basilico, E. Pagani, D. Monti et al., “A microtitre-based method for measuring the haem polymerization inhibitory activity (HPIA) of antimalarial drugs,” *Journal of Antimicrobial Chemotherapy*, vol. 42, no. 1, pp. 55–60, 1998.
- [17] T. S. Macedo, L. Colina-Vegas, M. Da Paixão et al., “Chloroquine-containing organoruthenium complexes are fast-acting multistage antimalarial agents,” *Parasitology*, vol. 143, no. 12, pp. 1543–1556, 2016.
- [18] S. Joshi, R. Munshi, G. Talele et al., “An experimental in vitro study to evaluate the antimalarial activity of select homeopathy preparations,” *International Journal of Medical and Health Research*, vol. 3, no. 7, pp. 65–68, 2017.
- [19] R. B. S. Lima, L. F. Rocha e Silva, M. R. S. Melo et al., “In vitro and in vivo anti-malarial activity of plants from the Brazilian Amazon,” *Malaria Journal*, vol. 14, no. 1, p. 508, 2015.
- [20] L. F. Rocha e Silva, K. L. Nogueira, A. C. da Silva Pinto et al., “In vivo antimalarial activity and mechanisms of action of 4-nerolidylcatechol derivatives,” *Antimicrobial Agents and Chemotherapy*, vol. 59, no. 6, pp. 3271–3280, 2015.
- [21] H. C. Upadhyay, B. S. Sisodia, J. Agrawal, A. Pal, M. P. Darokar, and S. K. Srivastava, “Antimalarial potential of extracts and isolated compounds from four species of genus *Ammannia*,” *Medicinal Chemistry Research*, vol. 23, no. 2, pp. 870–876, 2014.
- [22] M. F. Musila, S. F. Dossaji, J. M. Nguta, C. W. Lukhoba, and J. M. Munyao, “In vivo antimalarial activity, toxicity and phytochemical screening of selected antimalarial plants,” *Journal of Ethnopharmacology*, vol. 146, no. 2, pp. 557–561, 2014.
- [23] J. M. Nguta and J. M. Mbaria, “Brine shrimp toxicity and antimalarial activity of some plants traditionally used in treatment of malaria in Msambweni district of Kenya,” *Journal of Ethnopharmacology*, vol. 148, no. 3, pp. 988–992, 2013.
- [24] J. W. Gathirwa, G. M. Rukunga, P. G. Mwitari et al., “Traditional herbal antimalarial therapy in Kilifi district, Kenya,” *Journal of Ethnopharmacology*, vol. 134, no. 2, pp. 434–442, 2011.

- [25] V. Muñoz, M. Sauvain, G. Bourdy et al., "The search for natural bioactive compounds through a multidisciplinary approach in Bolivia. Part II. Antimalarial activity of some plants used by Mosaic Indians," *Journal of Ethnopharmacology*, vol. 69, no. 2, pp. 139–155, 2000.
- [26] E. J. Calabrese and L. A. Baldwin, "Hormesis: U-shaped dose responses and their centrality in toxicology," *Trends in Pharmacological Sciences*, vol. 22, no. 6, pp. 285–291, 2001.
- [27] B. A. Cunha, J. B. Comer, and M. Jonas, "The tetracyclines," *Medical Clinics of North America*, vol. 66, no. 1, pp. 293–302, 1982.
- [28] P. G. Welling, P. A. Koch, C. C. Lau, and W. A. Craig, "Bioavailability of tetracycline and doxycycline in fasted and nonfasted subjects," *Antimicrobial Agents and Chemotherapy*, vol. 11, no. 3, pp. 462–469, 1977.
- [29] Z. Guo, "Artemisinin anti-malarial drugs in China," *Acta Pharmaceutica Sinica B*, vol. 6, no. 2, pp. 115–124, 2016.
- [30] K. Karunamoorthi, S. Sabesan, K. Jegajeevanram, and J. Vijayalakshmi, "Role of traditional antimalarial plants in the battle against the global malaria burden," *Vector-Borne and Zoonotic Diseases*, vol. 13, no. 8, pp. 521–544, 2013.
- [31] A. C. Chou and C. D. Fitch, "Mechanism of hemolysis induced by ferriprotoporphyrin IX," *Journal of Clinical Investigation*, vol. 68, no. 3, pp. 672–677, 1981.



**KOMISI ETIK PENELITIAN  
FAKULTAS KEDOKTERAN HEWAN UNIVERSITAS AIRLANGGA  
*Animal Care and Use Committee (ACUC)***

**KETERANGAN KELAIKAN ETIK  
“ ETHICAL CLEARANCE ”**

**No : 2.KE.119.07.2018**

**KOMISI ETIK PENELITIAN (ANIMAL CARE AND USE COMMITTEE)  
FAKULTAS KEDOKTERAN HEWAN UNIVERSITAS AIRLANGGA SURABAYA,  
TELAH MEMPELAJARI SECARA SEKSAMA RANCANGAN PENELITIAN YANG  
DIUSULKAN, MAKA DENGAN INI MENYATAKAN BAHWA :**

**PENELITIAN BERJUDUL** : Aktivitas Profilaksis Antimalaria Ekstrak Etanol 96%  
Akar *Heliathus annuus* L Secara *In Vivo* Pada  
Mencit Balb/C

**PENELITI UTAMA** : Zelmira Amanda

**UNIT/LEMBAGA/TEMPAT  
PENELITIAN** : Departemen Farmakognosi dan Fitokimia  
Fakultas Farmasi Universitas Airlangga

**DINYATAKAN** : LAIK ETIK

Surabaya, 19 Juli 2018

Mengetahui,  
Dekan FKH-Unair,

Prof. Dr. Pudji-Srianto, M.Kes.,Drh.  
NIP. 195601051986011001

Ketua,

Dr. Nusdianto Triakoso, M.P.,Drh.  
NIP. 196805051997021001



## Evidence-based Complementary and Alternative Medicine

COUNTRY

United States

 Universities and research institutions in United States

PUBLICATION TYPE

Journals

SUBJECT AREA AND CATEGORY

Medicine  
└ Complementary and Alternative Medicine

ISSN

1741427X, 17414288

PUBLISHER

Hindawi Publishing Corporation

COVERAGE

2005-2021

H-INDEX

**100**


INFORMATION

[Homepage](#)

[How to publish in this journal](#)

Quartiles


Complementary and Alternative Medicine



SJR

2013	0.614
2014	0.747
2015	0.784
2016	0.747
2017	0.683
2018	0.590
2019	0.510
2020	0.552
2021	0.461

Total Documents





## Source details

[Feedback >](#) [Compare sources >](#)

### Evidence-based Complementary and Alternative Medicine

Open Access ⓘ

Scopus coverage years: **from 2005 to Present**

Publisher: Hindawi

ISSN: 1741-427X E-ISSN: 1741-4288

Subject area: Medicine: Complementary and Alternative Medicine

Source type: Journal

[View all documents >](#)

[Set document alert](#)

[Save to source list](#)

[Source Homepage](#)

CiteScore 2021

3.2 ⓘ

SJR 2021

0.461 ⓘ

SNIP 2021

0.901 ⓘ

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)



#### Improved CiteScore methodology

CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)



CiteScore 2021



$$3.2 = \frac{13,705 \text{ Citations 2018 - 2021}}{4,268 \text{ Documents 2018 - 2021}}$$

Calculated on 05 May, 2022


CiteScoreTracker 2022 ⓘ

$$2.8 = \frac{11,432 \text{ Citations to date}}{4,118 \text{ Documents to date}}$$

Last updated on 06 June, 2022 • Updated monthly


#### CiteScore rank 2021 ⓘ

Category	Rank	Percentile
Medicine Complementary and Alternative Medicine	#26/93	



Scimago Journal & Country Rank

also developed by scimago:



SCIMAGO INSTITUTIONS RANKINGS

Enter Journal Title, ISSN or Publisher Name

Home

Journal Rankings

Country Rankings

Viz Tools

Help

About Us

Medicine

Complementary and Alternative Medicine

United States




Journals

2019

Only Open Access Journals
  Only SciELO Journals
  Only WoS Journals

Display journals with at least 0 Citable Docs. (3years)  Apply

1 - 7 of 7

Title	Type	↓ SJR	H index	Total Docs. (2019)	Total Docs. (3years)	Total Refs. (2019)	Total Cites (3years)	Citable Docs. (3years)	Cites / Doc. (2years)	Ref. / Doc. (2019)	
1 Integrative Cancer Therapies	journal	0.708 <span style="background-color: #90EE90; padding: 2px;">Q1</span>	63	121	253	5541	636	244	2.46	45.79	
2 Complementary Therapies in Medicine	journal	0.576 <span style="background-color: #90EE90; padding: 2px;">Q1</span>	69	285	435	12040	1043	415	2.08	42.25	
3 Evidence-based Complementary and Alternative Medicine	journal	0.510 <span style="background-color: #90EE90; padding: 2px;">Q1</span>	100	786	2065	36451	4190	2022	1.83	46.38	
4 American Journal of Clinical Hypnosis	journal	0.266 <span style="background-color: #FFD700; padding: 2px;">Q2</span>	29	9	113	457	48	98	0.33	50.78	