


NATURAL PRODUCT COMMUNICATIONS

An International Journal for Communications and Reviews Covering All Aspects of Natural Products Research



Natural Product Communications



Editorial Board

[Hide All](#)

Editor-in-Chief

Dr. Pawan K. Agrawal

Natural Product, Inc

Honorary Editor

Professor Gerald Blunden

University of Portsmouth, UK

Editors

Professor Vladimir I. Kalinin

G.B. Elyakov Pacific Institute of Bioorganic Chemistry, Russia

Professor Phan Van Kiem

Vietnam Academy of Science and Technology (VAST), Vietnam

Professor Carmen Martin-Cordero

University of Seville, Spain

Professor Manfred G. Reinecke

Texas Christian University, USA

Professor Sergio Rosselli

Università Degli Studi di Palermo, Italy

Professor Tadaaki Satou

International University of Health and Welfare, Japan

Professor William N. Setzer

University of Alabama in Huntsville, USA

Professor Ping-Jyun Sung

National Museum of Marine Biology and Aquarium

Professor Yasuhiro Tezuka

Hokuriku University, Japan

Professor David E. Thurston

King's College London, UK

Coordinating Editor

Chandan Agrawal

Natural Product, Inc

Advisory Editorial Board

Professor Giovanni Appendino

Università del Piemonte Orientale, Italy

Professor Yoshinori Asakawa

Tokushima Bunri University, Japan

Professor Vassya Bankova

Bulgarian Academy of Science, Bulgaria

Professor Anna R. Bilia	University of Florence, Italy
Professor Maurizio Bruno	University of Palermo, Italy
Professor Geoffrey Cordell	Natural Products Inc.
Professor Fatih Demirci	Anadolu University, Turkey
Professor Francesco Epifano	D'Annunzio University of Chieti-Pescara, Rovigo, Italy
Professor Antonio Evidente	University of Naples "Federico II", Napoli, Italy
Professor Ana Cristina Figueiredo	Universidade de Lisboa, Portugal
Professor Mary J. Garon	The University of Queensland, Australia
Professor Cristina Gracia-Viguera	CEBAS-CSIC, Spain
Dr. Christopher Gray	University of New Brunswick, Canada
Professor Dominique Guillaume	Université de Reims-Champagne-Ardenne, France
Professor Hisahiro Hagiwara	Niigata University, Japan
Professor Judith Hohmann	University of Szeged, Hungary
Professor Tsukasa Iwashina	National Museum of Nature and Science, Japan
Professor Niel A. Koorbanally	University of KwaZulu-Natal, South Africa
Professor George A. Kraus	Iowa State University, USA
Professor Chiaki Kuroda	Rikkyo University, Japan
Professor M. A. Lacaille-Dubois	Université de Bourgogne, France
Professor Shoei-Sheng Lee	National Taiwan University
Professor Yoshihiro Mimaki	Tokyo University of Pharmacy and Life Sciences (TUPL), Japan
Professor Robert W. Owen	German Cancer Research Centre, Germany
Professor M. Soledade C. Pedras	University of Saskatchewan, Canada
Professor Luc Pieters	University of Antwerp, Belgium
Professor Phila Raharivelomanana	Université de la Polynésie Française, French Polynesia
Professor Jesús M. Rodilla	Universidade da Beira Interior, Covilhã, Portugal
Professor Stefano Serra	C.N.R.-Istituto di Chimica del Riconoscimento Molecolare, Italy
Dr. Bikram Singh	CSIR-Institute of Himalayan Bioresource, India
Professor Alexios-Leandros Skaltsounis	University of Athens, Greece
Professor John L. Sorensen	University of Manitoba, Canada
Professor Marina Stefova	Saints Cyril and Methodius University of Skopje, Macedonia
Professor Valentin Stonik	G.B. Elyakov Pacific Institute of Bioorganic Chemistry, Russia
Professor Félix Tomi	Université de Corse-CNRS, France
Professor Karin Valant-Vetschera	University of Vienna, Austria
Professor Johannes van Staden	University of KwaZulu-Natal, South Africa
Dr. Alessandro Venditti	Università di Roma "La Sapienza", Roma, Italy
Assoc. Professor Yasunori Yaoita	Tohoku Medical and Pharmaceutical University, Japan

Natural Product Communications

Table of Contents

Volume 6 Issue 3, March 2011

Article

 **Cytotoxic Bibenzyls, and Germacrane- and Pinguisane-type Sesquiterpenoids from Indonesian, Tahitian and Japanese Liverworts**

Ismiarni Komala, Takuya Ito, Fumihiro Nagashima, Yasuyuki Yagi, Yoshinori Asakawa

First Published March 1, 2011

[Abstract](#)
> [Preview](#)



 **Total Synthesis of the Bicyclic Marine Sesquiterpenoid Drechslerine B**

Hisahiro Hagiwara, Masakazu Fukushima, Kimihiko Kinugawa, Takuya Matsui, Takashi Hoshi, Toshio Suzuki

First Published March 1, 2011

[Abstract](#)
> [Preview](#)



 **Chemosystematics of *Porella* (Marchantiophyta, Porellaceae)**

Agnieszka Ludwiczuk, S. Rob Gradstein, Fumihiro Nagashima, Yoshinori Asakawa

First Published March 1, 2011

[Abstract](#)
> [Preview](#)



 **Melidianolic Acid A and B, New Antimalarial Acyclic Diterpenes from *Aphanamixis grandifolia***

Adil Astulla, Yusuke Hirasawa, Abdul Rahman, Idha Kusumawati, Wiwied Ekasari, Aty Widyawaruyanti, Noor Cholies Zaini, Hiroshi Morita

First Published March 1, 2011

[Abstract](#)
> [Preview](#)



Melidianolic Acid A and B, New Antimalarial Acyclic Diterpenes from *Aphanamixis grandifolia*

Adil Astulla^a, Yusuke Hirasawa^a, Abdul Rahman^b, Idha Kusumawati^b, Wiwied Ekasari^b, Aty Widayawaruyanti^b, Noor Cholies Zaini^b and Hiroshi Morita^{a,*}

^aFaculty of Pharmaceutical Sciences, Hoshi University, Ebara 2-4-41 Shinagawa-ku, Tokyo 142-8501, Japan

^bFaculty of Pharmacy, Airlangga University, Jalan Dharmawangsa Dalam, Surabaya 60286, Indonesia

moritah@hoshi.ac.jp

Received: December 13th, 2010; Accepted: January 12th, 2011


Two new acyclic diterpenes, melidianolic acids A (**1**) and B (**2**), have been isolated from the bark of *Aphanamixis grandifolia*. Their structures were elucidated on the basis of spectroscopic and chemical methods. Melidianolic acids A (**1**) and B (**2**) showed antimalarial activity against *Plasmodium falciparum* 3D7 with IC₅₀ of 6.1 and 7.3 μg/mL, respectively.

Keywords: *Aphanamixis grandifolia*, melidianolic acids A and B, acyclic diterpene, antimalarial activity, *Plasmodium falciparum* 3D7.

The Meliaceae is a flowering plant of mostly trees, and shrub and a few herbaceous plants, native to tropical and subtropical regions. Most species, comprising about 50 genera and 550 species, are evergreen, but some are deciduous [1]. Meliaceae is of particular interest because of the presence of abundance and structural diversity of the highly oxygenated tetranortriterpenoids, limonoids, with a range of biological activities such as insecticidal, insect antifeedant, antibacterial, antifungal, antimalarial, anticancer, and antiviral activities [1]. Some plants belonging to Meliaceae have traditionally been employed as fish or dart arrow poisons, and afforded a several toxic components [2,3].

Aphanamixis grandifolia is a timber tree occurring in Java, Indonesia, and produces poisonous fruits. The peel and seeds of *A. grandifolia* were known to contain sesquiterpenes [4] and limonoids [5-7], which have inhibited the growth of the P388 cell line [7]. Our continuing search for biogenetically interesting compounds from *A. grandifolia* led to isolation of two new acyclic diterpenes, melidianolic acids A (**1**) and B (**2**). This paper describes the isolation and structure elucidation of **1** and **2** as well as antimalarial activity against *Plasmodium falciparum* 3D7.

The bark of *A. grandifolia*, which were collected at Alas Purwo, Indonesia, was extracted with MeOH, and then partitioned between hexane, EtOAc, *n*-BuOH, and water. The EtOAc extract was chromatographed over silica gel eluted with hexane-EtOAc and then CHCl₃-MeOH. The




CHCl₃-MeOH fraction was further separated by using an ODS column and C₁₈ HPLC to give melidianolic acids A (**1**, 22.8 mg) and B (**2**, 6.5 mg).

The ESIMS of melidianolic acid A (**1**) showed the pseudomolecular ion peak at m/z 339 ($M + H$)⁺ and the molecular formula, C₂₀H₃₄O₄, was established by HRESIMS (m/z 339.2524 ($M + H$)⁺, Δ -1.1 mmu). IR absorptions implied the presence of hydroxyl (3410 cm⁻¹) and carbonyl (1695 cm⁻¹) functionalities. ¹H and ¹³C NMR data were shown in Table 1. The ¹³C NMR and HSQC spectra revealed 20 carbon signals due to one carbonyl carbon, three sp² quaternary carbons, three sp² methines, one sp³ quaternary carbon, one sp³ methine, six sp³ methylenes, and five methyl groups.


The gross structure of **1** was elucidated by analysis of 2D NMR data including ¹H-¹H COSY, HSQC, and HMBC spectra in Figure 1. ¹H-¹H COSY spectrum of **1** revealed three partial structures **a** (C-4 to C-6), **b** (C-8 to C-10), and

Table 1: ^1H and ^{13}C NMR Data for Melidianolic Acids A (**1**) and B (**2**) in CDCl_3 at 300 K^a.

Position	1		2	
	δ_{H} (int.; mult.; J (Hz))	δ_{C}	δ_{H}	δ_{C}
1		170.3		177.1
2a	5.69 (1H, s)	116.0	2.14 (1H, m)	34.9
2b	2.31 (1H, m)			
3		162.5	2.01 (1H, m)	25.1
4a	2.63 (1H, m)	33.2	1.25 (1H, m)	29.6
4b	2.73 (1H, m)		1.97 (1H, m)	
5	2.19 (2H, m)	26.3	2.10 (2H, m)	25.8
6	5.12 (1H, t, 8.0)	123.6	5.09 (1H, t, 8.0)	124.5
7		135.2		135.2
8	2.01 (2H, m)	39.3	2.06 (2H, m)	39.3
9	2.09 (2H, m)	25.8	2.09 (2H, m)	25.8
10	5.16 (1H, t, 8.0)	124.8	5.16 (1H, t, 8.0)	124.7
11		134.6		135.5
12a	2.06 (1H, m)	36.6	2.03 (1H, m)	36.9
12b	2.25 (1H, m)		2.28 (1H, m)	
13a	1.41 (1H, m)	29.7	1.42 (1H, m)	29.3
13b	1.59 (1H, m)		1.61 (1H, m)	
14	3.36 (1H, brd, 12.0)	78.1	3.37 (1H, brd, 12.0)	78.4
15		73.4		73.3
16	1.20 (3H, s)	26.3	1.22 (3H, s)	26.3
17	1.16 (3H, s)	23.2	1.17 (3H, s)	23.1
18	1.59 (3H, brs)	15.9	1.59 (3H, brs)	15.8
19	1.59 (3H, brs)	15.9	1.60 (3H, brs)	15.8
20	1.91 (3H, s)	25.3	0.97 (3H, d, 6.4)	19.5

^a δ in ppm.**Figure 1:** Selected 2D NMR Correlations for Melidianolic Acid A (**1**).

c (C-12 to C-14). Connection of the partial structures **a** and **b** through C-7 was unambiguously established by HMBC correlations for H-6 and H₃-19 to C-7 (δ_{C} 135.2) and H₃-19 to C-8 (δ_{C} 39.3). The partial structures **b** and **c** were connected to C-11 on the basis of HMBC correlations for H-10 and H₃-18 to C-11 (δ_{C} 134.6) and C-12 (δ_{C} 36.6). The isopropanol function was connected to C-14 on the basis of HMBC cross-peaks for H₃-17 and H₃-16 to C-15 (δ_{C} 73.4) and H₃-17 to C-14 (δ_{C} 78.1). The connectivity between among C-2 and C-4 through C-3 were implied by HMBC correlations for H-4 and H₃-20 to C-2 (δ_{C} 116.0). The presence of a conjugated carboxylic acid was verified by a HMBC correlation for H-2 to C-1 (δ_{C} 170.3). Methylation of **1** with trimethylsilyl diazomethane to afford a methylated derivative supported the presence of a carboxylic acid function. The geometry of the double bonds was determined as *6E* and *10E* by comparison of the ^1H and ^{13}C chemical shifts with similar compounds [8-10]. Remaining geometry at C-2 was elucidated as *Z* by the NOESY correlation of H₃-20/H-2. Thus, the gross structure of melidianolic acid A was assigned to be **1**.

**Figure 2:** Selected 2D NMR Correlations for Melidianolic Acid B (**2**).

Melidianolic acid B {**2** [α]²⁰_D-12 (*c* 1.0, MeOH)} showed the pseudomolecular ion peak at m/z 341 ($\text{M} + \text{H}$)⁺ in the ESIMS, and the molecular formula, $\text{C}_{20}\text{H}_{36}\text{O}_4$, was established by HRESIMS [m/z 341.2511, ($\text{M} + \text{H}$)⁺, Δ -1.1 mmu]. The molecular formula of melidianolic acid B (**2**) was larger than that of **1** by H_2 unit. IR absorptions implied the presence of hydroxyl (3400 cm^{-1}) and carbonyl (1710 cm^{-1}) functionalities. ^1H and ^{13}C NMR chemical shifts of **2** were analogous to those of **1** with four methyl groups at C-16, C-17, C-18, and C-19 although signals of trisubstituted olefin carbons (δ_{H} 5.69; δ_{C} 116.0 and 162.5) at C-2 and C-3 of **1** were not observed for **2** (Table 1). The gross structure of **2** was elucidated by 2D NMR (^1H - ^1H COSY, HSQC, and HMBC) data (Figure 2). The presence of a carboxylic group was verified by a HMBC correlation for H₂-2 to C-1 (δ_{C} 177.1), which was also supported by methylation of **2** according to the same procedure as in **1**. The geometry of the double bonds was determined as *6E* and *10E* by comparison of the chemical shifts of the ^1H and ^{13}C NMR spectra of similar compounds [8-10]. Thus **2** was elucidated to be melidianolic acid B, which was 2,3-dihydro from of **1**.

Melidianolic acids A (**1**) and B (**2**) showed in vitro antiplasmodial activity against *P. falciparum* 3D7 (IC₅₀ 6.1 and 7.3 µg/mL, respectively), whereas both of them showed no cytotoxicity (IC₅₀ > 100 µg/mL). An acyclic diterpene, glaucaic acid, from *Knema glauca* (Myristicaceae), did not show antimalarial activity against *P. falciparum* [11]. Although recently acyclic triterpenes from *Ekebergia capensis* (Meliaceae) showed in vitro and in vivo antimalarial activities [12], it is the first report on the presence of antimalarial acyclic diterpenes from Meliaceae.

Experimental

General Experimental Procedures: Optical rotations were measured on a JASCO DIP-1000 automatic digital polarimeter. UV spectra were obtained on an Ultrospec 2100 pro spectrophotometer, and IR spectra were recorded on a JASCO FT/IR-4100 spectrophotometer. 1D and 2D NMR spectra were recorded on a Bruker AV 400 spectrometer, and chemical shifts were referenced to the residual chloroform-*d* (δ_{H} 7.26 and δ_{C} 77.0). Standard pulse sequences were employed for the 2D NMR experiments. ¹H-¹H COSY spectrum was measured with spectral widths of both dimensions of 4800 Hz, and 32 scans with two dummy scans were accumulated into 1K data points for each of 256 *t*₁ increments. For HSQC spectra in the phase sensitive mode and HMBC spectra, a total of 256 increments of 1K data points were collected. For HMBC spectra with Z-axis PFG, a 50 ms delay time was used for long-range C-H coupling. Zero-filling to 1K for *F*₁ and multiplication with squared cosine-bell windows shifted in both dimensions were performed prior to 2D Fourier transformation. High-resolution ESI MS were obtained on a LTQ Orbitrap XL (Thermo Scientific).

Plant Material: *Aphanamixis grandifolia* (Meliaceae) were collected at Alas Purwo, East Java, Indonesia in 2007. The botanical identification was made by Ms. Suri, Purbodadi Botanical Garden, Indonesia. A voucher specimen (no. AP070905) is deposited at the Herbarium of Purbodadi Botanical Garden, Indonesia.

Extraction and Isolation: The bark of *A. grandifolia* was extracted with MeOH (10 L x 3). MeOH extracted (126.3 g) was partitioned between hexane, EtOAc, *n*-BuOH, and water. The EtOAc soluble materials (16.7 g) were subjected to a silica gel column (hexane/EtOAc, 1:0 → 5:5, and then CHCl₃/MeOH 20:1 → 0:1). The fraction eluted with CHCl₃/MeOH (9:1) was applied to an ODS column (10% MeOH→MeOH), in which a fraction eluted with 80% MeOH was purified with an ODS HPLC (55% CH₃CN/0.1% TFA) to afford melidianolic acids A (**1**, 22.8 mg, 0.0016% yield) and B (**2**, 6.5 mg, 0.0004% yield).

Melidianolic Acid A (**1**)

colorless solid.

$[\alpha]_{\text{D}}^{20}$: -18 (*c* 1.0, MeOH).

IR (neat): 3410, 2970, 1695, 1170 cm⁻¹.

¹H and ¹³C NMR: Table 1.

ESIMS (pos.): *m/z* = 339 [M + H]⁺.

HRESITOFMS: *m/z* [M + H]⁺ calcd for C₂₀H₃₅O₄: 339.2535; found: 339.2524.

Melidianolic Acid B (**2**)

colorless solid.

$[\alpha]_{\text{D}}^{20}$: -12 (*c* 1.0, MeOH).

IR (neat): 3400, 2930, 2855, 1710 cm⁻¹.

¹H and ¹³C NMR: Table 1.

ESIMS (pos.): *m/z* = 363 [M + Na]⁺.

HRESITOFMS: *m/z* [M + Na]⁺ calcd for C₂₀H₃₆O₄Na: 363.2511; found: 363.2504.

Antiplasmodial activity: Human malaria parasites were cultured according to the method of Trager et al. [13]. The antimalarial activity of the isolated compounds was determined by the procedure described by Budimulja et al. [14]. In brief, stock solutions of the samples were prepared in DMSO (final DMSO concentrations of < 0.5%) and were diluted to the required concentration with complete medium (RPMI 1640 supplemented with 10% human plasma, 25 mM HEPES and 25 mM NaHCO₃) until the final concentrations of samples in culture plate wells were 10; 1; 0.1; 0.01; 0.001 µg/mL. The malarial parasite *P. falciparum* 3D7 clone was propagated in a 24-well culture plates. Growth of the parasite was monitored by making a blood smear fixed with MeOH and stained with Geimsa stain. The antimalarial activity of each compound was expressed as an IC₅₀ value, defined as the concentration of the compound causing 50% inhibition of parasite growth relative to an untreated control. The percentage of growth inhibition was expressed according to following equation: Growth inhibition % = 100 - [(test parasitaemia/control parasitemia) × 100]. Chloroquine: IC₅₀ 0.011 µM.

Cytotoxic Activity: A549 (human lung adenocarcinoma epithelial cell line) was seeded onto 96-well microtiter plates at 5 × 10³ cells per well. Cells were preincubated for 24 h at 37 °C in a humidified atmosphere of 5% CO₂. Different concentrations of each compound (10 µL) were added to the cultures, and then the cells were incubated at 37 °C for 48 h. On the third day, 15 µL MTT solution (5 mg/mL) was added into each well of the cultured medium. After further 2 hours of incubation, 100 µL of 10% SDS-0.01 N HCl solution was added to each well and the formazan crystals in each well were dissolved by stirring with a pipette. The optical density measurements were made using a micropipette reader (Benchmark Plus microplate spectrometer, BIO-RAD) equipped with a two wavelengths system (550 and 700 nm). In each experiment, three replicate of wells were prepared for each sample. The ratio of the living cells was determined based on the difference of the absorbance between those of samples and controls. These differences are expressed in percentage and cytotoxic activity was indicated as an IC₅₀ value. Taxol: IC₅₀ 0.0014 µM.

Acknowledgments - This work was partly supported by a Grant-in-Aid for Scientific Research from the Ministry of

Education, Culture, Sports, Science, and Technology of Japan, and a grant from the Open Research Center Project.


References

- [1] Amit R, Shailendra S. (2006) Limonoids: Overview of Significant Bioactive Triterpenes Distributed in Plants Kingdom. *Biological & Pharmaceutical Bulletin*, **29**, 191-201.
- [2] Nishizawa M, Nishide H, Hayashi Y, Kosela S. (1982) The structure of Lansioside A : A novel triterpene glycoside with amino-sugar from *Lansium domesticum*. *Tetrahedron Letters*, **23**, 1349-1350.
- [3] Nishizawa M, Inoue A, Sastrapradja S, Hayashi Y. (1983) (+)-8-Hydroxycalamenene: A fish-poison principle of *Dysoxylum acutangulum* and *D. Alliaceum*. *Phytochemistry*, **22**, 2083-2085.
- [4] Nishizawa M, Inoue A, Hayashi Y, Sastrapradja S, Kosela S, Iwashita T. (1984) Structure of aphanamols I and II. *The Journal of Organic Chemistry*, **49**, 3660-3662.
- [5] Polonsky J, Varon Z, Arnoux B, Pascard C, Pettir GR, Schmid JH, Lange LM. (1978) Antineoplastic agents. 57. Isolation and structure of aphanastatin. *Journal of the American Chemical Society*, **100**, 2575-2576.
- [6] Polonsky J, Varon Z, Arnoux B, Pascard C, Pettir GR, Schmid JM. (1978) Antineoplastic agents. 61. Isolation and structure of amoorastatin. *Journal of the American Chemical Society*, **100**, 7731-7734.
- [7] Polonsky J, Varon Z, Marazano C, Arnoux B, Pettir GR, Schmid JM, Ochi M, Kotsuki H. (1979) The structure of amoorastatone and the cytotoxic limonoid 12-hydroxyamoorastatin. *Experientia*, **35**, 987-989.
- [8] Perez AL, Ortega A, Vivar AR. (1988) An acyclic diterpene and sesquiterpene lactones from *Tithonia pedunculata*. *Phytochemistry*, **27**, 3897-3901.
- [9] Guerreiro E. (1986) Heliangolides and acyclic diterpene from *Viguiera gilliesii*. *Phytochemistry*, **25**, 748-750.
- [10] Valls R, Banaigs B, Francisco C, Codomier L, Cave A. (1986) An acyclic diterpene from the brown alga *Bifurcaria bifurcata*. *Phytochemistry*, **25**, 751-752.
- [11] Rangkaew N, Suttisri R, Moriyasu M, Kawanishi K. (2009) A new acyclic diterpene acid and bioactive compounds from *Knema glauca*. *Archives of Pharmacal Research*, **32**, 685-692.
- [12] Murata T, Miyase T, Muregi FW, Naoshima-Ishibashi Y, Umehara K, Warashina T, Kanou S, Mkoji GM, Terada M, Ishih A. (2008) Antiplasmodial Triterpenoids from *Ekebergia capensis*. *Journal of Natural Products*, **71**, 167-174.
- [13] Trager W, Jensen JB. (1976) Human malaria parasites in continuous culture. *Science*, **193**, 673-675.
- [14] Budimulja AS, Syafruddin TP, Wilariat P, Marzuki S. (1977) The sensitivity of *Plasmodium* protein synthesis to prokaryotic ribosomal inhibitors. *Molecular and Biochemical Parasitology*, **84**, 137-141.

Natural Product Communications

COUNTRY

United States

 Universities and research institutions in United States

PUBLICATION TYPE

Journals

SUBJECT AREA AND CATEGORY

Agricultural and Biological Sciences
└ Plant Science

Medicine
└ Complementary and Alternative Medicine
└ Medicine (miscellaneous)

Pharmacology, Toxicology and Pharmaceutics
└ Drug Discovery
└ Pharmacology

ISSN

15559475, 1934578X

PUBLISHER

SAGE Publications Inc.

COVERAGE

2006-2022

H-INDEX

50

INFORMATION

- Homepage
- How to publish in this journal
- editorial@naturalproduct.us

Quartiles



SJR

Year	SJR
2007	0.247
2008	0.280
2009	0.311
2010	0.367
2011	0.451
2012	0.449
2013	0.412
2014	0.371

Total Documents



Bukti – Scopus Coverage, Publisher dan ISSN

Source details

Feedback > Compare sources >

Natural Product Communications

Open Access ⓘ

Scopus coverage years: from 2006 to Present

Publisher: SAGE

ISSN: 1934-578X E-ISSN: 1555-9475

Subject area: Medicine: Complementary and Alternative Medicine Agricultural and Biological Sciences: Plant Science
Pharmacology, Toxicology and Pharmaceutics: Pharmacology Pharmacology, Toxicology and Pharmaceutics: Drug Discovery

Source type: Journal

View all documents >

Set document alert

Save to source list

CiteScore 2021
2.0 ⓘ

SJR 2021
0.270 ⓘ

SNIP 2021
0.529 ⓘ

CiteScore CiteScore rank & trend Scopus content coverage

Improved CiteScore methodology

CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021

$$2.0 = \frac{2,491 \text{ Citations } 2018 - 2021}{1,262 \text{ Documents } 2018 - 2021}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022 ⓘ

$$2.1 = \frac{1,991 \text{ Citations to date}}{970 \text{ Documents to date}}$$

Last updated on 06 June, 2022 • Updated monthly

CiteScore rank 2021 ⓘ

Category	Rank	Percentile
Medicine └ Complementary and Alternative Medicine	#44/93	53rd
Agricultural and Biological Sciences └ Plant Science	#240/482	50th
Pharmacology, Toxicology and Pharmaceutics └ Pharmacology	#212/303	30th
Pharmacology, Toxicology and Pharmaceutics └ Drug Discovery	#114/154	26th