



Buletin Peternakan

(Bulletin of Animal Science)

OPEN ACCESS



pISSN : 0126-4400 | eISSN : 2407-876X

- Home
- About
- Login
- Register
- Search
- Current
- Archives
- Announcements
- Statistics
- Indexing & Abstracting
- Journal History
- Contact

Home > Vol 44, No 2 (2020)

Buletin Peternakan

Buletin Peternakan (*Bulletin of Animal Science*) with registered number ISSN 0126-4400 (print) ISSN 2407-876X (online) is a scientific open access journal published by **Faculty of Animal Science Universitas Gadjah Mada** jointly with **ISSTAP (Indonesian Society for Sustainable Tropical Animal Production)**. Buletin Peternakan has been accredited as a scientific journal by the Ministry of Research-Technology and Higher Education Republic of Indonesia: 36a/E/KPT/2016 PERIOD : 2016-2021. Buletin Peternakan (*Bulletin of Animal Science*) aims to publish original research results and reviews on farm tropical animals such as cattle, buffaloes, sheep, goats, pigs, horses, poultry, as well as non domesticated Indonesian endemic animals, such as deers, anoa, babirusa, etc. Buletin Peternakan encompasses a broad range of research topics in animal sciences: breeding and genetics, reproduction and physiology, nutrition, feed sciences, agrastology, animal products



- Focus & Scope
- Author Guidelines
- Author Fees
- How to Online Submission
- Publication Ethics
- Open Access Policy
- Screening For Plagiarism
- Editorial Board
- Copyright Notice
- Peer Reviewers

Buletin Peternakan have been indexed on:

1. DOAJ (Directory of Open Access Journals)
2. Google Scholar
3. CrossRef
4. SINTA (Sistem Indeksi dan Sitasi)
5. BASE (Bielefeld Academic Search Engine)
6. ROAD
7. Scilit
8. WorldCat
9. OpenAire
10. Sherpa/Romeo
11. Boston University
12. Livivo
13. Harvard Library
14. Stanford University Libraries
15. Berkeley Library Catalog
16. University of California Libraries
17. Western
18. Cabi.org
19. Indonesian One Search
20. Garuda (Garba Rujukan Digital)
21. Renmin University of China
22. University of Saskatchewan
23. DRJI
24. DAIJ
25. Realtime Impact Factor
26. JournalTOCS
27. Repositori Publikasi Badan Litbang Pertanian
28. DISNAKESWAN - Sulsel
29. Universal Impact Factor
30. USDA
31. Global Impact Factor
32. Indonesian Scientific Journal Database (ISJD)
33. FAO
34. University of Saskatchewan Library



CITATION ANALYSIS

- ▶ SCOPUS
- ▶ Google Scholar

TEMPLATE



REFERENCE MANAGEMENT TOOLS



USER

Username

Buletin Peternakan

Tidak aman | jurnal.ugm.ac.id/buletinpeternakan

Vol 44, No 2 (2020): BULETIN PETERNAKAN VOL. 44 (2) MAY 2020

Table of Contents

Articles

Evaluation of Different Techniques in Recovering of Oocytes and Storage Duration of Ovaries on the Quality and Quantity of Bovine In Vitro Maturation
 Zaituni Udin, Masrizal Masrizal, Hendri Hendri, Syafril Nanda
 10.21059/buletinpeternak.v44i2.52181 Abstract views : 312 | views : 186

Reproductive Performances of Friesian Holstein Dairy Cows in Different Agricultural Ecosystems
 Tendy Kusmayadi, Muhamad Hasan Hadiana, Ujang Hidayat Tanuwiria
 10.21059/buletinpeternak.v44i2.39917 Abstract views : 184 | views : 118

Identification of Glucogenic Amino Acids Content in *Gliricidia maculata* as an Alternative Energy Source for High-Yielding Periparturient Dairy Cows
 Sulvia Dwi Astuti SW, Budi Prasetyo Widyobroto, Ali Agus, Lies Mira Yusiati
 10.21059/buletinpeternak.v44i2.54103 Abstract views : 190 | views : 112

The Effect of Choline Chloride Supplementation on the Reproductive Performance of Simmental Bulls Fed Protected Protein in the Ration
 Bambang Waluyo Hadi Eko Prasetyono, Yon Soepri Ondho, Agung Subrata, Pramesti Kusuma Pratiwi, Mutiara Briliiana Zahra, Trisna Itmamulwafa, Tertyani Kartika Pratiwi, Muslichatun Nisa, Widiyanto Widiyanto
 10.21059/buletinpeternak.v44i2.55338 Abstract views : 202 | views : 145

Application of *Saccharomyces cerevisiae* as a Probiotic for Producing Low Cholesterol and Antibiotic-Free Broiler Meat
 Suci Wulandari, Theo Mahiseta Syahniar, Dadik Pantaya

Search
 Search Scope
 All
 Search

Browse
 By Issue
 By Author
 By Title
 Other Journals

Type here to search

Buletin Peternakan

Tidak aman | jurnal.ugm.ac.id/buletinpeternakan

Probiotic-Enhanced Liquid Acidifier
 Danung Nur Adli, Oskar Sjojjan
 10.21059/buletinpeternak.v44i2.54713 Abstract views : 284 | views : 199

Refeeding Postmolting Method to Improve Weekly Production Performance of Rejected Laying Hens with Low Mortality
 Claude Mona Airin, Amelia Hana, Pudji Astuti, Sarmin Sarmin, Trini Susmiati, Okti Herawati
 10.21059/buletinpeternak.v44i2.45508 Abstract views : 242 | views : 188

Resistance Profile of Extended Spectrum Beta Lactamase-Producing *Escherichia coli* Bacteria using Vitek® 2-Compact Method
 Freshinta Jellia Wibisono, Bambang Sumiarto, Tri Untari, Mustofa Helmi Effendi, Dian Ayu Permatasari, Adiana Mutamsari Witaningrum
 10.21059/buletinpeternak.v44i2.51347 Abstract views : 213 | views : 169

Physicochemical Characteristics and Microbiological Quality of the Topside and Longissimus Dorsi of Indonesian Local Buffalo Meat
 Noraimah Binti Sulaiman, Cece Sumantri, Irma Isnafia Arief, Cahyo Budiman
 10.21059/buletinpeternak.v44i2.45544 Abstract views : 189 | views : 93

Buletin Peternakan (Bulletin of Animal Science) Indexed by:

DOAJ
 Google Scholar
 Crossref
 SINTA
 BASE
 ROAD
 WorldCat
 SHERPA/RMEO
 BOSTON UNIVERSITY
 UNWU
 HARVARD LIBRARY
 WESTERN
 ONEsearch
 IPI
 UNIVERSITY OF SASKATCHEWAN

Type here to search

Editorial Team

Buletin Peternakan (*Bulletin of Animal Science*)

Editor in chief:

- **Muhlisin**, Department of Animal Nutrition and Feed Science, Faculty of Animal Sciences, Universitas Gadjah Mada, Indonesia. Scopus ID = 54391557700; h-index = 5.

Managing Editor

- **Bambang Ariyadi**, Department of Animal Production, Faculty of Animal Sciences, Universitas Gadjah Mada, Indonesia. Scopus ID = 55184028700; h-index = 5.
- **Tri Anggraeni Kusumastuti**, Faculty of Animal Sciences, Universitas Gadjah Mada, Indonesia. Scopus ID = 57194135275; h-index = 1.
- **Tri Satya Mastuti Widi**, Faculty of Animal Sciences, Universitas Gadjah Mada, Indonesia. Scopus ID = 57194418225; h-index = 2.
- **Tety Hartatik**, Faculty of Animal Sciences, Universitas Gadjah Mada, Indonesia. Scopus ID = 6505863692; h-index = 4.

International Editor:

- **Yuny Erwanto**, Department of Animal Products Technology, Faculty of Animal Sciences, Universitas Gadjah Mada, Indonesia. Scopus ID = 6507184764; h-index = 7.
- **Nafiatul Umami**, Department of Animal Nutrition and Feed Science, Faculty of Animal Sciences, Universitas Gadjah Mada, Indonesia. Scopus ID = 55515013800; h-index = 1.
- **Nanung Agus Fitriyanto**, Department of Animal Products Technology, Faculty of Animal Sciences, Universitas Gadjah Mada, Indonesia. Scopus ID = 36544733200; h-index = 2.
- **Nanung Danar Dono**, Department of Animal Nutrition and Feed Science, Faculty of Animal Sciences, Universitas Gadjah Mada, Indonesia. Scopus ID = 56131603300; h-index = 2.
- **Eka Meutia Sari**, Breeding dan Pemuliaan Ternak, Universitas Syiah Kuala, Indonesia. Scopus ID = 57190665652; h-index = 2.
- **Dyah Maharani**, Department of Animal Production, Faculty of Animal Sciences, Universitas Gadjah Mada, Indonesia. Scopus ID = 54881785100; h-index = 3.
- **Jamhari**, Department of Animal Products Technology, Teknologi Hasil Ternak, Universitas Gadjah Mada, Indonesia. Scopus ID = 35082056800; h-index = 2.
- **Sri Mukodiningih**, Nutrisi dan Makanan Ternak, Universitas Diponegoro, Indonesia. Scopus ID = 57189042929; h-index = 1.
- **Sutrisno Hadi Purnomo**, Sosial Ekonomi Peternakan, Universitas Negeri Sebelas Maret. Scopus ID = 55683342400; h-index = 3.
- **Trisiwi Wahyu Widayati**, Faculty of Animal Sciences, Universitas Negeri Papua, Indonesia. Scopus ID = 57191056182; h-index = 1.

JOURNAL CONTENT

Search

Search Scope

All

Search

Browse

- ▶ By Issue
- ▶ By Author
- ▶ By Title
- ▶ Other Journals

- **Yuny Erwanto**, Department of Animal Products Technology, Faculty of Animal Sciences, Universitas Gadjah Mada, Indonesia. Scopus ID = 6507184764; h-index = 7.
- **Nafiatul Umami**, Department of Animal Nutrition and Feed Science, Faculty of Animal Sciences, Universitas Gadjah Mada, Indonesia. Scopus ID = 55515013800; h-index = 1.
- **Nanung Agus Fitriyanto**, Department of Animal Products Technology, Faculty of Animal Sciences, Universitas Gadjah Mada, Indonesia. Scopus ID = 36544733200; h-index = 2.
- **Nanung Danar Dono**, Department of Animal Nutrition and Feed Science, Faculty of Animal Sciences, Universitas Gadjah Mada, Indonesia. Scopus ID = 56131603300; h-index = 2.
- **Eka Meutia Sari**, Breeding dan Pemuliaan Ternak, Universitas Syiah Kuala, Indonesia. Scopus ID = 57190665652; h-index = 2.
- **Dyah Maharani**, Department of Animal Production, Faculty of Animal Sciences, Universitas Gadjah Mada, Indonesia. Scopus ID = 54881785100; h-index = 3.
- **Jamhari**, Department of Animal Products Technology, Teknologi Hasil Ternak, Universitas Gadjah Mada, Indonesia. Scopus ID = 35082056800; h-index = 2.
- **Sri Mukodiningih**, Nutrisi dan Makanan Ternak, Universitas Diponegoro, Indonesia. Scopus ID = 57189042929; h-index = 1.
- **Sutrisno Hadi Purnomo**, Sosial Ekonomi Peternakan, Universitas Negeri Sebelas Maret. Scopus ID = 55683342400; h-index = 3.
- **Trisiwi Wahyu Widayati**, Faculty of Animal Sciences, Universitas Negeri Papua, Indonesia. Scopus ID = 57191056182; h-index = 1.
- **Abdul Razak Alimon**, Universiti Putra Malaysia, Malaysia. Scopus ID = 6603104909; h-index = 10.
- **Bounmy Keohavong**, Souphanouvong University, Laos. Scopus ID = 57205074640; h-index = 1.
- **Cheorun Jo**, Seoul National University, Korea. Scopus ID = 7006072985; h-index = 46.
- **Egil Robert Orskov**, International Feed Resources Unit, Aberdeen, UK. Scopus ID = 7005860217; h-index = 48.
- **Henning Otte Hansen**, University of Copenhagen, Denmark. Scopus ID = 56295438900; h-index = 2.
- **Liang Chou Hsia**, Pingtung University of Science and Technology, Taiwan. Scopus ID = 36722621700; h-index = 4.
- **Mohammad Taghi Beigi Nassiri**, Khuzwstan Ramin Agricultural and Natural Resources University, Iran. Scopus ID = 56994110200; h-index = 3.
- **Moriyama Hiromitsu**, Tokyo University of Agriculture and Technology, Japan.
- **Sang Jip Ohh**, Kangwon National University, Korea. Scopus ID = 6603464243; h-index = 10.
- **Peter Charles Wynn**, Charles Stuart University, Australia. Scopus ID = 7006216390; h-index = 25.
- **Tomoyaki Nakagawa**, Gifu University, Japan.
- **Wu Chean Ping**, National Chiayi University, Taiwan.
- **Zulkifli Idrus**, Universitas Putra Malaysia. Scopus ID = 56040494900; h-index = 24.

Editorial staffs:

- Galuh Adi Insani
- Sri Agtin Tejawati
- Adhitya Permadi

Doi: 10.21059/buletinpeternak.v44i2.51347

Resistance Profile of Extended Spectrum Beta Lactamase-Producing *Escherichia coli* Bacteria using Vitek® 2 Compact Method

Freshinta Jellia Wibisono¹, Bambang Sumiarto², Tri Untari³, Mustofa Helmi Effendi^{4*}, Dian Ayu Permatasari⁴, and Adiana Mutamsari Witaningrum⁴

¹Doctoral Program in Veterinary Science, Faculty of Veterinary Medicine, Universitas Gadjah Mada, Yogyakarta, 55281, Indonesia

²Department of Veterinary Public Health, Faculty of Veterinary Medicine, Universitas Gadjah Mada, Yogyakarta, 55281, Indonesia

³Department of Microbiology, Faculty of Veterinary Medicine, Universitas Gadjah Mada, Yogyakarta, 55281, Indonesia

⁴Department of Veterinary Public Health, Faculty of Veterinary Medicine, Airlangga University, Surabaya, 60115, Indonesia

ABSTRACT

This study aimed to determine the resistance profile and the nature of multidrug resistance in *Extended Spectrum Beta Lactamase* (ESBL)-producing *Escherichia coli* (*E.coli*) against several classes of antibiotics. Positive isolates of ESBL-producing *E.coli* were tested for antibiotic sensitivity using the VITEK® 2 compact method which then analyzed automatically. The results showed an antibiotic resistance profile against ESBL-producing *E.coli* showed the highest level of antibiotics in beta lactam, amoxicillin, ampicillin, cefazolin, cefotaxime, and ceftriaxone at 100%. Subsequent results found a relatively high level of resistance in the antibiotics aztreonam (86.36%), trimethoprim/sulfamethoxazole (77.27%), gentamicin (72.73%), and ciprofloxacin (68.18%). Antibiotics from carbapenem groups such as ertapenem and meropenem, and antibiotics from the aminoglycosides (amikacin) and tetracycline groups of tetracycline still showed a high sensitivity level of 100%. The most common resistance patterns found in ESBL-producing *E.coli* isolates are AM/AMP/KZ/CTX/CRO/ATM/GM/CIP as much as 22.73%, and AM/AMP/KZ/CTX/CRO/ATM/GM/CIP/SXT patterns of 18.2%. The results of multi-class antibiotic resistance showed that 86.36% had multidrug resistance. The highest multidrug resistance pattern in ESBL-producing *E.coli* occurred with a BL/AG/Q/SP pattern of 50%. Other patterns of multidrug resistance in ESBL-producing *E.coli* that can be found in this study are, the BL/AG/Q/SP pattern is 18.20%, the BL/AG/Q/SP pattern is 13.64%, and the BL/AG/Q pattern is 4.55%. The high profile of resistance and the nature of multidrug resistance in ESBL-producing *E.coli* has the potential to spread these resistant genes, thus risking the use of antibiotics as a public health therapy and animal health, therefore further evaluation and control are needed.

Keywords: ESBL-producing *Escherichia coli*, Multidrug resistance, Vitek® 2 Compact Method

Article history

Submitted: 10 November 2019

Accepted: 3 May 2020

* Corresponding author:

Telp. +628175111783

E-mail: mheffendi@yahoo.com

Introduction

Disease in commercial chicken is one of constraints causing the decline in economic value (Wiedosari and Wahyuwardani, 2015). Pathogenic microorganisms cause infectious diseases, which is one of a leading cause of death in animals or even human. The high prevalence of infectious disease is associated with the high use of antibiotic in health care (Noor and Peoleonga, 2005). A negative economic consequence in commercial chicken production can be caused by *Escherichia coli* (*E.coli*) infection. The existence of Extended Spectrum Beta Lactamase (ESBL)-producing *E.coli* on commercial chicken is highly

associated with *E.coli*, a normal microflora on gastrointestinal tract of commercial chicken. Feces and animal housing environment can facilitate the existence of *E.coli*, although, chicken show no symptoms at all of any diseases caused by the bacteria (Wibisono *et al.*, 2018).

Commercial chicken farm is one of sources of antibiotic resistance. The uncontrolled use of antibiotic causes the emergence of antibiotic resistance. Beta lactam antibiotic is the type of antibiotic commonly used in commercial chicken production. Resistance to beta lactam antibiotic can occur in ESBL-producing *E.coli* (Santos *et al.*, 2013; Hammerum *et al.*, 2014). Extended Spectrum Beta Lactamase is an enzyme that

causes resistance to wider spectrum on third generation of cephalosporins and mobobactams; thus, the antibiotic becomes ineffective. ESBL-producing bacteria can also be resistant to wide array of antibiotic class: aminoglycoside, tetracycline chloramphenicol, and sulfamethoxazole-trimethoprim (Brower *et al.*, 2017; Sudarwanto *et al.*, 2017). Incidence on multidrug resistance to third generation cephalosporins and other antibiotic classes are often found in ESBL-producing bacteria (Masrurh *et al.*, 2016). Multidrug resistance is a resistance to three or more different antibiotic classes (Handayani *et al.*, 2017).

Identification of ESBL bacteria is a problem in confirming a diagnosis at the laboratory, either in therapeutic approach or in the field to prevent its spread. Clinical and Laboratory Standards Institute (CLSI) states that ESBL examination must be performed routinely. According to CLSI recommendation, ESBL detection consists of two steps. The first is an initial screen test, a filter test to reduce the susceptibility to more than one cephalosporin indicators (cefotaxime, ceftriaxone, ceftazidime, cefpodoxime) and aztreonam. The susceptibility reduction from cephalosporins shows positive results. A positive result from screening test is followed by the second step, ESBL confirmation test. The second step aims to detect the hydrolytic potency of ESBL against antibiotic used in the screening test. The ESBL confirmation test deliver a picture of collaboration action between ceftazidime or cefotaxime, and clavulanate acid (Amelia *et al.*, 2016; CLSI, 2017; Biutifasari, 2018).

Methods of identification, antibiotic sensitivity test, and ESBL-producing bacteria confirmation test are categorized into phenotype and genotype examination methods, with their advantages and limitations. Both phenotypic bacteria identification and antibiotic sensitivity is relatively easier than genotypic examination. Phenotypic examination is performed by using agar testing method or fully automated automatic method as in VITEK[®] 2 compact automated system. The VITEK[®] 2 compact automated system from Biomerieux works on calorimetric principle for identification, through biochemical and antibiotic sensitivity tests. The accuracy of VITEK[®] 2 compact automated system ranges from 97.8% (O'Hara, 2005) to 98.02% (Duggal *et al.*, 2012). This study aimed to determine resistance profile and the nature of multidrug resistance in ESBL-producing *E.coli* against several classes of antibiotic. Antibiotic resistance is a problem occurring across the globe, including Indonesia, both in human and animal health.

Materials and Methods

This study used commercial chicken cloaca swab sample, collected from commercial chicken farms in Blitar Regency. 22 positive ESBL-producing *E.coli* isolate were identified using IMBIC test and confirmed as ESBL-

producing *E.coli* using Double Disc Synergy (DDST) test (CLSI, 2017; Effendi *et al.*, 2018). Identification of ESBL-producing *E.coli* isolate was performed using VITEK[®] 2 GN card, while VITEK[®] 2 AST cards were used to determine the resistance profile. 3 ml of sterile saline solution (0.45 – 0.50%; pH 4.5 – 7.0) was put into plastic tube aseptically to make a bacterial suspension. *E. coli* isolate was isolated on MacConkey Agar (MCA) medium, incubated at 35-37°C for 20-24 hours. *E. coli* isolate was put into sterile saline solution using sterile swab to create bacterial suspension. Bacterial suspensions were homogenized and 0.50 – 0.63 McFarland bacterial turbidity were made using VITEK[®] 2 DensiCHEK. These bacterial suspensions were inoculated to VITEK[®] 2 card no more than after 30 minutes. Bacterial suspension tubes and VITEK[®] 2 cards (GN and AST) were placed into special rack or cassette. Racks containing bacterial suspensions and cards were placed into vacuum chamber station for 30-60 seconds. Bacterial suspension was then transferred into wells. Transfer tubes were cut automatically. Cards were transferred to incubator room after 15 minutes, for approximately 8 hours at 35°C. Cards were then analyzed automatically. Antibiotic sensitivity test on ESBL-producing *E.coli* was carried out using VITEK[®] 2 compact, that consists of 18 antibiotics: amoxicillin, ampicillin, ampicillin/sulbactam, piperacillin/tazobactam, cefazolin, cefotaxime, ceftazidime, ceftriaxone, cefepime, aztreonam, ertapenem, meropenem, amikacin, gentamicin, ciprofloxacin, tigecycline, nitrofurantoin, dan trimethoprim/sulfamethoxazole. The results were automatically analyzed by system and interpreted as sensitive, intermediate, or resistant (Sugiartha, 2016; Biomerieux, 2017). ESBL-producing *E.coli* multidrug resistance in this study used six different antibiotic classes: beta lactam, aminoglycoside, quinolone, tetracycline, nitrofurantoin, and sulfonamide-potential.

Results and Discussion

The resistance test results of ESBL-producing *E.coli* from commercial chicken cloaca swabs against several classes of antibiotic (Figure 1) shows high level of resistance. The highest percentage of resistance was observed in amoxicillin, ampicillin, cefazolin, cefotaxime, and ceftriaxone, by 100%. This result is according to resistance study on ESBL-producing *E.coli* at RPHR of Bogor City that showed 100% level of resistance on penicillin and amoxicillin (Normaliska *et al.*, 2019). Amoxicillin, ampicillin, cefazolin, cefotaxime, and ceftriaxone are categorized as beta lactam class antibiotic, amoxicillin and ampicillin belong to penicillin, cefazolin belongs to first generation cephalosporins, cefotaxime and ceftriaxone is third generation cephalosporins class. *E.coli* which is often referred as ESBL-producing bacteria has ability to produce beta lactamase enzyme, an enzyme that able to inhibit beta lactam (Paterson and Bonomo, 2005). Beta lactam resistance

commonly occurs in gram negative bacteria, including *E.coli* that can be isolated from animal products. Poultry is known as ESBL-producing *E.coli* reservoir (Schmid *et al.*, 2013; Hammerum *et al.*, 2014). The high resistance level of ESBL-producing *E.coli* from commercial chicken cloaca swab on beta lactam class antibiotic is caused by the presence of ESBL enzyme in *E.coli*, which is not limited on digestive tract, but also in the cage, feces, and surrounding environment (Schroeder *et al.*, 2004). The enzyme is not only able to hydrolyze penicillin, but also third generation cephalosporins and monobactam (Paterson and Bonomo, 2005; Nuangmek *et al.*, 2018).

A relatively high level resistance of ESBL-producing bacteria was observed on aztreonam, trime thoprim/sulfamethoxazole, gentamicin, and ciprofloxacin, by 86.36%, 77.27%, 72.73%, and 68.18% consecutively, as the previous study with relatively high level of resistance of ciprofloxacin (50.0%), gentamicin (60.0%), trimethoprim-sulfamethoxazole (60%, and ciprofloxacin (40.0%) (Cormican *et al.*, 1996; *et al.*Kürekcı *et al.*, 2017; Sudarwanto *et al.*, 2017). Other antibiotics such as ertapenem and mempenem from carbapenem class, amikacin (aminoglycoside class), and tigecycline (tetracycline class) still show high level of sensitivity (100%), according to mechanism of action of ESBL which hydrolyze penicillin class antibiotics, first, second, and third generations of cephalosporin, and monobactam antibiotic class, but inactive on carbapenem class (imipenem, meropenem, ertapenem) (Lim *et al.*, 2013; Biutifasari, 2018). Nitrofurantoin has lower resistance level than trimethoprim-sulphamethoxazole and fluoroquinolone (Rank *et al.*, 2018).

The susceptibility result shows a nearly similar pattern for all isolates (Table 1). The most resistance pattern seen on ESBL-producing *E.coli* was AM/AMP/KZ/CTX/CRO/ATM/GM/CIP pattern (22.73% or 5/22), and AM/AMP/KZ/CTX/CRO/ATM/GM/CIP/SXT (18.2% or 4/22). The result of t multiclass antibiotic resistance shows 86.36% (19/22) had multidrug resistance (Table 2). Multidrug resistance is certain bacterial resistance to three or more different antibacterial class (Kurniawati *et al.*, 2015). Multidrug resistance is a problem that hard to resolve in disease treatment. This condition is caused by the practice of the antibiotic use as disease preventive measures in commercial chicken production. Antibiotic has been used to control the level of morbidity, mortality, and infection of *E.coli*. While antibiotics are mostly used as therapeutic, several antibiotic classes are used for sub therapeutic purpose to prevent the emergence of disease in animal production (Niasono *et al.*, 2019). The use of antibiotic at sub therapeutic level can lead to the emergence of multidrug resistance (Wang *et al.*, 2015).

The highest level of multidrug resistance pater of ESBL-producing *E.coli* was observed on BL/AG/Q/SP pattern (50% or 11/22). Other multidrug resistance patterns found in this study were BL/AG/Q/SP (18.20% or 4/22), BL/AG/Q/SP (13.64% or 3/22), and BL/AG/Q (4.55% or 1/22). This study demonstrates that the high incidence of multidrug resistance can indicate the high risk of transmission on other chicken at the farm, to cause a transmission of antibiotic resistance on multi class antibiotics. The condition can result in the high potential failure of antibiotic treatment




Figure 1. Resistance profiles of ESBL-producing *E.coli* using VITEK® 2 method (Amoxicillin = AM; Ampicillin = AMP; Ampicillin/sulbactam = SAM; Piperacillin/Tazobactam = TZP; Cefazolin = KZ; Cefotaxime = CTX; Cefotaxime = CAZ; Ceftriaxone = CRO; Cefepime = FEP; Aztreonam = ATM; Ertapenem = ERT; meropenem = MEM; amikacin = AK; gentamicin = GM; ciprofloxacin = CIP; tigecycline = TC; nitrofurantoin = NFT; trimethoprim/sulfamethoxazole = SXT; R = Resistance; I = Intermediate; S = Susceptible)

Table 1. Isolates belonging to individual antibiotic resistance profiles

Code	Antibiotic																Anti-microbial Resistance Profiles		
	A M	A M P	S A M	T Z P	K Z	C T X	C A Z	C R O	F E P	A T M	E R T	M E M	A K	G M	C I P	T C		N F T	S X T
Ec1	R	R	S	S	R	R	S	R	S	R	S	S	R	R	S	S	S	AM/AMP/KZ/CTX/CRO/ATM/GM/CIP	
Ec2	R	R	S	S	R	R	S	R	S	R	S	S	R	R	S	S	R	AM/AMP/KZ/CTX/CRO/ATM/GM/CIP/SXT	
Ec3	R	R	S	S	R	R	S	R	S	R	S	S	R	R	S	S	R	AM/AMP/KZ/CTX/CRO/ATM/GM/CIP/SXT	
Ec4	R	R	I	S	R	R	S	R	S	R	S	S	R	R	S	I	S	AM/AMP/KZ/CTX/CRO/ATM/GM/CIP	
Ec5	R	R	S	S	R	R	S	R	S	R	S	S	R	R	S	S	R	AM/AMP/KZ/CTX/CRO/ATM/GM/CIP/SXT	
Ec6	R	R	I	S	R	R	S	R	S	S	S	S	R	R	S	S	S	AM/AMP/KZ/CTX/CRO/ATM/GM/CIP	
Ec7	R	R	R	S	R	R	R	R	S	R	S	S	S	I	S	S	R	AM/AMP/SAM/KZ/CTX/CAZ/CRO/ATM/SXT	
Ec8	R	R	S	S	R	R	S	R	S	R	S	S	R	S	S	S	R	AM/AMP/KZ/CTX/CRO/ATM/GM/SXT	
Ec9	R	R	I	S	R	R	S	R	S	R	S	S	R	S	S	S	R	AM/AMP/KZ/CTX/CRO/ATM/GM/SXT	
Ec10	R	R	S	S	R	R	S	R	S	S	S	S	R	S	S	I	R	AM/AMP/KZ/CTX/CRO/GM/SXT	
Ec11	R	R	I	S	R	R	S	R	S	S	S	S	R	R	S	S	R	AM/AMP/KZ/CTX/CRO/GM/CIP/SXT	
Ec12	R	R	I	S	R	R	S	R	S	R	S	S	R	R	S	I	S	AM/AMP/KZ/CTX/CRO/ATM/GM/CIP	
Ec13	R	R	S	S	R	R	S	R	S	R	S	S	R	R	S	S	R	AM/AMP/KZ/CTX/CRO/ATM/GM/CIP/SXT	
Ec14	R	R	S	S	R	R	S	R	S	R	S	S	R	R	S	S	S	AM/AMP/KZ/CTX/CRO/ATM/GM/CIP	
Ec15	R	R	S	S	R	R	S	R	S	R	S	S	S	I	S	S	R	AM/AMP/KZ/CTX/CRO/ATM/SXT	
Ec16	R	R	I	S	R	R	R	R	R	R	S	S	R	R	S	S	R	AM/AMP/KZ/CTX/CAZ/CRO/FEP/ATM/GM/CIP/SXT	
Ec17	R	R	R	I	R	R	R	R	R	R	S	S	S	S	R	S	I	AM/AMP/SAM/KZ/CTX/CAZ/CRO/FEP/ATM/CIP/SXT	
Ec18	R	R	R	S	R	R	R	R	R	R	S	S	S	R	R	S	S	AM/AMP/SAM/KZ/CTX/CAZ/CRO/FEP/ATM/GM/CIP/SXT	
Ec19	R	R	R	S	R	R	R	R	R	R	S	S	S	S	R	S	I	AM/AMP/SAM/KZ/CTX/CAZ/CRO/FEP/ATM/CIP/SXT	
Ec20	R	R	R	I	R	R	R	R	R	R	S	S	S	I	S	S	I	AM/AMP/SAM/KZ/CTX/CAZ/CRO/FEP/ATM/SXT	
Ec21	R	R	R	S	R	R	S	R	S	R	S	S	R	S	S	S	R	AM/AMP/SAM/KZ/CTX/CRO/ATM/GM/SXT	
Ec22	R	R	I	S	R	R	R	R	R	R	S	S	S	S	R	S	S	AM/AMP/KZ/CTX/CAZ/CRO/FEP/ATM/CIP/SXT	
Total	R	22	22	6	0	22	22	7	22	6	19	0	0	0	1	1	0	0	1
I	0	0	7	2	0	0	0	0	0	0	0	0	0	0	1	2	0	6	0
S	0	0	9	20	0	0	15	0	16	3	2	2	2	5	5	2	1	5	

(Amoxicillin = AM; Ampicillin = AMP; Ampicillin/sulbactam = SAM; Piperacillin/Tazobactam = TZP; Cefazolin = KZ; Cefotaxime = CTX; Ceftriaxone = CAZ; Cefepime = FEP; Aztreonam = ATM; Ertapenem = ERT; meropenem = MEM; amikacin = AK; gentamicin = GM; ciprofloxacin = CIP; tigecycline = TC; nitrofurantoin = NFT; trimethoprim/sulfamethoxazole = SXT; R = Resistance; I = Intermediate; S = Susceptible)

Table 2. Multidrug resistance pattern of ESBL-producing *E. coli* isolate

Sample code	Beta lactam	Aminoglycoside	Quinolones	Tetracycline	Nitrofurantoin	Sulfonamide potential	MDR	Multidrug Resistance Pattern
Ec1	R	R	R	S	S	S	MDR	BL/AG/Q
Ec2	R	R	R	S	S	R	MDR	BL/AG/Q/SP
Ec3	R	R	R	S	S	R	MDR	BL/AG/Q/SP
Ec4	R	R	R	S	I	S	MDR	BL/AG/Q
Ec5	R	R	R	S	S	R	MDR	BL/AG/Q/SP
Ec6	R	R	R	S	S	S	MDR	BL/AG/Q/NF/SP
Ec7	R	S	I	S	S	R	NEGATIF	BL/SP
Ec8	R	R	S	S	S	R	MDR	BL/AG/SP
Ec9	R	R	S	S	S	R	MDR	BL/AG/SP
Ec10	R	R	S	S	I	R	MDR	BL/AG/SP
Ec11	R	R	R	S	S	R	MDR	BL/AG/Q/SP
Ec12	R	R	R	S	I	S	MDR	BL/AG/Q
Ec13	R	R	R	S	S	R	MDR	BL/AG/Q/SP
Ec14	R	R	R	S	S	S	MDR	BL/AG/Q
Ec15	R	S	I	S	S	R	NEGATIF	BL/SP
Ec16	R	R	R	S	S	R	MDR	BL/AG/Q/SP
Ec17	R	S	R	S	I	R	MDR	BL/AG/Q/SP
Ec18	R	R	R	S	S	R	MDR	BL/AG/Q/SP
Ec19	R	S	R	S	I	R	MDR	BL/AG/Q/SP
Ec20	R	I	S	S	I	R	NEGATIF	BL/SP
Ec21	R	R	S	S	S	R	MDR	BL/AG/SP
Ec22	R	S	R	S	S	R	MDR	BL/AG/Q/SP

(Beta lactam = BL; Aminoglycoside = AG; Quinolones = Q; Tetracycline = TC; Nitrofurantoin = NF; Sulfonamide potential = SP; R = Resistance; I = Intermediate; S = Susceptible).

during the existence of infectious disease at the farm. The different resistance pattern arises due to the varied antibiotic combinations used by farmers, as stated by Bywater *et al.* (2014) in which various type of antibiotics, geographic conditions, and production system are factors causing the different resistance pattern.

Conclusion

The highest antibiotic resistance of ESBL-producing *E.coli* was observed on amoxicillin, ampicillin, cefazolin, cefotaxime, and ceftriaxone (100%). Ertapenem and mempenem from carbapenem antibiotic class, and amikacin (aminoglycoside class), and tigecycline (tetracycline class) still display high sensitivity (100%). The most resistance pattern found in this study was AM/AMP/KZ/CTX/CRO/ATM/GM/CIP (22.73%), and AM/AMP/KZ/CTX/CRO/ATM/GM/CIP/SXT (18.2%). The result of multi class antibiotic resistance exhibits 86.36% are multidrug resistant, with the highest multidrug resistance occurred in BL/AG/Q/SP (50%).

Acknowledgment

This study was funded by *Dana Hibah Penelitian Mandat* from Universitas Airlangga, Indonesia in 2019.

References

- Amelia, A., A. Nugroho, and P. N. Harijanto. 2016. Diagnosis and management of infections caused by enterobacteriaceae producing extended spectrum Beta-Lactamase. *Acta Medica Indonesiana -The Indonesian Journal of Internal Medicine*. 48: 156–166.
- Biomerieux. 2017. AST and Resistance Detection, Antibiotic Susceptibility Testing bioMérieux Industry website. Available from: <http://www.biomerieux-industry.com/ast-and-resistance-detection>.
- Biutifasari, V. 2018. Extended Spectrum Beta – Lactamase (ESBL). *oceana Biomedicina Journal*. 1: 1–6.
- Brower, C. H., S. Mandal, S. Hayer, M. Sran, A. Zehra, S. J. Patel, R. Kaur, L. Chatterjee, S. Mishra, B. R. Das, P. Singh, R. Singh, J. P. S. Gill, and R. Laxminarayan. 2017. The prevalence of extended-spectrum beta-lactamase-producing multidrug-resistant *Escherichia coli* in poultry chickens and variation according to farming practices in Punjab, India. *Environmental Health Perspectives*. 125: 1–10. doi:10.1289/EHP292.
- Bywater, R., H. Deluyker, E. Deroover, A. de Jong, H. Marion, M. McConville, T. Rowan, T. Shryock, D. Shuster, V. Thomas, M. Vallé, and J. Walters. 2004. A European survey of antimicrobial susceptibility among zoonotic and commensal bacteria isolated from food-producing animals. *Journal of Antimicrobial Chemotherapy*. 54:744–754. doi:10.1093/jac/dkh422.
- Clinical and Laboratory Standards Institute (CLSI). 2017. M100 Performance Standards for Antimicrobial. 27th ed. Clinical and Laboratory Standards Institute, USA. Available from: www.clsi.org
- Cormican, M. G., S. A. Marshall, and R. N. Jones. 1996. Detection of extended-spectrum β -lactamase (ESBL)-producing strains by the Etest ESBL screen. *Journal of Clinical Microbiology*. 34: 1880–1884.
- Duggal, S., R. Gaiind, N. Tandon, M. Deb, and T. Das Chugh. 2012. Comparison of an automated system with conventional identification and antimicrobial susceptibility testing. *International Scholarly Research Network Microbiology*. 2012. doi:10.5402/2012/107203.
- Effendi, M. H., I. G. Bintari, E. B. Aksoro, and I. P. Hermawan. 2018. Detection of blaTem Gene of Klebsiella pneumoniae Isolated from swab of food-producing animals in East Java. *Tropical Animal Science Journal*. 41: 174–178. doi:10.5398/tasj.2018.41.3.174.
- Hammerum, A. M., J. Larsen, V. D. Andersen, C. H. Lester, T. S. S. Skytte, F. Hansen, S. S. Olsen, H. Mordhorst, R. L. Skov, F. M. Aarestrup, and Y. Agersø. 2014. Characterization of extended-spectrum β -lactamase (ESBL)-producing *Escherichia coli* obtained from Danish pigs, pig farmers and their families from farms with high or no consumption of third- or fourth-generation cephalosporins. *Journal of Antimicrobial Chemotherapy*. 69: 2650–2657. doi:10.1093/jac/dku180.
- Handayani, R. S., S. Siahaan, and M. J. Herman. 2017. Resistensi antimikroba dan penerapan kebijakan pengendalian di Rumah Sakit di Indonesia. *Jurnal Penelitian dan Pengembangan Pelayanan Kesehatan*. 1: 131–140.
- Kürekci, C., M. Aydin, M. Yipel, M. Katouli, and A. Gündogdu. 2017. Characterization of extended spectrum β -lactamase (ESBL)-producing *Escherichia coli* in Asi (Orontes) River in Turkey. *Journal of Water and Health*. 15: 788–798. doi:10.2166/wh.2017.257.
- Kurniawati, A. F., P. Satyabakti, and N. Arbianti. 2015. Perbedaan Risiko Multidrug Resistance Organisms (MDROS) Menurut Faktor Risiko dan Kepatuhan Hand Hygiene. *Jurnal Berkala Epidemiologi*. 3:277–289. doi:<http://dx.doi.org/10.20473/jbe.V3i32015.277-289>.
- Lim, C. L. L., W. Lee, A. L. C. Lee, L. T. T. Liew, S. C. Nah, C. N. Wan, M. P. Chlebicki, and A. L. H. Kwa. 2013. Evaluation of ertapenem use with impact assessment on extended-spectrum β -lactamases

- (ESBL) production and gram-negative resistance in Singapore general hospital (SGH). *BMC Infectious Diseases*. 13: 1–10. doi:10.1186/1471-2334-13-523.
- Masrurroh, C. A., M. B. Sudarwanto, and H. Latif. 2016. Tingkat kejadian *Escherichia coli* penghasil extended spectrum B - Lactamase yang diisolasi dari feses broiler di kota Bogor. *Jurnal Sain Veteriner*. 34: 42–49.
- Niasono, A. B., H. Latif, and T. Purnawarman. 2019. Resistensi antibiotik terhadap bakteri *Escherichia coli* yang diisolasi dari peternakan ayam pedaging di Kabupaten Subang, Jawa Barat. *Jurnal Veteriner*. 20: 187–195. doi:10.19087/jveteriner.2019.20.2.187.
- Noor, S. M. and M. Poeloenga. 2005. Pemakaian antibiotika pada ternak. Lokakarya Nasional Keamanan Pangan Produk Peternakan. Auditorium Balai Penelitian Veteriner. Bogor. 56–64.
- Normaliska, R., M. B. Sudarwanto, and H. Latif. 2019. Pola resistensi antibiotik pada *Escherichia coli* penghasil ESBL dari sampel lingkungan di RPH-R Kota Bogor. *Acta Veterinaria Indonesiana*. 7: 42–48. doi:10.29244/avi.7.2.42-48.
- Nuangmek, A., S. Rojanasthien, S. Chotinun, P. Yamsakul, P. Tadee, V. Thamlikitkul, N. Tansakul, and P. Patchanee. 2018. Antimicrobial resistance in ESBL-Producing *Escherichia coli* isolated from layer and pig farms in Thailand. *Acta Scientiae Veterinariae*. 46: 1538.
- O'Hara, C. M. 2005. Manual and automated instrumentation for identification of *enterobacteriaceae* and other aerobic gram-negative bacilli. *Clinical microbiology reviews*. 18: 147–162. doi:10.1128/CMR.18.1.147.
- Paterson, D. L. and R. A. Bonomo. 2005. Extended Spectrum Beta-Lactamases: a Clinical Update. *Clinical Microbiology Reviews*. 18: 657–686. doi:10.1128/CMR.18.4.657.
- Rank, E. L., T. Lodise, L. Avery, E. Bankert, E. Dobson, G. Dumyati, S. Hasset, M. Keller, M. Pearsall, T. Lubowski, and J. J. Carreno. 2018. Antimicrobial susceptibility trends observed in urinary pathogens obtained from New York state. *Open Forum Infectious Diseases*. 5: 1–6. doi:10.1093/ofid/ofy297.
- Santos, L. L. dos, R. A. Moura, P. Aguilar-Ramires, A. P. De Castro, and N. Lincopan. 2013. Current status of extended-spectrum β -lactamase (ESBL)-producing Enterobacteriaceae in animals. Microbial pathogens and strategies for combating them: science, technology and education. 1600–07.
- Schroeder, C. M., D. G. White, and J. Meng. 2004. Retail meat and poultry as a reservoir of antimicrobial-resistant *Escherichia coli*. *Food Microbiology*. 21:249–255. doi:10.1016/S0740-0020(03)00074-1.
- Schmid, A., S. Hörmansdorfer, U. Messelhäusser, A. Käsbohrer, C. Sauter-Louis, and R. Mansfeld. 2013. Prevalence of extended-spectrum β -lactamase-producing *Escherichia coli* on Bavarian dairy and beef cattle farms. *Applied and Environmental Microbiology*. 79: 3027–3032. doi:10.1128/AEM.00204-13.
- Sudarwanto, M. B., D. W. Lukman, T. Purnawarman, H. Latif, H. Pisestyani, and E. Sukmawinata. 2017. Multidrug resistance extended spectrum β -lactamase and AmpC producing *Escherichia coli* isolated from the environment of Bogor Slaughterhouse, Indonesia. *Asian Pacific Journal of Tropical Biomedicine*. 7: 708–711. doi:10.1016/j.apjtb.2017.07.012.
- Sugiartha, I. G. E. 2016. Perbandingan Hasil Identifikasi Metode Analytical Profile Index (API) dan Tes Kepekaan Antibiotika Konvensional dengan Metode Technical Dedicated Reasonable (TDR)-300B. Universitas Airlangga.
- Wang, X., C. Cao, H. Huan, L. Zhang, X. Mu, Q. Gao, X. Dong, S. Gao, and X. Liu. 2015. IsoEc22 R S R S S R MDR BL/AG/Q/SPIation, identification, and pathogenicity of O142 avian pathogenic *EscherichEc22 R S R S S R MDR BL/AG/Q/SPIa coli* causing black proventriculus and septicemia in broiler breeders. *Infection, Genetics and Evolution*. 32:23–29. doi:10.1016/j.meegid.2015.02.013.
- Wibisono, F. J., B. Sumiarto, and T. K. Kusumastuti. 2018. Economic losses estimation of pathogenic *Escherichia coli* infection in Indonesian Poultry Farming. *Buletin Peternakan* 42: 341–346. doi:10.21059/buletinpeternak.v42i4.37505.
- Wiedosari, E. and S. Wahyuwardani. 2015. A case study on the diseases of broiler chicken in Sukabumi and Bogor Districts. *Jurnal Kedokteran Hewan*. 9: 9–13.