

Application of chitosan scaffolds on vascular endothelial growth factor and fibroblast growth factor 2 expressions in tissue engineering principles

by Maretaningtias Dwi Ariani

Submission date: 04-Apr-2023 10:43AM (UTC+0800)

Submission ID: 2055226505

File name: 2015_Dental_Journal.pdf (510.42K)

Word count: 2832

Character count: 15853

Research Report

Application of chitosan scaffolds on vascular endothelial growth factor and fibroblast growth factor 2 expressions in tissue engineering principles

Ariyati Retno Pratiwi,¹ Anita Yulianti,² Istiati Soepribadi,³ and Maretaningtias Dwi Ariani⁴

¹Master Program in Dental Health Science

²Department of Dental Material

³Department of Oral Pathology and Maxillofacial

⁴Department of Prosthodontics

Faculty of Dental Medicine, Universitas Airlangga
Surabaya-Indonesia

ABSTRACT

Background: Tissue engineering has given satisfactory results as biological tissue substitutes to restore, replace, or regenerate tissues that have a defect. Chitosan is an organic biomaterial often used in the biomedical field. Chitosan has biocompatible, antifungal, and antibacterial properties. Chitosan is osteoconductive, suitable for bone regeneration applications. Bone defect healing begins with inflammatory phase as a response to the presence of vascular injury, so new vascularization is required. Vascular endothelial growth factor (VEGF) and basic fibroblast growth factor-2 (FGF2) are indicators of the beginning of bone regeneration process, playing an important role in angiogenesis. **Purpose:** This research was aimed to determine the effects of chitosan scaffold application on the expressions of VEGF and FGF2 in tissue engineering principles. **Method:** Chitosan was dissolved in CH_3COOH and NaOH to form a gel. Chitosan gel was then printed in mould to freeze dry for 24 hours. Those rats with defected bones were divided into two groups. Group 1 was the control group which defected bones were not administrated with chitosan scaffolds. Group 2 was the treatment group which defected bones were administrated with chitosan scaffolds. Those rats were sacrificed on day 14. Tissue preparations were made, and then immunohistochemical staining was conducted. Finally, a statistical analysis was conducted using Kruskal Wallis test. **Result:** There was no significant difference in the expressions of VEGF and FGF2 between the control group and the treatment group ($p > 0.05$). **Conclusion:** Chitosan scaffolds do not affect the expressions of VEGF and FGF2 during bone regeneration process on day 14 in tissue engineering principles

Keywords: chitosan; VEGF; FGF2; tissue engineering

Correspondence: Ariyati Retno Pratiwi, c/o: Peserta Program S2 Ilmu Kesehatan Gigi, Fakultas Kedokteran Gigi Universitas Airlangga, Jl. Mayjend. Prof. Dr. Moestopo no. 47 Surabaya 60132, Indonesia. E-mail: ariyatiretnop@gmail.com

INTRODUCTION

Tissue engineering is an application of the principles and methods of engineering used to restore, maintain, or improve tissue function. The principles of tissue engineering are conducted by providing appropriate materials to trigger the cells in order to regenerate or combine the performance of the cells in the body by using scaffold to trigger the growth of new tissue. Tissue engineering has given satisfactory results as biological tissue substitutes to

restore, replace, or regenerate tissue suffering from defects.¹ Cells, scaffold, and growth stimulating signals are generally called as tissue engineering triad, a major component of tissue engineering.

Scaffold is an important material in tissue engineering. Scaffold is a very porous artificial extracellular matrix used for cell accommodation, cell growth, and tissue regeneration.² Scaffold must meet some requirements, such as has interconnecting pores appropriate to support tissue integration and vascularity. It made from materials that have

certain properties, such as biodegradation or bioresorption. As a result, the tissue will eventually be replaced with scaffolds, which have a surface suitable for supporting cell attachment, cell differentiation, and cell proliferation, as well as have good mechanical properties, easily made into a variety of shape and size.³

Organic biomaterials are often used in the biomedical field, especially chitosan application. Chitosan has biocompatibility, antifungi, and antibacteria properties.⁴ Chitosan also is osteoconductive, so it is suitable for hard tissue regeneration application, but the mechanical properties and biological activities need to be improved.⁵ The biological properties of chitosan that very influential in the process of wound healing are biodegradability, biocompatibility, and antimicrobial activity. The high biodegradation rates can trigger chitosan to be increasingly inadequate in accelerating wound healing. Biodegradability of chitosan also affects its biocompatibility because high degradation will increase amino sugar accumulation and inflammatory response.⁶

Bone healing process is a series of molecular and cellular process as well as tissue transformation, from resorption to hard and soft tissue formation. Bone defect healing process begins with inflammatory phase, which is vascular response to injury, requiring new vascularization. Vascularization plays an important role in osteogenesis during the bone healing process.⁷⁻⁹ Increased formation of blood vessels in the area indicates that wound epithelialization process occurs faster.¹⁰

Bone regeneration can be affected by vascular endothelial growth factor (VEGF) directly or indirectly. VEGF plays an important role in the formation of new blood vessels serving to mobilize and recruit endothelial progenitor cell (EPC), as well as to differentiate and proliferate endothelial cells.⁸ VEGF induces angiogenic process through endothelial cells. Bone-forming precursor cells migrate through the bloodstream to the callus that will differentiate into osteoblasts.⁹ VEGF affects osteogenesis from day 14 to day 21 after the defect occurred.

In addition to VEGF, basic fibroblast growth factor 2 (FGF2) plays an important role in the process of vascularization. VEGF and FGF2 can also stimulate fibroblasts to migrate to the defect and trigger collagen synthesis. FGF2 expression occurs in the early phase of bone healing process to form osteoblasts.¹¹ FGF2 also plays a role in mitogenesis of mesenchymal cells, proliferating and differentiating into progenitor cells. Progenitor cells will differentiate into osteoblasts as bone formation cells.¹³

This research was aimed to determine the effects of chitosan scaffold application on VEGF and FGF2 expressions in tissue engineering principles.

MATERIALS AND METHOD

This research was a laboratory experimental research with post control group design. Animals used were male rats (*Rattus norvegicus*) aged 3 months old and weighed 250 grams. Materials used in this research were chitosan with deacetylation 81% (Sigma 93646, USA), a solution of acetic acid (CH₃COOH) and NaOH (Merck, Germany), VEGF polyclonal antibody (BIOSSUSA), and FGF2 polyclonal antibody (BIOSSUSA). Tools used in this research were a glass beaker, connicle tube, freezer (Royal Chest Freezer BD 195, China), spatula glass, scales (Pioneer, USA), magnetic stirrer (HANNA, USA), and freeze-dryer (Heto FD3, EN 87 164, Japan).

Chitosan scaffold was synthesized by dissolving chitosan powder into a solution of acetic acid (CH₃COOH 0.5M) and sodium hydroxide (NaOH 0.1M), and then centrifuged at a speed of 9000 rpm for 10 minutes. Supernatant in the form of chitosan gel was inserted into the mold. The mold already containing chitosan gel was frozen at a temperature of -20° C using deep freezer for 2 hours, and then freeze-drying was conducted for 24 hours to form a porous three-dimensional structure, known as scaffold.

Research procedure was performed into several phases. Those six rats were divided into two groups. Group 1 was the control group, while group 2 was the treatment group. Those rats were acclimatized for one week. The manufacture of bone defects was performed in both groups by drilling the rats' femoral and dextral areas. During the making of bone defects, irrigation was made using Ringer solution. In the defect area of the control group, placebo scaffold was applied to each defect using tweezers and escavator, and then sutured with 3/0 non-absorbable black-silk thread on muscles. In the defect area of the treatment group, chitosan scaffold was applied to each defect using tweezers and escavator, and then sutured with 3/0 non-absorbable black-silk thread on muscles.

On the 14th day after the closure of the defect, those rats were anesthetized using 10% ether as asphyxiation, and then sacrificed. The defected tissues of those rats were cut. The pieces of the tissues were fixed with 10% formalin for two days. They were decalcified using EDTA in order to soften the tissues to facilitate the process of cutting. The tissues were cut to a thickness of 0.3 mm. Dehydration process was carried out using alcohol. Those pieces of the tissues were put in xylol solution three times. Paraffin infiltration was conducted by melting solid paraffin. Embedding tissue was performed by pouring paraffin into molding devices. The tissues were taken using tweezers, and then implanted into the mold, which had been filled by paraffin. The mold, which had been filled by paraffin and tissue, was cooled to form paraffin blocks. Those rats that had been sacrificed were then buried properly.

Immunohistochemical preparations were performed by cutting the paraffin block using a rotary microtome. The pieces placed in the glass object was then deparaffinized using xylol solution. The preparations were dripped with normal serum evenly on a glass object, and then put in a preparation box (humidity chamber) with tissue paper, etched with PBS to keep the moisture. The preparations were put in an incubator at a temperature of 37° C for 45 minutes. The tissues were dripped with primary antibody against inducible nitric oxide synthase (iNOS) enzyme, and incubated at 4° C for one night. The tissues then were dripped with secondary antibody (biotinylation), and incubated at 37° C for 35 minutes. The administration of chromogen was performed by dripping a solution of 3,3'- diaminobenzidine (DAB), then incubated at 37° C for 35 minutes. Counter stain was conducted by dripping hematoxylin shed evenly, then left for 15 seconds and washed with running water. Dehydration was carried out with alcohol solution 70%, 80%, 90%, and 100%. Clearing with xylol (I, II, and III) and closing the preparat (mounting) then were carried out immediately using the cover glass.

Expressions of VEGF and FGF2 in each sample were assessed semiquantitatively using the modification of Remmele method. Remmele scale index (immuno reactive score/IRS) is the result of multiplying immunoreactive cells percentage score to color intensity score on the immunoreactive cells. Data then were obtained from the average value of the IRS in each sample observed in five different fields of view at 400x magnification. The whole of this examination was performed using light microscope (Nikon N600L) equipped with a 300 megapixel DS Fi2 digital camera and image processing software (Nikon Image System).

RESULTS

The VEGF and FGF2 distribution were show in Figure 1. The average of VEGF and FGF2 expression in the control and treatment groups were shown in Figure 2. Based on the results of the statistical Kruskal Wallis test, p value obtained was less than 0.05, which means that there was no significant difference in VEGF and FGF2 expressions between the control group and the treatment group.

DISCUSSION

Tissue engineering involves cells as a building block, scaffold as a template, and growth factor as a biochemical signal that indicates there has been a growth of tissue. The primary function of scaffold is as cell support, an artificial extracellular matrix. It is not only providing sufficient mechanical environment of cells, but also causing cell attachment, proliferation, differentiation, and metabolism signals.⁴ Selection of biomaterials for scaffold design is essential to cell growth and proliferation in three-dimensional matrix.

Chitosan is a natural polysaccharide which is similar to glycosaminoglycans and has a good interaction with cell membrane.⁶ Natural polysaccharides can stimulate the activity of growth factors, which can maintain cell phenotype in particular morphology and play an important role as scaffold component of soft tissue and hard tissue.⁷

Chitosan-containing N-acetyl-D-glucosamine can bind to receptors that recognize macrophages. Macrophages produce VEGF directly to stimulate endothelial cell proliferation.⁹ Chitosan may provide more amino groups




Figure 1. VEGF expressions on the control group (a) and the treatment group; (b) FGF2 expressions on the control group; (c) and the treatment group; (d) VEGF and FGF2 expressions were evenly distributed in the control and treatment groups.




Figure 2. The average of VEGF and FGF2 expression.

for the attachment and proliferation of endothelial cells because of the affinity between cations of the ammonium group of chitosan and the anion surface of the endothelial cell membrane.¹³

In this research, chitosan with a degree of deacetylation of 81% was used. Deacetylation degree can increase the attachment and stimulate fibroblast proliferation. Deacetylation degree of chitosan can also indicate free amine groups presented in the structure of chitosan. Cationic amine chitosan group provides a suitable environment for cell attachment. Cell attachment, is not only influenced by acetylation degree, but also by molecular weight and porosities. Chitosan, furthermore, can increase the proliferation of fibroblasts indirectly through the formation of poly-electrolyte complex with serum as heparin.¹³

The results of this research showed that there was no significant difference in VEGF expressions between the treatment group and the control group on day 14 during the process of bone regeneration. Chitosan could stimulate inflammatory cells and growth factors in the early phase of wound healing process. This is supported by a research conducted by Inan and Saraydin indicating that the expression of VEGF in post-administration of chitosan will decline on day 14.¹⁴ Chitosan could also stimulate the formation of granulation tissue. At the same time, chitosan could stimulate both fibroblasts to proliferate and extracellular matrix formation.

This study also showed that there was no significant difference in FGF2 expressions between the treatment group and the control group on day 14 during the process of bone regeneration. The number of FGF2 expressions on the 14th day is lower than on the 3rd day.¹⁴ This condition is possible because fibroblasts have produced collagen fiber on the 14th day. Consequently, fibroblasts will grow into inactive fibroblasts, known as fibrocyte. In this study, however, chitosan scaffold on the 14th day could still interact with the endothelial cells and fibroblasts although not optimal. Finally, it may be concluded that chitosan

scaffold cannot affect VEGF and FGF2 expressions on the 14th day during bone regeneration process in tissue engineering principles.

REFERENCES

- Karp JM, Langer R. Development and therapeutic applications of advanced biomaterials. *Curr Opin Biotechnol* 2007; 18(5): 454-9.
- Chan BP, Leong KW. Scaffolding in tissue engineering: general approaches and tissue-specific considerations. *Eur Spine J* 2008; 17(Suppl 4): 467-9.
- Sachlos E, Czernuszka JT. Making tissue engineering scaffolds work. Review: The application of solid free form fabrication technology to the production of tissue engineering scaffolds. *Euro Cell Mater* 2003; 5: 29-39.
- Shin JA, Choi JY, Kim ST, Kim CS, Lee YK, Cho KS, Chai JK, Kim CK, Choi SH. The effect of hydroxyapatite-chitosan membrane on bone regeneration in rat calvarial defects. *J Korean Acad Periodontol* 2009; 39: 213-4.
- Isikli C, Hasirci V, Hasirci N. Development of porous chitosan-gelatin/hydroxyapatite composite scaffold for hard tissue engineering application. *J Tissue Eng Reg Med* 2012; 6(2): 135-43.
- Aranaz I, Mengibar M, Harris R, Panos I, Miralles B, Acosta N, Galed G, Heras A. Functional characterization of chitin and chitosan. *Current Chemical Biology* 2009; 3: 203-30.
- Hankenson KD, Dishowitz M, Gray C, Schenker M. Angiogenesis in bone regeneration. *Injury* 2011; 42(6): 556-61.
- Saran U, Piperni SG, Chatterjee S. Role of angiogenesis in bone repair. *Arch Biochem and Biophys* 2014; 561: 109-17.
- Stegen S, Gastel NV, Carmeliet G. Bringing new life to damaged bone: The importance of angiogenesis in bone repair and regeneration. *Bone* 2014; 70: 19-25.
- Mackay D, Miller A. Nutritional support for wound healing. *Altern Med Rev* 2003; 8(4): 359-77.
- Dimitriou R, Tsiridis E, Giannoudis PV. Current concept of molecular aspects of bone healing. *Injury* 2005; 36(12): 1392-404.
- Arvidson K, Abdallah BM, Applegate LA, Baldini N, Cenni E, Gomez-Barrena E, Granchi D, Kassem M, Kontinen YT, Mustafa K, Pioletti DP, Sillat T, Finne-Wistrand A. Bone regeneration and stem cells. *J Cell Mol Med* 2011; 15(4): 718-46.
- Ma L, Gao C, Mao Z, Zhou J, Shen J, Hu X, Han C. Collagen/chitosan porous scaffold with improved biostability for skin tissue engineering. *Biomaterials* 2003; 24(26): 4833-41.
- Deniz I, Serpil S. Investigation of the wound healing effects of chitosan on FGF3 and VEGF immunolocalization in experimentally diabetic rats. *Int J Biomed Mat Research* 2013; 1: 1-8.

Application of chitosan scaffolds on vascular endothelial growth factor and fibroblast growth factor 2 expressions in tissue engineering principles

ORIGINALITY REPORT

19%

SIMILARITY INDEX

12%

INTERNET SOURCES

17%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Rini Devijanti Ridwan, Yuliati, Sidarningsih, Fitri Mar'atush Sholihah, Mohammed Ahmed Qasim Aljunaid, Dur Mohammad Lashari. "A study of the mucoadhesive patches loaded with mangosteen peel extract in periodontitis", Journal of Taibah University Medical Sciences, 2021
Publication 2%

- 2** www.dovepress.com
Internet Source 2%

- 3** Juan Chen, Lu Qian, Cuilian Chen, Xiaoling Wang. "The Characteristics of Fear of Recurrence and the Effect of Cognitive-Behavioral Stress Management Intervention in Patients after Radiofrequency Ablation of Atrial Fibrillation", Evidence-Based Complementary and Alternative Medicine, 2022
Publication 1%

4

Chao Li, Zhi-bin Wang. "Clinical application of compound Glycyrrhizin tablets in the treatment of patients with Simplex Henoch-Schonlein Purpura and its effect on immune function", Pakistan Journal of Medical Sciences

Internet Source

1 %

5

Ashish Agarwal, Narinder Dev Gupta, Avikal Jain. "Platelet rich fibrin combined with decalcified freeze-dried bone allograft for the treatment of human intrabony periodontal defects: a randomized split mouth clinical trail", Acta Odontologica Scandinavica, 2015

Publication

1 %

6

[researcherslinks.com](https://www.researcherslinks.com)

Internet Source

1 %

7

Kaviya Baskar, Saravanakarthykeyan Balasubramanian, Ishwarya Gurucharan, Sekar Mahalaxmi et al. "Eggshell derived nano - hydroxyapatite incorporated carboxymethyl chitosan scaffold for dentine regeneration: a laboratory investigation", International Endodontic Journal, 2021

Publication

1 %

8

Xu Zhang, Li Zhou, Min Cai, Naxin Cui, Guoyan Zou, Qian Wang. "Effects of photocatalysis using a photocatalytic concrete board on water qualities and microbial communities in

1 %

the aquaculture wastewater", Separation and Purification Technology, 2023

Publication

9

Yuan Xu, Hongbin Wen, Jie Li, Jing Yang, Kai Luo, Liying Chang. "The relationship between sleep disorders, anxiety, depression, and cognitive function with restless legs syndrome (RLS) in the elderly", Sleep and Breathing, 2021

Publication

1 %

10

www.mmj.eg.net

Internet Source

1 %

11

Amira M. M. Amin, Emad M. M. Ewais. "Chapter 3 Bioceramic Scaffolds", IntechOpen, 2017

Publication

1 %

12

www.omicsdi.org

Internet Source

1 %

13

S. Geetha Priya. "Skin Tissue Engineering for Tissue Repair and Regeneration", Tissue Engineering Part B Reviews, 03/2008

Publication

<1 %

14

www.sfn.org

Internet Source

<1 %

15

B. P. Chan. "Scaffolding in tissue engineering: general approaches and tissue-specific

<1 %

considerations", European Spine Journal,
12/2008

Publication

16

Tamara Yuanita, Ridzki A Oktavianti, Debby F Suryani, Mandojo Rukmo, Sri Kunarti, Andrie H Kusuma. "The Inhibitory Ability of Cocoa Pod Husk Extract on Enterococcus faecalis Glucosyltransferase Enzyme Activity", The Journal of Contemporary Dental Practice, 2020

Publication

17

journals.itb.ac.id

Internet Source

18

Richard da Costa Marques, Johanna Simon, Cyril d'Arros, Katharina Landfester, Kerstin Jurk, Volker Mailänder. "Proteomics reveals differential adsorption of angiogenic platelet lysate proteins on calcium phosphate bone substitute materials", Regenerative Biomaterials, 2022

Publication

19

pubmed.ncbi.nlm.nih.gov

Internet Source

20

www.science.gov

Internet Source

21

123dok.com

Internet Source

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

22 Anuradha Subramanian, Hsin-Yi Lin. <1 %
"Crosslinked chitosan: Its physical properties and the effects of matrix stiffness on chondrocyte cell morphology and proliferation", Journal of Biomedical Materials Research Part A, 2005
Publication

23 de.scribd.com <1 %
Internet Source

24 Sanna M. Peltola, Ferry P. W. Melchels, Dirk W. Grijpma, Minna Kellomäki. <1 %
"A review of rapid prototyping techniques for tissue engineering purposes", Annals of Medicine, 2009
Publication

25 Shoufeng Yang. <1 %
"The Design of Scaffolds for Use in Tissue Engineering. Part I. Traditional Factors", Tissue Engineering, 12/2001
Publication

26 Wookbong Kwon, Hyeng - Soo Kim, Jain Jeong, Yonghun Sung et al. <1 %
" Tet1 overexpression leads to anxiety - like behavior and enhanced fear memories the activation of calcium - dependent cascade through Egr1 expression in mice ", The FASEB Journal, 2017
Publication

27 openarchive.ki.se
Internet Source

<1 %

28

phcogj.com

Internet Source

<1 %

29

www.mdpi.com

Internet Source

<1 %

30

www.scribd.com

Internet Source

<1 %

31

www.tandfonline.com

Internet Source

<1 %

32

Nike Hendrijantini, Yonatan Christian Suisan, Rizko Wira Artha Megantara, Bambang Agustono Satmoko Tumali et al. "Bone Remodeling in Mandible of Wistar Rats with Diabetes Mellitus and Osteoporosis", European Journal of Dentistry, 2022

Publication

<1 %

33

crjim.ub.ac.id

Internet Source

<1 %

34

Junyu Xiang, Jianmei Li, Jian He, Xiangyu Tang et al. "Cerium Oxide Nanoparticle Modified Scaffold Interface Enhances Vascularization of Bone Grafts by Activating Calcium Channel of Mesenchymal Stem Cells", ACS Applied Materials & Interfaces, 2016

Publication

<1 %

35

Salmon Charles PT Siahaan, Budi Santoso, -
Widjiati. "Effectiveness of Moringa oleifera
Leaves on TNF- α Expression, Insulin Levels,
Glucose Levels and Follicle Count in Rattus
norvegicus PCOS Model", Diabetes, Metabolic
Syndrome and Obesity: Targets and Therapy,
2022

Publication

<1 %

36

hangtuah.ac.id

Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Application of chitosan scaffolds on vascular endothelial growth factor and fibroblast growth factor 2 expressions in tissue engineering principles

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4
