

## DAFTAR PUSTAKA

- Boneti, N. *et al.* (2018) ‘Does functional capacity , fall risk awareness and physical activity level predict falls in older adults in different age groups ?’, *Archives of Gerontology and Geriatrics*. Elsevier, 77(March), pp. 57–63. doi: 10.1016/j.archger.2018.04.002.
- Boongird, C. and Ross, R. (2015) ‘Views and Expectations of Community-Dwelling Thai Elderly in Reporting Falls to Their Primary Care Physicians : A Mixed-Methods Study’, *Shouthern Gerontological Society*, pp. 1–19. doi: 10.1177/0733464815606799.
- Brigham, J. C. (1991) *Social Psychology*. 2nd Editio. New York: Harper Collins Publishers. Inc.
- Darmojo, B. (2014) *Buku Ajar Boedhi-Darmojo: Ilmu Kesehatan Usia Lanjut*. 5th edn. Edited by H. H. Martono and K. Pranaka. Jakarta: Badan Penerbit FKUI.
- Dinas Kesehatan Pemerintah Kota Surabaya (2018) *Profil Kesehatan 2017*. Surabaya: Dinas Kesehatan Pemerintah Kota Surabaya.
- Duray, M. and Genc, A. (2017) ‘The relationship between physical fitness and falling risk and fear of falling in community-dwelling elderly people with different physical activity levels’, *Turkish Journal of Medical Sciences*, 47, pp. 455–462. doi: 10.3906/sag-1511-101.
- Goto, N. A. *et al.* (2019) ‘Accidental Falling in Community-Dwelling Elderly with Chronic Kidney Disease’, *International Urology and Nephrology*. Springer Netherlands, 51(1), pp. 119–127. doi: 10.1007/s11255-018-1992-9.
- Hu, J. *et al.* (2015) ‘Risk factors of indoor fall injuries in community-dwelling older women : A prospective cohort study’, *Archives Gerontology and Geriatrics*. Elsevier Ireland Ltd, 60(2), pp. 259–264. doi: 10.1016/j.archger.2014.12.006.
- Jitapunkul, S. *et al.* (1998) ‘Falls and their associated factors: a national survey of the Thai elderly’, *Journal of the Medical Association of Thailand = Chotmaihet Thangphaet*, 81(4), pp. 233–242.
- Junior, C. R. F. *et al.* (2018) ‘Factors associated with low levels of physical activity among elderly residents in a small urban area in the interior of the Brazilian Amazon’, *Gerontology and Geriatrics*, 75(November 2017), pp. 37–43. doi: 10.1016/j.archger.2017.11.007.
- Kementrian Kesehatan (2016) *Situasi lanjut usia (lansia) di Indonesia, 2016*.
- Kementrian Kesehatan (2017) ‘Analisis Lansia di Indonesia’, 2017.
- Klenk, J. *et al.* (2015) ‘Physical Activity and Different Concepts of Fall Risk Estimation in Older People – Results of the ActiFE-Ulm Study’, *Plos ONE*, (1), pp. 1–11. doi: 10.1371/journal.pone.0129098.
- Mendatu, A. (2010) *Pemulihan Trauma: Strategi Penyembuhan Trauma untuk Diri Sendiri, Anak, dan Orang Lain di Sekitar Anda*. Yogyakarta: Jalasutra.

- Mihaljcic, T. *et al.* (2014) ‘Development of a new self-awareness of falls risk measure ( SAFRM )’, *Archives of Gerontology and Geriatrics*. Elsevier Ireland Ltd, 59(2), pp. 249–256. doi: 10.1016/j.archger.2014.06.001.
- Mihaljcic, T. *et al.* (2015) ‘Self-Awareness of Falls Risk among Elderly Patients: Characterizing Awareness Deficits and Exploring Associated Factors’, *Physical Medicine and Rehabilitation*. Elsevier Ltd. doi: 10.1016/j.apmr.2015.08.414.
- Mihaljcic, T. *et al.* (2017) ‘Investigating the relationship between reduced self-awareness of falls risk , rehabilitation engagement and falls in older adults’, *Archives of Gerontology and Geriatrics*. Elsevier Ireland Ltd, 69, pp. 38–44. doi: 10.1016/j.archger.2016.11.003.
- Miller, C. A. (2012) *Nursing for Wellness in Older Adults*. Sixth. Edited by L. Williams and Wilkins. Hong Kong: Wolters Kluwer Health.
- Nursalam (2013) *Konsep Penerapan Metode Penelitian Ilmu Keperawatan*. Jakarta: Salemba Medika.
- Nursalam (2016) *Metodologi Penelitian Ilmu Keperawatan Edisi 4*. Jakarta: Penerbit Salemba Medika.
- Oliveira, C. C. *et al.* (2015) ‘Fear of falling in people with chronic obstructive pulmonary disease’, *Respiratory Medicine*. Elsevier Ltd, 109(4), pp. 483–489. doi: 10.1016/j.rmed.2015.02.003.
- Pengpid, S. and Peltzer, K. (2018) ‘Prevalence and Risk Factors Associated with Injurious Falls among Community-Dwelling Older Adults in Indonesia’, *Current Gerontology and Geriatrics Research*. Hindawi, 2018, pp. 1–8. doi: 10.1155/2018/5964305.
- Podsiadlo, D. and Richardson, S. (1991) ‘The Timed “Up & Go”: A Test of Basic Functional Mobility for Frail Elderly Person’, *Journal of American Geriatrics Society*, 39, pp. 142–148.
- Possamai, L. T. and Zogo, A. (2015) ‘Fitness for elders : A comparison between practioners and non-practioners of exercise’, *American Aching Association*, 37, pp. 1–8. doi: 10.1007/s11357-015-9772-2.
- Rolenz, E. and Reneker, J. C. (2016) ‘Validity of the 8-Foot Up and Go, Timed Up and Go, and Activities- Specific Balance Confidence scale in older adults with and without cognitive impairment’, *Journal of Rehabilitation and Development*, 53(4), pp. 511–518. doi: <http://dx.doi.org/10.1682/JRRD.2015.03.0042>.
- Russell, K. *et al.* (2017) ‘Measurement of Fall Prevention Awareness and Behaviours among Older Adults at Home’, *Canadian Journal on Aging*. doi: 10.1017/S0714980817000332.
- Sibley, K. M. *et al.* (2014) ‘Chronic disease and falls in community-dwelling Canadians over 65 years old : a population-based study exploring associations with number and pattern of chronic conditions’, *BMC Geriatrics*. Available

at: <http://www.biomedcentral.com/1471-2318/14/22>.

- Sugiyono (2012) *Metode Penelitian Kuantitatif, Kualitatif*. Bandung: Alfabeta.
- Sya'diyah, H. (2018) *Keperawatan Lanjut Usia*. Sidoarjo: Indomedia Pustaka.
- U.S. Department of Health and Human Services (2013) *Summary Health Statistics for the U . S . Population : National Health Interview Survey , 2012*. Washington, DC: U.S. Department of Health and Human Services.
- Uen Yea, Y. *et al.* (2016) 'Prevalence of falls among community-dwelling elderly and its associated factors : A cross-sectional study in Perak , Malaysia', *Malaysian Family Physician*, 11, pp. 7–14.
- Untari, I. (2018) *Buku Ajar Keperawatan Gerontik: Terapi Tertawa dan Senam Cegah Pikun*. Edited by W. Praptiani. Jakarta: EGC.
- Williams, J. S. *et al.* (2015) 'Prevalence , risk factors and disability associated with fall-related injury in older adults in low- and middle-income countries : results from the WHO Study on global AGEing and adult health ( SAGE )', *BMC Medicine*. BMC Medicine, pp. 1–12. doi: 10.1186/s12916-015-0390-8.
- World Health Organization (2011) 'Global Recommendations on Physical Activity for Health 65 years and above'. Available at: <http://www.who.int/dietphysicalactivity/pa/en/index.html>.
- World Health Organization (2015) *World Report on Ageing and Health*. Luxembourg.
- World Health Organization (2018a) 'Falls', pp. 1–4. Available at: <https://www.who.int/news-room/fact-sheets/detail/falls>.
- World Health Organization (2018b) *Noncommunicable Diseases Country Profiles 2018*. Geneva: World Health Organization.
- World Health Organization (2019a) 'Global Strategy on Diet, Physical Activity and Health', pp. 1–3. Available at: <https://www.who.int/dietphysicalactivity/pa/en/>.
- World Health Organization (2019b) 'Noncommunicable diseases', pp. 1–4. Available at: <https://www.who.int/en/news-room/fact-sheets/detail/noncommunicable-diseases>.