



# Journal of Advanced Pharmaceutical Technology & Research

An Official Publication of Society of Pharmaceutical Education & Research

Volume 11 / Issue 4 / October-December 2020

[www.japtr.org](http://www.japtr.org)

JAPTR



## Table of Contents



**October-December 2020**

Volume 11 | Issue 4

Page Nos. 157-237

Online since Saturday, October 10, 2020

Accessed 12,165 times.

### PDF access policy

Journal allows immediate open access to content in HTML + PDF

### EPub access policy

allows EPub access only to users from **INDIA**, developing countries and paid subscribers.

View issue as eBook

Issue citations

Issue statistics

RSS

### Feedback

Subscribe

Next Issue

Previous Issue

Most popular articles

Join as Reviewer

Email alerts

Recommend this journal to your library

[Show all abstracts](#) [Show selected abstracts](#) [Export selected to ▼](#) [Add to my list](#)

## ORIGINAL ARTICLES


**The coronavirus disease 2019 main protease inhibitor from *Andrographis paniculata* p. 157**

(Burm.f) Ness

Sukardiman, Martha Ervina, Mohammad Rizki Fadhil Pratama, Hadi Poerwono, Siswandono Siswodihardjo

DOI:10.4103/japtr.JAPTR\_84\_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Citations (1)] [Sword Plugin for Repository]<sup>Beta</sup>




**Anticancer evaluation of *N*-benzoyl-3-allylthiourea as potential antibreast cancer agent p. 163 through enhances HER-2 expression**

Tri Widiandani, Siswandono, Edy Meiyanto

DOI:10.4103/japtr.JAPTR\_77\_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]<sup>Beta</sup>




**Effects of lime (*Citrus aurantifolia*) peel to the expression of mRNA toll-like receptors 4 in p. 169 balb/c mice-infected *Salmonella typhi***

Vivien Novarina Kasim, Mochammad Hatta, Rosdiana Natzir, Veni Hadju, Ami Febriza, Hasta Handayani Idrus

DOI:10.4103/japtr.JAPTR\_48\_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]<sup>Beta</sup>




**Anti-aging potential and phytochemicals of *Centella asiatica*, *Nelumbo nucifera*, and *Hibiscus sabdariffa* extracts p. 174**

Monsicha Khuaneckaphan, Chanai Noysang, Warachate Khobjai

DOI:10.4103/japtr.JAPTR\_79\_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]<sup>Beta</sup>




**Proximate composition, vitamin and mineral composition, antioxidant capacity, and anticancer activity of *Acanthopanax trifoliatum* p. 179**

Aunyachulee Ganogpichayagrai, Chutinan Suksaard

DOI:10.4103/japtr.JAPTR\_61\_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]<sup>Beta</sup>




**Stabilization of eye drops containing autologous serum and recombinant human epidermal growth factor for dry eye syndrome p. 184**


Sriwidodo Sriwidodo, Insan Sunan Kurniawan Syah, Iman Permana Maksum, Toto Subroto, Ulvi Zasvia, Abd Kakhar Umar

DOI:10.4103/japtr.JAPTR\_72\_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]<sup>Beta</sup>



**Antidiabetic activity of thin film containing astaxanthin-loaded nanoemulsion using p. 189 carboxymethylcellulose sodium polymer on alloxan-induced diabetic rabbit**



Lusi Nurdianti, Taofik Rusdiana, Iyan Sopyan

DOI:10.4103/japtr.JAPTR\_55\_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]<sup>Beta</sup>**In silico docking and comparative ADMET profile of different glycogen synthase kinase 3 beta inhibitors as the potential leads for the development of anti-Alzheimer drug therapy** p. 194

Manisha Prajapat, Phulen Sarma, Nishant Shekhar, Hardeep Kaur, Sanjay Singh, Subodh Kumar, Harpinder


Kaur, Saniya Mahendiratta, Amit Raj Sharma, Sukhmandeep Kaur, Vidya M Mahalmani, Bikash Medhi

DOI:10.4103/japtr.JAPTR\_178\_19

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]<sup>Beta</sup>**Effect of implantation Ti-6Al-4V ELI in femoral bone defect regeneration of Sprague Dawley rat** p. 202

I Nyoman Jujur, Damisih, Kartika Devy, Shovy Suha, Boy Muchlis Bachtiar, Endang Winiati Bachtiar


DOI:10.4103/japtr.JAPTR\_74\_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]<sup>Beta</sup>**Development and validation of an ultra-performance liquid chromatography mass spectrometry/mass spectrometry method for simultaneous quantification of total and free mycophenolic acid and its metabolites in human plasma** p. 207

Napatsanan Tanathitiphuwarat, Pannee Tanudechpong, Pajaree Chariyavilaskul, Nantaporn Prompila, Nat

Tansrisawad, Apinya Tubtimrattana, Supeecha Wittayalertpanya


DOI:10.4103/japtr.JAPTR\_40\_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]<sup>Beta</sup>**The enhancement apoptosis of osteosarcoma mesenchymal stem cells co-cultivation with peripheral blood mononuclear cells sensitized by secretome and granulocyte macrophage colony-stimulating factor** p. 213

Ferdiansyah Mahyudin, Hizbillah Yazid, Mouli Edward, Mohammad Hardian Basuki, Yunus Abdul Bari, Fedik


Abdul Rantam

DOI:10.4103/japtr.JAPTR\_52\_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]<sup>Beta</sup>**Peel-off emulgel mask of *Cocos nucifera* L. Extract using gelling agent carbomer 940 as antiacne against *Propionibacterium acnes* ATCC 11827** p. 220

Dewi Melani Hariyadi, Isnaeni Isnaeni, Sisunandar Sudarma, Suciati Suciati, Noorma Rosita

DOI:10.4103/japtr.JAPTR\_51\_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]<sup>Beta</sup>**Accelerated wound healing ability of *Jatropha sap* by iota carrageenan-poly (vinyl alcohol) hydrogel film** p. 226

Nurain Thomas, Agus Rusdin, Madania Tulsyahra, Nasrul Wathoni, Asep Kuswandi


DOI:10.4103/japtr.JAPTR\_11\_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]<sup>Beta</sup>**Induction of cytotoxicity by *Bruguiera gymnorrhiza* in human breast carcinoma (MCF-7) cell line via activation of the intrinsic pathway** p. 233

Gul-e-Saba Chaudhry, Nurul Huda Rahman, Vigneswari Sevakumaran, Aziz Ahmad, Habsah Mohamad,

Muhammad Naveed Zafar, Yeong Yik Sung, Tengku Sifzizul Tengku Muhammad

DOI:10.4103/japtr.JAPTR\_81\_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]<sup>Beta</sup>



[Home](#) | [About JAPTR](#) | [Editorial board](#) | [Search](#) | [Ahead of print](#) | [Current issue](#) | [Archives](#) | [Submit article](#) | [Instructions](#) | [Subscribe](#) | [Advertise](#) | [Contacts](#) | [Login](#)

Users Online: 75    

## Editorial Board

### Editor-in-Chief

#### Dr. Upendra Nagaich

M.Pharm., Ph.D., FSPER

Secretary,

Society of Pharmaceutical Education & Research (SPER), India

&

Research Coordinator,

Amity Institute of Pharmacy, Amity University,

Noida (U.P.) India

### Executive Editors

#### Dr. Paul Wan Sia Heng

Department of Pharmacy,

National University of Singapore,

Singapore

#### Ms. Jaya Pandey

School of Studies in Pharmaceutical Sciences,

Jiwaji University,

Gwalior (M.P.) India

### Associate Editor

#### Dr. Vandana Chaudhary

School of Pharmacy,

Pharmacy Australia Centre of Excellence,

The University of Queensland,

Brisbane, Australia

### Consultant Editor

#### Prof. (Dr.) Laszlo Endrenyi

Professor Emeritus

Department of Pharmacology & Toxicology

University of Toronto, Toronto, Canada

### Section Editors (Alphabetical Order)

#### Dr. Ikoni Ogaji

Dept. of Pharmaceutics & Pharmaceutical Technology,

Faculty of Pharmaceutical Sciences,

University of Jos, Nigeria

#### Dr. Ke Li

J. David Gladstone Institutes,

San Francisco, California, USA

### Editorial Board Members (In Alphabetical Order)

#### Prof. (Dr.) A. N. Kalia

Vice-President, Society of Pharmacognosy,

Professor & Head

I.S.F. College of Pharmacy,

Moga (Punjab), India

#### Dr. Abdul Wahab

### Feedback

[Subscribe](#)

[About the journal](#)

[Most popular articles](#) 

[Join as Reviewer](#) 

[Email alerts](#) 

[Recommend this journal to your library](#) 

Department of Pharmacy  
Kohat University of Science & Technology  
Kohat, KPK, Pakistan

**Prof. (Dr.) Arun Dharmarajan**

Professor  
School of Biomedical Sciences,  
Faculty of Health Sciences,  
Curtin University,  
Perth, Australia

**Prof. (Dr.) Ashoke Kumar Ghosh**

Vice-Chancellor  
IFTM University,  
Moradabad (Uttar Pradesh), India

**Prof. (Dr.) B. Mishra**

Professor & Head  
Department of Pharmaceutics,  
Indian Institute of Technology,  
Banaras Hindu University,  
Varanasi (Uttar Pradesh), India

**Prof. B. G. Shivananda**

Secretary, APTI  
Bangalore (Karnataka), India

**Prof. (Dr.) Bharat B. Aggarwal**

Professor of Cancer Medicine &  
Chief, Cytokine Research Laboratory  
The University of Texas  
Texas, USA

**Dr. Bhaswat Sundar Chakraborty**

Senior VP & Chairman, R&D Core Committee  
Cadila Pharmaceuticals Ltd.  
Ahmedabad (Gujarat), India

**Prof. (Dr.) C. K. Kokate**

National Advisor, Society of Pharmacognosy,  
Vice-Chancellor  
KLE University,  
Belgaum (Karnataka), India

**Prof. (Dr.) Cedric Pearce**

Adjunct Professor  
University of North Carolina  
CEO, Mycosynthetix Inc.  
North Carolina, USA  
[Associate Editor, Journal of Natural Products]

**Prof. (Dr.) Ciddi Veeresham**

Professor  
University College of Pharmaceutical Sciences,  
Kakatiya University,  
Warangal (Andhra Pradesh), India

**Dr. Dâmaris Silveira**

Universidade de Brasília  
Brasilia, Brazil

**Prof. (Dr.) Dilipkumar Pal**

Professor  
Dept. of Pharmaceutical Sciences,  
Guru Ghasidas Vishwavidyalaya,  
Bilaspur (Chhattisgarh), India

**Prof. (Dr.) Divakar Goli**

Associate Secretary, IPA  
Secretary, Education Division, IPA  
Professor & Principal  
Acharya & B.M. Reddy College of Pharmacy,  
Bangalore (Karnataka), India

**Prof. (Dr.) G. K. Dash**

Professor  
Universiti Kuala Lumpur  
Perak Darul Ridzuan, Malaysia

**Prof. (Dr.) H. S. Buttar**

Adjunct Professor,  
School of Medicine,  
University of Ottawa,  
Ottawa, Canada

**Kaushik Desai**

Gen. Secretary, IPA  
Immediate Vice President, IPA  
Mumbai (Maharashtra), India

**Prof. (Dr.) Mainul Haque**

Professor  
Universiti Sultan Zainal Abidin  
Kuala Terengganu, Terengganu, Malaysia

**Dr. Muhammad Akhlaq**

University of Auckland  
Auckland, New Zealand

**Prof. (Dr.) N. Udupa**

Professor & Research Director [Health Sciences]  
Manipal University,  
Manipal (Karnataka), India

**Prof. (Dr.) Pornsak Sriamornsak**

Professor,  
Department of Pharmaceutical Technology &  
Pharmaceutical Biopolymer Group (PBiG),  
Faculty of Pharmacy, Silpakorn University,  
Nakhon Pathom, Thailand

**Dr. Rakesh Kumar Sharma**

Scientist 'F'  
Director,  
Defence Food Research Laboratory (DFRL),  
Mysuru (Karnataka), India

**Prof. (Dr.) Roop K. Khar**

Former Professor  
Department of Pharmaceutics,  
Jamia Hamdard (Hamdard University)  
New Delhi, India

**Prof. (Dr.) Roopa Karki**

Professor & Head  
Department of Pharmaceutics,  
Acharya & BM Reddy College of Pharmacy,  
Bangalore (Karnataka), India

**Prof. (Dr.) S. D. Tonpay**

Former Professor & Head  
Govt. Gajra Raja Medical College,  
Gwalior (Madhya Pradesh), India

**Prof. (Dr.) S. H. Ansari**

Professor & Head,  
Department of Pharmacognosy & Phytochemistry,  
Jamia Hamdard (Hamdard University)  
New Delhi, India

**Prof. (Dr.) S. H. Mishra**

Professor Emeritus  
Faculty of Pharmacy,  
The M. S. University of Baroda,  
Vadodara (Gujarat), India

**Prof. (Dr.) Satish Sardana**

Immediate Past President, APTI (Haryana)

Director,  
Amity Institute of Pharmacy,  
Amity University,  
Manesar (Haryana), India

**Prof. (Dr.) T. V. Narayana**

President, IPA  
Director  
Vikas Institute of Pharmaceutical Sciences  
Rajahmundry (Andhra Pradesh), India

**Prof. (Dr.) V. K. Dixit**

President, Society of Pharmacognosy,  
Former Professor  
Department of Pharmaceutical Sciences,  
Dr. H. S. Gour University,  
Sagar (M.P.), India

**Prof. (Dr.) Y. K. Agrawal**

Director  
Institute of Research & Development,  
Gujarat Forensic Sciences University,  
Gandhinagar (Gujarat), India

**Technical Board Members (In Alphabetical Order)**

Dr. Deenanath Jhade, Bhopal  
Dr. Inderbir Singh, Rajpura  
Dr. Mohd. Rageeb Mohd. Usman, Chopda  
Dr. Neeraj Sethiya, Vadodara  
Dr. Praveen Sharma, Indore  
Dr. Rakesh Pahwa, Kurukshetra  
Dr. Sanjib Bhattacharya, Kolkata  
Dr. Shweta Dixit, Pune  
Dr. Sunder Singh, Indore

# Peel-off emulgel mask of *Cocos nucifera* L. Extract using gelling agent carbomer 940 as antiacne against *Propionibacterium acnes* ATCC 11827

Dewi Melani Hariyadi,  
Isnaeni Isnaeni<sup>1</sup>,  
Sisunandar Sudarma<sup>2</sup>,  
Suciati Suciati<sup>3</sup>, Noorma Rosita

Departments of Pharmaceutics,  
<sup>1</sup>Pharmaceutical Chemistry and  
<sup>3</sup>Pharmacognosy and Phytochemsitry,  
Faculty of Pharmacy, Universitas  
Airlangga, Surabaya, <sup>2</sup>Department of  
Biology, Universitas Muhammadiyah  
Purwokerto, Purwokerto, Indonesia

*J. Adv. Pharm. Technol. Res.*

## ABSTRACT

*Cocos nucifera* Linn., which contain lauric acid has been known had antibacterial activity against *Propionibacterium acnes* that usually improve severe of pimple. Current study investigated optimum formula of emulgel mask based on the *C. nucifera* L. Extract from Kopyor coconut. Extract were tested for antibacterial against *P. acnes* ATCC 11827. In this research, *C. nucifera* L. extract of 1 and 5% were formulated as an active agent of peel off antiacne emulgel mask-containing carbomer 940 in various concentration (1% and 1.5%). The peel-off emulgel mask of *C. nucifera* L extract was then evaluated in terms of viscosity, pH, drying time, spreadability, and antibacterial activity. The selected formula was formula containing 5% of extract and 1% of carbomer 940. This formula had pH that suitable with skin pH 4.5–6.5, had good spreadability, and also produced highest antibacterial activity against *P. acnes*.

**Key words:** Antibacterial activity, carbomer 940, *Cocos nucifera* L. extract, *Propionibacterium acnes*

## INTRODUCTION

*Cocos nucifera* L is a multipurpose plant that has main component is water and soft flesh which has not been much researched as drug or cosmetic. Kopyor coconut is one type of it. To provide great benefits in developing potential as drug and cosmetic, kopyor need to be extracted and characterized. Muhammadiyah University Purwokerto Coconut Research Center Indonesia collaborated with the Faculty of Pharmacy Airlangga University to undergo research using kopyor

extract as active agent for cosmetic. This study investigated the potential of kopyor extract (*C. nucifera* L.) and tested its potential and activity as an antimicrobial and cosmetic.

## MATERIALS AND METHODS

### Materials

Materials used in the study had a degree of pharmaceutical-grade purity including, polyvinyl alcohol, Carbomer 940, Na-ethylenediaminetetraacetic acid, phenoxyethanol, propylene glycol, Tween 80, Span 80, TEA, BHT, NaCl. Kopyor coconut was obtained from Purwokerto Research Centre, and ATCC 11827 *Propionibacterium acnes* was obtained from the Surabaya Health Laboratory Center. Nutrient agar media was obtained from the Testing Service Unit (ULP) Faculty of Pharmacy Universitas Airlangga.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

**For reprints contact:** reprints@medknow.com

**How to cite this article:** Hariyadi DM, Isnaeni I, Sudarma S, Suciati S, Rosita N. Peel-off emulgel mask of *Cocos nucifera* L. Extract using gelling agent carbomer 940 as antiacne against *Propionibacterium acnes* ATCC 11827. *J Adv Pharm Technol Res* 2020;11:220-5.

### Address for correspondence:

Dr. Dewi Melani Hariyadi,  
Department of Pharmaceutics, Faculty of Pharmacy, Universitas  
Airlangga, Surabaya, Indonesia.  
E-mail: dewi-m-h@ff.unair.ac.id

Submitted: 24-Apr-2020

Revised: 21-Jul-2020

Accepted: 06-Aug-2020

Published: 10-Oct-2020

### Access this article online

#### Quick Response Code:



#### Website:

www.japtr.org

#### DOI:

10.4103/japtr.JAPTR\_51\_20



## Methods

### *Kopyor coconut extraction with maceration method*

A sample of 150 g was immersed in 96% ethanol with a ratio of 1: For 1 day and was repeated three times. Maserate was allowed to settle for sedimentation purposes. The oil sample was separated from the ethanol layer and the residue. Then, it was shaken out with a rotary evaporator to remove the remaining solvent. The yield was calculated, and phytochemical screening was analyzed.<sup>[1]</sup>

### Saponification test

#### *Volumetric method*

A total of 2.5 g of sample was weighed in a 250 mL erlenmyer. 25.0 mL KOH was added to ethanol 0.5 N then continued to reflux for 30 min while turning occasionally. Then, 1 mL PP indicator was added. The excess of KOH was titrated with 0.5 N HCl. The saponification value was calculated.

$$N \text{ HCl} = (1000 \times m \text{ KHP} \times \% \text{KHP} \times V_{t_2}) / (V_{t_1} \times Mr \text{ KHP} \times V_{\text{HCl}})$$

$$\text{Saponification value} = ([v \text{ blank titration} - v \text{-sample titration}] \times N \text{ HCl} \times Mr \text{ KOH}) / (m \text{ sample})$$

### Formulation of Peel-off Mask Emulgel

Polyvinyl alcohol and Carbomer 940 solution were prepared. Into Carbomer 940, TEA was added dropwise while stirring. Tween 80 was dissolved in water and then propylene glycol was added followed by addition of phenoxyethanol and Na-EDTA. Then the water phase was put into a gel base which has expanded and was stirred homogeneously. Kopyor (*Cocos nucifera* L.) extract was mixed with Span 80 and BHT, then PVA was added and the gel mixture was stirred until homogeneous. Formula can be seen in Table 1.

### Phytochemical screening analysis

#### *Physical evaluation of peel-off mask*

Evaluation of the peel-off mask of kopyor coconut extract was included the examination of the physical characteristics by determining pH, viscosity, dispersibility, particle size, particle size distribution, and zeta potential.<sup>[2]</sup>

#### *pH evaluation*

pH measurements used a pH meter.<sup>[3]</sup>

#### *Viscosity evaluation*

The viscosity was measured with a cone and plate DV-1 viscometer with a CP-41 type spindle. A sample of 2 g of peel-off mask was weighed. The sample must be ensured to be bubble-free and spread evenly on the surface of the cup. After that, the sample cup was reassembled on the viscometer. The viscometer was turned on at 4 rpm and then was left for a while until the reading was stable.<sup>[4]</sup>

#### *Spreadability evaluation*

One gram of peel-off mask was weighed and was placed in the middle of a glass plate and was covered with another glass plate. Additional weighting was carried out, starting from 5 g on the top glass plate. More weighting load was added out until the preparation showed no longer spreads. Deployment ability was measured by the slope of the distribution equation of the diameter of the dispersion versus weight load.<sup>[3,5]</sup>

### Drying time of peel-off mask

A total of 700 mg peel-off gel mask was applied to the glass slab. The plates were left at room temperature (25°C) and at the oven (37°C). Then the mask was left to dry by checking every 5 min.<sup>[6]</sup>

### Evaluation of antibacterial activity of peel-off mask

The *P. acnes* bacterial inoculum was prepared. Transmission of a bacterial inoculum was measured by a spectrophotometer at a wavelength of 580 nm until it received a 25% transmittance. The media base layer (30 mL) and seat layer (20 mL) were made with NA and then was sterilized at 121°C for 30 min. Base layer media was poured in a petri dish. It was then allowed to cool and solidify. After cooling and solidifying, the seat layer media was added to the 20 µL inoculum solution, followed by allowing cooling and solidifying. After cooling and solidifying, wells were made sterile to contain bacteria with a diameter of the center. A total well of 21 points consisted of 4 × 3 points for each sample (F1, F2, F3, and F4) including replication, 2 × 3 points for the positive control (contains 5% benzoyl peroxide concentration) and 1 × 3 points for negative control (dimethyl sulfoxide). Then, the petri dish was placed upside down and was incubated for 24 h at 37°C. After 24 h, the diameter of the zone was measured.

### Statistical analysis used

#### *Data analysis*

Measurement of a significant difference between the formula with different concentrations of extract and gel base Carbopol 940 with viscosity, dispersion, and inhibitory strength of kopyor extract preparations, was performed with statistical analysis with one-way variant analysis (ANOVA). Significant differences between formulas were identified by Tukey's honestly significant difference test with a 95% confidence level ( $\alpha = 0.05$ ). To find out the significant differences in the physical characteristics, namely viscosity before and after the thermal cycling stability test, particle size, and particle size distribution, a paired *t*-test was performed.

## RESULTS

Characterization of Kopyor (*C. nucifera* L.) extract was as follows:

**Table 1: Formula of peel-off mask**

Components	Concentration (%)					
	Formulas					
	1	2	3	4	5	6
Kopyor coconut extract	1	5	1	5	-	-
PVA	10	10	10	10	10	10
Carbopol 940	1	1	1.5	1.5	1	1.5
Propylenglycol	15	15	15	15	15	15
Na EDTA	0.1	0.1	0.1	0.1	0.1	0.1
BHT	0.1	0.1	0.1	0.1	0.1	0.1
Phenoxyethanol	1	1	1	1	1	1
Triethanolamine	0.5	0.5	0.5	0.5	0.5	0.5
Tween 80	4.5	4.5	4.5	4.5	4.5	4.5
Span 80	0.5	0.5	0.5	0.5	0.4	0.5
Aquadest	Ad 100	Ad 100	Ad 100	Ad 100	Ad 100	Ad 100

PVA: Polyvinyl alcohol, BHT: Butyl Hydroxy Toluene

**Table 2: Standardization of 0.5 N HCl**

m Na <sub>2</sub> CO <sub>3</sub> (99,9%); Mr= 106 (g)	Volume HCl (mL)	N HCl (mol/L)
264.8	6.30	0.7923
265.0	6.35	0.7866
Average of N HCl		0.7894

**Table 3: Standardization of 0.5 N KOH**

Volume KOH (mL)	Volume HCl (mL)	N HCl (mol/L)
25.0	15.80	0.4989
25.0	15.85	0.5005
Average N KOH		0.4997

$$\text{Saponification Index} = \frac{(v \text{ titrasi blanko} - v \text{ titrasi sampel}) \times 28.05}{m \text{ sampel}}$$

**Table 4: Saponification Index**

m sample (g)	Blank titration (mL)	Sample titration (mL)	Saponification value
1.2286	15.80	13.70	47.92
1.2916	15.85	13.65	47.75
Average			47.83
RPD			0.30

**Table 5: Minimum inhibitory concentration of extract**

Concentration (%)	Inhibition zone diameter (mm)±SD
30	7.16±0.05
20	6.58±0.11
10	6.40±0.21
5	5.90±0.42
2.5	5.80±0.12
1	5.50±0.23
0.5	0.0±0.0

SD: Standard deviation

**Phytochemical screening**

Alkaloid, saponin, and flavonoid test are shown in Figure 1.

**Saponification index**

Saponification index values results are as calculation in Tables 2-4.

Saponification index = 47.83

MIC of the extract against *Propionibacterium Acnes*

MIC of the extract is shown in Table 5.

**Antibacterial activity of extract**

Antibacterial activity of extract against *Propionibacterium Acnes* as shown in Table 6.

**pH test**

pH of the formula was demonstrated in Table 7.

**Viscosity test**

Viscosity of the formula was demonstrated in Table 8.

**Spreadability test**

Spradability of the formula was demonstrated in Table 9.

Slope of Spreadability is shown in Table 10.

**Drying time**

The drying time of the formula is shownin Table 11.

**Activity test of peel-off mask formula against propionibacterium acnes ATCC 11827**

Zone of Inhibition of formula is shown in Table 12.

**DISCUSSION**

Kopyor coconut has been investigated worldwide using some parts of plants have been used for the treatment of various pathological conditions such as antiinflammation, antioxidant, antihelminthic, antimalarial, and antiviral activity,<sup>[7-9]</sup> thus this current research studied antiacne

**Table 6: Antibacterial activity of extract against *Propionibacterium Acnes***

Sample	Concentration (%)	<i>Propionibacterium Acnes</i> inhibition zone diameter (mm)
Kopyor Extract	1	6.20
	2.5	6.50
	5	8.20
Control (+)	2.5	6.20
Control (-)		5.30
Neutrogena mask (contains benzoyl peroxide)	2.5	6.15

**Table 7: pH formula**

Formula	pH±SD
F1	4.89±0.02
F2	4.83±0.03
F3	5.45±0.02
F4	5.21±0.03
F5	5.79±0.07
F6	5.45±0.14

SD: Standard deviation

**Table 8: Viscosity of formula**

Formula	Viscosity (cps)±SD
F1	42,540±471.27
F2	46,320±559.42
F3	44,470±256.22
F4	48,789±925.84
F5	29,953±352.04
F6	33,190±115.30

formulation from the extract of *C. nucifera* L. from flesh part.

### Saponification index

Saponification index values results are shown as below calculation on Tables 2-4.

The saponification index value was 47.83 and consisted of lauric acid of 29.20%.


### Physical characteristics of peel-off mask formula

#### pH test

From these results, it showed that the pH of the peel-off mask of kopyor extract fulfilled the pH of the skin, namely pH 4.5–6.5 and was safe for use on the skin.<sup>[10]</sup> To ensure the difference in pH during the storage process, it was also advisable to look for free fatty acids that may be oxidized in the kopyor extract content that can allow an increase in free fatty acids when exposed to high temperatures.<sup>[11]</sup>

#### Viscosity test

The results of viscosity between formulas showed that an increase in the extract and gelling agent carbomer showed an increase in the viscosity of the preparation in line with similar studies.<sup>[12,13]</sup>



**Figure 1:** (1) Alkaloid test: Positive; (2) Saponin test: Negative; (3) Flavonoid test: Positive

#### Spreadability test

The spread diameter was shown between 4.37 and 7.70 for antiacne formulas contain extract. This was confirmed with other researchers for mask capacity of 5–7 cm.<sup>[14]</sup> Based on the value of the slope and the load to achieve maximum dispersal capacity, it can be concluded that the base formula without an active agent was more easily spread than the preparation formula. It was appropriate that an increase in the viscosity of the preparations will reduce the spreadability of the preparations.<sup>[13]</sup> Another explanation may be due to the effect of formula, which still contains a large amount of water content in each formula that can slow the evaporation and the formation of a film layer in a peel-off mask as similar to a study conducted by Phindo.<sup>[15]</sup>

#### Drying time

Based on the results of the drying time, the preparation of kopyor extract mask formulas 1–4 had a drying time of 40–50 min. The formula containing kopyor extract had a longer drying time compared to the two mask base formulas without active agent. This showed the drying time of the four formulas was longer than the peel-off mask drying time in general, which was <30 min.<sup>[16]</sup> This result was maybe due to kopyor extract in which there was oil content that can still resist the evaporation of water in the preparation therefore it took longer to dry.

#### Activity test of peel-off mask formula against *Propionibacterium acnes* ATCC 11827

Based on the results of the peel-off mask activity test, all formulations showed inhibition against *P. acnes* ATCC 11827. However, the comparison of inhibition diameter between formulas that had low *C. nucifera* L. extract concentrations,

**Table 9: Spreadability of formula**

Formula	Weight (g)	Spreadability diameter (cm)±SD
F1	-	4.37±0.15
	5	4.60±0.17
	10	4.83±0.15
	15	5.13±0.23
	20	5.23±0.23
F2	-	6.03±0.06
	5	6.27±0.21
	10	6.33±0.23
	15	6.47±0.21
	20	6.53±0.25
F3	-	4.63±0.15
	5	4.87±0.11
	10	5.07±0.11
	15	5.13±0.06
	20	5.20±0.10
F4	-	6.80±0.26
	5	7.23±0.29
	10	7.43±0.38
	15	7.67±0.58
	20	7.70±0.61
F5	-	3.83±0.05
	5	4.00±0.11
	10	4.10±0.11
	15	4.17±0.12
	20	4.20±0.10
F6	-	3.97±0.06
	5	4.20±0.17
	10	4.43±0.11
	15	4.57±0.15
	20	4.57±0.15

SD: Standard deviation

**Table 10: Spreadability slope of Formula**

Formula	F1	F2	F3	F4	F5	F6
Slope (cm <sup>2</sup> /g)	0.3406	0.2372	0.2168	0.5130	0.2102	0.2208

inhibitory activity was as large or not significantly different from positive control of benzoyl peroxide. The inhibitory activity against acne bacteria increased with increasing levels of kopyor extract. This can be influenced by the content of lauric acid, which was not maximum in the content of kopyor extract, and hence, it was necessary to optimize the extraction and selection of different types of kopyor coconut.<sup>[17]</sup>

## CONCLUSIONS

Kopyor (*C. nucifera* L.) extract had antibacterial activity against *P. acnes* ATCC 11827 at concentration of 1 and 5% and has been successfully formulated into peel-off mask formula using gelling agent carbomer 940 at a concentration of 1 and 1.5%. The variation of kopyor extract and gelling agent concentration affected the viscosity, pH, and spreadability.

**Table 11: Drying time of formula**

Formula	Drying time (min) Temperature (37°C)
F1	40:00
F2	40:05
F3	45:07
F4	50:03
F5	40:15
F6	50:00

**Table 12: Zone of inhibition of formula**

Sample	Inhibition zone diameter (mm)±SD
F1	10.05±0.13
F2	11.68±0.30
F3	9.90±0.05
F4	11.43±0.50
Control +	9.55±0.25
Control-	6.50±0.00

SD: Standard deviation

However, the antibacterial activity between formulas in masks with a low extract concentration of 1% did not show different results of its activity, therefore it was suggested to increase concentration of extract of higher than 5% or carbomer of more than 1.5%, it was predicted that there would be a significant increase in antibacterial activity.

Based on the physical evaluation and antibacterial activity of the mask preparations, a formula which has good characteristics was selected. The chosen formula was formula, which was a formula containing 5% extract and 1% carbomer produced formula with suitable skin pH, good dispersion, and sufficient viscosity. Moreover, it had a high antibacterial activity. Further recommendation of the peel-off mask formula included study the stability of peel-off mask for a longer period.

## Financial support and sponsorship

DRPM DIKTI for grant.

## Conflicts of interest

There are no conflicts of interest.

## REFERENCES

- Rao US, Abdurrazak M, Suryati MK. Phytochemical screening. Total flavonoid and phenolic content assays of various solvent extracts of tepal of *Musa paradisiaca*. *Malaysian J Analyt Sci* 2016;20:1181-90.
- Santoso I, Prayoga T, Agustina I, Rahayu W. Formulation of peel-off gel mask fruit of (*Aloe vera* l) with polyvinyl alcohol gelling agent. *J Riset Kefarmasian Indonesia* 2020;2:17-25.
- Zettira, Natalia Z. Antibactery Activity of Gel Rosella Extract (*Hibiscus sabdariffa* L.)-HPMC 6000 on *Staphylococcus aureus* ATCC 25923. Thesis. Faculty of Pharmacy, Universitas Airlangga; 2016.

4. Dantas MG, Reis SA, Damasceno CM, Rolim LA, Rolim-Neto PJ, Carvalho FO, *et al.* Development and evaluation of stability of a gel formulation containing the monoterpene borneol. *ScientificWorldJournal* 2016;2016:7394685.
5. Gayatri PA, Krisnawati K, Sahlan M, Pratami DK, Widayati R. Stability of zoledronate gel emulsion in virgin coconut oil. *Int J Applied Pharma* 2019;11:201-6.
6. Lestari PM, Sutyaningsih RB, Ruhimat. The Influence of Increase Concentration Poliviny Alcohol (PVA) as a Gelling Agent on Physical Properties of The Peef-off of Pineapple Juice (*Ananas comosus* L). *Asian Societies of Cosmetic Scientists Conference*; 2013.
7. Aggarwal B, Lamba HS, Sharma P, Ajeet PS. Various pharmacological aspects of *Cocos nucifera* – A review. *Am J Pharmacol Sci* 2017;5:25-30.
8. Lima EB, Sousa CN, Meneses LN, Ximenes NC, Santos Júnior MA, Vasconcelos GS, *et al.* *Cocos nucifera* (L.) (Arecaceae): A phytochemical and pharmacological review. *Braz J Med Biol Res* 2015;48:953-64.
9. Dwestiwati R, Sulistyowati E. The utilization of coconut husk extract (*Cocos nucifera* L.) As an antioxidant in the palm oil krengseng. *J Kimia Dasar* 2016;5:1-7.
10. Prakash C, Bhargava P, Tiwari S, Majumdar B, Bhargava RK. Skin surface pH in acne vulgaris: insights from an observational study and review of the literature. *J Clin Aesthet Dermatol* 2017;10:33-9.
11. Gharby S, Harhar H, Matthaus B, Bouzoubaa Z, Charrouf Z. The chemical parameters and oxidative resistance to heat treatment of refined and extra virgin Moroccan Picholine olive oil. *J Taibah Univer Sci* 2016;10:100-6.
12. Setyawati Y, Aladin A, Wiyani L, Modding B. Evaluation of Viscosity and pH on Emulsions of Virgin Coconut Oil Beverages. *IOP Conf. Series: Earth and Environmental Science*; 2018. p. 175.
13. Affianti HP, Murrukmihadi M. Effect of concentration of gelling agent HPMC on physical stability and antibacterial activity of gel ethanolic extract of Kemangi leaves. *Farmaseutik* 2015;11:307-15.
14. Garg A, Deepika A, Sanjay G, Anil KS. Spreading of semisolid formulations: An update. *Pharma Technol* 2002;26:84-105.
15. Phindo L. Formulation and Physical Evaluation of Mask Peel-Off of Ethanolic Extract of *Artocarpus Heterophyllus*. Lamk, Glycolic acid and Niacinamide. Thesis. UIN Syarif Hidayatullah. Jakarta; 2016.
16. Priani S, Irawati I, Gita CE. Formulation of mask gel peel-off skin of *Garcinia mangostana* Linn. *IJPST* 2015;2:90-5.
17. Novariantio H, Meity T. Lauric acid content on variety of coconut as materials of VCO. *Littri J* 2017;13:28-33.



# Journal of Advanced Pharmaceutical Technology and Research

<b>Country</b>	India -  SCIMAGO INSTITUTIONS RANKINGS
<b>Subject Area and Category</b>	Pharmacology, Toxicology and Pharmaceutics Pharmaceutical Science
<b>Publisher</b>	Wolters Kluwer Medknow Publications
<b>Publication type</b>	Journals
<b>ISSN</b>	01105558, 09762094
<b>Coverage</b>	2010-2020

# 29

H Index

**Scope** Journal of Advanced Pharmaceutical Technology & Research (JAPTR) is an Official Publication of Society of Pharmaceutical Education & Research™. It is an international journal published Quarterly. Journal of Advanced Pharmaceutical Technology & Research (JAPTR) is available in online and print version. It is a peer reviewed journal aiming to communicate high quality original research work, reviews, short communications, case report, Ethics Forum, Education Forum and Letter to editor that contribute significantly to further the scientific knowledge related to the field of Pharmacy i.e. Pharmaceutics, Pharmacology, Pharmacognosy, Pharmaceutical Chemistry. Articles with timely interest and newer research concepts will be given more preference.



[Homepage](#)

[How to publish in this journal](#)









[Contact](#)



[Join the conversation about this journal](#)

Quartiles





Journal of Advanced Pharmaceutical Technology...

Q2 Pharmaceutical Science  
best quartile

SJR 2019  
0.27

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```

Metrics based on Scopus® data as of April 2020

B **Beatriz G.** 7 months ago

Estimados, solicito de su ayuda como puedo saber cual es el factor de impacto de un articulo publicado en su revista ??

De antemano agradezco su respuesta

reply



**Melanie Ortiz** 7 months ago

SCImago Team

Dear Beatriz, thank you very much for your comment.

SCImago Journal and Country Rank uses Scopus data, our impact indicator is the SJR.

The next update will be made in June 2020. We suggest you consult the Journal Citation Report for other indicators (like Impact Factor) with a Web of Science data source. Best Regards, SCImago Team

E **eden fisha** 1 year ago

good

reply