



Journal of Advanced Pharmaceutical Technology & Research

An Official Publication of Society of Pharmaceutical Education & Research

Volume 11 / Issue 4 / October-December 2020

www.japtr.org

JAPTR



Editorial Board

Editor-in-Chief

Dr. Upendra Nagaich

M.Pharm., Ph.D., FSPER

Secretary,

[Society of Pharmaceutical Education & Research \(SPER\), India](#)

&

Research Coordinator,

Amity Institute of Pharmacy, Amity University,

Noida (U.P.) India

Executive Editors

Dr. Paul Wan Sia Heng

Department of Pharmacy,

National University of Singapore,

Singapore

Ms. Jaya Pandey

School of Studies in Pharmaceutical Sciences,

Jiwaji University,

Gwalior (M.P.) India

Associate Editor

Dr. Vandana Chaudhary

School of Pharmacy,

Pharmacy Australia Centre of Excellence,

The University of Queensland,

Brisbane, Australia

Consultant Editor

Prof. (Dr.) Laszlo Endrenyi

Professor Emeritus

Department of Pharmacology & Toxicology

University of Toronto, Toronto, Canada

Section Editors (Alphabetical Order)

Dr. Ikoni Ogaji

Dept. of Pharmaceutics & Pharmaceutical Technology,

Faculty of Pharmaceutical Sciences,

University of Jos, Nigeria

Dr. Ke Li

J. David Gladstone Institutes,

San Francisco, California, USA

Editorial Board Members (In Alphabetical Order)

Prof. (Dr.) A. N. Kalia

Vice-President, Society of Pharmacognosy,

Professor & Head

I.S.F. College of Pharmacy,

Moga (Punjab), India


Feedback

[Subscribe](#)

[About the journal](#)

[Most popular articles](#) 

[Join as Reviewer](#) 

[Email alerts](#) 

[Recommend this journal to your library](#) 

Dr. Abdul Wahab

Department of Pharmacy
Kohat University of Science & Technology
Kohat, KPK, Pakistan

Prof. (Dr.) Arun Dharmarajan

Professor
School of Biomedical Sciences,
Faculty of Health Sciences,
Curtin University,
Perth, Australia

Prof. (Dr.) Ashoke Kumar Ghosh

Vice-Chancellor
IFTM University,
Moradabad (Uttar Pradesh), India

Prof. (Dr.) B. Mishra

Professor & Head
Department of Pharmaceutics,
Indian Institute of Technology,
Banaras Hindu University,
Varanasi (Uttar Pradesh), India

Prof. B. G. Shivananda

Secretary, APTI
Bangalore (Karnataka), India

Prof. (Dr.) Bharat B. Aggarwal

Professor of Cancer Medicine &
Chief, Cytokine Research Laboratory
The University of Texas
Texas, USA

Dr. Bhaswat Sundar Chakraborty

Senior VP & Chairman, R&D Core Committee
Cadila Pharmaceuticals Ltd.
Ahmedabad (Gujarat), India

Prof. (Dr.) C. K. Kokate

National Advisor, Society of Pharmacognosy,
Vice-Chancellor
KLE University,
Belgaum (Karnataka), India

Prof. (Dr.) Cedric Pearce

Adjunct Professor
University of North Carolina
CEO, Mycosynthetix Inc.
North Carolina, USA
[Associate Editor, Journal of Natural Products]

Prof. (Dr.) Ciddi Veeresham

Professor
University College of Pharmaceutical Sciences,
Kakatiya University,
Warangal (Andhra Pradesh), India

Dr. Dâmaris Silveira

Universidade de Brasilia
Brasilia, Brazil

Prof. (Dr.) Dilipkumar Pal

Professor
Dept. of Pharmaceutical Sciences,
Guru Ghasidas Vishwavidyalaya,
Bilaspur (Chhattisgarh), India

Prof. (Dr.) Divakar Goli

Associate Secretary, IPA
Secretary, Education Division, IPA

Professor & Principal
Acharya & B.M. Reddy College of Pharmacy,
Bangalore (Karnataka), India

Prof. (Dr.) G. K. Dash

Professor
Universiti Kuala Lumpur
Perak Darul Ridzuan, Malaysia

Prof. (Dr.) H. S. Buttar

Adjunct Professor,
School of Medicine,
University of Ottawa,
Ottawa, Canada

Kaushik Desai

Gen. Secretary, IPA
Immediate Vice President, IPA
Mumbai (Maharashtra), India

Prof. (Dr.) Mainul Haque

Professor
Universiti Sultan Zainal Abidin
Kuala Terengganu, Terengganu, Malaysia

Dr. Muhammad Akhlaq

University of Auckland
Auckland, New Zealand

Prof. (Dr.) N. Udupa

Professor & Research Director [Health Sciences]
Manipal University,
Manipal (Karnataka), India

Prof. (Dr.) Pornsak Sriamornsak

Professor,
Department of Pharmaceutical Technology &,
Pharmaceutical Biopolymer Group (PBiG),
Faculty of Pharmacy, Silpakorn University,
Nakhon Pathom, Thailand

Dr. Rakesh Kumar Sharma

Scientist 'F'
Director,
Defence Food Research Laboratory (DFRL),
Mysuru (Karnataka), India

Prof. (Dr.) Roop K. Khar

Former Professor
Department of Pharmaceutics,
Jamia Hamdard (Hamdard University)
New Delhi, India

Prof. (Dr.) Roopa Karki

Professor & Head
Department of Pharmaceutics,
Acharya & BM Reddy College of Pharmacy,
Bangalore (Karnataka), India

Prof. (Dr.) S. D. Tonpay

Former Professor & Head
Govt. Gajra Raja Medical College,
Gwalior (Madhya Pradesh), India

Prof. (Dr.) S. H. Ansari

Professor & Head,
Department of Pharmacognosy & Phytochemistry,
Jamia Hamdard (Hamdard University)
New Delhi, India

Prof. (Dr.) S. H. Mishra

Professor Emeritus

Faculty of Pharmacy,
The M. S. University of Baroda,
Vadodara (Gujarat), India

Prof. (Dr.) Satish Sardana

Immediate Past President, APTI (Haryana)
Director,
Amity Institute of Pharmacy,
Amity University,
Manesar (Haryana), India

Prof. (Dr.) T. V. Narayana

President, IPA
Director
Vikas Institute of Pharmaceutical Sciences
Rajahmundry (Andhra Pradesh), India

Prof. (Dr.) V. K. Dixit

President, Society of Pharmacognosy,
Former Professor
Department of Pharmaceutical Sciences,
Dr. H. S. Gour University,
Sagar (M.P.), India

Prof. (Dr.) Y. K. Agrawal

Director
Institute of Research & Development,
Gujarat Forensic Sciences University,
Gandhinagar (Gujarat), India

Technical Board Members (In Alphabetical Order)

Dr. Deenanath Jhade, Bhopal
Dr. Inderbir Singh, Rajpura
Dr. Mohd. Rageeb Mohd. Usman, Chopda
Dr. Neeraj Sethiya, Vadodara
Dr. Praveen Sharma, Indore
Dr. Rakesh Pahwa, Kurukshetra
Dr. Sanjib Bhattacharya, Kolkata
Dr. Shweta Dixit, Pune
Dr. Sunder Singh, Indore



Table of Contents






October-December 2020
Volume 11 | Issue 4
Page Nos. 157-237

Online since Saturday, October 10, 2020

Accessed 15,690 times.

PDF access policy
Journal allows immediate open access to content in HTML + PDF

EPub access policy
allows EPub access only to users from **INDIA**, developing countries and paid subscribers.


-  [View issue as eBook](#)
-  [Issue citations](#)
-  [Issue statistics](#)
-  [RSS](#)

- Feedback**
- [Subscribe](#)
 - [Next Issue](#)
 - [Previous Issue](#)


- [Most popular articles](#) 
- [Join as Reviewer](#) 
- [Email alerts](#) 
- [Recommend this journal to your library](#) 

- [Show all abstracts](#)
- [Show selected abstracts](#)
- [Export selected to ▼](#)
- [Add to my list](#)


ORIGINAL ARTICLES




The coronavirus disease 2019 main protease inhibitor from *Andrographis paniculata* p. 157
(Burm.f) Ness POPULAR
Sukardiman , Martha Ervina, Mohammad Rizki Fadhil Pratama, Hadi Poerwono, Siswandono Siswodihardjo
DOI:10.4103/japtr.JAPTR_84_20
[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[Citations \(1\)\]](#) [\[Sword Plugin for Repository\]](#)^{Beta}




Anticancer evaluation of *N*-benzoyl-3-allylthiourea as potential antibreast cancer agent p. 163
through enhances HER-2 expression
Tri Widiandani, Siswandono , Edy Meiyanto
DOI:10.4103/japtr.JAPTR_77_20
[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[Sword Plugin for Repository\]](#)^{Beta}



Effects of lime (*Citrus aurantifolia*) peel to the expression of mRNA toll-like receptors 4 in p. 169
balb/c mice-infected *Salmonella typhi*
Vivien Novarina Kasim, Mochammad Hatta, Rosdiana Natzir, Veni Hadju, Ami Febriza, Hasta Handayani Idrus
DOI:10.4103/japtr.JAPTR_48_20
[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[Sword Plugin for Repository\]](#)^{Beta}



Anti-aging potential and phytochemicals of *Centella asiatica*, *Nelumbo nucifera*, and p. 174
Hibiscus sabdariffa extracts
Monsicha Khuaneckaphan, Chanai Noysang, Warachate Khobjai
DOI:10.4103/japtr.JAPTR_79_20
[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[Sword Plugin for Repository\]](#)^{Beta}



Proximate composition, vitamin and mineral composition, antioxidant capacity, and p. 179
anticancer activity of *Acanthopanax trifoliatum*
Aunyachulee Ganogpichayagrai, Chutinan Suksaard
DOI:10.4103/japtr.JAPTR_61_20
[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[Sword Plugin for Repository\]](#)^{Beta}



Stabilization of eye drops containing autologous serum and recombinant human p. 184
epidermal growth factor for dry eye syndrome
Sriwidodo Sriwidodo, Insan Sunan Kurniawan Syah, Iman Permana Maksum, Toto Subroto, Ulvi Zasvia, Abd Kakhar Umar
DOI:10.4103/japtr.JAPTR_72_20
[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[Sword Plugin for Repository\]](#)^{Beta}


Antidiabetic activity of thin film containing astaxanthin-loaded nanoemulsion using p. 189

carboxymethylcellulose sodium polymer on alloxan-induced diabetic rabbit

Lusi Nurdianti, Taofik Rusdiana, Iyan Sopyan

DOI:10.4103/japtr.JAPTR_55_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}



In silico docking and comparative ADMET profile of different glycogen synthase kinase 3 beta inhibitors as the potential leads for the development of anti-Alzheimer drug therapy p. 194

Manisha Prajapat, Phulen Sarma, Nishant Shekhar, Hardeep Kaur, Sanjay Singh, Subodh Kumar, Harpinder

Kaur, Saniya Mahendiratta, Amit Raj Sharma, Sukhmandeep Kaur, Vidya M Mahalmani, Bikash Medhi

DOI:10.4103/japtr.JAPTR_178_19

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}




Effect of implantation Ti-6Al-4V ELI in femoral bone defect regeneration of Sprague Dawley rat p. 202

I Nyoman Jujur, Damisih, Kartika Devy, Shovy Suha, Boy Muchlis Bachtiar, Endang Winiati Bachtiar

DOI:10.4103/japtr.JAPTR_74_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}




Development and validation of an ultra-performance liquid chromatography mass spectrometry/mass spectrometry method for simultaneous quantification of total and free mycophenolic acid and its metabolites in human plasma p. 207

Napatsanan Tanathitiphuwarat, Pannee Tanudechpong, Pajaree Chariyavilaskul, Nantaporn Prompila, Nat

Tansrisawad, Apinya Tubtimrattana, Supeecha Wittayalerpanya

DOI:10.4103/japtr.JAPTR_40_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}




The enhancement apoptosis of osteosarcoma mesenchymal stem cells co-cultivation with peripheral blood mononuclear cells sensitized by secretome and granulocyte macrophage colony-stimulating factor p. 213

Ferdiansyah Mahyudin, Hizbillah Yazid, Mouli Edward, Mohammad Hardian Basuki, Yunus Abdul Bari, Fedik

Abdul Rantam

DOI:10.4103/japtr.JAPTR_52_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}




Peel-off emulgel mask of *Cocos nucifera* L. Extract using gelling agent carbomer 940 as antiacne against *Propionibacterium acnes* ATCC 11827 p. 220

Dewi Melani Hariyadi, Isnaeni Isnaeni, Sisunandar Sudarma, Suciati Suciati, Noorma Rosita

DOI:10.4103/japtr.JAPTR_51_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}



Accelerated wound healing ability of *Jatropha sap* by iota carrageenan-poly (vinyl alcohol) hydrogel film p. 226

Nurain Thomas, Agus Rusdin, Madania Tulsyahra, Nasrul Wathoni, Asep Kuswandi

DOI:10.4103/japtr.JAPTR_11_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}




Induction of cytotoxicity by *Bruguiera gymnorhiza* in human breast carcinoma (MCF-7) cell line via activation of the intrinsic pathway p. 233

Gul-e-Saba Chaudhry, Nurul Huda Rahman, Vigneswari Sevakumaran, Aziz Ahmad, Habsah Mohamad,

Muhammad Naveed Zafar, Yeong Yik Sung, Tengku Sifzizul Tengku Muhammad

DOI:10.4103/japtr.JAPTR_81_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}



Peel-off emulgel mask of *Cocos nucifera* L. Extract using gelling agent carbomer 940 as antiacne against *Propionibacterium acnes* ATCC 11827

Dewi Melani Hariyadi,
Isnaeni Isnaeni¹,
Sisunandar Sudarma²,
Suciati Suciati³, Noorma Rosita

Departments of Pharmaceutics,
¹Pharmaceutical Chemistry and
³Pharmacognosy and Phytochemsitry,
Faculty of Pharmacy, Universitas
Airlangga, Surabaya, ²Department of
Biology, Universitas Muhammadiyah
Purwokerto, Purwokerto, Indonesia

J. Adv. Pharm. Technol. Res.

ABSTRACT

Cocos nucifera Linn., which contain lauric acid has been known had antibacterial activity against *Propionibacterium acnes* that usually improve severe of pimple. Current study investigated optimum formula of emulgel mask based on the *C. nucifera* L. Extract from Kopyor coconut. Extract were tested for antibacterial against *P. acnes* ATCC 11827. In this research, *C. nucifera* L. extract of 1 and 5% were formulated as an active agent of peel off antiacne emulgel mask-containing carbomer 940 in various concentration (1% and 1.5%). The peel-off emulgel mask of *C. nucifera* L extract was then evaluated in terms of viscosity, pH, drying time, spreadability, and antibacterial activity. The selected formula was formula containing 5% of extract and 1% of carbomer 940. This formula had pH that suitable with skin pH 4.5–6.5, had good spreadability, and also produced highest antibacterial activity against *P. acnes*.

Key words: Antibacterial activity, carbomer 940, *Cocos nucifera* L. extract, *Propionibacterium acnes*

INTRODUCTION

Cocos nucifera L is a multipurpose plant that has main component is water and soft flesh which has not been much researched as drug or cosmetic. Kopyor coconut is one type of it. To provide great benefits in developing potential as drug and cosmetic, kopyor need to be extracted and characterized. Muhammadiyah University Purwokerto Coconut Research Center Indonesia collaborated with the Faculty of Pharmacy Airlangga University to undergo research using kopyor

extract as active agent for cosmetic. This study investigated the potential of kopyor extract (*C. nucifera* L.) and tested its potential and activity as an antimicrobial and cosmetic.

MATERIALS AND METHODS

Materials

Materials used in the study had a degree of pharmaceutical-grade purity including, polyvinyl alcohol, Carbomer 940, Na-ethylenediaminetetraacetic acid, phenoxyethanol, propylene glycol, Tween 80, Span 80, TEA, BHT, NaCl. Kopyor coconut was obtained from Purwokerto Research Centre, and ATCC 11827 *Propionibacterium acnes* was obtained from the Surabaya Health Laboratory Center. Nutrient agar media was obtained from the Testing Service Unit (ULP) Faculty of Pharmacy Universitas Airlangga.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Hariyadi DM, Isnaeni I, Sudarma S, Suciati S, Rosita N. Peel-off emulgel mask of *Cocos nucifera* L. Extract using gelling agent carbomer 940 as antiacne against *Propionibacterium acnes* ATCC 11827. *J Adv Pharm Technol Res* 2020;11:220-5.

Address for correspondence:

Dr. Dewi Melani Hariyadi,
Department of Pharmaceutics, Faculty of Pharmacy, Universitas
Airlangga, Surabaya, Indonesia.
E-mail: dewi-m-h@ff.unair.ac.id

Submitted: 24-Apr-2020

Revised: 21-Jul-2020

Accepted: 06-Aug-2020

Published: 10-Oct-2020

Access this article online

Quick Response Code:



Website:

www.japtr.org

DOI:

10.4103/japtr.JAPTR_51_20

Methods

Kopyor coconut extraction with maceration method

A sample of 150 g was immersed in 96% ethanol with a ratio of 1: For 1 day and was repeated three times. Maserate was allowed to settle for sedimentation purposes. The oil sample was separated from the ethanol layer and the residue. Then, it was shaken out with a rotary evaporator to remove the remaining solvent. The yield was calculated, and phytochemical screening was analyzed.^[1]

Saponification test

Volumetric method

A total of 2.5 g of sample was weighed in a 250 mL erlenmyer. 25.0 mL KOH was added to ethanol 0.5 N then continued to reflux for 30 min while turning occasionally. Then, 1 mL PP indicator was added. The excess of KOH was titrated with 0.5 N HCl. The saponification value was calculated.

$$N \text{ HCl} = (1000 \times m \text{ KHP} \times \% \text{KHP} \times V_{t_2}) / (V_{t_1} \times Mr \text{ KHP} \times V_{\text{HCl}})$$

$$\text{Saponification value} = ([v \text{ blank titration} - v \text{-sample titration}] \times N \text{ HCl} \times Mr \text{ KOH}) / (m \text{ sample})$$

Formulation of Peel-off Mask Emulgel

Polyvinyl alcohol and Carbomer 940 solution were prepared. Into Carbomer 940, TEA was added dropwise while stirring. Tween 80 was dissolved in water and then propylene glycol was added followed by addition of phenoxyethanol and Na-EDTA. Then the water phase was put into a gel base which has expanded and was stirred homogeneously. Kopyor (*Cocos nucifera* L.) extract was mixed with Span 80 and BHT, then PVA was added and the gel mixture was stirred until homogeneous. Formula can be seen in Table 1.

Phytochemical screening analysis

Physical evaluation of peel-off mask

Evaluation of the peel-off mask of kopyor coconut extract was included the examination of the physical characteristics by determining pH, viscosity, dispersibility, particle size, particle size distribution, and zeta potential.^[2]

pH evaluation

pH measurements used a pH meter.^[3]

Viscosity evaluation

The viscosity was measured with a cone and plate DV-1 viscometer with a CP-41 type spindle. A sample of 2 g of peel-off mask was weighed. The sample must be ensured to be bubble-free and spread evenly on the surface of the cup. After that, the sample cup was reassembled on the viscometer. The viscometer was turned on at 4 rpm and then was left for a while until the reading was stable.^[4]

Spreadability evaluation

One gram of peel-off mask was weighed and was placed in the middle of a glass plate and was covered with another glass plate. Additional weighting was carried out, starting from 5 g on the top glass plate. More weighting load was added out until the preparation showed no longer spreads. Deployment ability was measured by the slope of the distribution equation of the diameter of the dispersion versus weight load.^[3,5]

Drying time of peel-off mask

A total of 700 mg peel-off gel mask was applied to the glass slab. The plates were left at room temperature (25°C) and at the oven (37°C). Then the mask was left to dry by checking every 5 min.^[6]

Evaluation of antibacterial activity of peel-off mask

The *P. acnes* bacterial inoculum was prepared. Transmission of a bacterial inoculum was measured by a spectrophotometer at a wavelength of 580 nm until it received a 25% transmittance. The media base layer (30 mL) and seat layer (20 mL) were made with NA and then was sterilized at 121°C for 30 min. Base layer media was poured in a petri dish. It was then allowed to cool and solidify. After cooling and solidifying, the seat layer media was added to the 20 µL inoculum solution, followed by allowing cooling and solidifying. After cooling and solidifying, wells were made sterile to contain bacteria with a diameter of the center. A total well of 21 points consisted of 4 × 3 points for each sample (F1, F2, F3, and F4) including replication, 2 × 3 points for the positive control (contains 5% benzoyl peroxide concentration) and 1 × 3 points for negative control (dimethyl sulfoxide). Then, the petri dish was placed upside down and was incubated for 24 h at 37°C. After 24 h, the diameter of the zone was measured.

Statistical analysis used

Data analysis

Measurement of a significant difference between the formula with different concentrations of extract and gel base Carbopol 940 with viscosity, dispersion, and inhibitory strength of kopyor extract preparations, was performed with statistical analysis with one-way variant analysis (ANOVA). Significant differences between formulas were identified by Tukey's honestly significant difference test with a 95% confidence level ($\alpha = 0.05$). To find out the significant differences in the physical characteristics, namely viscosity before and after the thermal cycling stability test, particle size, and particle size distribution, a paired *t*-test was performed.

RESULTS

Characterization of Kopyor (*C. nucifera* L.) extract was as follows:

Table 1: Formula of peel-off mask

Components	Concentration (%)					
	Formulas					
	1	2	3	4	5	6
Kopyor coconut extract	1	5	1	5	-	-
PVA	10	10	10	10	10	10
Carbopol 940	1	1	1.5	1.5	1	1.5
Propylenglycol	15	15	15	15	15	15
Na EDTA	0.1	0.1	0.1	0.1	0.1	0.1
BHT	0.1	0.1	0.1	0.1	0.1	0.1
Phenoxyethanol	1	1	1	1	1	1
Triethanolamine	0.5	0.5	0.5	0.5	0.5	0.5
Tween 80	4.5	4.5	4.5	4.5	4.5	4.5
Span 80	0.5	0.5	0.5	0.5	0.4	0.5
Aquadest	Ad 100	Ad 100	Ad 100	Ad 100	Ad 100	Ad 100

PVA: Polyvinyl alcohol, BHT: Butyl Hydroxy Toluene

Table 2: Standardization of 0.5 N HCl

m Na ₂ CO ₃ (99,9%); Mr= 106 (g)	Volume HCl (mL)	N HCl (mol/L)
264.8	6.30	0.7923
265.0	6.35	0.7866
Average of N HCl		0.7894

Table 3: Standardization of 0.5 N KOH

Volume KOH (mL)	Volume HCl (mL)	N HCl (mol/L)
25.0	15.80	0.4989
25.0	15.85	0.5005
Average N KOH		0.4997

$$\text{Saponification Index} = \frac{(v \text{ titrasi blanko} - v \text{ titrasi sampel}) \times 28.05}{m \text{ sampel}}$$

Table 4: Saponification Index

m sample (g)	Blank titration (mL)	Sample titration (mL)	Saponification value
1.2286	15.80	13.70	47.92
1.2916	15.85	13.65	47.75
Average			47.83
RPD			0.30

Table 5: Minimum inhibitory concentration of extract

Concentration (%)	Inhibition zone diameter (mm)±SD
30	7.16±0.05
20	6.58±0.11
10	6.40±0.21
5	5.90±0.42
2.5	5.80±0.12
1	5.50±0.23
0.5	0.0±0.0

SD: Standard deviation

Phytochemical screening

Alkaloid, saponin, and flavonoid test are shown in Figure 1.

Saponification index

Saponification index values results are as calculation in Tables 2-4.

Saponification index = 47.83

MIC of the extract against *Propionibacterium Acnes*

MIC of the extract is shown in Table 5.

Antibacterial activity of extract

Antibacterial activity of extract against *Propionibacterium Acnes* as shown in Table 6.

pH test

pH of the formula was demonstrated in Table 7.

Viscosity test

Viscosity of the formula was demonstrated in Table 8.

Spreadability test

Spradability of the formula was demonstrated in Table 9.

Slope of Spreadability is shown in Table 10.

Drying time

The drying time of the formula is shown in Table 11.

Activity test of peel-off mask formula against propionibacterium acnes ATCC 11827

Zone of Inhibition of formula is shown in Table 12.

DISCUSSION

Kopyor coconut has been investigated worldwide using some parts of plants have been used for the treatment of various pathological conditions such as antiinflammation, antioxidant, antihelminthic, antimalarial, and antiviral activity,^[7-9] thus this current research studied antiacne

Table 6: Antibacterial activity of extract against *Propionibacterium Acnes*

Sample	Concentration (%)	<i>Propionibacterium Acnes</i> inhibition zone diameter (mm)
Kopyor Extract	1	6.20
	2.5	6.50
	5	8.20
Control (+)	2.5	6.20
Control (-)		5.30
Neutrogena mask (contains benzoyl peroxide)	2.5	6.15

Table 7: pH formula

Formula	pH±SD
F1	4.89±0.02
F2	4.83±0.03
F3	5.45±0.02
F4	5.21±0.03
F5	5.79±0.07
F6	5.45±0.14

SD: Standard deviation

Table 8: Viscosity of formula

Formula	Viscosity (cps)±SD
F1	42,540±471.27
F2	46,320±559.42
F3	44,470±256.22
F4	48,789±925.84
F5	29,953±352.04
F6	33,190±115.30

formulation from the extract of *C. nucifera* L. from flesh part.

Saponification index

Saponification index values results are shown as below calculation on Tables 2-4.

The saponification index value was 47.83 and consisted of lauric acid of 29.20%.

Physical characteristics of peel-off mask formula

pH test

From these results, it showed that the pH of the peel-off mask of kopyor extract fulfilled the pH of the skin, namely pH 4.5–6.5 and was safe for use on the skin.^[10] To ensure the difference in pH during the storage process, it was also advisable to look for free fatty acids that may be oxidized in the kopyor extract content that can allow an increase in free fatty acids when exposed to high temperatures.^[11]

Viscosity test

The results of viscosity between formulas showed that an increase in the extract and gelling agent carbomer showed an increase in the viscosity of the preparation in line with similar studies.^[12,13]




Figure 1: (1) Alkaloid test: Positive; (2) Saponin test: Negative; (3) Flavonoid test: Positive

Spreadability test

The spread diameter was shown between 4.37 and 7.70 for antiacne formulas contain extract. This was confirmed with other researchers for mask capacity of 5–7 cm.^[14] Based on the value of the slope and the load to achieve maximum dispersal capacity, it can be concluded that the base formula without an active agent was more easily spread than the preparation formula. It was appropriate that an increase in the viscosity of the preparations will reduce the spreadability of the preparations.^[13] Another explanation may be due to the effect of formula, which still contains a large amount of water content in each formula that can slow the evaporation and the formation of a film layer in a peel-off mask as similar to a study conducted by Phindo.^[15]

Drying time

Based on the results of the drying time, the preparation of kopyor extract mask formulas 1–4 had a drying time of 40–50 min. The formula containing kopyor extract had a longer drying time compared to the two mask base formulas without active agent. This showed the drying time of the four formulas was longer than the peel-off mask drying time in general, which was <30 min.^[16] This result was maybe due to kopyor extract in which there was oil content that can still resist the evaporation of water in the preparation therefore it took longer to dry.

Activity test of peel-off mask formula against *Propionibacterium acnes* ATCC 11827

Based on the results of the peel-off mask activity test, all formulations showed inhibition against *P. acnes* ATCC 11827. However, the comparison of inhibition diameter between formulas that had low *C. nucifera* L. extract concentrations,

Table 9: Spreadability of formula

Formula	Weight (g)	Spreadability diameter (cm)±SD
F1	-	4.37±0.15
	5	4.60±0.17
	10	4.83±0.15
	15	5.13±0.23
	20	5.23±0.23
F2	-	6.03±0.06
	5	6.27±0.21
	10	6.33±0.23
	15	6.47±0.21
	20	6.53±0.25
F3	-	4.63±0.15
	5	4.87±0.11
	10	5.07±0.11
	15	5.13±0.06
	20	5.20±0.10
F4	-	6.80±0.26
	5	7.23±0.29
	10	7.43±0.38
	15	7.67±0.58
	20	7.70±0.61
F5	-	3.83±0.05
	5	4.00±0.11
	10	4.10±0.11
	15	4.17±0.12
	20	4.20±0.10
F6	-	3.97±0.06
	5	4.20±0.17
	10	4.43±0.11
	15	4.57±0.15
	20	4.57±0.15

SD: Standard deviation

Table 10: Spreadability slope of Formula

Formula	F1	F2	F3	F4	F5	F6
Slope (cm ² /g)	0.3406	0.2372	0.2168	0.5130	0.2102	0.2208

inhibitory activity was as large or not significantly different from positive control of benzoyl peroxide. The inhibitory activity against acne bacteria increased with increasing levels of kopyor extract. This can be influenced by the content of lauric acid, which was not maximum in the content of kopyor extract, and hence, it was necessary to optimize the extraction and selection of different types of kopyor coconut.^[17]

CONCLUSIONS

Kopyor (*C. nucifera* L.) extract had antibacterial activity against *P. acnes* ATCC 11827 at concentration of 1 and 5% and has been successfully formulated into peel-off mask formula using gelling agent carbomer 940 at a concentration of 1 and 1.5%. The variation of kopyor extract and gelling agent concentration affected the viscosity, pH, and spreadability.

Table 11: Drying time of formula

Formula	Drying time (min) Temperature (37°C)
F1	40:00
F2	40:05
F3	45:07
F4	50:03
F5	40:15
F6	50:00

Table 12: Zone of inhibition of formula

Sample	Inhibition zone diameter (mm)±SD
F1	10.05±0.13
F2	11.68±0.30
F3	9.90±0.05
F4	11.43±0.50
Control +	9.55±0.25
Control-	6.50±0.00

SD: Standard deviation

However, the antibacterial activity between formulas in masks with a low extract concentration of 1% did not show different results of its activity, therefore it was suggested to increase concentration of extract of higher than 5% or carbomer of more than 1.5%, it was predicted that there would be a significant increase in antibacterial activity.

Based on the physical evaluation and antibacterial activity of the mask preparations, a formula which has good characteristics was selected. The chosen formula was formula, which was a formula containing 5% extract and 1% carbomer produced formula with suitable skin pH, good dispersion, and sufficient viscosity. Moreover, it had a high antibacterial activity. Further recommendation of the peel-off mask formula included study the stability of peel-off mask for a longer period.

Financial support and sponsorship

DRPM DIKTI for grant.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Rao US, Abdurrazak M, Suryati MK. Phytochemical screening. Total flavonoid and phenolic content assays of various solvent extracts of tepal of *Musa paradisiaca*. *Malaysian J Analyt Sci* 2016;20:1181-90.
- Santoso I, Prayoga T, Agustina I, Rahayu W. Formulation of peel-off gel mask fruit of (*Aloe vera* l) with polyvinyl alcohol gelling agent. *J Riset Kefarmasian Indonesia* 2020;2:17-25.
- Zettira, Natalia Z. Antibactery Activity of Gel Rosella Extract (*Hibiscus sabdariffa* L.)-HPMC 6000 on *Staphylococcus aureus* ATCC 25923. Thesis. Faculty of Pharmacy, Universitas Airlangga; 2016.

4. Dantas MG, Reis SA, Damasceno CM, Rolim LA, Rolim-Neto PJ, Carvalho FO, *et al.* Development and evaluation of stability of a gel formulation containing the monoterpene borneol. *ScientificWorldJournal* 2016;2016:7394685.
5. Gayatri PA, Krisnawati K, Sahlan M, Pratami DK, Widayati R. Stability of zoledronate gel emulsion in virgin coconut oil. *Int J Applied Pharma* 2019;11:201-6.
6. Lestari PM, Sutyaningsih RB, Ruhimat. The Influence of Increase Concentration Poliviny Alcohol (PVA) as a Gelling Agent on Physical Properties of The Peef-off of Pineapple Juice (*Ananas comosus* L). *Asian Societies of Cosmetic Scientists Conference*; 2013.
7. Aggarwal B, Lamba HS, Sharma P, Ajeet PS. Various pharmacological aspects of *Cocos nucifera* – A review. *Am J Pharmacol Sci* 2017;5:25-30.
8. Lima EB, Sousa CN, Meneses LN, Ximenes NC, Santos Júnior MA, Vasconcelos GS, *et al.* *Cocos nucifera* (L.) (Arecaceae): A phytochemical and pharmacological review. *Braz J Med Biol Res* 2015;48:953-64.
9. Dwestiwati R, Sulistyowati E. The utilization of coconut husk extract (*Cocos nucifera* L.) As an antioxidant in the palm oil krengseng. *J Kimia Dasar* 2016;5:1-7.
10. Prakash C, Bhargava P, Tiwari S, Majumdar B, Bhargava RK. Skin surface pH in acne vulgaris: insights from an observational study and review of the literature. *J Clin Aesthet Dermatol* 2017;10:33-9.
11. Gharby S, Harhar H, Matthaus B, Bouzoubaa Z, Charrouf Z. The chemical parameters and oxidative resistance to heat treatment of refined and extra virgin Moroccan Picholine olive oil. *J Taibah Univer Sci* 2016;10:100-6.
12. Setyawati Y, Aladin A, Wiyani L, Modding B. Evaluation of Viscosity and pH on Emulsions of Virgin Coconut Oil Beverages. *IOP Conf. Series: Earth and Environmental Science*; 2018. p. 175.
13. Affianti HP, Murrukmihadi M. Effect of concentration of gelling agent HPMC on physical stability and antibacterial activity of gel ethanolic extract of Kemangi leaves. *Farmaseutik* 2015;11:307-15.
14. Garg A, Deepika A, Sanjay G, Anil KS. Spreading of semisolid formulations: An update. *Pharma Technol* 2002;26:84-105.
15. Phindo L. Formulation and Physical Evaluation of Mask Peel-Off of Ethanolic Extract of *Artocarpus Heterophyllus*. Lamk, Glycolic acid and Niacinamide. Thesis. UIN Syarif Hidayatullah. Jakarta; 2016.
16. Priani S, Irawati I, Gita CE. Formulation of mask gel peel-off skin of *Garcinia mangostana* Linn. *IJPST* 2015;2:90-5.
17. Novariantio H, Meity T. Lauric acid content on variety of coconut as materials of VCO. *Littri J* 2017;13:28-33.



Oxygen Transport Whitepaper

Download the whitepaper to learn more

GE Healthcare

DOWNLOAD

Journal of Advanced Pharmaceutical Technology and Research

COUNTRY

India



Universities and research institutions in India

SUBJECT AREA AND CATEGORY

Pharmacology, Toxicology and
Pharmaceutics
Pharmaceutical Science

PUBLISHER

Wolters Kluwer Medknow
Publications

H-INDEX

29

PUBLICATION TYPE

Journals

ISSN

01105558, 09762094

COVERAGE

2010-2020

INFORMATION

[Homepage](#)

[How to publish in this journal](#)

editor@japtr.org

SCOPE

Journal of Advanced Pharmaceutical Technology & Research (JAPTR) is an Official Publication of Society of Pharmaceutical Education & Research™. It is an international journal published Quarterly. Journal of Advanced Pharmaceutical Technology & Research (JAPTR) is available in online and print version. It is a peer reviewed journal aiming to communicate high quality original research work, reviews, short communications, case report, Ethics Forum, Education Forum and Letter to editor that contribute significantly to further the scientific knowledge related to the field of Pharmacy i.e. Pharmaceutics, Pharmacology, Pharmacognosy, Pharmaceutical Chemistry. Articles with timely interest and newer research concepts will be given more preference.



Join the conversation about this journal

Trade seperti biasa



Tambahkan hadiah ke dalam keuntungan Anda

OctaFX

Quartiles

Roulette Terbaru 2021

Sebuah permainan kecil yang bisa dimainkan kapan saja, dimana saja datang dan bersantai

777slot.io

B

Mitochondria-Targeted

The only industry-led meeting focused on end-to-end mitochondria-targeting.

mitochondria-targets.com

OPEN

FIND SIMILAR JOURNALS ?

1
**Brazilian Journal of
Pharmaceutical Sciences**
BRA

41%
similarity

2
**Saudi Pharmaceutical
Journal**
SAU







40%
similarity

3
**DARU, Journal of
Pharmaceutical Sciences**
GBR

39%
similarity

4
**Asian Journal of
Pharmaceutics**
IND

39%
similarity



Journal of Advanced Pharmaceutical Technology...

Pharmaceutical Science
best quartile

Q2

SJR 2019
0.27

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```

Trade seperti biasa

Tambahkan hadiah ke dalam keuntungan Anda

OctaFX

BUKA

Metrics based on Scopus® data as of April 2020

B **Beatriz G.** 9 months ago

Estimados, solicito de su ayuda como puedo saber cual es el factor de impacto de un articulo publicado en su revista ??

De antemano agradezco su respuesta

reply



Melanie Ortiz 9 months ago

SCImago Team

Dear Beatriz, thank you very much for your comment.

SCImago Journal and Country Rank uses Scopus data, our impact indicator is the SJR. The next update will be made in June 2020. We suggest you consult the Journal Citation Report for other indicators (like Impact Factor) with a Web of Science data source. Best Regards, SCImago Team

E **eden fisha** 1 year ago

good

reply



Melanie Ortiz 1 year ago

SCImago Team

Dear Eden, thanks for your participation! Best Regards, SCImago Team

A **Amina** 1 year ago

Dear sir

i would like to ask a question about this journal

why the ISSN numbers of this journal are different in the website than those in SJR website

in the journal website:

ISSN: Print -2231-4040, Online - 0976-2094

in SJR website:

ISSN: print 0110-5558 , online 0976-2094

please tell me your explanation please as soon as possible.

Regards..

reply



Melanie Ortiz 1 year ago

SCImago Team

Dear Amina,

SJR is a portal with scientometric indicators of journals indexed in Scopus. SJR has no authority over the data of the journals; they are the ones that Scopus sends us. The metadata are property of Scopus. SCImago has a signed agreement that limits our performance to the generation of scientometric indicators derived from the metadata sent in the last update. We suggest that you contact Scopus through this link to report the inconsistencies or modifications that you want to make in SCImago:

https://service.elsevier.com/app/answers/detail/a_id/14883/kw/scimago/supporthub/scopus/.

Best Regards, SCImago Team

A **Amina** 1 year ago

Hi

Can you send me the website of this journal as soon as possible plz.

Regards

reply



Melanie Ortiz 1 year ago

SCImago Team

Dear Amina, you can see the updated information just above. Best Regards, SCImago Team

M **Mohamed** 1 year ago

<http://www.japtr.org/>



Melanie Ortiz 1 year ago

SCImago Team

Dear Mohamed, thanks for your participation! Best Regards, SCImago Team

Leave a comment

Name

Email

(will not be published)

I'm not a robot reCAPTCHA
Privacy - Terms


Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2020. Data Source: Scopus®

EST MODUS IN REBUS

Horatio (Satire 1, 1, 106)