

DAFTAR PUSTAKA

- Abdel-Mottaleb M. M. A., Moulari B., Beduneau A., Pellequer Y., Lamprecht A., 2012. Nanoparticles enhance therapeutic outcome in inflamed skin therapy. *European Journal of Pharmaceutics and Biopharmaceutics* 82, p. 151–157
- Anonim, 2005. Showing Metabocard for 4-Methoxycinnamic acid. www.sigma-aldrich.com, diakses pada tanggal 10 November 2013
- Anonim, 2013. www.chemicaland21.com, diakses pada tanggal 10 November 2013
- De Vry, C.G., Valdez, M., Lazarov, M., Muhr, E., Buelow, R., Fong, T., Suhasini, I., 2005. Topical application of a novel immunomodulatory peptide, RDP58, reduces skin inflammation in the phorbol ester-induced dermatitis model. *J. Invest. Dermatol.* 125, p. 473–481.
- Dubey, A., Prabhu, P., Kamath, J.V., 2012. Nano Structured Lipid Carriers : A Novel Topical Drug Delivery System. *International Journal of Pharmatech Research* Vol. 4, No. 2, p. 705-714.
- Gard, Paul., 2001. *Human Pharmacology*, Chapter IX., 135. Taylor & Francis., London, New York.
- Garud, A., Singh D., Garud N., 2012. Solid Lipid Nanoparticles (SLN): Method, Characterization and Applications. *International Current Pharmaceutical Journal* 1(11), p. 384-393
- Hasanah, N.A., Nazaruddin F., Febrina, E., Zuhrotun, A., (2011). Analisis Kandungan Minyak Atsiri dan Uji Aktivitas Antiinflamasi Ekstrak Rimpang Kencur (*Kaempferia galanga* L.). *Jurnal Matematika & Sains*, Vol. 16 No. 3, p. 147-151
- Hou, DongZhi, ChangSheng, Xie., Kaijin, Huang, ChangHong, Zhu., 2003. The Production and Characteristics of Solid Lipid Nanoparticles (SLNs). *Biomaterials* Vol. 24, p. 1781-1785

- Jenning, V., Gohla, S.H., 2000. Comparison of Wax and Glyceride Solid Lipid Nanoparticles (SLN). *International Journal of Pharmaceutics*, p.219-222
- Jenning V, Gohla S., 2001. Encapsulation of retinoids in solid lipid nanoparticles (SLN). *Microencapsulation*, 18, p. 149-158
- Khurana,S., Bedi, P.M.S., Jain, K., 2013. Preparation and evaluation of solid lipid nanoparticles based nanogel for dermal delivery of meloxicam. *Chemistry and Physics of Lipids* 175-176, p. 65-72.
- Lachman, L., Lieberman, H.A., Kanig, J.L., 1986. *The Theory and Practice of Industrial Pharmacy* 3rd Edition. Philadelphia, p. 491-515
- Lademann J, Richter H, Teichmann A, Otberg N, Blume-Peytavi U, Luengo J, et al., 2007. Nanoparticles—an efficient carrier for drug delivery into the hair follicles. *Eur J Pharm Biopharm* 66, p.159–64.
- Lippacher A, Muller RH, Mader., 2002. Semisolid SLN dispersion for topical application: Influence of formulation and production parameters on viscoelastic properties. *European Journal of Pharmaceutics and Biopharmaceutics* 53, p. 155–160.
- Mei, Z., Chen, H., Weng, T., Yang, Y., Yang, X., 2003. Solid lipid nanoparticle and microemulsion for topical delivery of triptolide. *European Journal of Pharmaceutics and Biopharmaceutics* 56, p.189-196.
- Meidan VC, Bonner M, Michniak B., 2005. Transfollicular drug delivery-is it a reality? *Int J Pharm* 306, p. 1–14.
- Mukherjee, S., Ray, S., Thakur, R.S., 2009. Solid lipid nanoparticles (SLN): A Modern Formulation Approach in Drug Delivery System. *Indian Journal of Pharmaceutical Sciences* 71(4), p. 349-358
- Muller RH, Mader K, Gohla S., 2000. Solid lipid nanoparticles (SLN) for controlled drug delivery a review of the state of art. *Eur J Pharm Biopharm*50, p. 161-177.

- Müller R. H., Radtke M., Wissing S. A., 2002. Solid lipid nanoparticles (SLN) and nanostructure lipid carriers (NLC) in cosmetic and dermatological preparations. *Adv. Drug Deliv. Rev.* 54
- Muller, R.H., Pardeike, J., Hommoss, A., 2009. Lipid nanoparticles (SLN, NLC) in cosmetic and pharmaceutical dermal products. *International Journal of Pharmaceutics* Vol. 366, p. 170-184.
- Patel, M., Murugananthan, Gowda K,P., 2012. *In Vivo* Animal Models in Preclinical Evaluation of Anti-Inflammatory Activity- A Review. *International Journal of Pharmaceutical Research & Allied Sciences* Volume 1, issue 2, p. 01-05
- Rahmatika, Dini., 2012. *Pengaruh Hambatan Stratum Korneum Terhadap Penetrasi APMS Melalui Membran Kulit Tikus Sebagai Studi Praformulasi Sediaan Topikal*. Surabaya : Universitas Airlangga, p.69.
- Rahmawan, T.G., Rosita, N., Erawati, T., 2012. Characterisation of Solid Lipid Nanoparticles p-methoxycinnamic Acid (SLN-PMCA) Formulated With Different Lipid Component: Stearic Acid and Cetyl Alcohol. *PharmaScientia*, Vol.1, No.1.
- Rosita, N., Soeratri, W., 2013. *Pembuatan dan Karakterisasi Sistem Solid Lipid Nanopartikel Asam Para Metoksi Sinamat (SLN-APMS) dengan Basis: Berbagai Komposisi Beeswax dan Gliseril Monostearat (Sebagai Prototipe Sistem Penghantaran Obat Dermal)*. Laporan Akhir Penelitian Unggulan Perguruan Tinggi. Surabaya : Universitas Airlangga.
- Rowe, Raymond, C., Paul, J.S., Marian, E.Q., 2009. *Handbook of Pharmaceutical Excipients*, 6th Ed. London : The Pharmaceutical Press, p. 290-292, 549-553, 592-593, 779-780.
- R. Toll, U. Jacobi, H. Richter, J. Lademann, H. Schaefer, U. Blume-Peytavi., 2004. Penetration profile of microspheres in follicular targeting of terminal hair follicles. *J. Invest. Dermatol.* 123, p.168–176

- Sadono dan Hasmono, D. 2000. *Ketersediaan Hayati / Profil Farmakokinetik Kristal EPMS (Isolat Bahan Aktif Serbuk Rimpang Kencur) pada Hewan coba Kelinci*. Laporan Penelitian Universitas Airlangga.
- Santosa, M.H., 1997. *Metabolisme Metabolit Sekunder Tumbuhan ObatIndonesia pada Kultur Sel Hepatosit Tikus Terisolasi*. Laporan Penelitian Lembaga Penelitian Universitas Airlangga.
- Smith, J.B dan Mangkoewidjojo, S. 1988. *Pemeliharaan, Pembuaian & Penggunaan Hewan Percobaan Di daerah Tropis*. Jakarta : Universitas Indonesia
- Swanson, Jason R., 1996. Anatomy and Histology of Normal Skin. <http://www.meddean.luc.edu/lumen/MedEd/medicine/dermatology/melto/kinlsn/sknltn.htm>, diakses pada tanggal 29 Januari 2013.
- Tadicherla, S., Berman, B., 2006. Percutaneous dermal drug delivery for local pain control. *Therapeutics and Clinical Risk Management* (1), p. 99–113.
- Umar, M.I., Asmawi, M.Z., Sadikun, A., Atangwho, I.J., 2012. Bioactivity-Guided Isolation of Ethyl-*p*-methoxycinnamate, an Anti-inflammatory Constituent, from *Kaempferia galanga* L.Extracts. *Molecules*, p. 8720-8734
- Vega,E., Egeaa, M. E., Ramirez, M.L., Garciaa M.L.,2013. Flurbiprofen PLGA-PEG nanospheres: Role of hydroxy- β -cyclodextrinon ex vivo human skin permeation and in vivo topical anti-inflammatory efficacy. *Colloids and Surfaces B: Biointerfaces*110, p. 339– 346
- Vivek K., Reddy, Harrivardhan., Murthy, Ramachandra S.R., 2006. Investigations of Effect of the Lipid Matrix on Drug Entrapment, In Vitro Release, and Physiscal Stability of Olanzepine-Loaded Solid Lipid Nanoparticles. *AAPS PharmSciTech*, Vol. 8 No. 4.
- Wijayandaru, R.A. 2013. *Pengaruh Kombinasi Lipid Beeswax-Gliseril Monostearat Terhadap Aktivitas Antiinflamasi Sistem SLN-APMS*. SKRIPSI. Surabaya : Airlangga University Press.

- Wissing, S.A., Muller R.H., 2002. The influence of the crystallinity of lipid nanoparticles on their occlusive properties. *International Journal of Pharmaceutics* 242, p. 377–379.
- Wissing, Sylvia, A., Rainer H. Muller., 2003. The influence of solid lipid nanoparticles on skin hydration and viscoelasticity – in vivo study. *European Journal of Pharmaceutics and Biopharmaceutics* 56, p.67–72
- Zhai, H., Maibach, H.I., 2001. Effects of skin occlusion on percutaneous absorption: an overview. *Skin Pharmacol Appl Skin Physiol*, 14(1), p. 1-10.

