

**DAFTAR PUSTAKA**

- Bagshaw, SM, Brophy, PD, Cruz, D, Ronco, C. (2008). Fluid balance as a biomarker: impact of fluid overload on outcome in critically ill patients with acute kidney injury. *Crit Care*, 12: 169.
- Basappanavar, VS, Vadlamani, S, Shreedhar, SJ, Mohandas, BS, Singh, NG, Pandey, G. (2018). End-Tidal Carbon Dioxide as Marker of Cardiac Index after Weaning from Cardiopulmonary Bypass. *EC Anaesthesia*, 4(7), pp 225-231.
- Berglund, JE, Halden, E, Jakobson, S (1994) The effect of PEEP ventilation on cardiac function in closed chest pigs. *Upsala Journal of Medical Sciences*, 99:2, pp 167-78.
- Botero, M, dkk. (2004). Measurement of Cardiac Output Before and After Cardiopulmonary Bypass: Comparison Among Aortic Transit-Time Ultrasound, Thermodilution, and Noninvasive Partial CO<sub>2</sub> Rebreathing. *Journal of Cardiothoracic and Vascular Anesthesia*, Vol 18, No 5 (October), pp 563-572
- Boulain, T, Achard, JM, Teboul, JL, Richard, C, Perrotin, D, Ginies, G. (2002). Changes in BP Induced by Passive Leg Raising Predict Response to Fluid Loading in Critically Ill Patients. *CHEST*, 121: pp 1245-52.
- Bressack, MA, Raffin, TA (1987) Importance of Venous Return, Venous Resistance, and Mean Circulatory Pressure in the Physiology and Management of Shock. *CHEST*, 92(5), pp 906-12
- Caille, V, Jabot, J, Belliard, G, Charron, C, Jardin, F, Baron, AV (2008) Hemodynamic effects of passive leg raising: an echocardiographic study in patients with shock. *Intensive Care Med*.
- Carpio ALM, Mora JI. Positive End-Expiratory Pressure (PEEP) [Updated 2019 Jun 2]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2019 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK441904/>
- Charbonneau H, Riu B, Faron M (2014) Predicting preload responsiveness using simultaneous recordings of inferior and superior vena cavae diameters. *Crit Care*, vol 18, pp 473.

- Fathil, SM, Mahdi, SN, Cheman, Z, Hassan, A, Ahmad, Z, Ismail, AK (2010) A prospective study of tracheal intubation in an academic emergency department in Malaysia. *Int J Emerg Med*, 3, pp 233-37.
- Geerts, BF, Aarts, LPHJ, Groeneveld, AB, Jansen, JRC (2011) Predicting cardiac output responses to passive leg raising by a PEEP-induced increase in central venous pressure, in cardiac surgery patients. *British Journal of Anaesthesia*, 107(2), pp 150-6.
- Giraud, R, Siegenthaler, N, Gayet-Ageron, A, Combescure, C, Romand, JA, Bendjelid, K. (2011). ScvO<sub>2</sub> as a marker to define fluid responsiveness. *J Trauma*, 70: 802-07.
- Grubler, MR, Wigger, O, Berger, D, Biochlinger, S. (2017) Basic concepts of heart-lung interactions during mechanical ventilation. *Swiss Med Wkly*.
- Guarracino F, Ferro B, Forfori F (2014) Jugular vein distensibility predicts fluid responsiveness in septic patients. *Crit Care*, vol 18, pp 647.
- Guyton AC, Jones CE, Coleman TG (1973) Circulatory physiology: cardiac output and its regulation. *Saunders*, Philadelphia
- He HW, Liu DW. (2016). Passive Leg Raising in Intensive Care Medicine. *Chin Med J*; 129:1 pp 755-8.
- Irwin, R. S, Rippe, J. M, Lilly, CM. (2018). Irwin and Rippe's intensive care medicine (8th ed.). Philadelphia: Wolters Kluwer Health/Lippincott Williams & Wilkins.
- Jin, X, Weil, MH, Tang, W, Povoas, H, Pernat, A, Xie, J, Bisera, J. (2000) End-tidal carbon dioxide as a noninvasive indicator of cardiac index during circulatory shock. *Crit Care Med*, 28(7), pp 2415-19
- Jozwiak, M, Monnet, X, Teboul, JL (2018). Prediction of fluid responsiveness in ventilated patients. *Ann Transl Med*, vol 6(18) pp 352-63.
- Kang, WS, Kim, SH, Kim, SY, Oh, CS, Lee, SA, Kim, JS (2014) The influence of positive end-expiratory pressure on stroke volume variation in patients undergoing cardiac surgery: an observational study. *J Thorac Cardiovasc Surg*, 148, pp 3139-45.
- Kellum, John A. (2018). Abnormal Saline and The History of Intravenous Fluids. *Nature*, 14, pp. 358-60.
- Lafanechere, A, Pene, F, Goulenok, C, Delahaye, A, Mallet, V, Choukroun, G, Chiche, JD, Mira, JP, Cariou A. (2006). Changes in aortic blood flow induced by passive leg raising predict fluid responsiveness in critically ill patients. *Critical Care*, 10, pp 132-41.
- Lakhal, K, Nay, MA, Kamel, T, Lortat-Jacob, B, Ehrmann, S, Rozec, B, Boulain, T. (2017). Change In End-Tidal Carbon Dioxide Outperforms Other

- Surrogates For Change In Cardiac Output During Fluid Challenge. *British Journal of Anesthesia*, 118(3) pp 355-62.
- Lansdorp, B, Lemson, J, van Putten, MJ, de Keijzer, A, van der Hoeven, JG, Pickkers, P. (2011). Dynamic indices do not predict volume responsiveness in routine clinical practice. *British Journal of Anaesthesia*, 108(3), 395-401.
- Luecke T, Pelosi P. (2005) Clinical review: Positive end-expiratory pressure and cardiac output. *Critical Care*, 9, pp 607-621.
- Malbrain, M, Van Regenmortel, N, Saugel, B, De Tavernier, B, Van Gaal, PJ, Joannes-Boyau, O, Teboul, JL, Mythen, M, Monnet, X. (2018) Principles of fluid management and stewardship in septic shock: it is time to consider the four D's and the four phases of fluid therapy. *Ann. Intensive Care*, pp 8:66-87.
- Marino, Paul. (2014). Marino's The ICU Book (Fourth edition). Philadelphia: Wolters Kluwer Health/Lippincott Williams & Wilkins.
- Michard, F, Teboul, JL. (2000) Using heart–lung interactions to assess fluid responsiveness during mechanical ventilation. *Crit Care*, 4 pp 282-289
- Marik, PE, Cavallazzi, R. (2008) Does Central Venous Pressure Predict Fluid Responsiveness?. *CHEST*, 134:172-78
- Marik, PE, Cavallazzi, R, Vasu, T, Hirani, A. (2009) Dynamic changes in arterial waveform derived variables and fluid responsiveness in mechanically ventilated patients: A systematic review of the literature. *Crit Care Med*, vol 37(9)
- Marik, PE, Cavallazzi, R. (2013) Does the Central Venous Pressure Predict Fluid Responsiveness? An Updated Meta-Analysis and a Plea for Some Common Sense, *Crit Care Med*, 41, pp 1774-1781
- Marini JJ, Culver BH, Butler J (1981) Effect of positive end-expiratory pressure on canine ventricular function curves. *J Appl Physiol* 51, pp 1367–1374
- Mesquida, J, Gruartmoner, G, Ferrer, R. (2017) Passive leg raising for assessment of volume responsiveness: a review. *Curr Opin Crit Care*, vol 23(3), pp 237-43
- Michard, F, Lopes, MR, Auler, JO. (2007). Pulse pressure variation: beyond the fluid management of patients with shock. *Crit Care*, vol 11(3) pp 131-7.
- Michard, F, Chemla, D, Teboul, JL. (2015). Applicability of pulse pressure variation: how many shades of grey?. *Crit Care*, vol 19, pp 144.
- Monnet X, Bleibtreu A, Ferre A. (2012). Passive leg raising and end expiratory occlusion tests perform better than pulse pressure variation in patients with low respiratory system compliance. *Crit Care Med*, vol 40, pp 152-7.

- Monnet X, Dres M, Ferre, A. (2012) Prediction of fluid responsiveness by a continuous non-invasive assessment of arterial pressure in critically ill patients: comparison with four other dynamic indices. *Br J Anaesth*, vol 109, pp 330-8.
- Monnet X, Teboul JL. (2012) Passive leg raising. In: Pinsky M.R., Brochard L., Mancebo J., Antonelli M. (eds) *Applied Physiology in Intensive Care Medicine 2*. Springer, Berlin, Heidelberg
- Monnet, X, Bataille, A, Magalhaes, E, Barrois, J, Le Corre, M, Gosset, C, Guerin, L, Richard, C, Teboul, JL. (2012) End-tidal carbon dioxide is better than arterial pressure for predicting volume responsiveness by the passive leg raising test. *Intensive Care Med*, 39, pp 93–100
- Monnet, X, Teboul, JL. (2013). Assessment of volume responsiveness during mechanical ventilation: recent advances. *Critical Care*, vol 17 pp 217-31.
- Monnet, X, Teboul, JL (2015) Passive leg raising: five rules, not a drop of fluid. *Critical Care*, 19:18.
- Murphy, CV, Schramm, GE, Doherty, JA, Reichley, RM, Gajic, O, Afessa, B, Micek, ST, Kollef, MH. (2009) The Importance of Fluid management in acute lung injury secondary to septic shock. *CHEST*, 136: pp 102-9
- Nassar BS, Schmidt GA (2015) Capnography During Critical Illness. *CHEST*.
- Ortega, R, Connor, C, Kim, S, Djang, R, Patel, K. (2012) Monitoring Ventilation with Capnography. *N Engl J Med*, pp 367: e27
- Pantazopoulos, C, Xanthos, T, Pantazopoulos, I, Papalois, A, Kouskouni, E, Iacovidou, N. (2015) A Review of Carbon Dioxide Monitoring During Adult Cardiopulmonary Resuscitation. *Heart, Lung, and Circulation*, vol 24, pp 1053-61.
- Pinsky, MR (2007) Heart-lung interactions. *Curr Opin Crit Care* 13, pp 528–531.
- Pinsky, MR (2012) Heart lung interactions during mechanical ventilation. *Curr Opin Crit Care*, 18:3, pp 256–260
- Renner, J, Gruenewald, M, Meybohm, P, Hedderich, J, Steinfath, M, Scholz, J, Bein, B (2008) Effect of elevated PEEP on dynamic variables of fluid responsiveness in a pediatric animal model. *Pediatric Anesthesia*, 18, pp 1170-77.
- Rivers, EP dkk (2006) Fluid-management strategies in acute lung injury – liberal, conservative, or both?. *New Engl J Med*, 354 pp 2598-600
- Roger, C, Zieleskiwicz, L, Demattei, C, Lakhal, K, Piton, G, Louart, B, Constantini, JM, Chabanne, R, Faure, JS, Mahjoub, Y, Desmeulles, I, Quintard, H, Lefrant, JY, Muller, L. (2019) Time course of fluid responsiveness in sepsis: the fluid challenge revisiting (FCREV) study. *Critical Care*, 23:179.

- RSUD Dr. Soetomo Surabaya (2019) Laporan Evaluasi Kinerja Internal Tahun 2019 RSUD Dr. Soetomo Surabaya. Diunduh tanggal 28 Maret 2020 dari: <http://rsudrsoetomo.jatimprov.go.id/laporan-evaluasi-internal-tw-ii-2019-fix-1/>
- Shibutani, K, Muraoka, M, Shirasaki, S, Kubal, K, Sanchala, VT, Gupte, P. (1994) Do changes in end-tidal PCO<sub>2</sub> quantitatively reflect changes in cardiac output? *Anesthesia and Analgesia*, 79(5) pp 829-33.
- Shojaee M, Sabzghabaei A, Alimohammadi H, Derakhshanfar H, Amini A, Esmailzadeh B (2017) Effect of Positive End-Expiratory Pressure on Central Venous Pressure in Patients under Mechanical Ventilation. *Emergency*. 5(1): e1
- Singh, G, Winardi, M, Rumende, CM (2014) Profil Klinis dan Luaran Pasien Gawat Darurat Medis Dewasa di Rumah Sakit Cipto Mangunkusumo, Jakarta. *Ina J Chest Crit and Emerg Med*, 1(3), pp 108-12.
- Smith, R, Fletcher, R. (2000). Positive end-expiratory pressure has little effect on carbon dioxide elimination after cardiac surgery. *Anesth Analg*, 90: pp 85-8.
- Tam, AYB, Lau, FL (2001) A prospective study of tracheal intubation in an emergency department in Hong Kong. *European Journal of Emergency Medicine*, 8(4) pp 305-10
- Tittley, JG, Fremes, SE, Weisel, RD, Christakis, GT, Evans, PJ, Madonik, MM, Ivanov, J, Teasdale, SJ, Mickle, DA, McLaughlin, PR. (1985) Hemodynamic and Myocardial Metabolic Consequences of PEEP. *CHEST*, 88(4) pp 496-502
- Vignon P, Repesse X, Begot E (2017) Comparison of echocardiographic indices used to predict fluid responsiveness in ventilated patients. *Am J Respir Crit Care Med*, vol 195, pp 1022-32.
- Wahba, RW, Tessler, MJ (1996) Misleading end-tidal CO<sub>2</sub> tensions. *Can J Anaesth*, vol 43, pp 862-66.
- Walls, RM, Hockberger, RS, Gausche-Hill, M. (2018) Rosen's emergency medicine: Concepts and clinical practice (9th ed.). Philadelphia, PA: Elsevier/Saunders
- Weil, MH, Bisera, J, Trevino, RP, Rackow, EC. (1985) Cardiac output and end-tidal carbon dioxide, *Crit Care Med*, 13(11) pp 907-9
- Whitesell, R, Asiddao, C, Gollman, D. (1981) Relationship between arterial and peak expired carbon dioxide pressure during anesthesia and factors influencing the difference. *Anesth Analg*, vol 60, pp 508-12.

- Wiesenack, C, Fiegl, C, Keyser, A, Prasser, C, Keyl, C. (2005). Assessment of fluid responsiveness in mechanically ventilated cardiac surgical patients. *European Journal of Anaesthesiology*, 22: pp 658-65.
- Wong, E, Fong, YT, Ho, KK (2004) Emergency airway management – experience of a tertiary hospital in south-east Asia. *Resuscitation*, 61, pp 349-55.
- Xiao-ting, W, Hua, Z, Da-wei, L, Hong-min, Z, Huai-wu, H, Yun, L, Wen-zhao, C (2015) Changes in end-tidal CO<sub>2</sub> could predict fluid responsiveness in the passive leg raising test but not in the mini fluid challenge test: A prospective and observational study. *Journal of Critical Care*, 30, pp 1061-66.
- Yang, X, Du B. (2014). Does pulse pressure variation predict fluid responsiveness in critically ill patients? A systematic review and meta-analysis. *Crit Care*, vol 18, pp 650-9.
- Yang, ZL, Zhou JQ, Sun, BL, Qian, ZX, Zhao, H, Liu, WD (2012) The influence of positive end-expiratory pressure on central venous pressure in patients with severe craniocerebral injury. *Chinese Critical Care Medicine*, 24(5), pp 283-5.
- Yosefy, C, Hay, E, Nasri, Y, Magen, E, Reisin, E. (2004). End tidal carbon dioxide as a predictor of the arterial PCO<sub>2</sub> in the emergency department setting. *Emerg Med J*, vol 21, pp 557-59.
- Young, A, Marik, PE, Sibole, S, Grooms, D, Levitov, A. (2012). Changes in end-tidal carbon dioxide and volumetric carbon dioxide as predictors of volume responsiveness in hemodynamically unstable patients. *J of Cardiothoracic and Vasc Anes*, 70(2): pp 200-08
- Zochios V, Wilkinson J. (2011). Assessment of intravascular fluid status and fluid responsiveness during mechanical ventilation in surgical and intensive care patients. *Journal of Intensive Care Society*, 12:4, pp 295-300