

DAFTAR PUSTAKA

- Annegers, J. F. *et al.* (1980) 'The incidence, causes, and secular trends of head trauma in Olmsted county, Minnesota, 1935-1974', *Neurology*. Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology, 30(9), pp. 912–919. doi: 10.1212/wnl.30.9.912.
- Attella, M. J., Nattinville, A. and Stein, D. G. (1987) 'Hormonal state affects recovery from frontal cortex lesions in adult female rats', *Behavioral and Neural Biology*, 48(3), pp. 352–367. doi: 10.1016/S0163-1047(87)90918-6.
- Badan Pusat Statistik* (no date a). Available at: <https://www.bps.go.id/linkTableDinamis/view/id/1134> (Accessed: 13 July 2020).
- Badan Pusat Statistik (no date b) 'Sensus Penduduk 2010 - Topik Site'. Available at: <https://sp2010.bps.go.id/index.php/site/topik?kid=7&kategori=Ketenagakerjaan>.
- Badan Pusat Statistik Kota Blitar* (no date). Available at: <https://blitarkota.bps.go.id/statictable/2020/05/04/1097/13-8-pelanggaran-lalu-lintas-jalan-raya-menurut-jenisnya-2019> (Accessed: 13 July 2020).
- Baugnon, K. L. and Hudgins, P. A. (2014) 'Skull base fractures and their complications', *Neuroimaging Clinics of North America*. W.B. Saunders, pp. 439–465. doi: 10.1016/j.nic.2014.03.001.
- Bell, R. B. *et al.* (2004) 'Management of cerebrospinal fluid leak associated with craniomaxillofacial trauma', *Journal of Oral and Maxillofacial Surgery*. W.B. Saunders, 62(6), pp. 676–684. doi: 10.1016/j.joms.2003.08.032.
- Berne, J. D. *et al.* (2010) 'A multivariate logistic regression analysis of risk factors for blunt cerebrovascular injury', *Journal of Vascular Surgery*. Mosby, 51(1), pp. 57–64. doi: 10.1016/j.jvs.2009.08.071.

- Bruns, J. and Hauser, W. A. (2003) 'The Epidemiology of Traumatic Brain Injury: A Review', *Epilepsia*, 44(s10), pp. 2–10. doi: 10.1046/j.1528-1157.44.s10.3.x.
- Cannon, C. R. and Jahrsdoerfer, R. A. (1983) 'Temporal Bone Fractures: Review of 90 Cases', *Archives of Otolaryngology*. American Medical Association, 109(5), pp. 285–288. doi: 10.1001/archotol.1983.00800190007002.
- Carstensen, L. L. *et al.* (2011) 'Emotional Experience Improves With Age: Evidence Based on Over 10 Years of Experience Sampling', *Psychology and Aging*, 26(1), pp. 21–33. doi: 10.1037/a0021285.
- CDC (2019) *TBI-related Deaths | Concussion | Traumatic Brain Injury | CDC Injury Center*. Available at: <https://www.cdc.gov/traumaticbraininjury/data/tbi-deaths.html> (Accessed: 29 June 2020).
- Chong, V. F. H., Khoo, J. B. K. and Fan, Y. F. (2004) 'Imaging of the nasopharynx and skull base', *Neuroimaging Clinics of North America*. Elsevier, pp. 695–719. doi: 10.1016/j.nic.2004.07.014.
- Connor, S. E. J. and Flis, C. (2005) 'The contribution of high-resolution multiplanar reformats of the skull base to the detection of skull-base fractures', *Clinical Radiology*. W.B. Saunders, 60(8), pp. 878–885. doi: 10.1016/j.crad.2005.04.003.
- Cushman, J. G. *et al.* (2001) 'Practice Management Guidelines for the Management of Mild Traumatic Brain Injury: The EAST Practice Management Guidelines Work Group', *Journal of Trauma and Acute Care Surgery*, 51(5). Available at: https://journals.lww.com/jtrauma/Fulltext/2001/11000/Practice_Management_Guidelines_for_the_Management.34.aspx.
- Dublin, A. B., French, B. N. and Rennick, J. M. (1977) 'Computed tomography in

- head trauma', *Radiology*. The Radiological Society of North America , 122(2), pp. 365–369. doi: 10.1148/122.2.365.
- Dublin, A. B. and Poirier, V. C. (1976) 'Fracture of the sella turcica', *American Journal of Roentgenology*. American Roentgen Ray Society, 127(6), pp. 969–972. doi: 10.2214/ajr.127.6.969.
- Farin, A. *et al.* (2003) 'Sex-related differences in patients with severe head injury: greater susceptibility to brain swelling in female patients 50 years of age and younger', *Journal of Neurosurgery*, 98(1), pp. 32–36. doi: 10.3171/jns.2003.98.1.0032.
- Faul, M. *et al.* (2010) 'Traumatic brain injury in the United States; emergency department visits, hospitalizations, and deaths, 2002-2006'. Edited by D. of I. R. National Center for Injury Prevention and Control (U.S.). Dept. of Health and Human Services, Centers for Disease Control and Prevention, National Center for Injury Prevention and Control. Available at: <https://stacks.cdc.gov/view/cdc/5571>.
- Feiz-Erfan, I. *et al.* (2007) 'Incidence and pattern of direct blunt neurovascular injury associated with trauma to the skull base', *Journal of Neurosurgery*. American Association of Neurological Surgeons, 107(2), pp. 364–369. doi: 10.3171/JNS-07/08/0364.
- Flores, L. P., Almeida, C. S. and Casulari, L. A. (2000) 'Positive predictive values of selected clinical signs associated with skull base fractures', *Journal of neurosurgical sciences*. Edizioni Minerva Medica, 44(2), p. 77.
- Frijda, N. H. (1986) *The emotions*. Cambridge University Press.
- Gemmete, J. J., Ansari, S. A. and Gandhi, D. M. (2009) 'Endovascular Techniques for Treatment of Carotid-Cavernous Fistula', *Journal of Neuro-Ophthalmology*,

- 29(1), pp. 62–71. doi: 10.1097/WNO.0b013e3181989fc0.
- Gill, H. S. and As, A. B. (2008) ‘Tension pneumocranium in childhood trauma’, *World Journal of Pediatrics*. Springer, 4(3), pp. 231–233. doi: 10.1007/s12519-008-0043-5.
- Glenn, L. W. (2005) ‘Innovations in neuroimaging of skull base pathology’, *Otolaryngologic Clinics of North America*. Elsevier, pp. 613–629. doi: 10.1016/j.otc.2005.03.007.
- Goh, K. Y. C. C. *et al.* (1997) ‘Is routine computed tomographic (CT) scanning necessary in suspected basal skull fractures?’, *Injury*. Elsevier, 28(5–6), pp. 353–357. doi: 10.1016/S0020-1383(97)00024-7.
- Groswasser, Z., Cohen, M. and Keren, O. (1998) ‘Female TBI patients recover better than males’, *Brain Injury*, 12(9), pp. 805–808. doi: 10.1080/026990598122197.
- Jimenez, D. F., Sundrani, S. and Barone, C. M. (1997) ‘Posttraumatic anosmia in craniofacial trauma.’, *The Journal of cranio-maxillofacial trauma*, 3(1), pp. 8–15.
- Kadish, H. A. and Schunk, J. E. (1995) ‘Pediatric Basilar Skull Fracture: Do Children With Normal Neurologic Findings and No Intracranial Injury Require Hospitalization?’, *Annals of Emergency Medicine*. Mosby, 26(1), pp. 37–41. doi: 10.1016/S0196-0644(95)70235-0.
- Karasu, A. *et al.* (2008) ‘Traumatic epidural hematomas of the posterior cranial fossa’, *Surgical Neurology*. Elsevier, 69(3), pp. 247–251. doi: 10.1016/j.surneu.2007.02.024.
- Kementrian Kesehatan Republik Indonesia (2018) ‘Laporan_Nasional_RKD2018_FINAL.pdf’, *Badan Penelitian dan Pengembangan Kesehatan*, p. 198. Available at:

http://labdata.litbang.kemkes.go.id/images/download/laporan/RKD/2018/Laporan_Nasional_RKD2018_FINAL.pdf.

- Keskil, S. *et al.* (1998) 'Clinical significance of acute traumatic intracranial pneumocephalus', *Neurosurgical Review*. Springer Verlag, 21(1), pp. 10–13. doi: 10.1007/BF01111479.
- Kimura, D. (1987) 'Are men's and women's brains really different?', *Canadian Psychology/Psychologie canadienne*, 28(2), pp. 133–147. doi: 10.1037/h0079885.
- Kirkness, C. J. *et al.* (2004) 'Is there a Sex Difference in the Course Following Traumatic Brain Injury?', *Biological Research For Nursing*, 5(4), pp. 299–310. doi: 10.1177/1099800404263050.
- Lewin, W. (1954) 'Cerebrospinal fluid rhinorrha in closed head injuries', *BJS (British Journal of Surgery)*. John Wiley & Sons, Ltd, 42(171), pp. 1–18. doi: 10.1002/bjs.18004217102.
- Liang, W. *et al.* (2007) 'Traumatic Carotid Cavernous Fistula Accompanying Basilar Skull Fracture: a Study on the Incidence of Traumatic Carotid Cavernous Fistula in the Patients With Basilar Skull Fracture and the Prognostic Analysis About Traumatic Carotid Cavernous Fistula', *The Journal of Trauma: Injury, Infection, and Critical Care*, 63(5), pp. 1014–1020. doi: 10.1097/TA.0b013e318154c9fb.
- Liu-Shindo, M. and Hawkins, D. B. (1989) 'Basilar skull fractures in children', *International Journal of Pediatric Otorhinolaryngology*. Elsevier, 17(2), pp. 109–117. doi: 10.1016/0165-5876(89)90086-4.
- Madhusudan, G. *et al.* (2006) 'Nomenclature of Frontobasal Trauma: A New Clinicoradiographic Classification', *Plastic and Reconstructive Surgery*, 117(7), pp. 2382–2388. doi: 10.1097/01.prs.0000218794.28670.07.

- Manson, P. N. *et al.* (2009) 'Frontobasal Fractures: Anatomical Classification and Clinical Significance', *Plastic and Reconstructive Surgery*, 124(6), pp. 2096–2106. doi: 10.1097/PRS.0b013e3181bf8394.
- Mark K. Wax, M. *et al.* (1997) 'Contemporary Management of Cerebrospinal Fluid Rhinorrhea', <http://dx.doi.org/10.1016/S0194-59989770292-4>. SAGE PublicationsSage CA: Los Angeles, CA. doi: 10.1016/S0194-59989770292-4.
- Menon, D. K. *et al.* (2010) 'Position statement: Definition of traumatic brain injury', *Archives of Physical Medicine and Rehabilitation*. W.B. Saunders, 91(11), pp. 1637–1640. doi: 10.1016/j.apmr.2010.05.017.
- MINCY, J. E. (1966) 'POSTTRAUMATIC CEREBROSPINAL FLUID FISTULA OF THE FRONTAL FOSSA', *Journal of Trauma and Acute Care Surgery*, 6(5). Available at: https://journals.lww.com/jtrauma/Fulltext/1966/09000/POSTTRAUMATIC_CEREBROSPINAL_FLUID_FISTULA_OF_THE.7.aspx.
- Mundinger, G. S. *et al.* (2013) 'Blunt-mechanism facial fracture patterns associated with internal carotid artery injuries: Recommendations for additional screening criteria based on analysis of 4,398 patients', *Journal of Oral and Maxillofacial Surgery*. W.B. Saunders, 71(12), pp. 2092–2100. doi: 10.1016/j.joms.2013.07.005.
- Nicholson, B. and Dhindsa, H. (2010) 'Traumatic tension pneumocephalus after blunt head trauma and positive pressure ventilation', *Prehospital Emergency Care*. Taylor & Francis, 14(4), pp. 499–504. doi: 10.3109/10903120903564522.
- Oh, J.-W., Kim, S.-H. and Whang, K. (2017) 'Traumatic Cerebrospinal Fluid Leak: Diagnosis and Management', *Korean Journal of Neurotrauma*, 13(2), pp. 63–67. doi: 10.13004/kjnt.2017.13.2.63.

- Olabinri, E. O. *et al.* (2015) ‘Comparative analysis of clinical and computed tomography features of basal skull fractures in head injury in southwestern Nigeria’, *Journal of Neurosciences in Rural Practice*. Medknow Publications, 6(2), pp. 139–144. doi: 10.4103/0976-3147.153215.
- Osborn, A. G. *et al.* (1978) ‘Intracranial air on computerized tomography’, *Journal of Neurosurgery*. Journal of Neurosurgery Publishing Group, 48(3), pp. 355–359. doi: 10.3171/jns.1978.48.3.0355.
- Parmar, H. *et al.* (2009) ‘Imaging of the Anterior Skull Base’, *Neuroimaging Clinics of North America*. Elsevier, pp. 427–439. doi: 10.1016/j.nic.2009.06.011.
- Ratilal, B. O. *et al.* (2015) ‘Antibiotic prophylaxis for preventing meningitis in patients with basilar skull fractures’, *Cochrane Database of Systematic Reviews*. John Wiley and Sons Ltd. doi: 10.1002/14651858.CD004884.pub4.
- Rincon, S., Gupta, R. and Ptak, T. (2016) ‘Imaging of head trauma’, in *Handbook of Clinical Neurology*. Elsevier B.V., pp. 447–477. doi: 10.1016/B978-0-444-53485-9.00022-2.
- Roof, R. L., Duvdevani, R. and Stein, D. G. (1992) ‘Progesterone treatment attenuates brain edema following contusion injury in male and female rats.’, *Restorative neurology and neuroscience*. Netherlands, 4(6), pp. 425–427. doi: 10.3233/RNN-1992-4608.
- Saarni, C. *et al.* (2007) ‘Emotional Development: Action, Communication, and Understanding’, in *Handbook of Child Psychology*. Hoboken, NJ, USA: John Wiley & Sons, Inc. doi: 10.1002/9780470147658.chpsy0305.
- Samii, M. and Tatagiba, M. (2002) ‘Skull base trauma: Diagnosis and management’, *Neurological Research*. Taylor & Francis, 24(2), pp. 147–156. doi: 10.1179/016164102101199693.

- Savva, A., Taylor, M. J. and Beatty, C. W. (2003) 'Management of Cerebrospinal Fluid Leaks Involving the Temporal Bone: Report on 92 Patients', *The Laryngoscope*. John Wiley & Sons, Ltd, 113(1), pp. 50–56. doi: 10.1097/00005537-200301000-00010.
- Schlosser, R. J. and Bolger, W. E. (2004) 'Nasal Cerebrospinal Fluid Leaks: Critical Review and Surgical Considerations', *Laryngoscope*. John Wiley & Sons, Ltd, pp. 255–265. doi: 10.1097/00005537-200402000-00015.
- Servadei, F., Teasdale, G. and Merry, G. (2001) 'Defining Acute Mild Head Injury in Adults: A Proposal Based on Prognostic Factors, Diagnosis, and Management', *Journal of Neurotrauma*. Mary Ann Liebert, Inc., publishers, 18(7), pp. 657–664. doi: 10.1089/089771501750357609.
- Sivanandapanicker, J. *et al.* (2018) 'Analysis and clinical importance of skull base fractures in adult patients with traumatic brain injury', *Journal of Neurosciences in Rural Practice*. Wolters Kluwer Medknow Publications, 9(3), pp. 370–375. doi: 10.4103/jnrp.jnrp_38_18.
- Stein, D. G. (2001) 'Brain damage, sex hormones and recovery: a new role for progesterone and estrogen?', *Trends in Neurosciences*, 24(7), pp. 386–391. doi: 10.1016/s0166-2236(00)01821-x.
- Steudel, W. I. and Hacker, H. (1986) 'Prognosis, incidence and management of acute traumatic intracranial pneumocephalus - A retrospective analysis of 49 cases', *Acta Neurochirurgica*. Springer-Verlag, 80(3–4), pp. 93–99. doi: 10.1007/BF01812281.
- Stiver, S. I. (2012) 'Management of Skull Base Trauma', in *Schmidek and Sweet Operative Neurosurgical Techniques*. Elsevier, pp. 1559–1572.
- Tagliaferri, F. *et al.* (2006) 'A systematic review of brain injury epidemiology in

- Europe’, *Acta Neurochirurgica*, 148(3), pp. 255–268. doi: 10.1007/s00701-005-0651-y.
- Takeuchi, S. *et al.* (2012) ‘Traumatic posterior fossa subdural hematomas’, *The Journal of Trauma and Acute Care Surgery*, 72(2), pp. 480–486. doi: 10.1097/TA.0b013e31823c89b3.
- Tasdemiroglu, E. and Patchell, R. A. (2002) ‘Classification and Management of Skull Base Fractures’, *Neurosurgery Quarterly*, 12(1). Available at: https://journals.lww.com/neurosurgery-quarterly/Fulltext/2002/03000/Classification_and_Management_of_Skull_Base.4.aspx.
- Teasdale, G. *et al.* (2014) ‘Forty years on: updating the Glasgow Coma Scale’, *Nursing times*, 110, pp. 12–16.
- Thibault, L. E. and Gennarelli, T. A. (1985) ‘Biomechanics and Craniocerebral’, *Central Nervous System Trauma Status Report [1985]*, p. 379.
- Tomura, N. *et al.* (1998) ‘Comparison of MR imaging and CT in discriminating tumor infiltration of bone and bone marrow in the skull base’, *Computerized Medical Imaging and Graphics*. Elsevier Sci Ltd, 22(1), pp. 41–51. doi: 10.1016/S0895-6111(98)00008-1.
- Tunik, M. G. *et al.* (2016) ‘Clinical Presentations and Outcomes of Children With Basilar Skull Fractures After Blunt Head Trauma’, *Annals of Emergency Medicine*. Mosby Inc., 68(4), pp. 431-440.e1. doi: 10.1016/j.annemergmed.2016.04.058.
- Wang, H. *et al.* (2018) ‘Traumatic skull fractures in children and adolescents: A retrospective observational study’, *Injury*. Elsevier Ltd, 49(2), pp. 219–225. doi: 10.1016/j.injury.2017.11.039.
- Wani, A. A. *et al.* (2013) ‘Skull base fractures: An institutional experience with

review of literature’, *The Indian Journal of Neurotrauma*. Georg Thieme Verlag KG, 10(2), pp. 120–126. doi: 10.1016/j.ijnt.2013.05.009.

World Health Organization (2014) ‘Manual Keselamatan Jalan Untuk Pengambil Keputusan dan Praktisi’, p. 248.

Yellinek, S. *et al.* (2016) ‘Clinical significance of skull base fracture in patients after traumatic brain injury’, *Journal of Clinical Neuroscience*. Churchill Livingstone, 25, pp. 111–115. doi: 10.1016/j.jocn.2015.10.012.

Yilmazlar, S. *et al.* (2006) ‘Cerebrospinal fluid leakage complicating skull base fractures: Analysis of 81 cases’, *Neurosurgical Review*. Springer, pp. 64–71. doi: 10.1007/s10143-005-0396-3.

Ziu, M., Savage, J. G. and Jimenez, D. F. (2012) ‘Diagnosis and treatment of cerebrospinal fluid rhinorrhea following accidental traumatic anterior skull base fractures’, *Neurosurgical Focus*. American Association of Neurological Surgeons, 32(6), p. E3. doi: 10.3171/2012.4.FOCUS1244.