

DAFTAR PUSTAKA

- Adani, S. and Cepanec, M. (2019) ‘Sex differences in early communication development: Behavioral and neurobiological indicators of more vulnerable communication system development in boys’, *Croatian Medical Journal*, 60(2), pp. 141–149. doi: 10.3325/cmj.2019.60.141.
- Amemiya, É. E., Soares, A. D. and Chiari, B. M. (2016) ‘Communicative indicators, motor and cognitive development of hearing-impaired children’, *Journal of Human Growth and Development*, 26(1), pp. 54–60. doi: 10.7322/jhgd.113717.
- Am Zehnhoff-Dinnesen, A., Wiskirska-Woznica, B. and Nawka, T., (2020). *Phoniatrics I Fundamentals – Voice Disorders – Disorders Of Language And Hearing Development*. Berlin, Heidelberg: Springer, pp.870-875
- Andrade, C. (2016) ‘Use of acetaminophen (paracetamol) during pregnancy and the risk of attention-deficit/hyperactivity disorder in the offspring’, *Journal of Clinical Psychiatry*, 77(3), pp. e312–e314. doi: 10.4088/JCP.16f10721.
- Anne, S., Lieu, J. E. C. and Cohen, M. S. (2017) ‘Speech and Language Consequences of Unilateral Hearing Loss: A Systematic Review’, *Otolaryngology - Head and Neck Surgery (United States)*, 157(4), pp. 572–579. doi: 10.1177/0194599817726326.
- Aoki, S. et al. (2018) ‘Developmental outcomes of Japanese children born through Assisted Reproductive Technology (ART) in toddlerhood’, *Journal of Obstetrics and Gynaecology Research*, 44(5), pp. 929–935. doi: 10.1111/jog.13613.
- Araujo, D. M., Cabrera Santos, D. C. and Marconi Pinheiro Lima, M. C. (2020) ‘Cognitive, language and motor development of infants exposed to risk and protective factors’, *International Journal of Pediatric Otorhinolaryngology*. Elsevier B.V., 138(August), p. 110353. doi: 10.1016/j.ijporl.2020.110353.
- ASHA (2015) *Audiology Information Series AUDIOLOGY Information Series TYPES OF HEARING LOSS CONFIGURATION OF HEARING LOSS*. Available at: www.asha.org (Accessed: 26 June 2019).
- Bernstein, N., Akram, M. and Daniyal, M. (2020) ‘Is it safe to consume traditional medicinal plants during pregnancy ?’, *Phytotherapy Research*, pp. 1–17. doi: 10.1002/ptr.6935.
- Boonen, N., Kloots, H. and Gillis, S. (2020) ‘Rating the overall speech quality of hearing-impaired children by means of comparative judgements’, *Journal of Communication Disorders*. Elsevier, 83(November 2019), p. 105969. doi: 10.1016/j.jcomdis.2019.105969.
- Boyle, A. K. et al. (2017) ‘Preterm birth: Inflammation, fetal injury and treatment strategies’, *Journal of Reproductive Immunology*. Elsevier Ireland Ltd, 119, pp. 62–66. doi: 10.1016/j.jri.2016.11.008.
- Carruth, B. R. and Skinner, J. D. (2002) ‘Feeding Behaviors and Other Motor Development in Healthy Children (2–24 Months)’, *Journal of the American College of Nutrition*. Routledge, 21(2), pp. 88–96. doi:

10.1080/07315724.2002.10719199.

- Celikkiran, S., Bozkurt, H. and Coşkun, M. (2015) 'Denver Developmental Test Findings and their Relationship with Sociodemographic Variables in a Large Community Sample of 0-4-Year-Old Children.', *Noro psikiyatri arsivi*. Turkish Neuropsychiatric Society, 52(2), pp. 180–184. doi: 10.5152/npa.2015.7230.
- Collisson, B. A. et al. (2016) 'Risk and protective factors for late talking: An epidemiologic investigation', *Journal of Pediatrics*. Elsevier Inc., 172, pp. 168–174.e1. doi: 10.1016/j.jpeds.2016.02.020.
- Cupples, L. et al. (2018) 'Language and speech outcomes of children with hearing loss and additional disabilities: Identifying the variables that influence performance at 5 years of age', *International Journal Audiology*. doi: 10.1080/14992027.2016.1228127.Language.
- Cushing, S. L. and Papsin, B. C. (2015) 'Taking the History and Performing the Physical Examination in a Child with Hearing Loss', *Otolaryngologic Clinics of North America*. Elsevier Inc, 48(6), pp. 903–912. doi: 10.1016/j.otc.2015.07.010.
- Dionne, G. et al. (2008) 'Gestational diabetes hinders language development in offspring', *Pediatrics*, 122(5). doi: 10.1542/peds.2007-3028.
- Douniadakis, D. E. et al. (2002) 'Incidence of hearing loss among children presented with speech-language delay', *Scandinavian Audiology*, 30(1), pp. 204–205. doi: 10.1080/010503901300007533.
- Ebrahimi, A. A. et al. (2017) 'Postural control in deaf children', *Acta Medica Iranica*, 55(2), pp. 115–122.
- Fouché, L. C., Kritzinger, A. and Le Roux, T. (2019) 'Corrigendum: Gestational age and birth weight variations in young children with language impairment at an early communication intervention clinic (South African Journal of Communication Disorders, (2018), 10.4102/sajcd.v65i1.584)', *South African Journal of Communication Disorders*, 66(1). doi: 10.4102/sajcd.v65i1.584.
- Gheysen, F., Loots, G. and Van Waelvelde, H. (2008) 'Motor development of deaf children with and without cochlear implants', *Journal of Deaf Studies and Deaf Education*, 13(1), pp. 61–70. doi: 10.1093/deafed/enm053.
- Gregersen, T. L. and Ulrik, C. S. (2013) 'Safety of bronchodilators and corticosteroids for asthma during pregnancy: What we know and what we need to do better', *Journal of Asthma and Allergy*, (6), pp. 117–125. doi: 10.2147/JAA.S52592.
- Hadders-Algra, M. (2018) 'Early human motor development: From variation to the ability to vary and adapt', *Neuroscience and Biobehavioral Reviews*, 90(January), pp. 411–427. doi: 10.1016/j.neubiorev.2018.05.009.
- Halliday, L. F., Tuomainen, O. and Rosen, S. (2017) 'Language development and impairment in children with mild to moderate sensorineural hearing loss', *Journal of Speech, Language, and Hearing Research*, 60(6), pp. 1551–1567. doi: 10.1044/2016_JSLHR-L-16-0297.
- Hammer, C. S. et al. (2017) 'Late talkers: A population-based study of risk factors and school readiness consequences', *Journal of Speech, Language, and Hearing*

- Research*, 60(3), pp. 607–626. doi: 10.1044/2016_JSLHR-L-15-0417.
- Hayes, S. H. et al. (2013) *Anatomy and physiology of the external, middle and inner ear*. 1st edn, *Handbook of Clinical Neurophysiology*. 1st edn. © 2013, Elsevier B.V. All rights reserved. doi: 10.1016/B978-0-7020-5310-8.00001-6.
- Hogan, A. et al. (2011) ‘Communication and behavioural disorders among children with hearing loss increases risk of mental health disorders’, *Australian and New Zealand Journal of Public Health*, 35(4), pp. 377–383. doi: 10.1111/j.1753-6405.2011.00744.x.
- de Hoog, B. E. et al. (2016) ‘Linguistic profiles of children with CI as compared with children with hearing or specific language impairment’, *International Journal of Language and Communication Disorders*, 51(5), pp. 518–530. doi: 10.1111/1460-6984.12228.
- Hsieh, D. C. et al. (2019) ‘Implications of vaginal instrumental delivery for children’s school achievement: A population-based linked administrative data study’, *Australian and New Zealand Journal of Obstetrics and Gynaecology*, 59(5), pp. 677–683. doi: 10.1111/ajo.12952.
- Inoue, A. et al. (2013) ‘Effect of vestibular dysfunction on the development of gross motor function in children with profound hearing loss’, *Audiology and Neurotology*, 18(3), pp. 143–151. doi: 10.1159/000346344.
- Keilmann, A. et al. (2011) ‘Manifestation of speech and language disorders in children with hearing impairment compared with children with specific language disorders’, *Logopedics Phoniatrics Vocology*, 36(1), pp. 12–20. doi: 10.3109/14015439.2010.517550.
- King-Dowling, S. et al. (2015) ‘Reprint of “Co-occurring motor, language and emotional-behavioral problems in children 3-6years of age”’, *Human Movement Science*, 42, pp. 344–351. doi: 10.1016/j.humov.2015.06.005.
- Krakowiak, P. et al. (2012) ‘Maternal metabolic conditions and risk for autism and other neurodevelopmental disorders’, *Pediatrics*, 129(5). doi: 10.1542/peds.2011-2583.
- Kuschke, S. et al. (2020) ‘Profile of childhood hearing loss in the Western Cape, South Africa’, *International Journal of Pediatric Otorhinolaryngology*. Elsevier B.V., 137(July), p. 110248. doi: 10.1016/j.ijporl.2020.110248.
- Leigh, G. et al. (2015) ‘Factors affecting psychosocial and motor development in 3-year-old children who are deaf or hard of hearing’, *Journal of Deaf Studies and Deaf Education*. doi: 10.1093/deafed/env028.
- Libertus, K. and Violi, D. A. (2016) ‘Sit to talk: Relation between motor skills and language development in infancy’, *Frontiers in Psychology*, 7(MAR), pp. 1–8. doi: 10.3389/fpsyg.2016.00475.
- Lim, R. and Brichta, A. M. (2016) ‘Anatomical and physiological development of the human inner ear’, *Hearing Research*. Elsevier B.V, 338, pp. 9–21. doi: 10.1016/j.heares.2016.02.004.
- McCreery, R. W. et al. (2015) ‘Speech Recognition and Parent Ratings From Auditory Development Questionnaires in Children Who Are Hard of Hearing.’, *Ear and*

- hearing.* NIH Public Access, 36 Suppl 1(0 1), pp. 60S-75S. doi: 10.1097/AUD.0000000000000213.
- McFarland, D. H. (2015) 'Oropharyngeal-Articulatory System', in *Netter's Atlas of Anatomy for Speech, Swallowing and Hearing*. 2nd edn. Elsevier Inc., pp. 109–160.
- McLaughlin, M. R. (2011) 'Speech and Language Delay in Children', *American Family Physician*. American Academy of Family Physicians, 83(10), pp. 1183–1188. Available at: <https://www.aafp.org/afp/2011/0515/p1183.html> (Accessed: 29 May 2019).
- Molini-Avejona, D. R., Ferreira, L. V. and Amato, C. A. de L. H. (2017) 'Risk Factors for Speech-Language Pathologies in Children', in Fernandez, F. D. M. (ed.) *Advances in Speech-language Pathology*. InTech, p. 38. doi: 10.5772/intechopen.70107.
- Moneta, L. B. and Quintanilla-Dieck, L. (2017) 'Embryology and anatomy of the ear', *Operative Techniques in Otolaryngology-Head and Neck Surgery*, 28(2), pp. 66–71. doi: <https://doi.org/10.1016/j.otot.2017.03.011>.
- Moore, C. et al. (2019) 'Point, walk, talk: Links between three early milestones, from observation and parental report.', *Developmental Psychology*, 55(8), pp. 1579–1593. doi: 10.1037/dev0000738.
- Netten, A. P. et al. (2015) 'Early identification: Language skills and social functioning in deaf and hard of hearing preschool children', *International Journal of Pediatric Otorhinolaryngology*. Elsevier Ireland Ltd, 79(12), pp. 2221–2226. doi: 10.1016/j.ijporl.2015.10.008.
- Niu, K. et al. (2020) 'Risk factors and etiology of childhood hearing loss: a cohort review of 296 subjects', *Acta Oto-Laryngologica*. Taylor & Francis, 140(8), pp. 668–674. doi: 10.1080/00016489.2020.1757753.
- Oguz, H., Hess, M. and Klein, A. M. (2015) *Phoniatriics, BioMed Research International*. doi: 10.1155/2015/156014.
- Olesen, A. W., Olsen, J. and Zhu, J. L. (2015) 'Developmental milestones in children born post-term in the Danish National Birth Cohort: A main research article', *BJOG: An International Journal of Obstetrics and Gynaecology*, 122(10), pp. 1331–1339. doi: 10.1111/1471-0528.13237.
- Oliveira, A. C. de et al. (2018) 'Hearing, language, motor and social skills in the child development: a screening proposal', *Revista CEFAC*. CEFAC Saúde e Educação, 20(2), pp. 218–227. doi: 10.1590/1982-0216201820216617.
- Oudgenoeg-Paz, O., Volman, M. J. M. and Leseman, P. P. M. (2016) 'First steps into language? Examining the specific longitudinal relations between walking, exploration and linguistic skills', *Frontiers in Psychology*, 7(SEP), pp. 1–12. doi: 10.3389/fpsyg.2016.01458.
- Polidano, C., Zhu, A. and Bornstein, J. C. (2017) 'The relation between cesarean birth and child cognitive development', *Scientific Reports*. Springer US, 7(1), pp. 1–10. doi: 10.1038/s41598-017-10831-y.
- Psarommatis, I. M. et al. (2001) 'Hearing loss in speech-language delayed children',

- International Journal of Pediatric Otorhinolaryngology*, 58(3), pp. 205–210. doi: 10.1016/S0165-5876(01)00430-X.
- R., L. and N., B. (2013) ‘12 minute consultation: An evidence-based approach to the management of a child with speech and language delay’, *Clinical Otolaryngology*, 38(2), pp. 148–153. doi: 10.1111/coa.12082.
- Rich, J. N., Allen, S. T. and Rivera, A. L. (2013) ‘Middle Ear Physiology’, in *Encyclopedia of Otolaryngology, Head and Neck Surgery*. Springer Berlin Heidelberg. doi: 10.1007/978-3-642-23499-6.
- Rosenbaum, S. and Simon, P. (eds) (2016) *Speech and Language Disorders in Children: Implications for the Social Security Administration’s Supplemental Security Income Program*. Washington, D.C.: National Academies Press. doi: 10.17226/21872.
- Rubio-Codina, M. et al. (2016) ‘Concurrent Validity and Feasibility of Short Tests Currently Used to Measure Early Childhood Development in Large Scale Studies’, *PLOS ONE*. Edited by D. O. Carpenter. Public Library of Science, 11(8), p. e0160962. doi: 10.1371/journal.pone.0160962.
- Rydz, D. et al. (2005) ‘Developmental Screening’, *Journal of Child Neurology*, pp. 4–21. Available at: https://www.medscape.com/viewarticle/504722_6 (Accessed: 19 June 2019).
- SABRA M. AHMED, M.D., H. H. A. M. D. . (2018) ‘Risk Factors of Delayed Language Development among Preschool Children Attending Assiut University Hospitals’, *The Medical Journal of Cairo University*, 86(9), pp. 2279–2285. doi: 10.21608/mjcu.2018.57520.
- Sahli, A. S. (2019) ‘Developments of children with hearing loss according to the age of diagnosis, amplification, and training in the early childhood period’, *European Archives of Oto-Rhino-Laryngology*. Springer Berlin Heidelberg, (0123456789). doi: 10.1007/s00405-019-05501-w.
- Seikel, J. A., King, D. W. and Drumright, D. G. (2020) *Anatomy & physiology for speech, language, and hearing*. 6th edn. San Diego: Plural Publishing, Inc.
- Shperling, R. B. and Yoge, Y. (2020) ‘Adverse outcomes of measles infection during pregnancy and in the perinatal period’, *Journal of Maternal-Fetal and Neonatal Medicine*. Taylor & Francis, 0(0), pp. 1–6. doi: 10.1080/14767058.2020.1759536.
- Stene-Larsen, K. et al. (2014) ‘Communication impairments in early term and late preterm children: A prospective cohort study following children to age 36 months’, *Journal of Pediatrics*. Elsevier Inc, 165(6), pp. 1123–1128. doi: 10.1016/j.jpeds.2014.08.027.
- Stübner, C. et al. (2020) ‘Schoolchildren with unilateral or mild to moderate bilateral sensorineural hearing loss should be screened for neurodevelopmental problems’, *Acta Paediatrica, International Journal of Paediatrics*, 109(7), pp. 1430–1438. doi: 10.1111/apa.15088.
- Sumi, S. and Editor, A. M. (2015) ‘U . S . Preventive Services Task Force Screening for Speech and Language Delay and Disorders in Children Aged 5 Years or

- Younger ':, *American Family Physician*, 92(4).
- Taylor, C. L. et al. (2018) 'Prenatal and perinatal risks for late language emergence in a population-level sample of twins at age 2', *BMC Pediatrics*. BMC Pediatrics, 18(1), pp. 1–9. doi: 10.1186/s12887-018-1035-9.
- Toda, K. (2017) 'Is acetaminophen safe in pregnancy?', *Scandinavian Journal of Pain*. Elsevier B.V., 17, pp. 445–446. doi: 10.1016/j.sjpain.2017.09.007.
- Tokgöz-Yilmaz, S. et al. (2013) 'Evaluation of hearing and speech-language in preschool children: How important, why we should perform?', *Turkish Journal of Pediatrics*, 55(6), pp. 606–611.
- Tomblin, J. B. et al. (2015) 'Language Outcomes in Young Children with Mild to Severe Hearing Loss.', *Ear and hearing*. NIH Public Access, 36 Suppl 1(0 1), pp. 76S-91S. doi: 10.1097/AUD.0000000000000219.
- Tonn, C. R. and Grundfast, K. M. (2014) 'What an otolaryngologist should know about evaluation of a child referred for delay in speech development', *JAMA Otolaryngology - Head and Neck Surgery*, 140(3), pp. 259–265. doi: 10.1001/jamaoto.2013.6368.
- Udagatti, V. D. and Dinesh Kumar, R. (2016) 'Endoscopic Ear Surgery: Critical Review of Anatomy and Physiology of Normal and Reconstructive Middle Ear', *Indian Journal of Otolaryngology and Head and Neck Surgery*. Springer India, 68(2), pp. 157–162. doi: 10.1007/s12070-015-0927-3.
- Vidranski, T. and Farkaš, D. (2015) 'Motor Skills in Hearing Impaired Children with or without Cochlear Implant--A Systematic Review.', *Collegium antropologicum*, 39 Suppl 1, pp. 173–9. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26434027>.
- Visscher, C. et al. (2007) 'Motor Profile of Children With Developmental Speech and Language Disorders', *Pediatrics*, 120(1), pp. e158–e163. doi: 10.1542/peds.2006-2462.
- Visser-Bochane, M. I. et al. (2020) 'Identifying Milestones in Language Development for Young Children Ages 1 to 6 Years', *Academic Pediatrics*. Elsevier Inc., 20(3), pp. 421–429. doi: 10.1016/j.acap.2019.07.003.
- Wallander, J. L. et al. (2014) 'Developmental trajectories of children with birth asphyxia through 36months of age in low/low-middle income countries', *Early Human Development*, 90(7), pp. 343–348. doi: 10.1016/j.earlhumdev.2014.04.013.
- WHO (2017) 'WHO | Grades of hearing impairment', WHO. World Health Organization. Available at: https://www.who.int/pbd/deafness/hearing_impairment_grades/en/ (Accessed: 14 June 2019).
- WHO (2019) *Deafness and hearing loss*, World Health Organization. Available at: <https://www.who.int/news-room/fact-sheets/detail/deafness-and-hearing-loss> (Accessed: 14 June 2019).
- WHO Multicentre Growth Reference Study Group (2006) 'WHO Motor Development Study: Windows of achievement for six gross motor development milestones',

- Acta Paediatrica, International Journal of Paediatrics*, 95(SUPPL. 450), pp. 86–95. doi: 10.1080/08035320500495563.
- Wilar, R. et al. (2010) ‘Correlation between hyperbilirubinemia in term infants and developmental delay in 2-4 year-old children’, *Paediatrica Indonesiana*, 50(3), p. 154. doi: 10.14238/pi50.3.2010.154-8.
- Williams, A. et al. (2020) ‘Cognitive and Behavioral Functioning in Hearing-Impaired Children with and without Language Delay’, *Otolaryngology - Head and Neck Surgery (United States)*, 163(3), pp. 588–590. doi: 10.1177/0194599820915741.
- Wooles, N., Swann, J. and Hoskison, E. (2018) ‘Speech and language delay in children: a case to learn from.’, *The British journal of general practice : the journal of the Royal College of General Practitioners*. Royal College of General Practitioners, 68(666), pp. 47–48. doi: 10.3399/bjgp17X694373.
- World Health Organization (2016) *CHILDHOOD HEARING LOSS Strategies for prevention and care*, World Health Organization.
- Yasin, A. et al. (2017) ‘Speech and language delay in childhood: a retrospective chart review’, *ENT Updates*, 7(1), pp. 22–27. doi: 10.2399/jmu.2017001004.
- Yılmaz, D. et al. (2016) ‘Screening 5 and 6 year-old children starting primary school for development and language’, *Turkish Journal of Pediatrics*, 58(2), pp. 136–144. doi: 10.24953/turkjped.2016.02.003.
- Zengin-Akkuş, P. et al. (2018) ‘Speech delay in toddlers: Are they only “late talkers”?’, *Turkish Journal of Pediatrics*, 60(2), pp. 165–172. doi: 10.24953/turkjped.2018.02.008.
- Zhang, Z. (2016) ‘Mechanics of human voice production and control’, *The Journal of the Acoustical Society of America*, 140(4), pp. 2614–2635. doi: 10.1121/1.4964509.