

**DAFTAR PUSTAKA**

- Aaberg, M., Burch, D., Hud, Z. and Zacharias, M. (2008). Gender differences in the onset of diabetic neuropathy. *Journal of Diabetes and its Complications*, 22(2), pp.83-87.
- ADA. (2008). Standards of medical care in diabetes. *Diabetes Care*, 31(Suppl 1), pp.S12-S54.
- ADA. (2018). Classification and Diagnosis of Diabetes:Standards of Medical Care in Diabetes—2018. *Diabetes Care*, 41(Suppl 1), pp.S13-S2.
- Al-Rubeaan, K., Al Derwish, M., Ouizi, S., Youssef, A., Subhani, S., Ibrahim, H. and Alamri, B. (2015). Diabetic Foot Complications and Their Risk Factors from a Large Retrospective Cohort Study. *PLOS ONE*, 10(5), p.e0124446.
- Andatu, R.A. (2016). Hubungan Antara Durasi Menderita Diabetes Melitus (DM). Skripsi S1. Universitas Muhammadiyah Yogyakarta.
- Ashok, S., Ramu, M., Deepa, R., Mohan, V. (2002). Prevalence of neuropathy in type 2 diabetic patients attending a diabetes centre in South India. *J Assoc Physicians India*, 50, pp.546-550.
- Belmin, J., & Valensi, P. (1996). Diabetic Neuropathy in Elderly Patients. *Drugs & Aging*, 8(6), pp.416–429.
- Berkowitz, G., Lapinski, R., Wein, R. and Lee, D. (1992). Race/Ethnicity and Other Risk Factors for Gestational Diabetes. *American Journal of Epidemiology*, 135(9), pp.965-973.
- Betteng, R., Pangemanan, D., and Mayulu, N. (2014). Analisis faktor resiko penyebab terjadinya diabetes melitus Tipe 2 pada wanita usia produktif di Puskesmas Wawonasa. *E-Biomedik*, 2(2), pp.404–412.
- Black, J.M., & Hawks, J. (2009). *Medical-surgical nursing: Clinical Management for pPsitive Outcomes*, Eight edition. Singapore: Saunders Elsevier.
- Booth, M., Okely, A., Chey, T. and Bauman, A. (2002). The reliability and validity of the Adolescent Physical Activity Recall Questionnaire. *Medicine & Science in Sports & Exercise*, 34(12), pp.1986-1995.
- Booya, F., Bandarian, F., Larijani, B., Pajouhi, M., Nooraei, M. and Lotfi, J. (2005). Potential risk factors for diabetic neuropathy: a case control study. *BMC Neurology*, 5(1).
- Boulton, A., Malik, R., Arezzo, J. and Sosenko, J. (2004). Diabetic Somatic Neuropathies. *Diabetes Care*, 27(6), pp.1458-1486.

- Brancati, F., Whelton, P., Kuller, L. and Klag, M. (1996). Diabetes mellitus, race, and socioeconomic status a population-based study. *Annals of Epidemiology*, 6(1), pp.67-73.
- British Columbia Provincial Nursing Skin & Wound Care. (2012). Procedure: Monofilament Testing for Loss of Protective Sensation of Diabetic/Neuropathic Feet for Adults & Children [online]. Available at: <https://www.clwk.ca/buddydrive/file/procedure-monofilament-testing/?download=106%253Aprocedure-monofilament-testing-for-lops> (Accessed 15 June 2019).
- Brownlee, M. (2005). The Pathobiology of Diabetic Complications: A Unifying Mechanism. *Diabetes*, 54(6), pp.1615-1625.
- Bruschi, L.K.M., Rocha, D.A., Filho, E.L.G., Barboza, N.M.P., Frisanco, P.A.B., Callegaro, R.M., Sá, L.B.P.C. and Arbex, A.K. (2017) Diabetes Mellitus and Diabetic Peripheral Neuropathy. *Open Journal of Endocrine and Metabolic Diseases*, 7, 12-21.
- Callaghan, B., Gao, L., Li, Y., Zhou, X., Reynolds, E., Banerjee, M., Pop-Busui, R., Feldman, E. and Ji, L. (2018). Diabetes and obesity are the main metabolic drivers of peripheral neuropathy. *Annals of Clinical and Translational Neurology*, 5(4), pp.397-405.
- Chan, J., Rimm, E., Colditz, G., Stampfer, M. and Willett, W. (1994). Obesity, Fat Distribution, and Weight Gain as Risk Factors for Clinical Diabetes in Men. *Diabetes Care*, 17(9), pp.961-969.
- Cleveland Clinic. (2019). Neuropathy | Cleveland Clinic. [online] Available at: <https://my.clevelandclinic.org/health/diseases/14737-neuropathy> [Accessed 5 Jul. 2019].
- Colditz, G. (1995). Weight Gain as a Risk Factor for Clinical Diabetes Mellitus in Women. *Annals of Internal Medicine*, 122(7), p.481.
- Creatore, M., Moineddin, R., Booth, G., Manuel, D., DesMeules, M., McDermott, S. and Glazier, R. (2010). Age- and sex-related prevalence of diabetes mellitus among immigrants to Ontario, Canada. *Canadian Medical Association Journal*, 182(8), pp.781-789.
- Duarsa, M., Arimbawa, I. and Indrayani, I. (2019). Hipertensi Sebagai Faktor Risiko Nyeri Neuropati Diabetik Pada Pasien Diabetes Mellitus Tipe I Di RSUP Sanglah Denpasar. *Jurnal Medika Udayana*, 8(10).
- Ghaderpanahi, M., Sharifi, F., Badamchizade, Z., Mirarefin, M., Ebrahim, P., Ghoutbi, S., Nouri, M. and Larijani, B. (2011). Association of Physical Activity with Risk of Type 2 Diabetes. *Iran J Public Health*, 40(1), pp.86-93.

- Hassan, M., Curley, S., Li, D., Kaseb, A., Davila, M., Abdalla, E., Javle, M., Moghazy, D., Lozano, R., Abbruzzese, J. and Vauthey, J. (2010). Association of diabetes duration and diabetes treatment with the risk of hepatocellular carcinoma. *Cancer*, 116(8), pp.1938-1946.
- IDF. (2019). IDF diabetes atlas - 2017 Atlas. [online] . Available at: <https://diabetesatlas.org/resources/2017-atlas.html> [Accessed 12 May 2019].
- Kapoor, N., David, K., Saravanan, Bharathy. (2017). Approach to Diabetic Neuropathy. *Curr Med Issues*. 3(15), pp.189-199.
- Kahn, S., Cooper, M. and Del Prato, S. (2014). Pathophysiology and treatment of type 2 diabetes: perspectives on the past, present, and future. *The Lancet*, 383(9922), pp.1068-1083.
- Kautzky-Willer, A., Harreiter, J. and Pacini, G. (2016). Sex and Gender Differences in Risk, Pathophysiology and Complications of Type 2 Diabetes Mellitus. *Endocrine Reviews*, 37(3), pp.278-316.
- Khana, R. (2016). Gambaran Neuropati Perifer Pada Diabetesi di Wilayah Kerja Puskesmas Kedungmundu Semarang. Skripsi S1. Fakultas Kedokteran Universitas Diponegoro.
- Kluding, P.M., Bareiss, S.K., Hastings, M., Marcus, R., Sinacore, D. and Mueller, M. (2017). Physical Training and Activity in People With Diabetic Peripheral Neuropathy: Paradigm Shift. *Physical Therapy*, Volume 97, Issue 1, January 2017, pp.31–43.
- Kominfo (2015). Masih Tinggi, Prevalensi Diabetes di Jatim | Dinas Komunikasi dan Informatika Provinsi Jawa Timur. [online] Available at: <http://kominfo.jatimprov.go.id/read/umum/masih-tinggi-prevalensi-diabetes-di-jatim-> [Accessed 12 May 2019].
- Lathifah, N. L. (2017). The Relationship Between Duration Disease and Glucose Blood Related to Subjective Compliance in Diabetes Mellitus. *Jurnal Berkala Epidemiologi*, 5, pp.218-230.
- Li, L., Chen, J., Wang, J., and Cai, D. (2015). Prevalence and risk factors of diabetic peripheral neuropathy in Type 2 diabetes mellitus patients with overweight/obese in Guangdong province, China. *Primary Care Diabetes*, 9(3), pp.191-195.
- Lipsky, B.A., Berendt, A.R., Deery, H.G., Embil, J.M., Joseph, W.S., Karchmer, A.W., LeFrock, J.L., Lew, D.P., Mader, J.T., Norden, C. and Tan, J.S. (2004). Diagnosis and Treatment of Diabetic Foot Infections : Clinical Infectious Diseases. Oxford University Press, 39(7) pp. 885-910.
- Maksimovic, M. (2013). Whether Obesity is Associated with Peripheral Arterial Disease. *Angiology: Open Access*, [online] 02(01). Available at:

<https://www.longdom.org/open-access/Whether-Obesity-is-Associated-with-Peripheral-Arterial-Disease-2329-9495.1000e104.pdf>

- Mildawati, M., Diani, N., and Wahid, A. (2019). Hubungan Usia, Jenis Kelamin, dan Lama Menderita Diabetes dengan Kejadian Neuropati Perifer Diabetik. *Caring Nursing Journal*, 3(2), pp.31-37.
- Nasution, N. D. (2017). Hubungan Indeks Masa Tubuh dengan Neuropati Diabetik di Puskesmas Sentosa Baru Medan. Medan: Fakultas Kedokteran Universitas Sumatera Utara.
- Nisar, M. U., Asad, A., Waqas, A., Ali, N., Nisar, A., Qayyum, M., Maryam, H., Javaid, M. and Jamil, M. (2015). Association of Diabetic Neuropathy with Duration of Type 2 Diabetes and Glycemic Control. *Cureus*, 7(8), e302.
- Obrosova, I., Ilnytska, O., Lyzogubov, V., Pavlov, I., Mashtalir, N., Nadler, J. and Drel, V. (2007). High-Fat Diet Induced Neuropathy of Pre-Diabetes and Obesity: Effects of "Healthy" Diet and Aldose Reductase Inhibition. *Diabetes*, 56(10), pp.2598-2608.2
- Ozougwu, J., Obimba, K.C., Belonwu, C.D. and Unakalamba, C.B. (2013). The pathogenesis and pathophysiology of type 1 and type 2 diabetes mellitus. *Academic Journals*, 4, pp.46-57.
- Perkeni. (2015). Pengelolaan dan Pencegahan Diabetes Melitus Tipe 2 di Indonesia [online] . Available at: <https://pbperkeni.or.id/wp-content/uploads/2019/01/4.-Konsensus-Pengelolaan-dan-Pencegahan-Diabetes-melitus-tipe-2-di-Indonesia-PERKENI-2015.pdf> [Accessed: 13 May 2019].
- Perkins, B., Olaleye, D., Zinman, B. and Bril, V. (2001). Simple Screening Tests for Peripheral Neuropathy in the Diabetes Clinic. *Diabetes Care*, 24(2), pp.250-256.
- Pop-Busui, R., Boulton, A., Feldman, E., Bril, V., Freeman, R., Malik, R., Sosenko, J. and Ziegler, D. (2016). Diabetic Neuropathy: A Position Statement by the American Diabetes Association. *Diabetes Care*, 40(1), pp.136-154.
- Prawira, D.A. (2013). Hubungan Durasi Diabetes Melitus Tipe 2 dengan Kejadian Diabetik Neuropati di Poliklinik Penyakit Dalam Rumah Sakit Muhammadiyah Palembang. Skripsi S1. Universitas Muhammadiyah Palembang.
- Quasthoff, S., & Hartung, H. P. (2002). Chemotherapy-induced peripheral neuropathy. *Journal of Neurology*, 249(1), pp.9–17.
- Sa'adah, Nur., (2016). Hubungan keyakinan kemampuan diri (self-efficacy) terhadap perilaku perawatan kaki pada pasien diabetes melitus. Thesis. Fakultas Kedokteran dan Ilmu Kesehatan Universitas Muhammadiyah Yogyakarta.

- Scheen, A. (2003). Pathophysiology Of Type 2 Diabetes. *Acta Clinica Belgica*, 58(6), pp.335-341.
- Scott, R., Fall, T., Pasko, D., Barker, A., Sharp, S., Arriola, L., Balkau, B., Barricarte, A., Barroso, I., Boeing, H., Clavel-Chapelon, F., Crowe, F., Dekker, J., Fagherazzi, G., Ferrannini, E., Forouhi, N., Franks, P., Gavrila, D., Giedraitis, V., Grioni, S., Groop, L., Kaaks, R., Key, T., Kuhn, T., Lotta, L., Nilsson, P., Overvad, K., Palli, D., Panico, S., Quiros, J., Rolandsson, O., Roswall, N., Sacerdote, C., Sala, N., Sanchez, M., Schulze, M., Siddiq, A., Slimani, N., Sluijs, I., Spijkerman, A., Tjonneland, A., Tumino, R., van der A, D., Yaghootkar, H., McCarthy, M., Semple, R., Riboli, E., Walker, M., Ingelsson, E., Frayling, T., Savage, D., Langenberg, C. and Wareham, N. (2014). Common Genetic Variants Highlight the Role of Insulin Resistance and Body Fat Distribution in Type 2 Diabetes, Independent of Obesity. *Diabetes*, 63(12), pp.4378-4387.
- Sherwood, L. (2014). Fisiologi Manusia dari Sel ke Sistem (Edisi 8). Jakarta: *EGC*. pp.697-826,
- Smeltzer, S. C., Bare, B. G., Hinkle, J. and Cheever, K. H. (2010). Brunner and Suddarth's: Textbook of medical-surgical nursing (12th ed.). Philadelphia :Lippincott Williams & Walkins.
- Soewondo, P., Soegondo, S., Suastika, K., Pranoto, A., Soeatmadji, D.W. and Tjokroprawiro, A. (2010). The DiabCare Asia 2008 study - Outcomes on control and complications of type 2 diabetic patients in Indonesia. *Med J Indones*. 19, pp.235-244.
- Soheilykhah, S., Rashidi, M., Dehghan, F., Shojaaddiny, A. and Rahimi-Saghand, S. (2013). Prevalence of Peripheral Neuropathy in Diabetic Patients. *Iranian Journal Of Diabetes And Obesity*, 5(3), pp.107-113.
- Stino, A. and Smith, A. (2017). Peripheral neuropathy in prediabetes and the metabolic syndrome. *Journal of Diabetes Investigation*, 8(5), pp.646-655.
- Suastika, K., Dwipayana, P., Siswadi, M., and Tuty, R. (2012). Age is an Important Risk Factor for Type 2 Diabetes Mellitus and Cardiovascular Diseases. *Glucose Tolerance* [online]. Available at : <https://www.intechopen.com/books/glucose-tolerance/age-is-an-important-risk-factor-for-type-2-diabetes-mellitus-and-cardiovascular-diseases> [Accessed 13 May 2019].
- Suri, M. H., Haddani, H., and Sinulingga, S. (2015). Hubungan Karakteristik, Hiperglikemi, dan Kerusakan Saraf Pasien Neuropati Diabetik di RSMH Palembang Periode 1 Januari 2013 Sampai Dengan 30 November 2014. *Jurnal Kedokteran Dan Kesehatan*, 2(3), pp.305-310.
- Tesfaye, S. and Selvarajah, D. (2012). Advances in the epidemiology, pathogenesis and management of diabetic peripheral neuropathy. *Diabetes/Metabolism Research and Reviews*, 28, pp.8-14.

- Tomic, M., Poljicanin, T., Pavlic-Renar, I. (2003). Obesity A Risk Factor for Microvascular and Neuropathic Complications in Diabetes?. Croatia: Diabetologia Croatia.
- Valensi, P., Giroux, C., Seeboth-Ghalayini, B. and Attali, J. (1997). Diabetic peripheral neuropathy: Effects of age, duration of diabetes, glycemic control, and vascular factors. *Journal of Diabetes and its Complications*, 11(1), pp.27-34.
- van Schie, C., Vermigli, C., Carrington, A. and Boulton, A. (2004). Muscle Weakness and Foot Deformities in Diabetes: Relationship to neuropathy and foot ulceration in Caucasian diabetic men. *Diabetes Care*, 27(7), pp.1668-1673.
- Wannamethee, S., Shaper, A. and Perry, I. (2001). Smoking as a Modifiable Risk Factor for Type 2 Diabetes in Middle-Aged Men. *Diabetes Care*, 24(9), pp.1590-1595.
- Wheat, L. (1980). Infection and Diabetes Mellitus. *Diabetes Care*, 3(1), pp.187-197.
- White, L. (2008). Screening of the Diabetic Foot How to use of a 10g Monofilament [online]. Available at: [https://www.northdevonhealth.nhs.uk/wp-content/uploads/2014/06/how\\_to\\_use\\_a\\_10\\_monofilament.pdf](https://www.northdevonhealth.nhs.uk/wp-content/uploads/2014/06/how_to_use_a_10_monofilament.pdf) [Accessed 15 June 2019].
- WHO (2018). Diabetes. [online] Available at: <https://www.who.int/news-room/fact-sheets/detail/diabetes> [Accessed 12 May 2019].
- Windasari, N. N., (2015). Pendidikan kesehatan dalam meningkatkan kepatuhan merawat kaki pada pasien diabetes melitus tipe II. *Muhammadiyah journal of nursing*, 2.1(79)
- Yang, W., Lu, J., Weng, J., Jia, Weiping., Ji, Linong., Xiao, Jianzhong., Shan, Zhongyan., Liu, Jie., Tian, Haoming., Ji, Qiuhe., Zhu, Dalong., Ge, Jiapu., Lin, Lixiang., Chen, Li., Guo, Xiaohui., Zhao, Zhigang., Li, Qiang., Zhou, Zhiguang., Shan, Guangliang. and He, Jiang. (2010). Prevalence of Diabetes among Men and Women in China. *New England of Journal Medicine*, 362(25), pp.2425 – 2426 [online]. Available at: <https://www.nejm.org/doi/full/10.1056/NEJMoa0908292> [ Accessed 13 May 2019]