



## EDITORIAL BOARD

sains malaysiana

### ✦ Editor-In-Chief

**Rusli Daik**

### ✦ Editors

Bioscience and Biotechnology  
**Endom Ismail/ Roohaida Othman/ Mohd Firdaus Mohd Raih/ Noor Liyana Sukiran**

Chemistry

Engineering Geology  
**Goh Thian Lai**

Environmental Science and Natural Resources  
**Choong Chee Yen/ Mohd Talib Latif/ Wee Suk Ling/ Marlia Mohd Hanafiah**

Earth Science  
**Norbert Simon**

Food Science and Nutrition  
**Wan Aida Wan Mustapha**

Health Science  
**Siti Balkis Budin/ Nur Azlina Mohd Fahami/ Dalia Abdullah/ Haliza Katas/ Dayang Fredalina Basri**

Materials Science  
**Chia Chin Hua/ Yap Chi Chin**

Mathematics  
**Anuar Mohd Ishak**

Physics

Statistics  
**Noriszura Ismail**

### ✦ International Advisory Board

Biodiversity  
**Abdul Latiff Mohamad**  
Universiti Kebangsaan Malaysia, Malaysia

Environmental Science  
**P. Brimblecombe**  
University of East Anglia, UK

Materials Science  
**M. P. Laborie**  
University of Freiburg, Germany

Materials Science  
**D. Hui**  
University of New Orleans, USA

Materials Chemistry  
**M. Oyama**  
Kyoto University, Japan

Mathematics  
**I. Pop**  
University of Cluj, Romania

Physics  
**P. Quentin**  
France

Health Sciences  
**K.L. Tucker**  
Northeastern University, USA

Food Science and Technology  
**L. Wicker**  
USA

## CONTENT, ABSTRACT AND REFERENCES

sains malaysiana

❖ Volume 49 ♦ Number 6 ♦ June 2020	Page
Phenomenon of Macrophyte Differentiation in a Small Lake <i>Sender Joanna, Danuta Urban &amp; Różańska-Boczula Monika</i>   Abstract and References   Full Text PDF (1.42MB)	1209 - 1222
Breeding and Evaluation of <i>Pleurotus giganteus</i> (Berk.) Karunarathna & K.D. Hyde Hybrids via Intraspecific Mating <i>Ha Kwan Yun, Chua Su Boon, Tan Yee Shin &amp; Vikineswary Sabaratnam</i>   Abstract and References   Full Text PDF (880KB)	1223 - 1236
Effect of <i>Lactobacillus casei</i> and <i>Lactobacillus acidophilus</i> in Laying Hens Challenged by <i>Escherichia coli</i> Infection <i>Widya Paramita Lokapirnasari, Adriana Monica Sahidu, Lilik Maslachah, Emy Koestanti Sabdongingrum &amp; Andreas Berny Yulianto</i>   Abstract and References   Full Text PDF (512KB)	1237 - 1244
Identification and Analysis of microRNAs Responsive to Abscisic Acid and Methyl Jasmonate Treatments in <i>Persicaria minor</i> <i>Abdul Fatah A. Samad, Nazaruddin Nazaruddin, Jaeyres Jani &amp; Ismanizan Ismail</i>   Abstract and References   Full Text PDF (1.91MB)	1245 - 1272
Ameliorative Effect of Apple Pomace and Mango Peels against Hyperlipidemia and Lipid Peroxidation Induced by Hyperlipidemic Diet <i>Huma Umbreen, Muhammad Umair Arshad, Razia Noreen &amp; Kiran Aftab</i>   Abstract and References   Full Text PDF (864KB)	1273 - 1282
Effect of Ethanol and Methanol on the Total Phenolic Content and Antioxidant Capacity of Chia Seeds ( <i>Salvia hispanica</i> L.) <i>Morales-Olán Gema, Rojas-López Marlon, Díaz-Reyes Joel, Rosas-Cárdenas Flor de Fátima &amp; Luna-Suárez Silvia</i>   Abstract and References   Full Text PDF (1.17MB)	1283 - 1292
Antimicrobial and Antioxidant Activities in 'Beluntas' ( <i>Pluchea indica</i> ), Turmeric ( <i>Curcuma longa</i> ) and Their Mixtures <i>'Atiqah Ruqayyah Zabidi, Mohd Azmi Mohd Razif, Sayyidah Nafisah Ismail, Muhammad Widus Sempo &amp; Nazariyah Yahaya</i>   Abstract and References   Full Text PDF (840KB)	1293 - 1302
Anti-inflammatory Effect of Tamarind Seed Coat Extract against LPS-Induced	1303 - 1312

## RAW264.7 Macrophages

*Jirapa Puntarut, Wipawadee Sianglum, Supita Tanasawet, Pennapa Chonpathompikunlert & Wanida Sukketsiri*

| Abstract and References | Full Text PDF (1MB) |

Trihoney Suppresses Soluble Adhesion Molecules (ICAM-1 and VCAM-1 in Hypercholesterolemic Atherosclerotic Rabbits: A Comparative Study with Atorvastatin

1313 - 1322

*Hamad Abdul Salam Hamad Alfarisi, Muhammad Ibrahim, Zenab b. Hamad Mohamamed, Asmah Hanim Hamdan & Che Anuar Che Mohamad*

| Abstract and References | Full Text PDF (712KB) |

Effect of Neuro-Dynamic Technique on Repetitive Inward Ankle Rolls among Young Malaysian Athletes- A Randomized Controlled Trial

1323 - 1332

*Vinodhkumar Ramalingam, Viswanath Sundar & Saju Joseph*

| Abstract and References | Full Text PDF (1.13MB) |

Fenitrothion Impaired Sexual Behaviour and Reproductive Performance in Male Sprague-Dawley Rats

1333 - 1344

*Nur Afizah Yusoff, Mahaneem Mohamed, Siti Balkis Budin & Izatus Shima Taib*

| Abstract and References | Full Text PDF (680KB) |

Disseminated Peritoneal Leiomyomatosis After 10 Years Post TAHBSO: A Case Report

1345 - 1350

*Seyed Jalaledin Tahsildar Tehrani, Thean Yean Kew & Mohd Amir Kamaruzzaman*

| Abstract and References | Full Text PDF (528KB) |

Role of Novel Therapeutic Agents in Modulating Invadopodia Formation in Metastatic Breast Cancer

1351 - 1358

*Siti Nor Aini Harun, Nurul Akmaryanti Abdullah, Noraina Muhamad Zakuan, Hafizah Abdul Hamid, Muhammad Zulfadli Mehat & Nur Fariesha Md Hashim*

| Abstract and References | Full Text PDF (412KB) |

Zerumbone Induces Cytotoxicity and Inhibits Cell Migration of Human Colon Cancer Cells

1359 - 1370

*Tan Min Jien, Siti Nur Parvin Ab Hamid, Nur Fariesha Md Hashim, Nurdin Armania, Hasni Idayu Saidi & Noraina Muhamad Zakuan*

| Abstract and References | Full Text PDF (2.57MB) |

Vonoprazan and Proton Pump Inhibitors in *Helicobacter pylori* Eradication Therapy: A Systematic Review

1371 - 1380

*Mohd Fahmi Nur Azlina & Che Roos Nur Aishah*

| Abstract and References | Full Text PDF (644KB) |

The Use of Fall Risk-Increasing Drugs among Older People in Nursing Homes

1381 - 1388

*Lee Shiow Ling, Marhanis Omar, Adliah Mhd-Ali & Mohd Makmor-Bakry*

| Abstract and References | Full Text PDF (600KB) |

Enhancing Effect of Vitexin on Osteogenic Activity of Murine Pre-Osteoblastic MC3T3-E1 Cells

1389 - 1400

*Xiaohan Yuan, Haiyan Han, Zhaohui Luo, Qiuxue Wang, Peijia Tang, Zhihui Zhang, Yujie Fu & Chengbo Gu*

| Abstract and References | Full Text PDF (976KB) |

The Differences between the Expression Levels of *axe-txe* Genes in Chloramphenicol-Sensitive and Penicillin-Resistant *Enterococcus faecium*

1401 - 1410

## Isolates

*Sri Indra Wahyuni Mohd Irmal, Suresh Kumar Subbiah, Vasantha Kumari Neela, Niazlin Mohd Taib, Azmiza Syawani Jasni, Maha Abdullah, Mohamad Khairil*

*Radzali & Rukman Awang Hamat*

| Abstract and References | Full Text PDF (800KB) |

Bone Density and Marginal Bone Loss around Implants Post Ultraviolet A and Ultraviolet C Irradiation

1411 - 1420

*Naauman Zaheer, Zainul Ahmad bin Rajion, Maliha Shahbaz, Usman Zaheer, Muhammad Qasim Saeed & Noor Hayati Abdul Razak*

| Abstract and References | Full Text PDF (716KB) |

Cytotoxic Effects of Organotin(IV) Dithiocarbamate Compounds with Different Functional Groups on Leukemic Cell Line, K-562

1421 - 1430

*Asmah Hamid, Mohd Azam Azmi, Nor Fadilah Rajab, Normah Awang & Nurul Farhana Jufri*

| Abstract and References | Full Text PDF (1MB) |

Accuracy of Contact Lens Method by Spherical and Aspheric Rigid Gas Permeable Lenses on Corneal Power Determination in Normal Eyes

1431 - 1437

*Md Muziman Syah Md Mustafa, Haliza Abdul Mutalib, Noorhazayti Ab. Halim & Mohd Radzi Hilmi*

| Abstract and References | Full Text PDF (732KB) |

Determination of the Radiological Risk from the Natural Radioactivity in Irrigation at Selected Areas of Peninsular Malaysia

1439 - 1450

*Khoirul Solehah Abdul Rahim, Zalita Zainuddin, Mohd Idzat Idris, Wahmisari Priharti & Murtadha Sh. Aswood*

| Abstract and References | Full Text PDF (852KB) |

*Ab-initio* Calculations of the Structural, Electronic and Optical Properties of (Cdse)<sub>2</sub> Clusters

1451 - 1460

*A.I.A. Alselaawe, MHH Jumali, G. Gopir & M.M. Anas*

| Abstract and References | Full Text PDF (1.25MB) |

Chemical Bonding Composition and Growth Mechanism of a-CN<sub>x</sub> Thin Films by Low-Temperature rf-PECVD Technique (Komposisi Ikatan Kimia dan Mekanisme Pertumbuhan Filem Nipis a-CN<sub>x</sub> oleh Teknik rf-PECVD Suhu Rendah)

1461 - 1470

*Siti Aisyah Abd Aziz & Rozidawati Awang*

| Abstract and References | Full Text PDF (556KB) |

A Security Upgrade on the GGH Lattice-based Cryptosystem

1471 - 1478

*Arif Mandangan, Hailiza Kamarulhaili & Muhammad Asyraf Asbullah*

| Abstract and References | Full Text PDF (0.99MB) |

[Home](#)

[Editorial Board](#)

[Guide for Authors](#)

[Contents, Abstract and References](#)

[Malay](#)

Copyright Reserved © 2007 Sains Malaysiana

## Effect of *Lactobacillus casei* and *Lactobacillus acidophilus* in Laying Hens Challenged by *Escherichia coli* Infection

(Kesan *Lactobacillus casei* dan *Lactobacillus acidophilus* dalam Ayam Bertelur yang Dijangkiti *Escherichia coli*)

WIDYA PARAMITA LOKAPIRNASARI\*, ADRIANA MONICA SAHIDU, LILIK MASLACHAH, EMY KOESTANTI  
SABDONINGRUM & ANDREAS BERNY YULIANTO

### ABSTRACT

This study aimed to prove the potential of *Lactobacillus casei* and *Lactobacillus acidophilus* probiotics as alternative substitutes of antibiotic growth promoters in laying hens challenged by *Escherichia coli* infection in order to enhance their growth performance and hen day production. The study used a total of 120 laying hens aged 25 weeks, divided into 3×2 treatments with each comprising 4 replications, and each replication consisted of 5 hens. The study used a completely randomised factorial design; factor a was the feed additive (control, antibiotics growth promoters /AGP, probiotic), whereas factor b was the *E. coli* infection (non-infection and *E. coli* infection). The results showed that there were significant differences ( $p < 0.05$ ) between the treatment of feed additive (factor a), and *E. coli* infection (factor b), and interaction ( $p < 0.05$ ) between the feed and the infection for the egg weight, hen day production, feed conversion ratio and feed efficiency. The probiotic use of 0.5% *L. casei* + 0.5% *L. acidophilus* in hens either infected or uninfected with *E. coli* still produced the highest egg weight, hen day production, feed efficiency and reduced feed conversion ratio compared to all treatments. Based on the results, it can be concluded that the use of probiotics 0.5% *L. casei* and 0.5% *L. acidophilus* act as alternative substitutes for antibiotic growth promoters in laying hens challenged by *E. coli* infection.

**Keywords:** *E. coli*; growth performance; hen day production; *Lactobacillus acidophilus*; *Lactobacillus casei*

### ABSTRAK

Kajian ini bertujuan membuktikan potensi probiotik *Lactobacillus casei* dan *Lactobacillus acidophilus* sebagai pengganti alternatif penggalak pertumbuhan antibiotik (AGP) dalam ayam bertelur yang dijangkiti *Escherichia coli* terhadap prestasi pertumbuhan dan pengeluaran telur harian. Kajian ini menggunakan 120 ekor ayam bertelur berumur 25 minggu yang dibahagikan kepada rawatan 3×2, dengan setiap rawatan terdiri daripada 4 replikasi dan setiap replikasi terdiri daripada 5 ekor ayam bertelur. Kajian ini menggunakan reka bentuk faktor rawak lengkap; faktor a adalah makanan tambahan (kawalan, AGP, probiotik), manakala faktor b adalah faktor jangkitan *E. coli* (bukan jangkitan dan jangkitan *E. coli*). Hasil kajian menunjukkan terdapat perbezaan yang signifikan ( $p < 0.05$ ) antara rawatan aditif makanan (faktor a) dan rawatan jangkitan *E. coli* (faktor b), serta interaksi antara faktor makanan tambahan dan jangkitan pada berat telur, pengeluaran telur harian, nisbah penukaran makanan dan kecekapan suapan. Penggunaan probiotik 0.5% *L. casei* + 0.5% *L. acidophilus* dalam ayam sama ada dijangkiti atau tidak dijangkiti *E. coli* masih memberikan berat telur tertinggi, pengeluaran telur harian dan kecekapan suapan yang tertinggi dan menurunkan nisbah penukaran makanan berbanding semua rawatan. Berdasarkan hasil kajian, dapat disimpulkan bahawa penggunaan probiotik 0.5% *L. casei* + 0.5% *L. acidophilus* dapat digunakan sebagai pengganti alternatif bagi promoter pertumbuhan antibiotik dalam ayam petelur yang dijangkiti *E. coli*.

**Kata kunci:** *E. coli*; *Lactobacillus acidophilus*; *Lactobacillus casei*; pengeluaran telur harian; prestasi pertumbuhan

### INTRODUCTION

Avian Pathogenic *Escherichia coli* (APEC) is an extra intestinal bacterium responsible for poultry colibacillosis. It affects chicken of all age groups leading to an increase in morbidity and mortality rate. In hens and broilers, pathogenic *E. coli* could generate numerous systemic infections that can cause pericarditis, perihepatitis,

salpingitis, salpingoperitonitis, colisepticaemia and airsacculitis infections, thereby, leading to decreases in egg production and economic losses (Dou et al. 2016; Paixao et al. 2016; Paudel et al. 2016). There are two forms of colibacillosis in laying hens, the salpingitis/peritonitis/salpingoperitonitis (SPS) and the *E. coli* peritonitis syndrome (EPS) (Landman & van Eck 2017).

The pathogenic and commensal *E. coli* strains colonise the mammalian intestine by interacting metabolically and physically with other microbiota in the mucus layer of the cecal and colonic epithelium (Mokrzycki et al. 2018). The stressor factor faced is the level of acidity in the stomach and its ability to survive in these conditions, owing to its stationary phase. In this condition, the bacteria are able to impede the system resistance or protection against acidity (Conway & Cohen 2015; Foster 2004) and after reaching the colon, they use nutrients to exit lag phase and grow. The failure in transition from the lag to logarithmic phase leads to the elimination of *E. coli* bacteria (Conway & Cohen 2015).

The use of unsuitable antibiotic growth promoters (AGP) in poultry farming induces bacteria resistant to antibiotics leading to the accumulation of antibiotic residues in the poultry products. This poses a threat to those consuming these poultry products, hence, many countries have prohibited to use them in poultry farms. Probiotics are considered as natural feed additives which can have a positive effect similar to AGP. Its supplementation in poultry feed has been reported to enhance growth performance (Awad et al. 2009; Tang et al. 2017), nutritional retention and caecal microbial balance (Mookiah et al. 2014; Mountzouris et al. 2010).

Probiotics are classified as the zootechnical feed additives and can be in the form of enzymes or microorganisms. They are non-pathogenic living microbes capable of improving the health of its host by increasing the microbial balance in intestine (Anadón 2006; Fuller 2001). The probiotic lowers the risk of gastro-intestinal diseases by stimulating the growth of beneficial microbes, improve immune modulatory, modulate the activity of epithelial cells, dendritic cells, natural killers, and the enhancement of nutrients bioavailability (Chiang & Pan 2012; Untoo et al. 2018; Yahfoufi et al. 2018). Probiotics could protect host against pathogens by colonisation in the gastrointestinal tract (Getachew 2016). Probiotics can be used as an alternative substitute for antibiotic growth promoters, owing to its beneficial effect on the health of its host in chicken's productivity, serum lipids and the intestinal morphology (Khaksefidi & Ghoorchi 2006; Nayeppor et al. 2007; Willis et al. 2007). The probiotic functions in disease prevention can be through competitively excluding gut pathogens, modulating gastrointestinal immune responses and producing metabolites such as bacteriocins that inhibit or kill pathogenic bacteria. For supporting growth performance, probiotics can counteract dysbiosis, maintain and replenish normal microorganism balance, and contribute in optimising nutrient absorption (Park et al. 2016).

There is still a limited number of research that has been conducted on the use of probiotics with the combination of *L. casei* and *L. acidophilus* in laying hens. The aim of this study was to determine the potential of *L. casei* and *L. acidophilus* probiotic as an alternative substitute of antibiotic growth promoters in laying hens

challenged by *E. coli* infection on egg weight, hen day production, feed conversion ratio and feed efficiency.

## MATERIALS AND METHODS

### ETHICAL APPROVAL

An approval from the Faculty of Veterinary Medicine Universitas Airlangga was obtained before the experimental trial, No: 1.KE.166.09.2018. The isolates were obtained from collections belonging to Widya Paramita Lokapirnasari, Animal Husbandry Department, Faculty of Veterinary Medicine, Universitas Airlangga. The source of laying hens (ISA 2019) was from commercial farm.

### EXPERIMENTAL DESIGN

The experimental design used was a factorial Completely Randomised Design (CRD). In the first factor, feed additives, consisted of 3 subfactors namely control, AGP and probiotic were utilised, while the second factor (*E. coli* infection), consisted of 2 subfactors namely the control treatment for hens not infected with *E. coli* and those infected. The isolates were obtained from the collection belonging to Emy Koestanti Sabdoningrum, Animal Husbandry Department, Faculty of Veterinary Medicine, Universitas Airlangga). The isolates *L. casei* and *L. acidophilus* were obtained from collections belonging to Widya Paramita Lokapirnasari, Animal Husbandry Department, Faculty of Veterinary Medicine, Universitas Airlangga (Yulianto & Lokapirnasari 2018).

### LAYERS AND HOUSING

ISA Brown laying hens (120) aged 25 weeks' old obtained from a commercial farm were used. They were randomly divided into 3×2 groups, with each group consisted of 4 replications and 5 layer chickens and placed in individual battery cages with a size of 20 × 35 × 35 cm with *ad libitum* drinking water and feeding. The climatic conditions and lighting programs were operated following the recommendations of Institut de Sélection Animale BV of ISA Brown throughout the experiment (ISA 2019).

### FEED, PROBIOTIC AND *E. COLI* TREATMENTS

The feed was given twice a day as much as 150 g/hen/day. Laying hens were fed a commercial layer diet with the following specifications: dry matter: 91.97%, ash: 9.28%, crude protein: 20.71%, extract ether 6.36%, crude fiber: 7.43%; nitrogen free extract (NFE): 48.18% and energy metabolism (EM:2938.60 Mcal/kg). Drinking water was given *ad libitum* during the treatment.

The induced *E. coli* bacteria containing  $1 \times 10^8$  CFU/mL *E. coli* was given orally through a disposable sterile syringe at  $10^8$  CFU/mL/hen to 28 weeks' old laying hens. Then, the clinical symptoms such as diarrhoea and decreased feed intake were observed.

The AGP treatment at 0.01% of feed was given every day for 4 weeks to laying hens at 26 to 30 weeks old. The probiotic administration contained 0.5% *L. casei* and 0.5% *L. acidophilus* at  $4.77 \times 10^8$  CFU/mL was given orally through drinking water. The probiotic administration was given for 4 weeks starting from 26 to 30 weeks old. Production performance data were carried out starting from the treatment of *E. coli* induction till the end of the study (28 to 30 weeks old). The egg weight, hen day production, Feed Conversion Ratio (FCR) and feed efficiency were obtained to analyse growth performance. For egg weight, the eggs were collected daily and the weight was recorded to calculate the mean egg weight using an electronic digital balance. FCR was calculated by dividing the feed intake by egg production. Feed intake was determined by subtracting the remaining feed from the original amount of feed at the end of each week. Feed efficiency (%) was calculated by dividing the egg production by feed intake and multiplying by 100. Hen day production was calculated by dividing number of eggs per day by number of laying hens per day and multiplying by 100 (Lokapirnasari et al. 2019a; Rattanawut et al. 2018). Eggs were extracted from 120 hens (3×2 treatments, 4 replications and each replication consisted of 5 laying hens) and weighed daily from the 26<sup>th</sup> to 30<sup>th</sup> week. The treatments used were as follow:

Control treatment (T0): negative control (not treated with AGP or probiotics and chicken not infected with *E. coli*); Treatment 1 (T1): 0.1% AGP and chicken not

infected with *E. coli*; Treatment 2 (T2): 0.5% *L. casei* and 0.5% *L. acidophilus* and chicken not infected with *E. coli*; Treatment 3 (T3): positive control (not given antibiotics or probiotics and chicken infected with *E. coli*); Treatment 4 (T4): 0.1% AGP and chicken infected with *E. coli*; and Treatment 5 (T5): 0.5% probiotics *L. casei* and 0.5% *L. acidophilus* and chicken infected with *E. coli*.

#### STATISTICAL ANALYSIS

Data of egg weight, hen day production, FCR and feed efficiency were compared between treatment groups using univariate general linear model statistics. If the results obtained were significantly different ( $p < 0.05$ ), it would be followed by subsequent Duncan multiple range tests.

#### RESULTS AND DISCUSSION

##### EGG WEIGHT

The results of the statistical analysis showed significant differences ( $p < 0.05$ ) between treatment factors a (feed additives) with the use of additive feeds on the egg weight. The treatment of factor b (*E. coli* infection) also showed significant differences ( $p < 0.05$ ) between the infected chicken and *E. coli* infection against egg weight. Between factors a and b, there was an interaction ( $p < 0.05$ ) on the additive feed on the egg weight (Table 1).

TABLE 1. Average egg weight for different treatments

Factor a × factor b	Control without feed additive (b0)	0.1% AGP (b1)	Probiotic (b2)
Non infected (a0)	58.500 <sup>b</sup> ± 0.01	58.900 <sup>c</sup> ± 0.01	61.225 <sup>e</sup> ± 0.09
Infected (a1)	58.300 <sup>a</sup> ± 0.01	58.300 <sup>a</sup> ± 0.01	60.925 <sup>d</sup> ± 0.09

a, b, c, d, e The different superscripts showed the significant differences ( $p < 0.05$ ) among the treatments

The lowest egg weight was produced in the control treatment i.e. without AGP administration and probiotics in *E. coli* infected chickens. No difference was noted with the treatment of AGP in *E. coli*-infected laying hens. The highest egg weight was found in the treatment of probiotics *L. casei*, and *L. acidophilus* in laying hens not infected with *E. coli* with the value of 61.225 (g/egg). The size of the eggs produced were in the range of 58.30 to 61.225 g. This result was supported by other studies in which the probiotic containing *L. casei*, and *L. rhamnosus* at  $1.2 \times 10^8$  CFU/g could increase the egg weight, egg length and egg width of quail eggs (Lokapirnasari et al. 2019b).

Similarly, the weight of chicken eggs treated with the probiotics, *L. casei*, and *L. acidophilus* in this research gave higher results than Tang et al. (2017), where the use of probiotic increased the egg weight in laying hens to 55.07 g compared to control (37.68 g) with medium size at 20 to 36 weeks. The size distribution of small (42.50 to 49.59 g), medium (49.60- to 6.69 g), large (56.70 to 63.78 g), extra large (63.79 to 70.87 g) and jumbo size eggs (> 70.88 g) was determined based on specified weights according to the United States Department of Agriculture (USDA) (Tang et al. 2017). It was shown that the administration of probiotics, *L. casei*, and *L.*

*acidophilus*, gave favourable results as they did not lead to a decrease in egg weight despite the fact that the chicken was infected with *E. coli*.

#### HEN DAY PRODUCTION

The Hen Day Production (HDP) is a comparison of the number of eggs produced with the number of laying

hens multiplied by 100%. The statistical analysis results showed significant differences ( $p < 0.05$ ) between treatment factors a using HDP feed additives. The treatment of factor b also showed significant differences ( $p < 0.05$ ) between the infected chicken and *E. coli* infection against HDP (Table 2).

TABLE 2. Average hen day production (HDP) for different treatments

Factor a × factor b	Control without feed additive (b0)	0.1% AGP (b1)	Probiotic (b2)
Non infected (a0)	94.700 <sup>d</sup> ± 0.05	93.300 <sup>c</sup> ± 0.05	97.000 <sup>e</sup> ± 1.288
Infected (a1)	69.300 <sup>a</sup> ± 0.05	89.300 <sup>b</sup> ± 0.05	96.000 <sup>c</sup> ± 1.061

<sup>a, b, c, d, e</sup> The different superscripts showed the significant differences ( $p < 0.05$ ) among the treatments

The results showed that there were interactions between the probiotics, *L. casei* and *L. acidophilus* or AGP with

*E. coli* infection. The lowest HDP results were found in the positive control treatment, i.e. without AGP or probiotics and *E. coli* infection which was 69.3%. The egg production in chicken treated with AGP and *E. coli* infection was higher (89.3%) than the others. This indicated that hens not treated with AGP and *E. coli* infection tended to show a decrease in egg production. Previous study had shown that the use of probiotics resulted in an increase of egg production at 76.51% compared to control (69.29%) (Tang et al. 2017). The HDP obtained as an effect of the use of probiotics, *L. casei* and *L. acidophilus*, showed a higher result. The highest HDP was found in the treatment of probiotics, *L. casei* and *L. acidophilus*, in laying hens not infected with *E. coli* (97.00%) and not significantly different from the treatment of probiotics in *E. coli* infected chicken (96.00%). The results indicated that when the probiotics were given to laying hens, no decrease in HDP was observed even though the chicken were infected with *E. coli*. The treatment of AGP in *E. coli*-infected laying hens showed a decrease in HDP of 4.28% compared to non-infected laying hens. In a positive control treatment where there was no addition of AGP or probiotics and the presence of *E. coli* infection, a decrease in HDP by 26.82% was observed compared to those not infected with *E. coli*. *E. coli* infection

reduces the percentage of egg production (Landman & van Eck 2017), but the use of probiotics, *L. casei* and *L. acidophilus*, were able to inhibit pathogenic *E. coli* hence not affecting the decrease in the host health since the probiotic strains inhibit the growth and colonisation of other pathogenic microorganisms (Shokryazdan et al. 2014). Lactic acid produced by *Lactobacillus* causes the environment to become acidic compared to its ability to inhibit the growth of pathogenic bacteria (Bernardeau et al. 2008). This result was in line with Raka et al. (2014) who reported that supplementing Liquid Probiotics Mixed Culture (LPMC) with *Lactobacillus* and *Bacillus* species showed the highest egg weight and hen day production in layers' chicken.

#### FEED CONVERSION RATIO

Feed Conversion Ratio (FCR) is the ratio between the amounts of feed consumed and the weight of the eggs produced. The statistical analysis results showed significant differences ( $p < 0.05$ ) between the treatment in factors a, using additive feeds on the FCR. The treatment of factor b also showed significant differences ( $p < 0.05$ ) between infected chicken and *E. coli* infection against FCR, with an interaction ( $p < 0.05$ ) between the factors of infection and the factor feed additive (Table 3).

TABLE 3. Average of feed conversion ratio (FCR) for different treatments

Factor a × factor b	Control without feed additive (b0)	0.1% AGP (b1)	Probiotic (b2)
Non infected (a0)	2.100 <sup>c</sup> ± 0.001	2.100 <sup>c</sup> ± 0.001	1.925 <sup>a</sup> ± 0.05
Infected (a1)	2.900 <sup>c</sup> ± 0.001	2.200 <sup>d</sup> ± 0.001	1.975 <sup>b</sup> ± 0.05

<sup>a, b, c, d, e</sup> The different superscripts showed the significant differences ( $p < 0.05$ ) among the treatments



The best feed conversion ratio value was found in the treatment of probiotics, *L. casei* and *L. acidophilus*, in *E. coli* infected chicken. The highest FCR value was produced in the control treatment, i.e. in hens not given AGP or probiotics and infected with *E. coli*. The result was found to be 2.900, followed by the treatment of *E. coli*-infected hens using AGP, which amounted to 2.200. The outcome indicated that the administration of probiotics, *L. casei* and *L. acidophilus*, in hens provided a good feed conversion value compared to AGP, therefore, the probiotics helped in increasing feed consumption, owing to the enhanced digestibility of feed in the digestive tract of the livestock. The enhanced feed consumption has an impact on growth and improves the feed conversion ratio (Bidarkar et al. 2014). The positive effect of *L. casei* and *L. acidophilus* probiotics in feed conversion ratio and feed efficiency might be due to the ability of the probiotics to induce physical changes in the gut structure, particularly in the development of crypt depth ratio and villus height in the ileum (Awad et al. 2008), thus increasing nutrient digestibility (Park et al. 2016).

The administration of *L. casei* and *L. acidophilus* probiotics to hens treated with *E. coli* infection, gave the best FCR value which was 1.925. This is because chickens in healthy conditions do not consume much feed and produce highest number of eggs. It also showed a better FCR value (1.975) than the use of AGP in non-infected and infected hens (2.100) and (2.200), respectively. The FCR value of negative control and the use of AGP gave an FCR value of 2.100. The highest FCR

value was produced in the positive control treatment, i.e. in hens not given AGP or probiotics but infected with *E. coli* which increased to 2.900. The results of the study done by Tang et al. (2017) showed that the use of probiotics improved FCR results (2.32) compared to controls (2.55). Probiotics can help maintain the health of the digestive tract by providing beneficial microbial balance. A healthy and functioning digestive and intestinal tract can reduce digestive disorders resulting in better nutrient utilisation and feed conversion to produce increased growth and productivity (Fuller 2001). The previous results were in agreement with the present experiment (Table 3), that the reduction in FCR by probiotic supplementation related to its promoting positive effects on metabolic processes of nutrient digestion and nutrients utilisation and increasing the host health status. The probiotic supplementation could enrich the diversity of *Lactobacillus* spp. in jejunum and cecum by increasing the abundance and prevalence of *Lactobacillus* spp. inhabiting the intestine, maintaining the natural stability of indigenous microbial and restoring the microbial flora balance (Hayirli et al. 2005).

#### FEED EFFICIENCY

The statistical analysis showed that there were differences between the feed additive factors ( $p < 0.05$ ) on the feed efficiency. The same results were also shown in the infection factors ( $p < 0.05$ ) with the factorial test analysis indicating an interaction between the feed additive and the infection for feed efficiency (Table 4).

TABLE 4. Average feed efficiency for different treatments

Factor a x factor b	Control without feed additive (b0)	0.1% AGP (b1)	Probiotic (b2)
Non infected (a0)	51.3800 <sup>b</sup> ± 0.001	51.7600 <sup>c</sup> ± 0.001	53.8700 <sup>e</sup> ± 0.133
Infected (a1)	51.1400 <sup>a</sup> ± 0.001	51.2800 <sup>b</sup> ± 0.001	53.1975 <sup>d</sup> ± 0.101

<sup>a, b, c, d, e</sup>The different superscripts showed the significant differences ( $p < 0.05$ ) among the treatments

The use of *L. casei* and *L. acidophilus* probiotics was able to inhibit colonisation of *E. coli* to compete in obtaining the nutrients needed. This was supported by several studies where the success of *E. coli* colonisation depends on competition with other micro biota (Khaksefidi & Ghoorchi 2006), penetration of the mucus layer (Møller et al. 2003), and its ability to avoid host defenses (Bergstrom et al. 2012; McGuckin et al. 2011). The use of probiotic strains helps to maintain microbial balance in gastrointestinal tract and make changes in the composition of the intestinal micro flora by increasing beneficial bacteria and suppressing harmful

pathogenic bacteria like *E. coli* (Fuller 2001; Srinu et al. 2013). This is due to the existence of competitive exclusion to compete for nutrition and attachment to the intestinal epithelial wall, or the ability of probiotic strains to produce antimicrobial substances or synergism between the two mechanisms (Mountzouris et al. 2010; Sohail et al. 2011). Other study also reported that probiotics have good effects in improving digestion, absorption and nutrient availability (Endens 2003). The use of probiotics containing *Lactobacilli*, *Bifidobacterium thermophilum* and *Enterococcus faecium* increases the jejunal villus height (Chichlowski et al. 2007), that it can

increase nutrient absorption and be used for growth and production. The supplementation of *L. sporogenes* at 100 mg/kg diet ( $6 \times 10^8$  spores) significantly increased the egg production and feed efficiency (Panda et al. 2008).

The effect of both antibiotics and probiotics on animals seem to be similar. In this study, the effect of giving AGP and probiotics containing *L. casei* and *L. acidophilus* showed the beneficial effect in decreasing FCR, and increasing feed efficiency, hen day production and egg weight. The positive impact of giving antibiotics on laying hens are related to changes of the microbial community in the gastrointestinal tract both in the ileum and the caecum (Choi et al. 2018) especially toward short-chain fatty acid producers (Banerjee et al. 2018), but also to the increase of amino acid metabolites, fatty acids, nucleosides, and vitamins (Gadde et al. 2018). Antibiotics could improve performance of chicken through an anti-inflammatory effect mediated by the intestinal epithelium (Niewold 2007).

#### CONCLUSION

The use of probiotics, 0.5% *L. casei* and 0.5% *L. acidophilus*, can be used as alternative substitutes for antibiotic growth promoters. They can provide better results for the laying hens performance to increase egg production (egg weight, hen day production, feed conversion ratio and feed efficiency) compared to the control and use of AGP, in both infected and uninfected laying chickens with *E. coli*. The use of *L. casei* and *L. acidophilus* probiotics would avoid the use of antibiotics in poultry industry.

#### ACKNOWLEDGEMENTS

The research team is grateful to the Ministry of Research and Technology of Higher Education (*Kemenristekdikti*), Chair of the Research and Innovation Institute, Rector Universitas Airlangga for funding this research (PTUPT 2019).

#### REFERENCES

- Anadón, A. 2006. WS14 The EU ban of antibiotics as feed additives alternatives and consumer safety. *Veterinary Pharmacology and Therapeutics* 29: 41-44.
- Awad, W., Ghareeb, K. & Böhm, J. 2008. Intestinal structure and function of broiler chickens on diets supplemented with a synbiotic containing *Enterococcus faecium* and oligosaccharides. *International Journal of Molecular Sciences* 9(11): 2205-2216.
- Awad, W.A., Ghareeb, K., Abdel-Raheem, S. & Böhm, J. 2009. Effects of dietary inclusion of probiotic and synbiotic on growth performance, organ weights, and intestinal histomorphology of broiler chickens. *Poultry Science* 88(1): 49-56.
- Banerjee, S., Sar, A., Misra, A., Pal, S., Chakraborty, A. & Dam, B. 2018. Increased productivity in poultry birds by sub-lethal dose of antibiotics is arbitrated by selective enrichment of gut microbiota, particularly short chain fatty acid producers. *Microbiology* 164(2): 142-153.
- Bergstrom, K.S., Sham, H.P., Zarepour, M. & Vallance, B.A. 2012. Innate host responses to enteric bacterial pathogens: A balancing act between resistance and tolerance. *Cellular Microbiology* 14(4): 475-484.
- Bernardeau, M., Vernoux, J.P., Dubernet, S.H. & Guéguen M. 2008. Safety assessment of dairy microorganisms: The *Lactobacillus* genus. *International Journal of Food Microbiology* 126: 278-285.
- Bidarkar, V.K., Swain, P.S., Ray, S. & Dominic, G. 2014. Probiotics: Potential alternative to antibiotics in ruminant feeding. *Trends in Veterinary and Animal Sciences* 1: 1-4.
- Chiang, S.S. & Pan, T.M. 2012. Beneficial effects of *Lactobacillus paracasei* subsp. *paracasei* NTU 101 and its fermented products. *Applied Microbiology and Biotechnology* 93(3): 903-916.
- Chichlowski, M., Croom, W.J., Edens, F.W., McBride, B.W., Qiu, R., Chiang, C.C. & Koci, M.D. 2007. Microarchitecture and spatial relationship between bacteria and ileal, cecal, and colonic epithelium in chicks fed a direct-fed microbial, PrimaLac, and salinomycin. *Poultry Science* 86(6): 1121-1132.
- Choi, J.H., Lee, K., Kim, D.W., Kil, D.Y., Kim, G.B. & Cha, C.J. 2018. Influence of dietary avilamycin on ileal and cecal microbiota in broiler chickens. *Poultry Science* 97(3): 970-979.
- Conway, T. & Cohen, P.S. 2015. Commensal and pathogenic *Escherichia coli* metabolism in the gut. *Microbiology Spectrum* 3(3): 1-24.
- Dou, X., Gong, J., Han, X., Xu, M., Shen, H., Zhang, D. & Zou, J. 2016. Characterization of avian pathogenic *Escherichia coli* isolated in Eastern China. *Gene* 576(1): 244-248.
- Edens, F.W. 2003. An alternative for antibiotic use in poultry: Probiotics. *Brazilian Journal of Poultry Science* 5(2): 75-97.
- Foster, J.W. 2004. *Escherichia coli* acid resistance: Tales of an amateur acidophile. *Nature Reviews Microbiology* 2(11): 898.
- Fuller, R. 2001. The chicken gut microflora and probiotic supplements. *The Journal of Poultry Science* 38(3): 189-196.
- Gadde, U.D., Oh, S., Lillehoj, H.S. & Lillehoj, E.P. 2018. Antibiotic growth promoters virginiamycin and bacitracin methylene disalicylate alter the chicken intestinal metabolome. *Scientific Reports* 8(1): 3592.
- Getachew, T. 2016. A review on effects of probiotic supplementation in poultry performance and cholesterol levels of egg and meat. *Journal of World's Poultry Research* 6(1): 31-36.
- Hayirli, A., Esenbuğa, N., Macit, M., Yörük, M.A., Yıldız, A. & Karaca, H. 2005. Nutrition practice to alleviate the adverse effects of stress on laying performance, metabolic profile and egg quality in peak producing hens: II. the probiotic supplementation. *Asian-Australasian Journal of Animal Sciences* 18(12): 1752-1760.
- ISA. 2019. ISA Brown. Cage production systems, management report. <https://www.isa-poultry.com/en/product/isa-brown/>. Accessed on 27 March 2019.

- Khaksefidi, A. & Ghoorchi, T. 2006. Effect of probiotic on performance and immunocompetence in broiler chicks. *The Journal of Poultry Science* 43(3): 296-300.
- Landman, W.J.M. & van Eck, J.H.H. 2017. The efficacy of inactivated *Escherichia coli* autogenous vaccines against the *E. coli* peritonitis syndrome in layers. *Avian Pathology* 46(6): 658-665.
- Lokapimasari, W.P., Pribadi, T.B., Al Arif, A., Soeharsono, S., Hidanah, S., Harijani, N., Najwan, R., Huda, K., Wardhani, H.C.P., Rahman, N.F.N. & Yulianto, A.B. 2019a. Potency of probiotics *Bifidobacterium* spp. and *Lactobacillus casei* to improve growth performance and business analysis in organic laying hens. *Veterinary World* 12(6): 860-867.
- Lokapimasari, W.P., Al Arif, A., Soeharsono, S., Fathinah, A., Najwan, R., Wardhani, H.C.P., Noorrahman, N.F., Huda, K., Ulfah, N. & Yulianto, A.B. 2019b. Improves in external and internal egg quality of Japanese quail (*Coturnix coturnix japonica*) by giving lactic acid bacteria as alternative antibiotic growth promoter. *Iranian Journal of Microbiology* 11(5): 406-411.
- McGuckin, M.A., Lindén, S.K., Sutton, P. & Florin, T.H. 2011. Mucin dynamics and enteric pathogens. *Nature Reviews Microbiology* 9(4): 265-278.
- Mokszycycki, M.E., Leatham-Jensen, M., Steffensen, J.L., Zhang, Y., Krogfelt, K.A., Caldwell, M.E., Conway, T. & Cohen, P.S. 2018. A simple *in vitro* gut model for studying the interaction between *Escherichia coli* and the intestinal commensal microbiota in cecal mucus. *Applied and Environmental Microbiology* 84(24): e02166-18.
- Møller, A.K., Leatham, M.P., Conway, T., Nuijten, P.J., de Haan, L.A., Krogfelt, K.A. & Cohen, P.S. 2003. An *Escherichia coli* MG1655 lipopolysaccharide deep-rough core mutant grows and survives in mouse cecal mucus but fails to colonize the mouse large intestine. *Infection and Immunity* 71(4): 2142-2152.
- Mookiah, S., Sieo, C.C., Ramasamy, K., Abdullah, N. & Ho, Y.W. 2014. Effects of dietary prebiotics, probiotic and synbiotics on performance, caecal bacterial populations and caecal fermentation concentrations of broiler chickens. *Journal of the Science of Food and Agriculture* 94(2): 341-348.
- Mountzouris, K.C., Tsitsrikos, P., Palamidi, I., Arvaniti, A., Mohnl, M., Schatzmayr, G. & Fegeros, K. 2010. Effects of probiotic inclusion levels in broiler nutrition on growth performance, nutrient digestibility, plasma immunoglobulins, and caecal microflora composition. *Poultry Science* 89(1): 58-67.
- Nayebpor, M., Farhomand, P. & Hashemi, A. 2007. Effects of different levels of direct fed microbial (Primalac) on growth performance and humoral immune response in broiler chickens. *Journal of Animal and Veterinary Advance* 6(11): 1308-1313.
- Niewold, T.A. 2007. The nonantibiotic anti-inflammatory effect of antimicrobial growth promoters, the real mode of action? A hypothesis. *Poultry Science* 86(4): 605-609.
- Paixao, A.C., Ferreira, A.C., Fontes, M., Themudo, P., Albuquerque, T., Soares, M.C. & Corrêa de Sá, M.I. 2016. Detection of virulence-associated genes in pathogenic and commensal avian *Escherichia coli* isolates. *Poultry Science* 95(7): 1646-1652.
- Panda, A.K., Rama Rao, S.S., Raju, M.V. & Sharma, S.S. 2008. Effect of probiotic (*Lactobacillus sporogenes*) feeding on egg production and quality, yolk cholesterol and humoral immune response of White Leghorn layer breeders. *Journal of the Science of Food and Agriculture* 88(1): 43-47.
- Park, J.W., Jeong, J.S., Lee, S.I. & Kim, I.H. 2016. Effect of dietary supplementation with a probiotic (*Enterococcus faecium*) on production performance, excreta microflora, ammonia emission, and nutrient utilization in ISA brown laying hens. *Poultry Science* 95(12): 2829-2835.
- Paudel, S., Stessl, B., Hess, C., Zloch, A. & Hess, M. 2016. High genetic diversity among extraintestinal *Escherichia coli* isolates in pullets and layers revealed by a longitudinal study. *BMC Veterinary Research* 12(1): 221.
- Raka, P.S., Sjojfan, O. & Radiati, L.E. 2014. Effect of liquid probiotics mixed culture supplements through drinking water on laying hens performance and yolk cholesterol. *Journal of World's Poultry Research* 4(1): 05-09.
- Rattanawut, J., Pimpa, O. & Yamauchi, K.E. 2018. Effects of dietary bamboo vinegar supplementation on performance, eggshell quality, ileal microflora composition, and intestinal villus morphology of laying hens in the late phase of production. *Animal Science Journal* 89(11): 1572-1580.
- Shokryazdan, P., Sieo, C.C., Kalavathy, R., Liang, J.B., Alitheen, N.B., Faseleh Jahromi, M. & Ho, Y.W. 2014. Probiotic potential of *Lactobacillus* strains with antimicrobial activity against some human pathogenic strains. *BioMed Research International* 2014: 927268.
- Shahail, M.U., Rahman, Z.U., Ijaz, A., Yousaf, M.S., Ashraf, K., Yaqub, T. & Rehman, H. 2011. Single or combined effects of mannan-oligosaccharides and probiotic supplements on the total oxidants, total antioxidants, enzymatic antioxidants, liver enzymes, and serum trace minerals in cyclic heat-stressed broilers. *Poultry Science* 90(11): 2573-2577.
- Shrinu, B., Rao, T.M., Reddy, P.M. & Reddy, K.K. 2013. Evaluation of different lactic acid bacterial strains for probiotic characteristics. *Veterinary World* 6(10): 785-788.
- Tang, S.G.H., Sieo, C.C., Ramasamy, K., Saad, W.Z., Wong, H.K. & Ho, Y.W. 2017. Performance, biochemical and haematological responses, and relative organ weights of laying hens fed diets supplemented with prebiotic, probiotic and synbiotic. *BMC Veterinary Research* 13(1): 248.
- Untoo, M., Banday, M.T., Afzal, I., Adil, S., Baba, I.A. & Khurshid, A. 2018. Potential of probiotics in poultry production. *Journal of Entomology and Zoology Studies* 6(3): 1293-1300.
- Willis, W.L., Isikhuemhen, O.S. & Ibrahim, S.A. 2007. Performance assessment of broiler chickens given mushroom extract alone or in combination with probiotics. *Poultry Science* 86(9): 1856-1860.
- Yahfoufi, N., Mallet, J.F., Graham, E. & Matar, C. 2018. Role of probiotics and prebiotics in immunomodulation. *Current Opinion in Food Science* 20: 82-91.
- Yulianto, B. & Lokapimasari, W.P. 2018. Isolation and identification of lactic acid bacteria from the digestive tract of kampung chicken (*Gallus gallus domesticus*). *The Philippine Journal of Veterinary Medicine* 55(1): 67-72.

Widya Paramita Lokapirnasari\* & Emy Koestanti Sabdongrum  
Department of Animal Husbandry  
Faculty of Veterinary Medicine  
Universitas Airlangga  
Jl. Mulyorejo, Kampus C, Surabaya, 60115  
Indonesia

Adriana Monica Sahidu  
Department of Marine  
Faculty of Fisheries and Marine  
Universitas Airlangga  
Jl. Mulyorejo, Kampus C, Surabaya, 60115  
Indonesia

Lilik Maslachah  
Department of Basic Medicine  
Veterinary Pharmacy Laboratory  
Faculty of Veterinary Medicine  
Universitas Airlangga  
Jl. Mulyorejo, Kampus C, Surabaya, 60115  
Indonesia

Andreas Berny Yulianto  
Sains Veteriner  
Faculty of Veterinary Medicine  
Universitas Airlangga  
Jl. Mulyorejo, Kampus C, Surabaya, 60115  
Indonesia

\*Corresponding author; email: [widya-p-l@fkh.unair.ac.id](mailto:widya-p-l@fkh.unair.ac.id)

Received: 6 April 2019

Accepted: 14 February 2020



SJR

Scimago Journal &amp; Country Rank

Enter Journal Title, ISSN or Publisher Name

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)

Ads by Google

[Report this ad](#) [Why this ad?](#)

# Sains Malaysiana

**Country** [Malaysia](#) -  SCIMAGO INSTITUTIONS RANKINGS**Subject Area and Category** [Multidisciplinary](#)  
[Multidisciplinary](#)**Publisher** [Penerbit Universiti Kebangsaan Malaysia](#) -  SCIMAGO INSTITUTIONS RANKINGS**Publication type** [Journals](#)**ISSN** [01266039](#)**Coverage** [2006-2020](#)

**Scope** Sains Malaysiana is a refereed journal committed to the advancement of scholarly knowledge and research findings of the several branches of science and technology. It contains articles on Earth Sciences, Health Sciences, Life Sciences, Mathematical Sciences and Physical Sciences. The journal publishes articles, reviews, and research notes whose content and approach are of interest to a wide range of scholars. Sains Malaysiana is published by the UKM Press and its autonomous Editorial Board are drawn from the Faculty of Science and Technology, Universiti Kebangsaan Malaysia. In addition, distinguished scholars from local and foreign universities are appointed to serve as advisory board members and referees.

[Homepage](#)[How to publish in this journal](#)[Contact](#)[Join the conversation about this journal](#)

# 26

H Index

# Indexed in SCOPUS and CPCI

A forum for researchers, practitioners to present and discuss research (Age 0+)

OpenPublish.eu

OPEN


# Indexed in SCOPUS and CPCI

A forum for researchers, practitioners to present and discuss research (Age 0+)


OpenPublish.eu

OPEN


Quartiles




SJR



Citations per document





Total Cites Self-Cites



External Cites per Doc



Cites per Doc

% International Collaboration



Citable documents    Non-citable documents    +

Cited documents    Uncited documents    +



Sains Malaysiana

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com" data-bbox="243 431 391 443">
```

## Free English Writing Tool

Grammarly makes sure everything you type is effective and mistake-free. Try now

Grammarly

DOWNLOAD

Metrics based on Scopus® data as of April 2020

Hani 1 week ago

what is the citescore for Sains Malaysiana

reply