

## DAFTAR PUSTAKA

- Afaq, F., Adhami, V. M., Ahmad, N., & Mukhtar, H. (2004). Health Benefits of Tea Consumption. In T. Wilson & N. J. Temple (Eds.), *Beverages in Nutrition and Health* (pp. 143–152). New York: Humana Press Inc.
- Ahmed, S., & Stepp, J. R. (2013). Green Tea: The Plants, Processing, Manufacturing and Production. In *Tea, Tea Drinking and Varieties* (pp. 19–31). USA: Elsevier.
- Al-Ghafari, A. B., Shorbaji, A. M., AL-Sarori, L. A., Baduwailan, E. O., Basaar, A. A., Doghaither, H. A. Al, Al-Marzouki, H.A., Omar, U. M. (2016). Phenolic Contents and Antioxidant Activities of Green Tea with and without Lemon. *Natural Science*, 08, 247–255.
- Alam, M. N., Bristi, N. J., & Rafiquzzaman, M. (2013). Review on in vivo and in vitro methods evaluation of antioxidant activity. *Studi Pharmaceutical Journal*, 21, 143–152.
- Alappat, B., Sarna, J. A., & Truong, C. (2015). Anticancer and Antioxidant Properties of Flavored Green Tea Extracts. *Journal of Agriculture and Life Sciences*, 2(1), 15–24.
- Amić, D., Davidović-Amić, D., Bešlo, D., & Trinajstić, N. (2003). Structure-radical scavenging activity relationships of flavonoids. *Croatica Chemica Acta*, 76(1), 55–61.
- Ananingsih, V. K., Sharma, A., & Zhou, W. (2013). Green tea catechins during food processing and storage: A review on stability and detection. *Food Research International*, 50, 469–479.
- Apak, R., Capanoglu, E., & Shahidi, F. (2018). *Measurement of antioxidant activity and capacity*. USA: John Wiley & Sons Ltd.

- Badan Standarisasi Nasional. 2016. Standar Nasional Indonesia: Teh Hijau. SNI 3945: 2016. Jakarta: Badan Standarisasi Nasional.
- Badan Standarisasi Nasional. 2016. Standar Nasional Indonesia: Teh Hitam. SNI 1902:2016. Jakarta: Badan Standarisasi Nasional, hal. 8.
- Bazinet, L., Araya-Farias, M., Doyen, A., Trudel, D., & Tetu, B. (2010). Effect of process unit operations and long-term storage on catechin contents in EGCG-enriched tea drink. *Food Research*, 43, 1692–1701.
- Bouayed, J., & Bohn, T. (2010). Exogenous antioxidant-double edged swords in cellular redox state. *Oxidative Medicine and Cellular Longevity*, 3(4), 228–237.
- Bruno, R. S., Bomser, J. A., & Ferruzzi, M. G. (2014). Antioxidant Capacity of Green Tea (*Camellia sinensis*). In *Processing and Impact on Antioxidants in Beverages* (pp. 33–39). USA: Elsevier Inc.
- Carocho, M., & Ferreira, I. C. F. R. (2013). A review on antioxidants, prooxidants and related controversy: Natural and synthetic compounds, screening and analysis methodologies and future perspectives. *Food and Chemical Toxicology*, 51, 15–25.
- Chambial, S., Dwivedi, S., Shukla, K. K., John, P. J., & Sharma, P. (2013). Vitamin C in Disease Prevention and Cure: An Overview. *Ind. J. Clin. Biochem*, 28(4), 314–328.
- Chen, G.-L., Chen, S.-G., Xie, Y.-Q., Chen, F., Zhao, Y.-Y., Luo, C.-X., & Gao, Y.-Q. (2015). Total phenolic, flavonoid and antioxidant activity of 23 edible flowers subjected to in vitro digestion. *Journal of Functional Foods*, 17, 243–259.
- Chen, G., Chen, S., Xiao, Y., & Fu, N. (2018). Antioxidant capacities and total phenolic contents of 30 flowers. *Industrial Crops & Products*, 111(October 2017), 430–445.
- Chen, M., Zhu, Y., Liu, B., Chen, Z., Zheng, J., Guan, M., Shi, H., Wang, Y., Yang, W. (2017). Changes in the volatiles, chemical components,

- and antioxidant activities of Chinese jasmine tea during the scenting processes. *International Journal of Food Properties*, 20(3), 681–693.
- Chen, Z. Y., Zhu, Q. Y., Tsang, D., & Huang, Y. (2001). Degradation of green tea catechins in tea drinks. *Journal of Agricultural and Food Chemistry*, 49(1), 477–482.
- Clifford, M. N., & Crozier, A. (2012). Phytochemicals in Teas and Tisanes and Their Bioavailability. In *Teas, Cocoa and Coffee: Plant Secondary Metabolites and Health* (pp. 45–54). USA: Blackwell Publishing.
- Departemen Kesehatan, R. (1989). *Materia Medika Indonesia* (5th ed.). Jakarta: Departemen Kesehatan Republik Indonesia.
- Dev, C., & Nidhi, S. R. R. S. (2016). Basketful Benefit of Citrus Limon. *International Research Journal of Pharmacy*, 7(6), 1–4.
- Dong, J., Ye, J., Lu, J., Zheng, X., & Liang, Y. (2011). Isolation of Antioxidant Catechins From Green Tea And Its Decaffeination. *Food and Bioproducts Processing*, 89(1), 62–66.
- Dong, X., Hu, Y., Li, Y., & Zhou, Z. (2019). The maturity degree , phenolic compounds and antioxidant activity of Eureka lemon [ Citrus limon ( L.) Burm . f .]: A negative correlation between total phenolic content , antioxidant capacity and soluble solid content. *Scientia Horticulturae*, 243(2018), 281–289.
- Duarte, A., Luis, A., Oleastro, M., & Domingues, F. C. (2016). Antioxidant properties of coriander essential oil and linalool and their potential to control Campylobacter spp . *Food Control*, 61, 115–122.
- Fadda, A., Serra, M., Molinu, M. G., Azara, E., Barberis, A., & Sanna, D. (2014). Reaction Time and DPPH Concentration Influence Antioxidant Activity and Kinetic Parameters of Bioactive Molecules and Plant Extracts in the Reaction With DPPH Radical. *Journal of Food Composition and Analysis*, 35(2), 112–119.

- Friedman, M., Kim, S.-Y., Lee, S.-J., Han, G.-P., Han, J.-S., Lee, K.-R., & Kozukue, N. (2005). Distribution of catechins, theaflavins, caffeine, and theobromine in 77 teas consumed in the united states. *Food Chemistry and Toxicology*, 70, 550–559.
- Furuse, M., Adachi, N., Tomonaga, S., Yamane, H., & D.M.Denbow. (2009). Central Functions of Green Tea Components. In *Handbook of Green Tea and Health Research* (pp. 1–22). New York: Nova Science.
- Gupta, D. (2015). Methods for determination of antioxidant capacity: A review. *International Journal of Pharmaceutical Sciences and Research*, 6(2), 546–566.
- Hajimahmoodi, M., Moghaddam, G., Mousavi, S. M., Sadeghi, N., Oveisi, M. R., & Jannat, B. (2014). Total Antioxidant Activity, and Hesperidin, Diosmin, Eriocitrin and Quercetin Contents of Various Lemon Juices. *Tropical Journal of Pharmaceutical Research*, 13(6), 951–956.
- Halliwell, B. (2002). Food-Derived Antioxidants: How to Evaluate Their Importance in Food and In Vivo. In E. Cadenas & L. Packer (Eds.), *Handbook of Antioxidants Revised and Expanded* (2nd ed., pp. 1–45). New York: Marcel Dekker.
- Hampton, M. G. (1992). Production of Black Tea. In K.C.Willson & M. N. Clifford (Eds.), *Tea: Cultivation to Consumption* (pp. 459–512). London: Chapman and Hall.
- Hara, Y. (2001). *Green Tea: Health Benefits And Applications*. New York: Marcel Dekker.
- Hardani, D. A., Hidayat, N., & Dewi, I. A. (2016). Ekstraksi Minyak Melati (Jasminum sambac) (Kajian Jenis Pelarut dan Lama Ekstraksi). *Industria*, 4(2), 82–88.
- He, F., Qian, Y. L., & Qian, M. C. (2018). Flavor and chiral stability of lemon-flavored hard tea during storage. *Food Chemistry*, 239, 622–630.

- Horzic, D., Komes, D., Belscak, A., Ganic, K. K., Ivezkovic, D., & Karlovic, D. (2009). The composition of polyphenols and methylxanthines in teas and herbal infusions. *Food Chemistry*, 115, 441–448.
- Indonesia, K. K. R. (2014). *Farmakope Indonesia* (5th ed.). Jakarta: Kementerian Kesehatan Republik Indonesia.
- Jabir, M. S., Taha, A. A., & Sabib, U. I. (2018). Antioxidant activity of linalool. *Engineering and Technology Journal*, 36, 64–67.
- Jagdale, A. D., Kamble, S. P., Nalawade, M. L., & Arvindekar, A. U. (2015). Citronellol: a potential antioxidant and aldose reductase inhibitor from Cymbopogon citratus. *International Journal of Pharmacy and Pharmaceutical Sciences*, 7(3), 203–209.
- Jeszka-skowron, M., Krawczyk, M., & Zgoła-grzeskowiak, A. (2015). Determination of antioxidant activity , rutin , quercetin , phenolic acids and trace elements in tea infusions : Influence of citric acid addition on extraction of metals. *Journal of Food Composition and Analysis*, 40, 70–77.
- Jiang, H., Engelhardt, U. H., Thräne, C., Maiwald, B., & Stark, J. (2015). Determination of flavonol glycosides in green tea , oolong tea and black tea by UHPLC compared to HPLC. *Food Chemistry*, 183, 30–35.
- Jun, M., Jeong, W.-S., & Ho, C.-T. (2006). Health Promoting Properties of Natural Flavor Substances. *Food Science and Biotechnology*, 15(3), 329–338.
- Kedare, S. B., & Singh, R. P. (2011). Genesis and development of DPPH method of antioxidant assay. *Journal Food Science Technology*, 48(4), 412–422.
- Khidzir, K. M., Cheng, S. F., & Chuah, C. H. (2015). Interspecies variation of chemical constituents and antioxidant capacity of extracts from Jasminum sambac and Jasminum multiflorum grown in Malaysia. *Industrial Crops and Products*, 74, 635–641.

- Kim, Y., Goodner, K. L., Park, J.-D., Choi, J., & Talcott, S. T. (2011). Changes in antioxidant phytochemicals and volatile composition of camellia sinensis by oxidation during tea fermentation. *Food Chemistry*, 129, 1331–1342.
- Komes, D., Horzic, D., Belscak, A., Kovacevic Ganic, K., & Vulic, I. (2010). Green tea preparation and its influence on the content of bioactive compounds. *Food Research International*, 43, 167–176.
- Kusmiyati, M., Sudaryat, Y., Lutfiah, I. A., Rustamsyah, A., & Rohdiana, D. (2016). Aktivitas antioksidan , kadar fenol total , dan flavonoid total dalam teh hijau ( Camellia sinensis ( L .) O . Kuntze ) asal tiga perkebunan Jawa Barat. *Jurnal Penelitian Teh Dan Kina*, 18(2), 101–106.
- Li, H., Luo, L., Ma, M., & Zeng, L. (2018). Characterization of volatile compounds and sensory analysis of jasmine scented black tea produced by different scenting processes. *Journal of Food Science*, 83(11), 1–15.
- Li, X., Jiang, Q., Wang, T., Liu, J., & Chen, D. (2016). Comparison of the antioxidant effects o quercitrin and isoquercitrin: Understanding the role of the 6'' -OH group. *Molecules*, 21, 1246.
- Lopez-Alarcon, C., & Lissi, E. (2009). Handbook of Green Tea and Health Research. In H. McKinley & M. Jamieson (Eds.), *Assessment of The Antioxidant Capacity of Green Teas: A Critical Review*. New York: Nova Science.
- Mabel, T.-G. D., Javier, S.-A. E., Cynthia, G.-P. I., Jorge, W.-P., Romeo, R., & Guadalupe, M.-A. G. C. (2017). Antioxidant profile of different types of herbal infusions and teas commercially available in mexico. *American Scientific Research Journal for Engineering, Technology, and Sciences (ASRJETS)*, 31(1), 67–77.

- Mathew, B. B., Jatawa, S. K., & Tiwari, A. (2012). Phytochemical analysis of Citrus limonum pulp and peel. *International Journal of Pharmacy and Pharmaceutical Sciences*, 4(2), 369–371.
- McKay, D. L., Miller, M. G., & Blumberg, J. B. (2012). Teas, Tisanes and Health. In A. Crozier, H. Ashihara, & F. Tomas-Barberan (Eds.), *Teas, Cocoa and Coffee: Plant Secondary Metabolites and Health* (pp. 109–116). USA: Blackwell Publishing.
- Moharram, H. ., & Youssef, M. M. (2014). Methods for Determining the Antioxidant Activity : A Review. *Journal Food Science & Technology*, 11(1), 31–42.
- Molyneux, P. (2004). The Use of the Stable Free Radical Diphenylpicrylhydrazyl (DPPH) for Estimating Antioxidant Activity. *Songklanakarin Journal of Science and Technology*, 26(December 2003), 211–219.
- Moosavy, M. H., Hassanzadeh, P., Mohammadzadeh, E., Mahmoudi, R., Khatibi, S. A., & Mardani, K. (2017). Antioxidant and Antimicrobial Activities of Essential Oil of Lemon (Citrus limon) Peel in Vitro and in a Food Model. *Journal of Food Quality and Hazards Control*, 4, 42–48.
- Mulja, M., & Suharman. (1995). *Analisis Instrumental*. Surabaya: Airlangga University Press.
- Namita, P., Mukesh, R., & Vijay, K. J. (2012). Camellia sinensis (Green Tea): A Review. *Global Journal of Pharmacology*, 6(2), 52–59.
- Naveed, M., Bibi, J., Kamboh, A. A., Suheryani, I., Kakar, I., Fazlani, S. A., FangFang, X., Kalhoro, S. A., Yunjuan, L., Kakar, M. U., El-Hack, M. E. A., Noreldin, A. E., Zhixiang, S., LiXia, C., XiaoHui, Z. (2018). Pharmacological values and therapeutic properties of black tea (Camellia sinensis ): A comprehensive overview. *Biomedicine & Pharmacotherapy*, 100, 521–531.

- Nimse, S. B., & Pal, D. (2015). Free radicals, natural antioxidants, and their reaction mechanisms. *RSC Advances*, 5, 27986–28006.
- Padayatty, S. J., Katz, A., Wang, Y., Eck, P., Kwon, O., Lee, J. H., Chen, S., Corpe, C., Dutta, A., Dutta, S. K., Levine, M. (2003). Vitamin C as an Antioxidant: Evaluation of Its Role in Disease Prevention. *Journal of the American College of Nutrition*, 22(1), 18–35.
- Permata, A. N., Kurniawati, A., & Lukiat, B. (2018). Screening fitokimia, aktivitas antioksidan dan antimikroba pada buah jeruk lemon (Citrus limon) dan jeruk nipis (Citrus aurantiifolia). *Jurnal Ilmiah Ibnu Sina*, 3(1), 64–76.
- Pisoschi, A. M., & Negulesce, G. P. (2011). Methods for total antioxidant activity determination: A review. *Biochemistry & Analytical Biochemistry*, 1(1), 1–10.
- Pratiwi, L., Fudholi, A., Martien, R., & Pramono, S. (2016). Ethanol Extract , Ethyl Acetate Extract , Ethyl Acetate Fraction , and n-Heksan Fraction Mangosteen Peels ( Garcinia mangostana L .) As Source of Bioactive Substance Free-Radical Scavengers Ekstrak etanol , Ekstrak etil asetat , Fraksi etil asetat , dan F. *Journal of Pharmaceutical Science and Clinical Research*, 1, 71–82.
- Prior, R. L., Wu, X., & Schaich, K. (2005). Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplement. *Agricultural and Food Chemistry*, 53, 4290–4302.
- Rajendran, N. K., George, B. P., Chandran, R., Tynga, I. M., Hourelid, N., & Abrahamse, H. (2019). The influence of light on reactive oxygen species and NF-kB in disease progression. *Antioxidants*, 8(12), 640.
- Rijken, P. J., Wiseman, S. A., Weisgerber, U. M., Mierlo, C. A. J. van, Quinlan, P. T., & Put, F. van de. (2002). Antioxidant and Other Properties of Green and Black Tea. In E. Cadenas & L. Packer (Eds.), *Handbook of Antioxidants* (pp. 371–399). New York: Marcel Dekker.

- Roy, M. K., Koide, M., Rao, T. P., Okubo, T., Ogasawara, Y., & Juneja, L. R. (2010). ORAC and DPPH assay comparison to assess antioxidant capacity of tea infusions: Relationship between total polyphenol and individual catechin content. *International Journal of Food Sciences and Nutrition*, 61(2), 109–124.
- Schaich, K. M., Tian, X., & Xie, J. (2015). Hurdles and pitfalls in measuring antioxidant efficacy: A critical evaluation of ABTS, DPPH, and ORAC assays. *Journal of Functional Food*, 14, 111–125.
- Scoboda, P., VIckova, H., & Novakova, L. (2015). Development and validation of UHPLC/MS-MS method for determination of eight naturally occurring catechin derivatives in various tea samples and the role of matrix effects. *Pharmaceutical and Biomedical Analysis*, 114, 62–70.
- Setyawati, A. (2015). Budidaya Tanaman Melati (Jasminum spp.). *Iptek Hortikultura*, 11, 1–4.
- Sharma, P. J., & Singh, R. P. (2013). Evaluation of antioxidant activity in foods with special References to TEAC method. *American Journal of Food Technology*, 8, 83–101.
- Shen, J. X., Rana, M. M., Liu, G. F., Ling, T. J., Gruber, M. Y., & Wei, S. (2017). Differential contribution of jasmine floral volatiles to the aroma of scented green tea. *Journal of Food Quality*, 2017, 1–10.
- Shimamura, T., Sumikura, Y., Yamazaki, T., Tada, A., Kashiwagi, T., Ishikawa, H., Matsui, T., Sugimoto, N., Akiyama, H., Ukeda, H. (2014). Applicability of the DPPH Assay for Evaluating the Antioxidant Capacity of Food Additives – Inter-laboratory Evaluation Study –. *Analytical Sciences*, 30, 717–721.

- Singh, P., Kesharwani, R. K., & Keservani, R. . (2017). Antioxidants and Vitamins: Roles in Cellular Function and Metabolism. In *Sustained Energy for Enhanced Human Functions and Activity* (pp. 385–407). UK: Elsevier Inc.
- Sochor, J., Ryvolova, M., Krystofova, O., Salas, P., Hubalek, J., Adam, V., Trnkova, L., Havel, L., Beklova, M., Zehnalek, J., Provaznik, I., Kizek, R. (2010). Fully Automated Spectrometric Protocols for Determination of Antioxidant Activity: Advantages and Disadvantages. *Molecules*, 15, 8618–8640.
- Sutherland, B. A., Rahman, R. M. A., & Appleton, I. (2006). Mechanisms of action of green tea catechins , with a focus on ischemia-induced neurodegeneration. *Journal of Nutritional Biochemistry*, 17, 291–306.
- Suyanti, Prabawati, S., & Sjaifullah. (2004). Karakterisasi Fisiko-Kimia Bunga Melati Putih. *Jurnal Hortikultura*, 14(2), 121–126.
- Takeo, T. (1992). Green and Semi-Fermented Teas. In K. C. Willson & M. N. Clifford (Eds.), *Tea: Cultivation to Consumption* (pp. 413–457). London: Chapman and Hall.
- Tao, W., Zhou, Z., Zhao, B., & Wei, T. (2016). Simultaneous determination of eight catechins and four theaflavins in green , black and oolong tea using new HPLC – MS – MS method. *Journal of Pharmaceutical and Biomedical Analysis*, 131, 140–145.
- Tariq, A. L., & Reyaz, A. L. (2013). Original Research Article Antioxidant activity of Camellia sinensis leaves. *Internasioanl Journal of Current Microbiology and Applied Sciences*, 2(5), 40–46.
- Tjitrosoepomo, G. (2005). *Morfologi Tumbuhan*. Yogyakarta: Gajah Mada University Press.

- Villano, D., Fernandez-Pachon, M. S., Moya, M. L., Troncoso, A. M., & Garcia-Parrilla, M. C. (2007). Radical scavenging ability of polyphenolic compounds towards DPPH free radical. *Talanta*, 71, 230–235.
- Wang, H.-F., Yih, K.-H., Yang, C.-H., & Huang, K.-F. (2017). Anti-oxidant activity and major chemical component analyses of twenty-six commercially available essential oil. *Journal of Food and Drug Analysis*, 25, 881–889.
- Watson, D. G. (2005). *Pharmaceutical Analysis: A Text Book For Pharmacy Students and Pharmaceutical Chemists*. New York: Harcourt.
- Weerawatanakorn, M., Hung, W.-L., Pan, M.-H., Li, S., Li, D., Wan, X., & Ho, C.-T. (2015). Chemistry and health beneficial effects of oolong tea and theasinensins. *Food Science and Human Wellness*, 4, 133–146.
- Widowati, W., Janeva, W., Nadya, S., Amalia, A., Arumwardana, S., Kusuma, H. S. W., & Arinta, Y. (2018). Antioxidant and antiaging activities of Jasminum sambac extract, and its compounds. *Journal of Reports in Pharmaceutical Sciences*, 7(3), 270–285.
- Xi, W., Lu, J., Qun, J., & Jiao, B. (2017). Characterization of phenolic profile and antioxidant capacity of different fruit part from lemon ( Citrus limon Burm .) cultivars. *Journal of Food Science and Technology*, 54(5), 1108–1118.
- Xie, J., & Schaich, K. M. (2014). Re-evaluation of the 2,2-diphenyl-1-picrylhydrazylfree radical (DPPH) assay for antioxidant activity. *Agricultural and Food Chemistry*, 62, 4251–4260.
- Yang, J., Guo, J., & Yuan, J. (2008). In vitro antioxidant properties of rutin. *Food Science and Technology*, 41, 1060–1066.

- Yang, S., Jeon, S., Lee, E., Shim, C.-H., & Lee, I.-S. (2010). Comparative study of the chemical composition and antioxidant activity of six essential oils and their components. *Natural Product Research*, 24(2), 140–151.
- Younis, A., Mehdi, A., & Riaz, A. (2011). Supercritical Carbon Dioxide Extraction and Gas Chromatography Analysis of Jasminum Sambac Essential Oil. *Pak. j. Bot.*, (43), 163–168. Retrieved from
- Zhang, P., & Zhou, Z. (2019). Postharvest ethephon degreening improves fruit color, flavor quality and increases antioxidant capacity in ‘Eureka’ lemon ( Citrus limon ( L. ) Burm . f .). *Scientia Horticulturae*, 248, 70–80.
- Zhang, Y., Li, Q., Xing, H., Lu, X., Zhao, L., Qu, K., & Bi, K. (2013). Evaluation of antioxidant activity of ten compounds in different tea samples by means of an on-line HPLC – DPPH assay. *Food Research International*, 1–10.
- Zhao, C. N., Tang, G. Y., Cao, S. Y., Xu, X. Y., Gan, R. Y., Liu, Q., Mao, Q. Q., Shang Ao., Li, H. Bin. (2019). Phenolic profiles and antioxidant activities of 30 tea infusions from green, black, oolong, white, yellow and dark teas. *Antioxidants*, 8(7).
- Zou, Z., Xi, W., Hu, Y., Nie, C., & Zhou, Z. (2016). Antioxidant activity of Citrus fruits. *Food Chemistry*, 196, 885–896.