

## [PJI] Copyediting Review Request

From: Editor in Chief (pji@ub.ac.id)

To: hanikhidayati@yahoo.com

Date: Friday, January 3, 2020 at 11:36 AM GMT+7

Ariani Rahayu Setianti:

Your submission "Effectiveness of Triamcinolone and Lidocaine in Patient with Piriformis Syndrome – A Case Series" for Pharmaceutical Journal of Indonesia has been through the first step of copyediting, and is available for you to review by following these steps.

1. Click on the Submission URL below.
2. Log into the journal and click on the File that appears in Step 1.
3. Open the downloaded submission.
4. Review the text, including copyediting proposals and Author Queries.
5. Make any copyediting changes that would further improve the text.
6. When completed, upload the file in Step 2.
7. Click on METADATA to check indexing information for completeness and accuracy.
8. Send the COMPLETE email to the editor and copyeditor.

Submission URL:

<https://pji.ub.ac.id/index.php/pji/author/submissionEditing/169>

Username: arianirahayu

This is the last opportunity to make substantial copyediting changes to the submission. The proofreading stage, that follows the preparation of the galleys, is restricted to correcting typographical and layout errors.

If you are unable to undertake this work at this time or have any questions, please contact me. Thank you for your contribution to this journal.

hananditia rachma pramestutie  
Department of Pharmacy, Faculty of Medicine, Brawijaya University  
Phone 081334624865  
[hananditia@gmail.com](mailto:hananditia@gmail.com)  
M.Farm.Klin., Apt

---

Pharmaceutical Journal of Indonesia

<http://pji.ub.ac.id/index.php/pji>

## [PJI] Editor Decision

From: Hananditia RAchma Pramestutie (hananditia@ub.ac.id)

To: arianirahayus@gmail.com

Cc: hanikhidayati@yahoo.com

Date: Tuesday, October 8, 2019 at 07:56 PM GMT+7

Ariani Rahayu Setianti:

We have reached a decision regarding your submission to Pharmaceutical Journal of Indonesia, "Effectiveness of Triamcinolone and Lidocaine in Patient with Piriformis Syndrome – A Case Series".

Our decision is to: please revision your article before 19 October 2019

hananditia rachma pramestutie  
Department of Pharmacy, Faculty of Medicine, Brawijaya University  
[hananditia@gmail.com](mailto:hananditia@gmail.com)

M.Farm.Klin., Apt

---

Pharmaceutical Journal of Indonesia  
<http://pji.ub.ac.id/index.php/pji>

## [PJI] Editor Decision

From: Editor in Chief (pji@ub.ac.id)

To: hanikhidayati@yahoo.com

Cc: arianirahayu@gmail.com

Date: Tuesday, October 22, 2019 at 06:28 PM GMT+7

Ariani Rahayu Setianti:

We have reached a decision regarding your submission to Pharmaceutical Journal of Indonesia, "Effectiveness of Triamcinolone and Lidocaine in Patient with Piriformis Syndrome – A Case Series".

Our decision is to: your article accepted to Pharmaceutical Journal of Indonesia

hananditia rachma pramestutie  
Department of Pharmacy, Faculty of Medicine, Brawijaya University  
Phone 081334624865  
[hananditia@gmail.com](mailto:hananditia@gmail.com)

---

Pharmaceutical Journal of Indonesia  
<http://pji.ub.ac.id/index.php/pji>



## Effectiveness of Triamcinolone and Lidocaine in Patient with Piriformis Syndrome – A Case Series

### Keywords:

Lidocaine,  
Piriformis  
Syndrome,  
Triamcinolone,  
Visual Analogue  
Scale (VAS)

### ABSTRACT

Piriformis syndrome (PS) has been documented as a contributory cause for sciatica, buttock, and low back pain. The worldwide incidence of PS would be about 2,4 million per year. The management of piriformis syndrome includes injection of the piriformis muscle with local anesthetic and steroid, however, the study for the effectiveness of the combination is lacking especially in Indonesian population. This case illustrates the effectiveness reduction of triamcinolone in combination with lidocaine measured by Visual Analogue Scale (VAS) in patient suffered from PS.

### Introduction

Piriformis syndrome (PS) first introduced in 1928 known as *wallet sciatica* or *fat wallet syndrome*.<sup>1,2</sup> PS is a peripheral neuritis of the sciatic nerve caused by an abnormal condition of the piriformis muscle.<sup>3</sup> PS has been recorded as a devoted cause for sciatica, buttock, and low back pain. Pain and localized tenderness around the gluteal region at the area of the piriformis muscle are the clinical manifestation of this syndrome and usually characterized as a deep, aching type of pain with or without signs and symptoms of sciatica.<sup>4</sup> PS may contribute to 0,3% to 6% of all cases of low back pain and/or sciatica with an estimated amount of new cases of low back pain at 40 million annually, the incidence of PS would be roughly at 2,4 million per year.<sup>5</sup> PS has been accused as the potential pain generator and affect individuals in both heavy activities population, such as athletes, and general population of all occupations with many kinds of activity levels.<sup>3,6</sup>

PS remains a controversial and nebulous diagnosis for back and buttock pain.<sup>7</sup> Delay in diagnosing PS may lead to pathologic condition of the sciatic nerve, chronic somatic dysfunction, and compensatory changes resulting in pain, paraesthesia, hyperesthesia, and muscle weakness.<sup>3</sup> Currently, gold standart therapy in PS has not been found yet.<sup>8</sup>

The first-line treatment for PS based on conservative pharmacotherapy with nonsteroidal anti-inflammatory drugs (NSAID), muscle relaxant, and neuropathic pain agents such as tricyclic antidepressants (TCAs) or anticonvulsants.<sup>9,10</sup> Local injection of lidocaine, steroids, or the newest agent, botulinum toxin can be used if the first line failed to reduce the pain.<sup>11</sup> Triamcinolone is one of corticosteroid used to treat piriformis syndrome in clinical settings, however, the research about effectiveness of lidocaine and triamcinolone is lacking especially in Indonesia populations.

The purpose of this case report is to determine the effectiveness of triamcinolone and lidocaine combination

as one of the therapy choices in PS. This case series will illustrate some patients experience after injected by both agents.

### Case Presentation

#### Case 1

Patient ND, 57 years old female and weighed 71 kg. Patient referred from Bunda Hospital diagnosed with low back pain suspect piriformis syndrome. Patient suffered from pain in both buttocks and hamstring since 5 years ago, radiating to both leg. Patient felt pain especially when sitting and walking, and also thick and tingling in both soles. There was no history of trauma, diabetes mellitus denied, and there was a history of high cholesterol.

FAIR position negative, Beatty test positive, and Freiberg test negative. Peripheral Extremities USG showed that the thickening of piriformis muscle was unclear. Patient was given paracetamol 500mg/24 hours and vitamin B12 50 mcg/12 hours, and vitamin B6 10 mg/12 hours. The pain persisted so the doctor decided to give Trilac® (Triamcinolone 10 mg/mL) 2 mL and lidocaine 2% 1 mL. The initial pain scale was 7 (severe) and after treatment the pain decreased into 2 (mild).

#### Case 2

Patient MB, 35 years old male, 159 cm and weighed 60 kg. Patient referred from PHC Hospital with PS. Patient suffered with pain from buttock radiating to right leg since one year ago, throbbing pain and increased if patient was standing, walking, or bowing. Patient felt his right sole thick and tingling. He ~~couldn't~~ ~~could not~~ stand for more than 5 minutes and walk for 100 meters because of the pain. Pain decreased if the patient sits or lie down. There ~~is~~ ~~was~~ a history of trauma due to fell in sitting

**Commented [WU1]:** The paper not only consisted of one case but reviewed some cases. Is there reason/method in choosing those cases? Mention here, along with one or two points that we can conclude from those cases.

**Commented [WU2]:** Use initial for the hospital name

**Formatted:** Strikethrough

**Formatted:** Strikethrough

position. There was no tumor, fever, half-body weakness, or stroke. The patient got oral analgesic as the therapy but the pain remained.

USG performed in the right extremities and there was a thickening of right sciatic nerve ± 0,53 cm with a tenderness and probe positive. Then he got injection with Trilac® (Triamcinolone 10 mg/mL) 2 mL and lidocaine 2% 2 mL. Patient felt a bit dizzy for about 5 minutes after injection. Patient's pain measured with Visual Analogue Scale (VAS), which the initial VAS was 8 (severe) and decreased into 6 (moderate). Pain reduction was obtained two days after injection.

### Case 3

Patient NS, 64 years old female, 157 cm and weighed 62 kg. Patient referred from Universitas Airlangga Hospital with PS. Patient has had history of heart disease and hypertension. Patient suffered from pain in left buttock since 2 months ago. Pain radiated to left hamstring, weakness in limbs denied. There was no tingling feeling. FAIR position tested positif.

Peripheral Extremities USG showed that the right piriformis muscle was thicker compared with left piriformis muscle. Patient got oral analgesics combination in capsules to relieve the pain. However, pain decreased but didn't did not disappear. Patient then got injected with Trilac® (Triamcinolone 10 mg/mL) and lidocaine 2% injection, 2 mL and 1 mL, respectively. Patient couldn't could not stand for about 10 minutes right after injection and needed to rest for about 15 minutes. The initial pain scale before injection was 4 (moderate) and after injection was 2 (mild). Pain decreased two days after injection.

### Discussion

Piriformis syndrome has been recorded as one of responsible factor of sciatica, buttock, and low back pain occurrence with relatively rare prevalence, about 0,3% to 6% of all cases of low back pain and sciatica.<sup>4,5</sup> This syndrome should be considered in the differential diagnosis of all patients with persistent pain in the gluteal area predominantly when there are no findings from the lumbar spine or the hip joints.<sup>7</sup>

There are some etiologies of piriformis syndrome, including gluteal trauma in the sacroiliac or gluteal areas (possibly several years previously), predisposing anatomic variants, myofascial trigger points, hypertrophy and spasm of the piriformis muscle. PS divided into two types of cause, primary and secondary. Primary cause happens when sciatic nerve entrapment occurs due to intrinsic abnormality within the muscle itself whereas the second one occurs as a result of direct or blunt trauma to the piriformis muscle.<sup>11</sup> Other researchers classify PS cause into trauma, muscle hypertrophy, sitting for prolonged periods, and anatomic anomalies.<sup>3,5</sup>

The piriformis muscle is flat, oblique, and pyramidal-shaped. It originates anterior to the vertebrae (S2 to S4),

the superior margin of the greater sciatic foramen, and the sacrotuberous ligament. This muscle receives innervation from nerve branches coming off L5, S1, and S2. Irritation of the neighboring sciatic nerve which is very close to the center of the muscle, could appear when piriformis muscle is overworked, irritated, or inflamed.<sup>5</sup> Some symptoms of PS occurs as the consequence of local inflammation and congestion due to muscular compression of small nerves and vessels, including the pudendal nerve and blood vessels, which exited at the medial inferior border of the piriformis muscle.<sup>3</sup>

The components added to the clinical presentation are somatic and neuropathic. Somatic component underlying PS is a myofascial pain syndrome whereas neuropathic component refers to compression and irritation of the sciatic nerve as it courses through the infrapiriform foramen.<sup>12</sup>

The symptoms suggesting PS are clinical, bringing together the patient's painful functional impairments and the signs reproduced by physical maneuvers soliciting the piriformis muscle and the enviroing anatomical structures.<sup>13</sup> Physical findings thought to be more specific for PS. Three onymous tests are the Freiberg test (forceful internal rotation of the hip with the patient supine), the Pace test (reproduction of buttock pain with resisted hip abduction), and Beatty test (reproduction of buttock pain with the abduction of the thigh against gravity with patient in lateral decubitus position). The other suggestive test is FAIR (Flexed, Adducted, Internally Rotated) position.<sup>11</sup> The patients presented in this study have been tested by those criteria to diagnose the PS. Beatty test was positive in patient 1 and FAIR position test was positive in patient 3.

Diagnostic imaging is invaluable in figuring out other probable causes of sciatic nerve irritation such as lumbar disk disease and radiculopathy.<sup>14</sup> CT and MRI are reportedly valuable for diagnosing PS by revealing enlargement of the piriformis muscle, however, adequate sensitivity remains questionable.<sup>15</sup> Zhang *et al* demonstrated ~~U~~ltrasound as one of technique for the diagnosis of PS. In PS patients, the piriformis muscle was enlarged and the echo intensity of this muscle was enhanced. Moreover, the thickness was significantly increased in the symptomatic side compared with that on the asymptomatic side.<sup>16</sup> In three patient presented, the result of USG musculoskeletal showed thickening of piriformis muscle in two patients while the other one is unclear.

Piriformis muscle injection is usually suggested to the patient as part of multimodal therapy.<sup>12</sup> Nonsteroidal anti-inflammatory drugs, analgesics, muscle relaxant, and neuropathic pain agents (gabapentin, nortriptyline, and carbamazepine) may relieve pain and spasm in this condition.<sup>7,9</sup> In addition, the management of piriformis syndrome includes injection of the piriformis muscle with local anesthetic and steroid or with botulinum toxin. Some investigators also inject dilute local anesthetic and steroid in the area of the sciatic nerve. The recommendation for local injection therapy may be both

Formatted: Strikethrough

Commented [WU3]: Use initial for hospital name

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

\* Corresponding author: Ariani Rahayu Setianti, Master of Clinical Pharmacy Programme, Faculty of Pharmacy, University of Airlangga, Surabaya, Indonesia .

E-mail: arianiarahayus@gmail.com

diagnostic and therapeutic.<sup>11,17</sup> Corticosteroid with local anesthetic injection aim is to reduce the irritation by decreasing inflammation in the area and temporarily stopping piriformis muscle spasm.<sup>18</sup>

Triamcinolone is one of corticosteroids injection approved by Food and Drugs Administration (FDA). Corticosteroids have both anti-inflammatory and immunosuppressive effect but the exact mechanisms are complex. Corticosteroids act directly on nuclear steroid receptor and interrupt the inflammatory and immune cascade at several levels. They decrease vascular permeability and inhibit the accumulation of inflammatory cells and prevent the synthesis and secretion of several inflammatory mediators.<sup>13</sup> Triamcinolone is used empirically for intra articular injection because of low solubility and longer duration of action whereas bethamethasone for soft tissue injection due to high solubility, shorter duration of action, and fewer cutaneous adverse effect. Triamcinolone common concentration used is 10-40 mg with approximate duration of action 14 days.<sup>19</sup>

Lidocaine is an amide-type local anesthetic that exerts its pharmacological action through the block of sodium channels in neural tissues, thereby interrupting neuronal transmission.<sup>20</sup> The analgesic effect of lidocaine can be due to blockage of the neuronal transmission at the site of injury, attenuating the neurogenic response, and by the intrinsic systemic anti-inflammatory properties. Moreover, lidocaine can reduce cytokine-induced cellular damage through a mechanism that involves mitochondrial adenosine triphosphate (ATP)-gated potassium channels.<sup>21</sup> The speed of onset of lidocaine is 1 to 5 minutes after local infiltration and 5 to 15 minutes after peripheral nerve blockade. Lidocaine has been shown to cross the placenta and blood-brain barrier by simple passive diffusion.<sup>22</sup>

Previous study conducted by Fishman *et al* reported that patient who got 1,5 mL 2% lidocaine combined with 20 mg triamcinolone generally feel immediate relieved after injection. Patient who conferred with two of three clinical signs, 468 out of 537 patients after 10 months follow up, 79% attained at least 50% improvement in symptoms. Patients with less than two clinical criteria, 75,2% improved 50% or more.<sup>23, 24, 25</sup>

A study performed by Ciurylo and Gray in patients with PS who got steroid injection showed that from 26 samples, 84,6% resulted in diagnostic response and average pain reduction was 65,6%. In this research, triamcinolone dose used was 20 mg mixed with 2 mL 1% lidocaine and 2 mL of 0,25% bupivacaine. In a randomized trial conducted in 2015, addition of corticosteroid to lidocaine therapy ~~didn't~~ did not give any beneficial compared with lidocaine alone in patient with piriformis syndrome. Fifty-seven patients between the ages 18 and 70 were enrolled in this research but 7 patients dropped out due to adverse effects?. Fifty patients were randomized and divided into two groups, first group was given (lidocaine 2% 5 mL only,) and second group was given(lidocaine 4% + 1 mL

betamethasone). Compared with the baseline, there was a significant improvement in NRS (statistically significant? P value?), however, there was no significant difference in pain reduction between each group.<sup>26</sup>

All of the patient presented in this paper received got triamcinolone and lidocaine injection to treat PS (same doses?)~~this~~ condition. The patients experienced a reduction in pain severity, however, the pain reduction was variedly among the patients. The first patient experienced 5 scales reduction, while the second and the third patient were 2 scales reduction. There was a delayed onset of pain reduction. Patient 2 and 3's pain reduced two days after injection. Review done by Brinks *et al* found that increased or persistent pain after injection or pain at the site of injection was described in 19 studies.<sup>27</sup> This phenomenon was likely the most possible cause of the delayed effect.

#### Conclusion

Triamcinolone and Lidocaine could be used as a therapy choice in patients with piriformis syndrome. Those agents could reduce pain scale from 2 up to 5 scales two days after injection, but longer observation is needed to achieve maximum pain reduction. However, a larger study is needed to determine the effectiveness in bigger population.

#### BIBLIOGRAPHY

1. Roy BA. Piriformis Syndrome. ACSMs Health Fit J [Internet]. 2014;18(4). Available from: [https://journals.lww.com/acsm-healthfitness/Fulltext/2014/07000/Piriformis\\_Syndrome.3.aspx](https://journals.lww.com/acsm-healthfitness/Fulltext/2014/07000/Piriformis_Syndrome.3.aspx)
2. Waldman S. Atlas of common pain syndrome. Philadelphia: Elsevier; 2019. 350-4 p.
3. A Boyajian-O'Neill L, McClain R, K Coleman M, Thomas P. Diagnosis and Management of Piriformis Syndrome: An Osteopathic Approach. J Am Osteopath Assoc. 2008 Dec 1;108:657-64.
4. Kean Chen C, Nizar AJ. Prevalence of Piriformis Syndrome in Chronic Low Back Pain Patients. A Clinical Diagnosis with Modified FAIR Test. Pain Pract [Internet]. 2013 Apr 1;13(4):276-81. Available from: <https://doi.org/10.1111/j.1533-2500.2012.00585.x>
5. Hicks B, Varacallo M. Piriformis Syndrome. In: StatPearls (Internet) [Internet]. StatPearls Publishing; 2019. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK448172>
6. Singh U, Meena R, Singh CA, Singh AKJ, Singh Am, Langshong R. Prevalence of piriformis syndrome among the cases of low back/buttock pain with sciatica: A prospective study. J Med Soc. 2013;27(2):94.

\* Corresponding author: Ariani Rahayu Setiandi, Master of Clinical Pharmacy Programme, Faculty of Pharmacy, University of Airlangga, Surabaya, Indonesia .

E-mail: [arianirahayus@gmail.com](mailto:arianirahayus@gmail.com)

7. Papadopoulos EC, Khan SN. Piriformis syndrome and low back pain: a new classification and review of the literature. *Orthop Clin North Am* [Internet]. 2004;35(1):65–71. Available from: <http://www.sciencedirect.com/science/article/pii/S0030589803001056>
8. Michel F, Decavel P, Toussiro E, Tatu L, Aleton E, Monnier G, et al. Piriformis muscle syndrome: Diagnostic criteria and treatment of a monocentric series of 250 patients. *Ann Phys Rehabil Med* [Internet]. 2013;56(5):371–83. Available from: <http://www.sciencedirect.com/science/article/pii/S1877065713000675>
9. Masala S, Crusco S, Meschini A. Piriformis Syndrome: Long-Term Follow-up in Patients Treated with Percutaneous Injection of Anesthetic and Corticosteroid Under CT Guidance. *Cardiovasc Interv Radiol* [Internet]. 2012;35(375). Available from: <https://doi.org/10.1007/s00270-011-0185-z>
10. Nishikawa N, Nomoto M. Management of neuropathic pain. *J Gen Fam Med* [Internet]. 2017 Apr 12;18(2):56–60. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29263992>
11. Miller T, White K, Ross D. The Diagnosis and Management of Piriformis Syndrome: Myths and Facts. *Can J Neurol Sci*. 2012;39(5):577–83.
12. Jankovic D, Peng P, van Zundert A. Brief review: Piriformis syndrome: etiology, diagnosis, and management. *Can J Anesth Can d'anesthésie* [Internet]. 2013;60(10):1003–12. Available from: <https://doi.org/10.1007/s12630-013-0009-5>
13. Ayhan E, Kesmezacar H, Akgun I. Intraarticular injections (corticosteroid, hyaluronic acid, platelet rich plasma) for the knee osteoarthritis. *World J Orthop* [Internet]. 2014 Jul 18;5(3):351–61. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25035839>
14. Miller TA, White KP, Ross DC. The Diagnosis and Management of Piriformis Syndrome: Myths and Facts. *Can J Neurol Sci*. 2012;29:577–83.
15. Niu C-C, Lai P-L, Fu T-S, Chen L-H, Chen W-J. Ruling out Piriformis syndrome before diagnosing lumbar radiculopathy. *Chang Gung Med J*. 2009 Mar 1;32:182–7.
16. Zhang W, Luo F, Sun H, Ding H. Ultrasound appears to be a reliable technique for the diagnosis of piriformis syndrome. *Muscle Nerve* [Internet]. 2019/02/27. 2019 Apr;59(4):411–6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30663080>
17. Benzon M.D. HT, Katz M.D. JA, Benzon B.A. HA, Iqbal M.D. MS. Piriformis Syndrome: Anatomic Considerations, a New Injection Technique, and a Review of the Literature. *Anesthesiol J Am Soc Anesthesiol*. 2003 Jun 1;98(6):1442–8.
18. Ciurylo W, Gray TK. Therapeutic Response Rate for Those with Piriformis by Fluoroscopy Guided Botulinum Toxin Injection. *J Anesth Pain Med*. 2017;2(2):1–5.
19. Stephens MB, Beutler AI, O'Connor FG. Musculoskeletal injections: A review of the evidence. *Am Fam Physician*. 2008;78(8):971–6.
20. Eipe N, Gupta S, Penning J. Intravenous lidocaine for acute pain: an evidence-based clinical update. *BJA Educ* [Internet]. 2016 Apr 12;16(9):292–8. Available from: <https://doi.org/10.1093/bjaed/mkw008>
21. de Oliveira CMB, Issy AM, Sakata RK. Intraoperative Intravenous Lidocaine. *Brazilian J Anesthesiol* [Internet]. 2010;60(3):325–33. Available from: <http://www.sciencedirect.com/science/article/pii/S0034709410700416>
22. Weinberg L, Peake B, Tan C, Nikfarjam M. Pharmacokinetics and pharmacodynamics of lignocaine: A review. *World J Anesthesiol*. 2015;4(2):17–29.
23. Cassidy L, Walters A, Bubb K, Shoja MM, Shane Tubbs R, Loukas M. Piriformis syndrome: implications of anatomical variations, diagnostic techniques, and treatment options. *Surg Radiol Anat* [Internet]. 2012;34(6):479–86. Available from: <https://doi.org/10.1007/s00276-012-0940-0>
24. Fishman LM, Dombi GW, Michaelsen C, Ringel S, Rozbruch J, Rosner B, et al. Piriformis syndrome: Diagnosis, treatment, and outcome—a 10-year study. *Arch Phys Med Rehabil* [Internet]. 2002 Mar 1;83(3):295–301. Available from: <https://doi.org/10.1053/apmr.2002.30622>
25. Jeong HS, Lee GY, Lee EG, Joe EG, Lee JW, Kang HS. Long-term assessment of clinical outcomes of ultrasound-guided steroid injections in patients with piriformis syndrome. *Ultrason (Seoul, Korea)* [Internet]. 2015/01/23. 2015 Jul;34(3):206–10. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25672769>
26. Akgun K, Palamar D, Misirlioglu T, Erden MG, Erbilir T. Piriformis Syndrome: Comparison of the Effectiveness of Local Anesthetic and Corticosteroid Injections: A Double-Blinded, Randomized Controlled Study. *Pain Physician*. 2015;18(2):163–71.
27. Brinks A, Koes BW, Volkens ACW, Verhaar JAN, Bierma-Zeinstra SMA. Adverse effects of extra-articular corticosteroid injections: a systematic review. *BMC Musculoskelet Disord* [Internet]. 2010

\* Corresponding author: Ariani Rahayu Setiandi, Master of Clinical Pharmacy Programme, Faculty of Pharmacy, University of Airlangga, Surabaya, Indonesia.

E-mail: [arianirahayus@gmail.com](mailto:arianirahayus@gmail.com)

Setiati and Hidayati: Effectiveness of Triamcinolone and Lidocaine in Patient With Piriformis Syndrome: A Case Series

Sep 13;11:206. Available from:  
<https://www.ncbi.nlm.nih.gov/pubmed/20836867>

---

\* Corresponding author: Ariani Rahayu Setiati, Master of Clinical Pharmacy Programme, Faculty of Pharmacy, University of Airlangga, Surabaya, Indonesia .

E-mail: [arianirahayus@gmail.com](mailto:arianirahayus@gmail.com)