

DAFTAR PUSTAKA

- Abdellatif, A. and Ali, M. (2013) 'Prevention of sevofluran related emergence agitation in children undergoing adenotonsillectomy: A comparison of dexmedetomidine and propofol', *Saudi Journal of Anaesthesia*, 7(3), pp. 296. doi: 10.4103/1658-354X.115363.
- Abu-Shahwan, I. (2008) 'Effect of propofol on emergence behavior in children after sevoflurane general anesthesia', *Pediatric Anesthesia*, 18(1), pp. 55–59. doi: 10.1111/j.1460-9592.2007.02376.x.
- Andriyanto, L. et al. (2019) 'Incidence of Emergence Agitation in Pediatric Patient after General Anesthesia', *Folia Medica Indonesiana*, 55(1), pp. 25. doi: 10.20473/fmi.v55i1.12546.
- Aouad, M. T. et al. (2007) 'A Single Dose of Propofol at the End of Surgery for the Prevention of Emergence Agitation in Children Undergoing Strabismus Surgery during Sevofluran Anesthesia', *Anesthesiology*, 107(5), pp. 733–738. doi: 10.1097/01.anes.0000287009.46896.a7.
- Bajwa, S. A., Costi, D. and Cyna, A. M. (2010) 'A comparison of emergence delirium scales following general anesthesia in children', *Pediatric Anesthesia*, 20(8), pp. 704–711. doi: 10.1111/j.1460-9592.2010.03328.x.
- Bedirli, N., Akçabay, M. and Emik, U. (2017) 'Tramadol vs dexmedetomidine for emergence agitation control in pediatric patients undergoing adenotonsillectomy with sevofluran anesthesia: Prospective randomized controlled clinical study', *BMC Anesthesiology*. *BMC Anesthesiology*, 17(1), pp. 1–7. doi: 10.1186/s12871-017-0332-4.
- Berghmans, J. M. et al. (2015) 'Does the Child Behavior Checklist predict levels of preoperative anxiety at anesthetic induction and postoperative emergence delirium? A prospective cohort study.', *Minerva anesthesiologica*, 81(2), pp. 145–56. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24861721>.
- Bilgen, S. et al. (2014) 'Effect of ketamine versus alfentanil following midazolam in preventing emergence agitation in children after sevofluran anaesthesia: A prospective randomized clinical trial', *Journal of International Medical Research*, 42(6), pp. 1262–1271. doi: 10.1177/0300060514543039.
- Blasiolo, B. and Davis, P. J. (2017) 'Intravenous Agents', in Davis, P. and Cladis, F. (eds) *Smith's Anesthesia for Infants and Children*. 9th edn. Elsevier Ltd, pp. 186–199.
- Bortone, L. et al. (2014) 'The effect of fentanyl and clonidine on early postoperative negative behavior in children: a double-blind placebo

- controlled trial', *Pediatric Anesthesia*. Edited by A. Bosenberg, 24(6), pp. 614–619. doi: 10.1111/pan.12388.
- Büttner, W. and Finke, W. (2000) 'Analysis of behavioural and physiological parameters for the assessment of postoperative analgesic demand in newborns, infants and young children: a comprehensive report on seven consecutive studies.', *Paediatric anaesthesia*, 10(3), pp. 303–18. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10792748>.
- Chandler, J. R. et al. (2013) 'Emergence delirium in children: A randomized trial to compare total intravenous anesthesia with propofol and remifentanyl to inhalational sevofluran anesthesia', *Paediatric Anaesthesia*, 23(4), pp. 309–315. doi: 10.1111/pan.12090.
- CHEN, J. et al. (2010) 'Emergence agitation after cataract surgery in children: a comparison of midazolam, propofol and ketamine', *Pediatric Anesthesia*, 20(9), pp. 873–879. doi: 10.1111/j.1460-9592.2010.03375.x.
- Chidambaran, V. and Costandi, A. (2016) 'PROPOFOL: A REVIEW OF ITS ROLE IN PEDIATRIC ANESTHESIA AND SEDATION', 29(7), pp. 543–563. doi: 10.1007/s40263-015-0259-6.PROPOFOL.
- Cho, E. J. et al. (2014) 'Comparison of the Effects of 0.03 and 0.05 mg/kg Midazolam with Placebo on Prevention of Emergence Agitation in Children Having Strabismus Surgery', *Anesthesiology*, 120(6), pp. 1354–1361. doi: 10.1097/ALN.0000000000000181.
- Choi, E. K. et al. (2019) 'Postoperative emergence agitation and intraoperative sevoflurane sedation under caudal block in children: a randomized comparison of two sevoflurane doses', *Anesthesia and Pain Medicine*, 14(4), pp. 434–440. doi: 10.17085/apm.2019.14.4.434.
- Cohen, I. T., Drewsen, S. and Hannallah, R. S. (2002) 'Propofol or midazolam do not reduce the incidence of emergence agitation associated with desflurane anaesthesia in children undergoing adenotonsillectomy', *Pediatric Anesthesia*, 12(7), pp. 604–609. doi: 10.1046/j.1460-9592.2002.00903.x.
- Cohen, I. T. et al. (2002) 'The effect of fentanyl on the emergence characteristics after desflurane or sevofluran anesthesia in children.', *Anesthesia and analgesia*, 94(5), p. 1178–81, table of contents. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11973185>.
- Cole, J. W. et al. (2002) 'Emergence behaviour in children: defining the incidence of excitement and agitation following anaesthesia.', *Paediatric anaesthesia*, 12(5), pp. 442–7. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12060332>.
- Costi, D. et al. (2014) 'Effects of sevofluran versus other general anaesthesia on emergence agitation in children', *Cochrane Database Syst Rev*, 9(9), p. Cd007084. doi: 10.1002/14651858.CD007084.pub2.

- Costi, D. et al. (2015) 'Transition to propofol after sevofluran anesthesia to prevent emergence agitation: A randomized controlled trial', *Paediatric Anaesthesia*, 25(5), pp. 517–523. doi: 10.1111/pan.12617.
- Cravero, J., Surgenor, S. and Whalen, K. (2000) 'Emergence agitation in paediatric patients after sevofluran anaesthesia and no surgery: a comparison with halothane.', *Paediatric anaesthesia*, 10(4), pp. 419–24. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10886700>.
- Creeley, C. et al. (2013) 'Propofol-induced apoptosis of neurones and oligodendrocytes in fetal and neonatal rhesus macaque brain', *British Journal of Anaesthesia*, 110, pp. i29–i38. doi: 10.1093/bja/aet173.
- Crosby, G. and Culley, D. J. (2011) 'Surgery and Anesthesia', *Anesthesia & Analgesia*, 112(5), pp. 999–1001. doi: 10.1213/ANE.0b013e3182160431.
- Dahmani, S. et al. (2010) 'Pharmacological prevention of sevofluran- and desflurane-related emergence agitation in children: a meta-analysis of published studies', *British Journal of Anaesthesia*, 104(2), pp. 216–223. doi: 10.1093/bja/aep376.
- Dahmani, S. et al. (2010) 'Pharmacological prevention of sevoflurane- and desflurane-related emergence agitation in children: a meta-analysis of published studies', *British Journal of Anaesthesia*, 104(2), pp. 216–223. doi: 10.1093/bja/aep376.
- Davis, P. J. et al. (1999) 'Recovery characteristics of sevofluran and halothane in preschool-aged children undergoing bilateral myringotomy and pressure equalization tube insertion.', *Anesthesia and analgesia*, 88(1), pp. 34–8. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9895062>.
- Eckenhoff, J. E., KNEALE, D. H. and Dripps, R. D. (1961) 'The incidence and etiology of postanesthetic excitement. A clinical survey.', *Anesthesiology*, 22, pp. 667–73. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/13889092>.
- Fleisher, L. E. E. A., Schwartz, A. J. A. Y. and Buck, D. (2014) 'Pediatric Anesthesiology', *Anesthesiology Clinics*, 32(1), pp. iii–vi. doi: 10.1016/S1932-2275(13)00110-9.
- Frawley, G. and Davidson, A. (2017) 'Inhaled anesthetic Inhaled anesthetic agents', in Peter Davis and Cladis, F. (eds) *Smith's Anesthesia for Infants and Children*. 9th edn. Elsevier Ltd, pp. 200–213.
- Frederick, H. J. et al. (2016) 'A Randomized Controlled Trial to Determine the Effect of Depth of Anesthesia on Emergence Agitation in Children.', *Anesthesia and analgesia*, 122(4), pp. 1141–6. doi: 10.1213/ANE.0000000000001145.
- Gaynor, J. and Ansermino, J. M. (2016) 'Paediatric total intravenous anaesthesia', *BJA Education*, 16(11), pp. 369–373. doi: 10.1093/bjaed/mkw019.

- Gowing, L. et al. (2016) 'Alpha 2 -adrenergic agonists for the management of opioid withdrawal', *Cochrane Database of Systematic Reviews*. doi: 10.1002/14651858.CD002024.pub5.
- Guo, J. et al. (2017) 'Emergence and Recovery Characteristics of Five Common Anesthetics in Pediatric Anesthesia: a Network Meta-analysis', *Molecular Neurobiology*. *Molecular Neurobiology*, 54(6), pp. 4353–4364. doi: 10.1007/s12035-016-9982-3.
- Hadi, S. M. et al. (2015) 'The effect of KETODEX on the incidence and severity of emergence agitation in children undergoing adenotonsillectomy using sevofluran based-anesthesia', *International Journal of Pediatric Otorhinolaryngology*, 79(5), pp. 671–676. doi: 10.1016/j.ijporl.2015.02.012.
- Hai, R. et al. (2020) 'Parental Presence Promoted Children Anesthesia Recovery: A Randomized Controlled Study', *Yangtze Medicine*, 04(01), pp. 54–61. doi: 10.4236/ym.2020.41005.
- Hosny, M. R. et al. (2019) 'Effect of IV Midazolam Premedication on the Recovery of Pediatric Patients after Isoflurane Anesthesia for Cochlear Implant Surgery', *The Open Anesthesia Journal*, 13(1), pp. 47–52. doi: 10.2174/2589645801913010047.
- Inomata, S. et al. (2010) 'Effects of fentanyl infusion on tracheal intubation and emergence agitation in preschool children anaesthetized with sevofluran', *British Journal of Anaesthesia*, 105(3), pp. 361–367. doi: 10.1093/bja/aeq168.
- Jalili, S. et al. (2019) 'Comparison of effects of propofol and ketofol (Ketamine-Propofol mixture) on emergence agitation in children undergoing tonsillectomy', *African Health Sciences*, 19(1), p. 1736. doi: 10.4314/ahs.v19i1.50.
- Jiang, S. et al. (2015) 'The efficacy of propofol on emergence agitation - a meta-analysis of randomized controlled trials', *Acta Anaesthesiologica Scandinavica*, 59(10), pp. 1232–1245. doi: 10.1111/aas.12586.
- Kain, Z. N. et al. (2006) 'Preoperative Anxiety, Postoperative Pain, and Behavioral Recovery in Young Children Undergoing Surgery', *Pediatrics*, 118(2), pp. 651–658. doi: 10.1542/peds.2005-2920.
- Kain, Z. N. et al. (2004) 'Preoperative Anxiety and Emergence Delirium and Postoperative Maladaptive Behaviors', *Anesthesia & Analgesia*, pp. 1648–1654. doi: 10.1213/01.ANE.0000136471.36680.97.
- Kanaya, A. (2016) 'Emergence agitation in children: risk factors, prevention, and treatment', *Journal of Anesthesia*, 30(2), pp. 261–267. doi: 10.1007/s00540-015-2098-5.
- Kanaya, A. et al. (2014) 'Lower incidence of emergence agitation in children after propofol anesthesia compared with sevofluran: A meta-analysis of

- randomized controlled trials', *Journal of Anesthesia*, 28(1), pp. 4–11. doi: 10.1007/s00540-013-1656-y.
- Kazmierski, J. et al. (2013) 'Cortisol levels and neuropsychiatric diagnosis as markers of postoperative delirium: a prospective cohort study', *Critical Care*, 17(2), p. R38. doi: 10.1186/cc12548.
- Kim, M.-S. et al. (2013) 'Comparison of propofol and fentanyl administered at the end of anaesthesia for prevention of emergence agitation after sevofluran anaesthesia in children', *British Journal of Anaesthesia*, 110(2), pp. 274–280. doi: 10.1093/bja/aes382.
- Kim, N. et al. (2017) 'Effects of intravenous fentanyl around the end of surgery on emergence agitation in children: Systematic review and meta-analysis', *Paediatric Anaesthesia*, 27(9), pp. 885–892. doi: 10.1111/pan.13181.
- Lee, C. J. et al. (2010) 'The effect of propofol on emergence agitation in children receiving sevofluran for adenotonsillectomy', *Korean Journal of Anesthesiology*, 59(2), p. 75. doi: 10.4097/kjae.2010.59.2.75.
- Lehmann, V. et al. (2015) 'Postoperative Agitation im Kindesalter', *Der Anaesthetist*, 64(5), pp. 373–380. doi: 10.1007/s00101-015-0026-6.
- Li, J. et al. (2011) 'Sufentanil reduces emergence agitation in children receiving sevofluran anesthesia for adenotonsillectomy compared with fentanyl.', *Chinese medical journal*, 124(22), pp. 3682–5. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22340224>.
- Li, J. et al. (2011) 'Sufentanil reduces emergence agitation in children receiving sevoflurane anesthesia for adenotonsillectomy compared with fentanyl.', *Chinese medical journal*, 124(22), pp. 3682–5. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22340224>.
- Liang, P. et al. (2014) 'Single-dose sufentanil or fentanyl reduces agitation after sevofluran anesthesia in children undergoing ophthalmology surgery', *Pakistan Journal of Medical Sciences*, 30(5). doi: 10.12669/pjms.305.4483.
- Malarbi, S. et al. (2011) 'Characterizing the behavior of children emerging with delirium from general anesthesia', *Pediatric Anesthesia*, 21(9), pp. 942–950. doi: 10.1111/j.1460-9592.2011.03646.x.
- Malherbe, S. et al. (2010) 'Total intravenous anesthesia and spontaneous respiration for airway endoscopy in children - a prospective evaluation', *Pediatric Anesthesia*, 20(5), pp. 434–438. doi: 10.1111/j.1460-9592.2010.03290.x.
- Mason, K. P. (2017) 'Paediatric emergence delirium: A comprehensive review and interpretation of the literature', *British Journal of Anaesthesia*, 118(3), pp. 335–343. doi: 10.1093/bja/aew477.
- Oofuvong, M. et al. (2013) 'Comparison the Incidence of Emergence Agitation between Sevo fl urane and Des fl urane after Pediatric Ambulatory Urologic Surgery', 96(11), pp. 1470–1475.

- Pickard, A. et al. (2014) 'Systematic review and meta-analysis of the effect of intraoperative $\alpha 2$ -adrenergic agonists on postoperative behaviour in children', *British Journal of Anaesthesia*, 112(6), pp. 982–990. doi: 10.1093/bja/aeu093.
- Pieters, B. J. et al. (2010) 'Emergence delirium and postoperative pain in children undergoing adenotonsillectomy: A comparison of propofol vs sevofluran anesthesia', *Paediatric Anaesthesia*, 20(10), pp. 944–950. doi: 10.1111/j.1460-9592.2010.03394.x.
- Sessler, C. N. et al. (2002) 'The Richmond Agitation–Sedation Scale', *American Journal of Respiratory and Critical Care Medicine*, 166(10), pp. 1338–1344. doi: 10.1164/rccm.2107138.
- Shi, F. et al. (2015) 'Effects of fentanyl on emergence agitation in children under sevofluran anesthesia: Meta-analysis of randomized controlled trials', *PLoS ONE*, 10(8), pp. 1–18. doi: 10.1371/journal.pone.0135244.
- Somainsi, M. et al. (2016) 'Emergence delirium or pain after anaesthesia - How to distinguish between the two in young children: A retrospective analysis of observational studies', *British Journal of Anaesthesia*. The Author(s), 116(3), pp. 377–383. doi: 10.1093/bja/aev552.
- Sriswasdi, P. et al. (2019) 'Predisposing Factors of Emergence Agitation in Pediatric Anesthesia', *J Med Assoc Thai*, 102(10), pp. 1065–1072.
- Stamper, M. J. et al. (2014) 'Identifying Pediatric Emergence Delirium by Using the PAED Scale: A Quality Improvement Project', *AORN Journal*. Elsevier Ltd, 99(4), pp. 480–494. doi: 10.1016/j.aorn.2013.08.019.
- Uezono, S. et al. (2000) 'Emergence agitation after sevoflurane versus propofol in pediatric patients', *Anesthesia and Analgesia*, 91(3), pp. 563–566. doi: 10.1213/00000539-200009000-00012.
- Van Hoff, S. L. et al. (2015) 'Does a prophylactic dose of propofol reduce emergence agitation in children receiving anesthesia? A systematic review and meta-analysis', *Paediatric Anaesthesia*, 25(7), pp. 668–676. doi: 10.1111/pan.12669.
- Vlajkovic, G. P. and Sindjelic, R. P. (2007) 'Emergence delirium in children: Many questions, few answers', *Anesthesia and Analgesia*, 104(1), pp. 84–91. doi: 10.1213/01.ane.0000250914.91881.a8.
- Vittinghoff, M. et al. (2018) 'Postoperative pain management in children: Guidance from the pain committee of the European Society for Paediatric Anaesthesiology (ESPA Pain Management Ladder Initiative)', *Pediatric Anesthesia*, 28(6), pp. 493–506. doi: 10.1111/pan.13373.
- Vlajkovic, G. P. and Sindjelic, R. P. (2007) 'Emergence delirium in children: Many questions, few answers', *Anesthesia and Analgesia*, 104(1), pp. 84–91. doi: 10.1213/01.ane.0000250914.91881.a8.
- Voepel-Lewis, T., Malviya, S. and Tait, A. R. (2003) 'A prospective cohort study of emergence agitation in the pediatric postanesthesia care unit.',

Anesthesia and analgesia, 96(6), p. 1625–30, table of contents. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12760985>.

Ydemann, M. et al. (2018) ‘Intraoperative clonidine for prevention of postoperative agitation in children anaesthetised with sevofluran (PREVENT AGITATION): a randomised, placebo-controlled, double-blind trial’, *The Lancet Child and Adolescent Health*, 2(1), pp. 15–24. doi: 10.1016/S2352-4642(17)30127-X.

Ydemann, M. et al. (2016) ‘Effect of clonidine to prevent agitation in children after sevofluran anaesthesia: a randomised placebo controlled multicentre trial.’, *Danish medical journal*, 63(6). Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27264945>.

Zainuddin, K. et al. (2016) ‘Abstract PR281’, *Anesthesia & Analgesia*, 123, pp. 362–363. doi: 10.1213/01.ane.0000492678.76530.18.

Zhong, Q., Qu, X. and Xu, C. (2018) ‘Effect of preoperative visiting operation room on emergence agitation in preschool children under sevofluran anesthesia’, *International Journal of Pediatric Otorhinolaryngology*. Elsevier, 104(September 2017), pp. 32–35. doi: 10.1016/j.ijporl.2017.10.038.