



Source details

Hypertension in Pregnancy

Formerly known as: Clinical and Experimental Hypertension - Part B Hypertension in Pregnancy
[Open Access](#) ⓘ

Scopus coverage years: from 1982 to Present

Publisher: Taylor & Francis

ISSN: 1064-1955 E-ISSN: 1525-6065

Subject area: Medicine: Obstetrics and Gynecology Medicine: Internal Medicine

Source type: Journal

CiteScore 2022

3.6 ⓘ

SJR 2022

0.451 ⓘ

SNIP 2022

0.638 ⓘ

[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology ×

CiteScore 2022 counts the citations received in 2019-2022 to articles, reviews, conference papers, book chapters and data papers published in 2019-2022, and divides this by the number of publications published in 2019-2022. [Learn more >](#)

CiteScore 2022 ▾

$$3.6 = \frac{549 \text{ Citations 2019 - 2022}}{154 \text{ Documents 2019 - 2022}}$$

Calculated on 05 May, 2023

CiteScoreTracker 2023 ⓘ

$$3.1 = \frac{428 \text{ Citations to date}}{138 \text{ Documents to date}}$$

Last updated on 05 October, 2023 • Updated monthly

CiteScore rank 2022 ⓘ

Category	Rank	Percentile
Medicine		
Obstetrics and Gynecology	#75/197	62nd
Medicine		
Internal Medicine	#74/140	47th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)

SJR Scimago Journal & Country Rank Enter Journal Title, ISSN or Publisher Name

Home Journal Rankings Country Rankings Viz Tools Help About Us

Hypertension in Pregnancy

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
United States Universities and research institutions in United States Media Ranking in United States	Medicine <ul style="list-style-type: none"> Internal Medicine Obstetrics and Gynecology 	Informa Healthcare	50
PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Journals	10641955, 15256065	1982-2022	Homepage How to publish in this journal

SCOPE


Hypertension in Pregnancy is a refereed journal in the English language which publishes data pertaining to human and animal hypertension during gestation. Contributions concerning physiology of circulatory control, pathophysiology, methodology, therapy or any other material relevant to the relationship between elevated blood pressure and pregnancy are acceptable. Published material includes original articles, clinical trials, solicited and unsolicited reviews, editorials, letters, and other material deemed pertinent by the editors.

Join the conversation about this journal

Quartiles

FIND SIMILAR JOURNALS options

1 Pregnancy Hypertension NLD 86% similarity	2 Journal of Maternal-Fetal and Neonatal Medicine GBR 62% similarity	3 Journal of Pregnancy USA 55% similarity	4 Journal of Obstetrics and Gynaecology GBR 53% similarity	5 Journal of Obstetrics and Gynaecology Research AUS 52% similarity
---	--	---	--	---



Metrics based on Scopus® data as of April 2023

[Home](#) ▶ [All Journals](#) ▶ [Hypertension in Pregnancy](#) ▶ [List of Issues](#) ▶ [Volume 39, Issue 3](#)

Hypertension in Pregnancy

[Submit an article](#) ▼[About this journal](#) ▼[Browse all articles & issues](#) ▼[Alerts & RSS feed](#) ▼

Taylor & Francis Online

[Log in](#)[Register](#)[Cart](#)

HYPERTENSION
IN
PREGNANCY

VOLUME 39
NUMBER 3
2020

Browse this journal

- [▶ Current issue](#)
- [▶ List of issues](#)
- [▶ Most read articles](#)
- [▶ Most cited articles](#)

Sample our
Medicine, Dentistry, Nursing
& Allied Health Journals

>> [Sign in here](#) to start your access
to the latest two volumes for 14 days

Hypertension in Pregnancy, Volume 39, Issue 3 (2020)

Hypertension in Pregnancy

◀ Issue Issue Issue Iss ▶

Submit an
article ▼

About this
journal ▼

Browse all articles &
issues ▼

Alerts & RSS
feed ▼



Browse by section (All) ▼



Display order (Default) ▼



Download citations

Download PDFs

Download issue

Research Article



Article

Magnesium intoxication in women with preeclampsia with severe features treated with magnesium sulfate >

Muhammad Ilham Aldika Akbar, Daniel Yoseph, Aditiawarman -, Muhammad Adrianes Bachnas, Erry Gumilar Dachlan, Gustaaf Albert Dekker & Ernawati

Pages: 221-227

Published online: 25 Apr 2020

335

Views

3

CrossRef citations

5

Altmetric



Article

Decreased placental IL9 and IL9R in preeclampsia impair trophoblast cell proliferation, invasion, and angiogenesis >

Yi Sun, Siyu Liu, Rong Hu, Qiongjie Zhou & Xiaotian Li

Pages: 228-235

Published online: 24 Apr 2020

186

Views

5

CrossRef citations

5

Altmetric



Article

Factors contributing to late stillbirth among women with

Hypertension in Pregnancy

Published online: 12 May 2020

Submit an
article ▾

About this
journal ▾

Browse all articles &
issues ▾

Alerts & RSS
feed ▾



Article

The diagnostic value and regulatory mechanism of miR-200a targeting *ZEB1* in pregnancy-induced hypertension >

Gui-Mei Wu, Yan Jin, Yan-Min Cao & Ji-Yun Li

Pages: 243-251

Published online: 29 Apr 2020

81

Views

7

CrossRef citations

Altmetric



Article

Antenatal and postpartum comparison of HD-OCT findings of macula, retinal nerve fiber layer, ganglion cell density between severe preeclampsia patients and healthy pregnant woman >

Abdullah Tok & Abdullah Beyoğlu

Pages: 252-259

Published online: 28 Apr 2020

136

Views

1

CrossRef citations

Altmetric



Article

Serum levels of uric acid may have a potential role in the management of immediate delivery or prolongation of pregnancy in severe preeclampsia >

Dan Liu, Chunfang Li, Pu Huang, Jing Fu, Xin Dong, Yunhui Tang, Xuelan Li & Qi Chen

Pages: 260-266

Published online: 29 Apr 2020

Hypertension in Pregnancy

[Submit an article](#) ▼[About this journal](#) ▼[Browse all articles & issues](#) ▼[Alerts & RSS feed](#) ▼

Repeated episodes of whole-body hypoxic preconditioning in rat model of preeclampsia >

Yan Li, Chunyun Wang, Jing Wang & Leisi Tao

Pages: 267-282

Published online: 13 May 2020

63

Views

1

CrossRef citations

Altmetric



Article

Sham surgical embryo transfer affects offspring neurodevelopment and manifestation of hypertensive phenotype in ISIAH rats >

Irina Rozkova, Eugeny Brusentsev, Tatyana Igonina, Diana Ragaeva, Olga Petrova, Maria A. Tikhonova, Anna A. Akopyan, Tamara G. Amstislavskaya, Yegor V. Antonov & Sergey Amstislavsky

Pages: 283-294

Published online: 13 May 2020

92

Views

2

CrossRef citations

Altmetric



Article

Impact of physical activity and traumatic exposure on occurrence of gestational hypertension: a survey of pregnant women in an armed-conflict region in Nigeria >

Aliyu Lawan, Cecelia Apeyemi, Muhammad Chutiyami, Umar Muhammad Bello, Dauda Salihu, Buhari Abdullahi Tafida, Umar Abubakar & Adamu Ahmad Rufa'i

Pages: 295-301

Published online: 18 May 2020

166

Views

0

CrossRef citations

Altmetric

Hypertension in Pregnancy

Article

[Fetal gender, serum human chorionic gonadotropin, and](#)

Submit an
article ▾

About this
journal ▾

Browse all articles &
issues ▾

Alerts & RSS
feed ▾

Taha

Pages: 302-307

Published online: 14 May 2020

100

Views

5

CrossRef citations

Altmetric

Article

[Cystatin C as an indicator of renal damage in pre-eclampsia](#) >

Hellen Cássia Dos Santos Gomes, Antônio Carlos Vieira Cabral, Silvia Passos Andrade, Henrique Vítor Leite, Patrícia Gonçalves Teixeira, Paula Peixoto Campos & Jaqueline Amorim Arantes Gomes

Pages: 308-313

Published online: 19 May 2020

108

Views

4

CrossRef citations

Altmetric

Article

[The comparison of side effects of methyldopa, amlodipine, and metoprolol in pregnant women with chronic hypertension](#) >

Masoumeh Ghafarzadeh, Amir Shakarami, Fatemeh Yari & Parsa Namdari

Pages: 314-318

Published online: 18 May 2020

496

Views

6

CrossRef citations

Altmetric

Article

[Carbetocin versus oxytocin for prevention of postpartum](#)

Hypertension in Pregnancy

Pages: 319-325

[Submit an article](#) ▼

[About this journal](#) ▼

[Browse all articles & issues](#) ▼

[Alerts & RSS feed](#) ▼

[Views](#) | [CrossRef citations](#) | [Altmetric](#)


Article

Predisposition to superimposed preeclampsia in women with chronic hypertension: endothelial, renal, cardiac, and placental factors in a prospective longitudinal cohort >

Kate Bramham, Pia M. Villa, Jennifer R. Joslin, Hannele Laivuori, Esa Hämäläinen, Eero Kajantie, Katri Räikkönen, Anukatriina Pesonen, Paul Seed, R Neil Dalton, Charles Turner, Max Wong, Peter Von Dadelszen, James M Roberts, Lucilla Poston & Lucy C Chappell

Pages: 326-335

Published online: 01 Jun 2020

372

Views

10

CrossRef citations

Altmetric



Article

An insertion-deletion polymorphism in angiotensin-converting enzyme is associated with a reduced risk of preeclampsia: an evidence-based meta-analysis from 44 studies >

Ting Huang, Yu Yan, Jianxin Li, Hong Chen & Zhen Chen

Pages: 336-347

Published online: 02 Jun 2020

30

Views

2

CrossRef citations

Altmetric



Article

Relationship between ABO blood groups and preeclampsia >

Nanhathai Mahasub & Dittakarn Boriboonhirunsarn

Home ▶ All Journals ▶ Hypertension in Pregnancy ▶ List of Issues ▶ Volume 39, Issue 3

Hypertension in Pregnancy

Submit an article ▼

About this journal ▼

Browse all articles & issues ▼

Alerts & RSS feed ▼

○ Association between intrahepatic cholestasis of pregnancy and risk of gestational diabetes and preeclampsia: a systematic review and meta-analysis >

Ahmed Arafa & Jia-Yi Dong

Pages: 354-360

Published online: 24 Apr 2020

415

Views

20

CrossRef citations

Altmetric

○ Complications and outcomes of SARS-CoV-2 in pregnancy: where and what is the evidence? >

Ayssa Teles Abrao Trad, Eniola R. Ibirogbá, Amro Elrefaei, Kavita Narang, Gabriele Tonni, Olivier Picone, Anna Suy, Elena Carreras Moratonas, Mark D. Kilby & Rodrigo Ruano

Pages: 361-369

Published online: 26 May 2020

1624

Views

15

CrossRef citations

Altmetric

Explore articles

Latest

Most cited

Trending

Explore the most recently published articles

Women with preeclampsia may have reduced risk of breast cancer: a



[Home](#) ▶ [All Journals](#) ▶ [Hypertension in Pregnancy](#) ▶ [List of Issues](#) ▶ [Volume 39, Issue 3](#)

Hypertension in Pregnancy

Genome-wide association study reveals HSF2, CIA1 and TRIM36 as

[Submit an article](#) ▼

[About this journal](#) ▼

[Browse all articles & issues](#) ▼

[Alerts & RSS feed](#) ▼

study in Hanjin, China >

Zhenhua Cao et al.

Article | Published online: 21 Sep 2023



[Cardiorespiratory fitness in women after severe pre-eclampsia](#) >

Lasse Gronningsaeter et al.

Article | Published online: 9 Aug 2023



[First-trimester maternal renin-angiotensin-aldosterone system activation and the association with maternal telomere length after natural and IVF/ICSI conceived pregnancies: the Rotterdam periconception cohort](#) >

D. Aoulad Fares et al.

Article | Published online: 21 Jul 2023





Hypertension in Pregnancy

[Submit an article](#) ▼[About this journal](#) ▼[Browse all articles & issues](#) ▼[Alerts & RSS feed](#) ▼

Ready to submit?

Start a new manuscript submission or continue a submission in progress

[Go to submission site](#) ↗

Submission information

[Instructions for authors](#)[Editorial policies](#) ↗

Editing services

[Editing services site](#) ↗

About this journal

[Journal metrics](#)[Aims & scope](#)[Journal information](#)[Special issue information](#)[Editorial board](#)[News & call for papers](#)[Advertising information](#)

Editorial board

Editor

Lana McClements, University of Technology Sydney, Sydney, Australia

Statistics Editor

Natasa M. Milic, Institute for Medical Statistics and Informatics, University of Belgrade, Serbia

Associate Editors

Stefan Hanson, Lund University, Lund, Sweden

Babbette LaMarca, University of Mississippi Medical Center, Mississippi, USA

Sonja Suvakov, Mayo Clinic, Rochester, USA

Editorial Board

Dr Bhavisha Bakrania, University of Queensland, Australia

Mathias Fasshauer, University of Leipzig, Germany

Annemarie Hennessy, University of Western Sydney, Penrith, Australia


Wolfgang Holzgreve, Institute of Advanced Study, Germany

Ananth Karumanchi, Harvard Medical School, Boston, USA

Dr Natasa Orlic, Ob/Gyn Clinic Narodni front, School of Medicine, University of Belgrade, Serbia

Holger Stepan, University of Leipzig, Germany

Ravi Thadhani, Harvard Medical School, Boston, USA



Magnesium intoxication in women with preeclampsia with severe features treated with magnesium sulfate

Muhammad Ilham Aldika Akbar, Daniel Yoseph, Aditiawarman -,
Muhammad Adrianes Bachnas, Erry Gumilar Dachlan, Gustaaf Albert
Dekker & Ernawati

To cite this article: Muhammad Ilham Aldika Akbar, Daniel Yoseph, Aditiawarman -, Muhammad Adrianes Bachnas, Erry Gumilar Dachlan, Gustaaf Albert Dekker & Ernawati (2020): Magnesium intoxication in women with preeclampsia with severe features treated with magnesium sulfate, Hypertension in Pregnancy, DOI: [10.1080/10641955.2020.1754851](https://doi.org/10.1080/10641955.2020.1754851)

To link to this article: <https://doi.org/10.1080/10641955.2020.1754851>



Published online: 25 Apr 2020.



Submit your article to this journal [↗](#)






View related articles [↗](#)



View Crossmark data [↗](#)



Magnesium intoxication in women with preeclampsia with severe features treated with magnesium sulfate

Muhammad Ilham Aldika Akbar ^{a,b,c}, Daniel Yoseph^b, Aditiawarman^{a,b}, Muhammad Adrianes Bachnas ^d, Erry Gumilar Dachlan^{a,b,c}, Gustaaf Albert Dekker^{a,e}, and Ernawati ^{a,b,c}

^aDepartment Obstetrics and Gynecology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia; ^bDepartment Obstetrics and Gynecology, DR. Soetomo General Academic Hospital, Surabaya, Indonesia; ^cDepartment Obstetrics and Gynecology, Universitas Airlangga Hospital, Surabaya, Indonesia; ^dDepartment Obstetrics and Gynecology, Dr. Moewardi General Hospital, Faculty of Medicine Universitas Sebelas Maret, Solo, Indonesia; ^eDepartment Obstetrics and Gynecology, Lyell-McEwin Hospital, The University of Adelaide, Adelaide, Australia

ABSTRACT

Objective: To evaluate the maternal-neonatal outcome in magnesium (Mg)-intoxicated women with preeclampsia with severe features (PESF) treated with magnesium sulfate (MgSO₄).

Methods: A total of 19 Mg intoxicated PESF women (cases) were compared with 166 PESF women without signs of intoxication (controls).

Results: Mg serum levels of cases was higher compared to control group (12.36 ± 3.54 mg/dl versus 2.69 ± 0.83 mg/dl). 3 women died and 3 had major maternal morbidity in cases group compared with zero in the control group (P = 0.009). Mg intoxication was also significantly associated with perinatal deaths and low Apgar scores at 1 and 5 minutes.

Conclusion: Mg intoxication is associated with a increased risk of maternal and perinatal mortality and morbidity.

ARTICLE HISTORY

Received 13 November 2019
Accepted 7 April 2020

KEYWORDS

Magnesium intoxication;
hypermagnesemia;
magnesium sulfate;
preeclampsia with severe
features

Introduction

Preeclampsia (PE) is a life-threatening syndrome in pregnancy characterized by the new onset of hypertension after 20 weeks of gestational age and accompanied by proteinuria or other signs of organ involvement (1). PE is one of the main problems in pregnancy, causing maternal mortality and morbidity in low and middle-income countries (Duley et al., 1992). About 50,000–63,000 women die each year of PE/eclampsia. Indonesia, the fourth most populous country in the world with a population of 267 million, is a middle-income country with PE incidence of 5–7%. The maternal (2.2%) and perinatal mortality rate (12%) of PE are still high (2). A large segment of the Indonesian population still faces poverty, has lack of access to health care, and often receives inadequate treatment in primary health-care centers. The wrong financial incentives under the current Indonesian universal health coverage system are partially to blame for too late referrals (3).

Magnesium sulfate (MgSO₄) is recommended unambiguously by all major guidelines as a first-choice agent to prevent eclamptic seizures and is used worldwide (4). The Magpie trial, the large international multicenter randomized controlled trial (RCT),

reported that the use of MgSO₄ reduces the risk of eclampsia about 58% in women with PE compared with PE women receiving placebo (5). MgSO₄ as an anticonvulsant agent works through multiple mechanisms: inhibits seizure excitability in the cerebral cortex, inhibits the N-methyl-D-aspartate receptor in the hippocampus, and calcium metabolism in the neuromuscular junction. MgSO₄ also works as a potent vasodilator, especially in the cerebral vasculature, potentially reducing brain ischemia-hypoxia in preeclamptic women (6). Although MgSO₄ has been proven to be effective in preventing eclampsia, its use still contains a small but definitive risk of magnesium intoxication or hypermagnesemia (7–9).

Mg intoxication is clinically evident when serum Mg levels exceed 12 mg/dL (hypermagnesemia), which almost exclusively occurs in PE women with marked renal involvement or in women receiving higher MgSO₄ infusion rates (10). The main symptoms of Mg intoxication are the result of its general CNS depressant effects, peripheral depression affecting muscle contractility, and central neuromuscular transmission blockade. Clinical manifestation of Mg

intoxication seldom occurs until magnesium levels reach a total dosage beyond 12 mg/dL in maternal blood. Clinical manifestations of Mg intoxication include general weakness, double vision, low blood pressure, loss of conscious and respiratory distress. The lethal manifestation, cardiac arrest, may happen if Mg concentration rises above 30 mg/dL (8,11,12).

Hypertension in pregnancy, especially PESF is in the top three diagnosis of the sickest high-risk obstetrics cases in Dr. Soetomo General Academic Hospital, the main tertiary referral hospital in Surabaya, Indonesia (13,14). Intravenous MgSO₄ is administrated routinely in PESF without measurements of Mg levels unless there are clinical indications (suspicion Mg intoxication) due to lack of adequate hospital funding under the Indonesian national insurance system.

In this study, we evaluated the incidence of Mg intoxication in women with PESF receiving MgSO₄ and investigated its association with adverse maternal and perinatal outcomes.

Material and methods

This study was conducted in Dr. Soetomo Hospital, the major tertiary referral center in East-Java, Indonesia, from January 2014 – December 2018; all women with PESF receiving Mg with clinical signs and symptoms of Mg intoxication and confirmed hypermagnesemia (cases n = 19) were compared to 166 PESF patients (8 controls per case) also treated with Mg but without signs of Mg intoxication (control group). The historical control group was recruited randomly with 33–34 patients each year to achieve a balanced distribution (2014–2018). Since Mg levels are not routinely checked in our hospital in PESF patients receiving MgSO₄, Mg levels in this study were only measured in women with signs or symptoms of Mg intoxication. In the control group, 24 random women out the 166 also had a Mg level measured. Mg level was measured using the Calmagite Colorimetric method (Hardness reagen set).

PE was defined using ISSHP classification: hypertension developing after 20 weeks gestation and the existence of one or more of the following: proteinuria, other maternal organ dysfunction (renal insufficiency, liver involvement, neurological complications, hematological complications), and uteroplacental dysfunction. HELLP syndrome was defined as a combination of hemolysis (lactic dehydrogenase >1000 U/L, schistocytes in blood smear), elevated liver enzymes, and thrombocytopenia (platelet count < 100.000 u/L) in PE women. PESF was defined as PE with any of the following findings: systolic blood pressure > 160 mmHg or diastolic blood pressure > 110 mmHg on two

occasions at least 4 hours apart, thrombocytopenia, elevated liver enzymes (AST > 45 µ/L, ALT > 35 µ/L), progressive renal insufficiency (BUN > 20 mg/dL, serum kreatinin > 1.1 mg/dL), pulmonary edema, and cerebral or visual disturbances. Pulmonary edema was defined based on symptoms of shortness of breath/difficulty breathing, physical examination, and confirmed with a chest X-ray (15). Hypertensive crisis was defined as systolic blood pressure more than 180 mmHg and/or diastolic blood pressure more than 120 mmHg (1,4,16,17).

Every patients with PESF in our hospital received MgSO₄, using the Zuspan regimen consisting of a loading dose 4 g in 20 ml (20% solution) administered iv over 15–20 minutes, followed by a maintenance dose of 1 g/hour iv infusion using syringe pump from admission until 12 hours or 24 hours after delivery in case of eclampsia (18). In preterm PESF women managed conservatively, MgSO₄ is administered the first 24 hours following admission. In these women, MgSO₄ is discontinued during their observation period and started again at the time of delivery. On the other hand, the regimen mostly used in primary health-care centers consists of a modified Pritchard scheme; loading dose of 4 gram 20% iv MgSO₄ in 10–15 minutes and 10 gram IM (5 gram in each buttock), followed by maintenance dose 5 gram IM MgSO₄ in alternate buttock for every 6 hours (19).

Cases of Mg intoxication were identified as women who received MgSO₄ treatment and had documented clinical signs of magnesium intoxication in addition to serum magnesium levels >12 mg/dL. Clinical signs and symptoms included muscles weakness, loss of deep tendon reflexes, respiratory paralysis, ECG changes (prolonged PR interval and widened QRS complex), SA or AV node block, loss of consciousness or cardiac arrest.

The primary outcomes of the study were maternal and perinatal outcomes among the 19 cases compared with the 166 controls. The maternal parameters evaluated included maternal age, body mass index (BMI), parity, referral origin, antenatal history, PE type. Maternal outcomes included mode of delivery, gestational age at delivery, laboratory results, maternal complications, and maternal death. PE complications included any of the following: eclampsia, HELLP syndrome, pulmonary edema, hypertensive crisis, acute kidney injury. The definition of maternal death in this study was any death during the treatment in hospital, and not necessarily a direct effect of Mg intoxication. Perinatal outcomes included fetal sex, birth weight, Apgar scores (minutes 1 & 5), SGA (<10th population birth weight centile), and perinatal death.

The results were analyzed using chi-square test, independent t-test, Mann Whitney test, and Fisher exact test where appropriate. Statistical measurement was performed using IBM SPSS Statistics for Windows, Version 25.0. Armonk, NY: IBM Corp. Released 2017.

Results

Incidence of Mg intoxication in women with PESF

The incidence of PESF during the period 2014–2018 was 1743 cases out of a total of 6823 total deliveries (25.54%); Mg intoxication was diagnosed in 19 patients (1.09%). The annual incidence of Mg intoxication over this 5-year period varied between 0.6% and 1.5%.

Maternal characteristics

The overall characteristics between the two groups were quite similar (Table 1) except the origin of the cases. All Mg intoxicated cases were referred from other hospitals or primary health-care centers compared with 80.2% in the control group: 19.8% women in the control group had regular antenatal care in our hospital (booked case). Twelve (63.2%) cases with Mg intoxication were referred from distant rural areas. There was no significant difference in maternal age, BMI, parity, and PE type between both groups.

Maternal outcomes

The clinical manifestations of Mg intoxication were varied: 42.1% of women demonstrated muscle weakness (loss of patellar reflex), 10.5% respiratory depression, and 42.1% loss of consciousness. The maternal outcomes of cases versus controls are presented in Table 2. The mode of delivery between both groups was not different,

most patients in both groups were delivered by cesarean section.

The rate of maternal mortality and severe morbidities was higher among cases compared with the controls; 14 out of the 19 cases (73%) had a major complication, including 3 maternal deaths compared with 30.7% major complication rate in the control group (OR 2.85; 95% CI 2.12–3.82). Eclampsia and HELLP syndrome were the main complications found in the case group. Interestingly we found no pulmonary edema among the cases compared with 13 (7.8%) in the control group.

Importantly, the three maternal deaths were not directly caused by the actual Mg intoxication but were related to other complications such as septic shock, thyroid crisis, and intracerebral hemorrhage. Laboratory manifestations reflecting disease severity were significantly worse in the case group (protein urine, LDH, ALT, AST, BUN, SK, and Albumin).

A subgroup analysis was performed in PESF control women who had their serum Mg levels measured (Table 3). The Mg level among the cases was markedly elevated compared with controls (12.36 ± 3.45 versus 2.69 ± 0.83 mg/dL; $p < 0.001$). The minimum and maximum level of both groups were as follows: cases (8.9–25.6 mg/dL) versus controls (1.5–4.3 mg/dL). In addition, we found a significant difference in the total Mg dose, length, and method of Mg administration between the groups. A significantly higher proportion of cases received >24 g of Mg, and had Mg administered for >24 hours, and had Mg administered by intramuscular injection. As high as 21.1% of cases had oliguria.

Perinatal outcomes

Maternal Mg intoxication had a significant association with worse perinatal outcomes: lower 1 and 5 minute

Table 1. Maternal characteristics of PESF patients treated with Mg with (cases) and without Mg intoxication (controls).

	Cases n = 19	Controls n = 166	<i>p</i> value	OR (95% CI)
Maternal Age	28.83 + 6.191	31.96 + 6.769	0.415	NA
BMI	30.9875 + 8.694	30.6698 + 7.210	0.973	NA
Parity				
Primipara	4 (21.2%)	51 (31.1%)	0.349	NA
Multipara	15 (78.9%)	115 (68.9%)		
Referral origin				
Country	12 (63.2%)	42 (25.5%)	0.001*	5.02
The City	7 (36.8%)	124 (74.5%)		(1.85–13.59)
Booked Case				
No	19 (100%)	134 (80.2%)	0.005*	9.72
Yes	0	32 (19.8%)		(0.57–165.03)
PE type				
Early onset PE	14 (73.6%)	104 (62.7%)	0.314	NA
Late onset PE	5 (26.4%)	62 (37.3%)		

Table 2. Maternal outcomes of cases versus controls.

	Cases n = 19	Controls n = 166	p value	OR (95% CI)
GA at delivery	31.63 ± 3.51	33.81 ± 3.33	0.014*	NA
Mode of delivery				
CS	11 (57.9%)	110 (66.3%)	0.473	NA
Vaginal delivery	8 (42.1%)	56 (33.7%)		
Complication				
Yes	14 (73.7%)	51 (30.7%)	0.0002*	2.85
No	5 (26.3%)	115 (69.3%)		(2.12–3.82)
Complication type	14	51	0.0113*	NA
Eclampsia	6 (42.9%)	9 (17.6%)		
HELLP Syndrome	3 (21.4%)	11 (21.6%)		
Pulmonary edema	0	13 (25.5%)		
IUGR	1 (7.1%)	4 (7.8%)		
IUFD	3 (21.5%)	1 (2%)		
Multiple complication	0	9 (17.6%)		
Emergency Hypertension	0	1 (2%)		
Acute Kidney Injury	1 (7.1%)	3 (5.9%)		
Maternal death				
Yes	3 (15.8%)	0	0.009*	
No	16 (84.2%)	166 (100%)		
BUN	33.75 ± 20.56	10.93 ± 8.84	0.000*	
Serum Creatinine	2.61 ± 1.68	0.65 ± 0.07	0.000*	
Serum Albumin	2.64 ± 0.38	2.58 ± 0.55	0.000*	
ALT	59.58 ± 94.79	43.35 ± 12.72	0.057*	
AST	96.68 ± 203.56	21.25 ± 3.8	0.000*	
Protein Urine			0.024*	NA
(-)	0	3 (1.8%)		
+1	1 (5.3%)	20 (12.1%)		
+2	1 (5.3%)	56 (33.9%)		
+3	10 (52.6%)	55 (33.3%)		
≥4	7 (36.8%)	31 (18.8%)		

Table 3. Subgroup analysis of cases vs controls group.

	Cases n = 19	Controls n = 24	p value	OR (95% CI)
Mg level (g)	12.36 ± 3.45	2.69 ± 0.83	<0.001*	NA
Urine output (ml/hour)				
<30	4 (21.1%)	0	0.031*	2.6
≥30	15 (78.9%)	24 (100%)		(0.9–7.2)
Methods of MgSO ₄ administration				
im	10 (52.6%)	5 (20.8%)	0.004*	2.07
iv	9 (47.4%)	19 (79.2%)		(1.56–19.4)
Total Mg Dose (g)				
≥ 24	14 (73.7%)	11 (45.8%)	0.027*	2.01
< 24	5 (26.3%)	13 (54.2%)		(1.51–1.75)
Length of administration (hour)				
≥ 24	14 (73.7%)	11 (45.8%)	0.027*	2.01
< 24	5 (26.3%)	13 (54.2%)		(1.51–1.75)

Apgar scores, lower birth weights, and a higher SGA and perinatal death rate (Table 4). The perinatal death rate in the case group was 36.8% compared with 6.6% in the control group.

Discussion

The results of this study confirm that while the incidence of Mg intoxication in women with PESF treated with MgSO₄ is quite low (1.09%), Mg intoxication is associated with significantly increased adverse maternal and perinatal outcomes. The incidence of Mg intoxication in a large systematic review involving 9556 women,

was 1.3–1.6% (8). Duley et al., in another large meta-analytic study, also found that Mg intoxication was rare and only occurred in around 1% of women receiving MgSO₄ (20). Importantly, Mg intoxication was not seen in patients receiving the “Magpie protocol” (5).

Three maternal deaths occurred among the 19 Mg intoxicated women, but the cause of maternal death was not directly related to the Mg level. The cause of death in these three cases was: septic shock, thyroid crisis, and intracerebral hemorrhage. Serum Mg level in these three cases had normalized a couple of days before they died. Lowe et al. reported no maternal death in their large systematic review, while Duley

Table 4. Perinatal outcomes of cases versus controls.

	Cases n = 19	Controls n = 166	<i>p</i> value	OR (95% CI)
Baby sex				
Male	13 (68.4%)	79 (50.3%)	0.131	NA
Female	6 (31.6%)	78 (49.7%)		
Baby Birthweight	1516.67 + 565.015	1872.32 + 662.860	0.005*	NA
Apgar Score				
Minutes 1	2.75 + 2.179	5.57 + 2.250	0.000*	NA
Minutes 5	5.42 + 1.782	7.34 + 1.573	0.000*	NA
SGA				
Yes	6 (31.6%)	14 (8.4%)	0.002*	3.74
No	13 (68.4%)	152 (91.6%)		(2.66–11.38)
Perinatal Death				
Yes	7 (36.8%)	11 (6.6%)	0.000*	5.56
No	12 (63.2%)	155 (93.4%)		(2.29–5.94)

et al. found two trials reported maternal deaths (8,20). The highest serum Mg level found among the cases in the current study was 25.6 mg/dL, this level is not considered to be high enough to directly cause maternal death. With serum levels of 20–34 mg/dL, Mg intoxication will be manifest as hypoventilation, acidosis, loss of tendon reflexes, and general weakness. Severe and life-threatening complications like respiratory depression and cardiac arrest will occur with the serum levels of 48–72 mg/dL (8). The clinical signs and symptoms in these Mg intoxication cases were general muscle weakness (loss of patellar reflex), loss of consciousness, and respiratory distress. The overall incidence of loss of patellar reflexes, loss of consciousness, and respiratory depression among all 1743 patients receiving MgSO₄ treatment was, respectively, 0.4% (8 cases), 0.5% (9 cases), and 0.1% (2 cases). The side effects caused by MgSO₄ treatment found in this study were significantly lower compared with the large systematic review by Lowe et al. (8). This review is comparable to our study since the research sample was only taken from middle- and low-income countries. The overall incidence of loss of patellar reflexes and respiratory distress in this systematic review was 1.2% and 1.3% (8). Unfortunately, this review did not evaluate the clinical sign of loss of consciousness. Perhaps this sign is not commonly interpreted as one of the major manifestations of Mg intoxication. The possible explanation of the much higher incidence of side effects reported in Lowe et al. (8) review compared with our study may be related to the regimen choices. Only eight studies used the Zuspan regimen (which is the same regimen used in our hospital) (18), while the other studies used the Pritchard (19) or the Dhaka regimens (21) with a relatively higher total dose of MgSO₄. Administration of MgSO₄ longer than 24 hours and a total dose of more than 24

grams would be potentially hazardous for the maternal outcome, as seen in the current study.

Interestingly, intramuscular administration of MgSO₄ was associated with a doubling of the rate of Mg intoxication; 52.6% of patients in the case group received MgSO₄ intramuscular versus 21.8% in the control group (*p* = 0.004). Similar findings have been reported by Kanti et al. (21); these authors reported a higher rate of signs and symptoms indicative of Mg intoxication and local site complications to be associated with intramuscular Mg administration. The Magpie trial also showed that intravenous administration had fewer side effects compared with the intramuscular route (5).

In the current study, the majority of the cases received intramuscular MgSO₄, since most of these patients were referred from primary health centers or primary health care or rural areas. This is in line with the regional referral guidelines on the management of PESF. Our regional policy for community-based care (primary health care) is to give a loading dose following the Pritchard regimen as initial management of PESF, before transferring the patients to a tertiary center. This approach has been implemented in many low- and middle-income countries such as Bangladesh, and India (22). The majority of smaller Indonesian medical centers prefer Pritchard IM administration regimen because of ease of administration, non-availability of infusion set/syringe pump, lack of nursing staff, and the fact that intramuscular administration is more cost-effective compared with iv regimen (23). Encouraging the government to provide syringe-pump availability throughout the nation could potentially improve outcomes in Indonesian women with PESF.

In our tertiary referral center, MgSO₄ is given via an intravenous route (syringe pump infusion) following the Zuspan regimen (18,22). Maintaining the therapeutic level of Mg while monitoring for sign of Mg

intoxication is important. Simple clinical assessment is adequate in most PESF women treated with MgSO₄, and more suitable in low- and middle-income countries (5,12,22), as is also demonstrated by the low overall rate of just over 1% in the current series.

The incidence of eclampsia and HELLP syndrome was significantly higher in the case group, while unexpectedly acute pulmonary edema, a common PE complication in Indonesia (15) was not seen among the 19 Mg intoxication cases. The presence of these complications may represent a significant risk factor of Mg intoxication due to the clinical necessity for a more prolonged MgSO₄ administration. Among the cases, more women received MgSO₄ for more than 5 days compared with the controls, although the average length of hospital stay between both groups did not differ significantly (cases versus controls: 10.21 ± 8.43 versus 11.29 ± 15.20, *p* = 0.783). The study results appear to indicate that giving MgSO₄ for more than 24 hours is not advisable unless Mg levels can be checked at regular intervals.

Another significant risk factor that could be contributing to the occurrence of Mg intoxication is oliguria and renal insufficiency. In this series, Mg-intoxicated patients had a worse renal function, reflected by higher BUN and serum creatinine level. Almost all of the serum magnesium is cleared by renal excretion, so any problems of urine production will significantly increase the risk of magnesium intoxication (6).

All patients in the Mg intoxicated group received calcium gluconate therapy as immediate management in line with our national protocol. This routine intervention is not found in the large systematic review by Lowe et al. The use of calcium gluconate in their review was extremely rare (0.18%). There is no clear explanation of this finding. However, Lowe et al. assumed that the low use of calcium gluconate reflects an infrequent need for its use as an antidote, rather than the unavailability of the drugs in the hospital (8,24).

The case had a higher rate of perinatal death, SGA, and low Apgar scores. Seven perinatal deaths (4 fetal demise) (36.8%) occurred in Mg-intoxicated group compared with 11 deaths (6.6%) in the control group. The higher rate of SGA in the case group reflects disease severity. About one-third of the cases were delivered of neonates who developed respiratory distress syndrome (RDS), significantly different from the 4% in the control group. Low Apgar scores were also more prominent among cases, in line with the strong correlation between maternal and fetal Mg levels resulting in respiratory muscle suppression and general muscle weakness. Previously, the use of antenatal MgSO₄ was discouraged by the majority of neonatologists

because of a perceived risk of neonatal respiratory depression (25). However, recent studies with a large number of subjects have shown that particular concerns regarding the use of antenatal MgSO₄ for adverse neonatal implications were only encountered after long duration of Mg administration (26). We also discourage the prolonged use of antenatal MgSO₄ in PESF cases, not only because of the aforementioned neonatal concern but also because of the association with an increased risk of maternal and perinatal complications. The meta-analytic study by Duley et al. did not find any difference in the risk of stillbirth or neonatal death, but there was a small increase in the overall risk of perinatal mortality associated with MgSO₄ use (RR 1.04) (27).

Conclusion

Mg intoxication was found in just over 1% of PESF women treated with MgSO₄, as was found to be associated with a significant risk to the mother and fetus. While Mg serum levels cannot be measured as a routine in many developing countries, serial monitoring for the simple clinical signs and symptoms indicative of Mg intoxication is effective as a monitoring tool. In these countries, Mg serum level monitoring is indicated in patients requiring longer duration and higher accumulated doses of MgSO₄, both carry a high risk of Mg intoxication.


Acknowledgments

We thank Prof. Dr. Hendy Hendarto dr. SpOG(K) as the head of department who supported and gave permission to perform this study; we thank all staff in Department Obstetrics & Gynecology Faculty of Medicine Universitas Airlangga - RSUD Dr. Soetomo - RS Universitas Airlangga for their genuine support for this study; we thank our resident in training and the research team for their hard work in collecting data in this study; and we would like to thank all the patients who were involved in this study, without their participation the study could never be completed.

Disclosure statement

The authors declare no conflict of interest.

ORCID

Muhammad Ilham Aldika Akbar  <http://orcid.org/0000-0002-2003-9282>

Muhammad Adrianes Bachnas  <http://orcid.org/0000-0002-1710-3909>

Ernawati  <http://orcid.org/0000-0002-9344-3606>

References

- [1] Tranquilli AL, Dekker G, Magee L, et al. The classification, diagnosis and management of the hypertensive disorders of pregnancy: A revised statement from the ISSHP. *Pregnancy Hypertens.* 2014;4(2):97–104.
- [2] DULEY L. Maternal mortality associated with hypertensive disorders of pregnancy in Africa, Asia, Latin America and the Caribbean. *BJOG An Int J Obstet Gynaecol.* 1992;99(7):547–553.
- [3] Ilham M, Akbar A, Adrianes Bachnas M, et al. The massive problem of preeclampsia in indonesia: in need of a redesigned national health care system. *Pregnancy Hypertens.* 2018;13(1):S16
- [4] ACOG. Hypertension in pregnancy. *Obs Gynecol.* 2013;122(5):1122–1131.
- [5] Altman D, Carroli G, Duley L, et al. Do women with pre-eclampsia, and their babies, benefit from magnesium sulphate? The Magpie trial: a randomised placebo-controlled trial. *Lancet.* 2002;359(9321):1877–90.
- [6] Tannirandorn Y. Is magnesium sulfate for prevention or only therapeutic in preeclampsia? *J Med Assoc Thail.* 2005;88(7):1003–10
- [7] Sibai BM, Gary Cunningham F. Prevention of preeclampsia and eclampsia [Internet]. Fourth Edn. In: Chesley's Hypertens. Disord. Pregnancy. Elsevier Inc.; 2009. DOI:10.1016/B978-0-12-407866-6.00012-2
- [8] Lowe RF, Smith JM, Felker-Kantor E, et al. An integrative review of the side effects related to the use of magnesium sulfate for pre-eclampsia and eclampsia management. *BMC Pregnancy Childbirth.* 2013;13:1–11.
- [9] Tukur J. The use of magnesium sulphate for the treatment of severe pre-eclampsia and eclampsia. *Ann Afr Med.* 2009;8(2):76.
- [10] Gröber U, Schmidt J, Kisters K. Magnesium in prevention and therapy. *Nutrients.* 2015;7(9):8199–8226.
- [11] Jackson CB, Drobatz KJ. Iatrogenic magnesium overdose: 2 Case reports. *J Vet Emerg Crit Care.* 2004;14(2):115–123.
- [12] Grissinger M. Preventing magnesium toxicity in obstetrics. *Pharm Ther* [Internet]. 2009;34. Available from. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2799127/pdf/ptj34_8p403.pdf
- [13] Ernawati, Gumilar E, Kuntoro, Soeroso J, Dekker G. Expectant management of preterm preeclampsia in Indonesia and the role of steroids. *J Matern Neonatal Med.* 2016;29:1736–1740.
- [14] Kartika AR, Aldika Akbar MI, Umiastuti P. Risk factor of severe preeclampsia in Dr. Soetomo Hospital Surabaya in 2015. *Maj Obstet Ginekol.* 2017;25(1).
- [15] Wardhana MP, Dachlan EG, Dekker G. Pulmonary edema in preeclampsia: an Indonesian case-control study. *J Matern Neonatal Med.* 2018;31(6):689–695.
- [16] Cunningham FG, Roberts JM, Lindheimer MD. The clinical spectrum of preeclampsia [Internet]. Fourth Edn. In: Chesley's Hypertens. Disord. Pregnancy. Elsevier Inc.; 2009. DOI:10.1016/B978-0-12-407866-6.00002-X
- [17] Muiesan ML, Salvetti M, Lorenza M, et al. An update on hypertensive emergencies and urgencies Pietro Scicchitano i, Enrico Agabiti Rosei a, Roberto Pedrinelli n, on behalf of the Working Group on Hypertension, Prevention, Rehabilitation of. 2015;
- [18] Zuspan FP. Problems encountered in the treatment of pregnancy-induced hypertension. A point of view. *Am J Obstet Gynecol.* 1978;131(6):591–597.
- [19] Pritchard JA, Cunningham FG, Pritchard SA. The Parkland memorial hospital protocol for treatment of eclampsia: evaluation of 245 cases. *Am J Obstet Gynecol.* 1984;148(7):951–963.
- [20] Duley L, Am G, Dj H Magnesium sulphate and other anticonvulsants for women with pre-eclampsia (Review). *Cochrane Collab.* 2003;
- [21] Begum R, Begum A, Johanson R, et al. A low dose (“Dhaka”) magnesium sulphate regime for eclampsia. *Acta Obstet Gynecol Scand.* 2001;80(11):998–1002.
- [22] Gordon R, Magee LA, Payne B, et al. Magnesium sulphate for the management of preeclampsia and eclampsia in low and middle income countries: a systematic review of tested dosing regimens. *J Obstet Gynaecol Canada* [Internet]. 2014;36(2):154–163.
- [23] Kanti V. Comparison between intramuscular and intravenous regimen of magnesium sulfate in management of severe pre-eclampsia and eclampsia. *Int J Reprod Contraception Obstet Gynecol.* 2015;4:1.
- [24] Ridge AL, Bero LA, Hill SR. Identifying barriers to the availability and use of magnesium sulphate injection in resource poor countries: A case study in Zambia. *BMC Health Serv Res.* 2010;10(1). DOI:10.1186/1472-6963-10-340
- [25] Levene M, Blennow M, Whitelaw A, et al. Acute effects of two different doses of magnesium sulphate in infants with birth asphyxia. *Arch Dis Child.* 1995;73(3):174–177.
- [26] M R, P PG. A review of the role for magnesium sulphate in preterm labour [Internet]. *BJOG An Int J Obstet Gynaecol.* 2005;84–88. Available from <http://www.embase.com/search/results?subaction=viewrecord&from=export&id=L40313989>
- [27] Duley L, Farrell B, Armstrong N, et al. The Magpie Trial: A randomised trial comparing magnesium sulphate with placebo for pre-eclampsia. Outcome for children at 18 months. *BJOG An Int J Obstet Gynaecol.* 2007;114:289–299.