

Two novel coumarins bearing an acetophenone derivative from the leaves of Melicope Quercifolia

by Mulyadi Tanjung

Submission date: 16-Dec-2020 02:23PM (UTC+0800)

Submission ID: 1476556332

File name: 3._Two_novel_coumarins.pdf (848.18K)

Word count: 2975

Character count: 13736



Two novel coumarins bearing an acetophenone derivative from the leaves of *Melicope Quercifolia*

Ratih Dewi Saputri, Tjitjik Srie Tjahjandarie & Mulyadi Tanjung

3
To cite this article: Ratih Dewi Saputri, Tjitjik Srie Tjahjandarie & Mulyadi Tanjung (2019): Two novel coumarins bearing an acetophenone derivative from the leaves of *Melicope Quercifolia*, Natural Product Research, DOI: 10.1080/14786419.2019.1644634

To link to this article: 51
<https://doi.org/10.1080/14786419.2019.1644634>



View supplementary material [↗](#)




Published online: 22 Jul 2019.



Submit your article to this journal [↗](#)



View Crossmark data [↗](#)



Two novel coumarins bearing an acetophenone derivative from the leaves of *Melicope Quercifolia*

Ratih Dewi Saputri, Tjitjik Srie Tjahjandarie and Mulyadi Tanjung

Natural Products Chemistry Research Group, Organic Chemistry Division, Faculty of Science and Technology, Department of Chemistry, Universitas Airlangga, Surabaya, Indonesia

ABSTRACT

Meliquercifolins A (1), and B (2), two new coumarins are bearing an acetophenone derivative were isolated from the leaves of *Melicope quercifolia* along with three known compounds, melico-denines E (3), F (4) and I (5). Structures of two new compounds were identified based on spectroscopic analyses (UV, HR-ESI-MS, 1-D and 2D NMR). Cytotoxic activities of compounds (1–5) towards three human cancer cells (HeLa, MCF-7, P-388), compounds 1, 4 and 5 showed very potent activity against Hela cells with IC₅₀ values 2.6; 0.8; 1.1 μM, respectively.

ARTICLE HISTORY

Received 5 June 2019
Accepted 13 July 2019

KEYWORDS


Cytotoxicity; *Melicope quercifolia*; Meliquercifolins A and B



1. Introduction

Melicope quercifolia (Rutaceae) is a small tree and found as an endemic plant in West Java, Indonesia. The leaves of *M. quercifolia* have been used to treat skin diseases (Appelhans et al. 2018). Acetophenones (Nguyen et al. 2016), alkaloids (George et al. 2017), coumarins (Xu et al. 2016), flavonoids (Saputri et al. 2018), and phenylpropa-noids (Nakashima et al. 2012) are phenolic compounds from the *Melicope* plants. Some of the phenolic compounds showed various Diels-Alder adduct and [2 + 2]

CONTACT Mulyadi Tanjung  mulyadi-t@fst.unair.ac.id

 Supplemental data for this article can be accessed at <https://doi.org/10.1080/14786419.2019.1644634>

© 2019 Informa UK Limited, trading as Taylor & Francis Group

cycloaddition (Nakashima et al. 2012). Melicodenines E (3), F (4) and I (5) formed a [2 + 2] cycloaddition type, and a Diels-Alder adduct type. In the ongoing phytochemical investigation of *M. quercifolia*, we report two new compounds, meliquercifolins A (1), and B (2) from the leaves of *M. quercifolia*. These structures of two new compounds are coumarins bearing an acetophenone moiety. Heterodimer compounds (1 and 2) are the first time found skeleton structure from *Melicope*. The cytotoxic properties against three human cancer cells (P-388, MCF-7, and HeLa) of isolated compounds will also report.

2. Result and discussion

Meliquercifolin A (1) obtained a colorless oil in which showed an optical rotation was inactive. Based on the HRESIMS measurement of 1 gave a positive ion peak $[M + H]^+$ at m/z 479.1729 correspondings for a molecular formula $C_{27}H_{27}O_8$. The IR absorptions of 1 displayed vibrations the presence of conjugated C=O (1636 cm^{-1}), aromatic (1608 and 1579 cm^{-1}) and ether (1119 cm^{-1}) groups. Two maximum absorptions at λ_{max} 280 (3.86) and 332 (3.88) nm very like with benzoyl and cinnamoyl chromophore. Four signals of a 1,2,3,4-tetrasubstituted cyclobutane ring showed at δ_{H} 3.05 (1H, dd, $J=9.8$; 5.8 Hz, H-3), δ_{H} 3.96 (1H, t, $J=2.2$ Hz, H-4), δ_{H} 4.71 (1H, t, $J=6.9$ Hz, H-6) and δ_{H} 5.42 (1H, ddd, $J=8.2$; 6.2; 2.2 Hz, H-7). The signal at δ_{H} 5.42 implied an oxymethine attached in the cyclobutane ring. Compound 1 also demonstrated the presence of two proton signals of two aromatics at δ_{H} 5.98 (1H, s, H-8), δ_{H} 6.21 (1H, s, H-9) and a pair of *cis* vinylic of chromen-2-one ring at δ_{H} 6.01 (1H, d, $J=9.8$ Hz, H-3), δ_{H} 7.82 (1H, d, $J=9.8$ Hz, H-4). Based on the ^1H NMR spectrum, compound 1 likewise showed the presence of three methyl signals at δ_{H} 2.17 (3H, s, H-10), δ_{H} 1.19 (3H, s, H-11), δ_{H} 1.61 (3H, s, H-12) as well as three methoxyl signals at δ_{H} 4.04 (3H, s, 5-OCH₃), δ_{H} 3.67 (3H, s, 5'-OCH₃), and δ_{H} 3.59 (3H, s, 7'-OCH₃). The ^{13}C NMR analysis (APT experiment), compound 1 showed 27 carbon signals that are completely separated, consisting of six methyl carbons, eight methine carbons, and 13 quaternary carbons. Three of them (δ_{C} 161.9; δ_{C} 139.5 and δ_{C} 109.7) are characteristic of C-2, C-3, and C-4 of the chromen-2-one ring (coumarin). Furthermore, two signals of the acetyl group (δ_{C} 32.3; δ_{C} 202.5) are characteristic of the acetophenone structure (Nakashima et al. 2012). The signal of a *cis* vinylic of chromen-2-one at δ_{H} 7.82 (H-4) in the HMBC spectrum exhibited correlation with two oxyaryl carbons, C-5 (δ_{C} 152.7), C-9a (δ_{C} 156.7) and a lactone carbonyl at C-2 (δ_{C} 161.9) as well as the methoxyl signal at δ_{H} 4.04 correlated to C-5 (δ_{C} 152.7) unambiguously located the methoxyl group at C-5. Another proton signal of a *cis* vinylic at δ_{H} 6.01 (H-3) correlated to C-2 (δ_{C} 161.9) and a quaternary carbon at C-4a (δ_{C} 108.1). Furthermore, a signal of aromatic at δ_{H} 6.21 correlated to two oxyaryl carbons at C-9a (δ_{C} 156.7); C-8a (δ_{C} 168.5) and two quaternary carbons at C-4a (δ_{C} 108.1); C-5a (δ_{C} 104.7) showed that the signal of δ_{H} 6.21 at H-9. A signal of an oxymethine at δ_{H} 5.42 correlated to C-8a (δ_{C} 168.5), and C-4' (δ_{C} 34.7) revealed that a part of the heterodimer is bergapten (coumarin) (Nakashima et al. 2012). The HMBC spectrum, a signal of methine at δ_{H} 4.71 (H-6) correlated to C-3' (δ_{C} 45.5), C-4' (δ_{C} 34.7) and C-8a (δ_{C} 168.5) reinforced the structure of the bergapten experiencing a [2 + 2] cycloaddition reaction on furano ring (Nakashima et al. 2012). A methine signal (cyclobutane ring) at




Figure 1. Chemical structures of isolated compounds from *M. quercifolia*.

⁵²
 δ_{H} 3.96 (H-4) correlated to δ_{C} 44.1 (C-6), δ_{C} 84.9 (C-7), δ_{C} 156.4 (C-8a) and δ_{C} 106.6 (C-4a). Two methyl signals of pyrano ring at δ_{H} 1.19 (H-11) and δ_{H} 1.61 (H-12) correlated to a methine carbon at C-3' (δ_{C} 45.5) and an ether cyclic carbon at C-2' (δ_{C} 74.0). Besides, a signal of aromatic isolated at δ_{H} 5.98 (H-8) correlated to an oxyaryl carbon at δ_{C} 156.8 (C-7), and two quaternary carbons at δ_{C} 106.6 (C-4a); 117.8 (C-6). A methoxyl signal at δ_{H} 3.67 (7-OCH₃) correlated to δ_{C} 156.8 (C-7). Another a methoxyl signal at δ_{H} 3.59 (5'-OCH₃) correlated to δ_{C} 155.7 (C-5). A methyl signal of acetyl at δ_{H} 2.17 (H-10) correlated to δ_{C} 202.5 (C-9). These correlations showed that a part of evodionol (Nakashima et al. 2012). This result implied that compound **1** is a reaction of [2 + 2] cycloaddition between bergapten-evodionol. Therefore, the structure of meliquercifolin A established as **1**.

Meliquercifolin B (**2**) also acquired as a colorless oil. The UV absorption bands (λ_{max} 269, 335), and IR (1620, 1601, 1571 and 1110 cm⁻¹) absorptions very semblable with **1**. The molecular formula of compound **2** was assigned to be C₂₈H₂₉O₉ exhibited positive ion peak [M + H]⁺ at *m/z* 509.1802 (calcd for 509.1812) by HRESIMS measurement. The ¹H and ¹³C NMR spectrum of **2** had very semblable with **1**. The main difference, the ¹H and ¹³C NMR of **2** showed a methoxyl group at C-9 (δ_{H} 3.98; δ_{C} 58.6). The HMBC and HMQC experiments confirmed the location of a methoxyl group at C-9. With the addition, a methoxyl group attached in the aromatic ring of coumarin skeleton indicated is [2 + 2] cycloaddition between isopimpinellin-evodionol. Based on UV, IR, HRESIMS, 1D, and 2D NMR measurements, the structure of meliquercifolin B assigned as **2**. The chemical structures of isolated compounds shown in (Figure 1).

Three known compounds, melicodenine E (**3**), melicodenine F (**4**) and melicodenine I (**5**) from HRESIMS, 1D and 2D NMR data showed chemical formula, and the chemical

Table 1. Cytotoxic activities of isolated compounds (1–5).

| Compound | IC ₅₀ (μM) | | |
|-------------|-----------------------|------------|-------------|
| | HeLa | MCF-7 | P-388 |
| 1 | 2.6 ± 0.10 | >100 | 78.9 ± 1.20 |
| 2 | >100 | >100 | >100 |
| 3 | >100 | >100 | 11.9 ± 0.42 |
| 4 | 0.8 ± 0.01 | >100 | 38.3 ± 0.87 |
| 5 | 1.1 ± 0.05 | >100 | >100 |
| Artonin E | – | – | 1.33 ± 0.07 |
| Doxorubicin | 0.9 ± 0.04 | 0.8 ± 0.02 | – |

shift very resemblant with published previously (Nakashima et al. 2012; George et al. 2017). Compound **3** is a form of a [2 + 2] cycloaddition reaction of *N*-methylflindersin-melicodin A, and compound **4** is *N*-methylflindersin-bergapten. Compound **5** is a [4 + 2] cycloaddition (a Diels-Alder adduct type) two *N*-methylflindersins (George et al. 2017; Nakashima et al. 2012).

Those cytotoxic activities of isolated compounds, compounds **1**, **4** and **5** showed high activity against HeLa cells while compounds **2** and **3** were inactive. However, all of the isolated compounds were inactive against MCF-7 and P-388 cells (Table 1). For coumarin bearing an acetophenone derivative, compound **1** more active than compound **2** against HeLa cells. The influence of a methoxyl group at C-9 suggested as a critical element to decrease the cytotoxic effect towards HeLa cells. The melicodenine E (**4**) more active than meliquercifolins A (**1**). However, compound **4** slightly active than compound **5**. The *N*-methylflindersin factor is a crucial element to enhance the cytotoxic effect of HeLa cells. In addition to the *N*-methylflindersin factor tend to increase cytotoxic activity against human colon cancer cells (DLD-1) (Nakashima et al. 2012).

38 3. Experimental

3.1. Plant material

The fresh leaves of *M. quercifolia* obtained from Cianten Farm, Cigudek District, Bogor, West Java, Indonesia on Nov. 2017. The plant identified by Mr. Ismail Rachman, a botanist senior from the Herbarium Bogoriense, Bogor, Indonesia. A specimen (MQ 20171104) deposited as a reference.

16 3.2. Extraction and isolation

The air-dried leaves of *M. quercifolia* (2.0 kg) were extracted with MeOH three times (6 L, each for two days) at room temperature. Evaporation of the solvent with evaporator gave a MeOH extract (450 g) and then partitioned with *n*-hexane three times. Furthermore, the MeOH extract was added with H₂O (4:1 v/v) and partitioned with EtOAc three times gave the crude extract (5 g). The EtOAc extract (4.8 g), fractionated by CC on silica gel, eluted with *n*-hexane-EtOAc (from 9:1 to 3:7) providing two fractions, A and B. Fraction A (1.3 g) was fractionated by CC chromatography on polyamide, eluted with *n*-hexane-CHCl₃ (from 9:1 to 1:4) gave two subfractions, A₁ and A₂. Compounds **3** (13 mg) and **4** (15 mg) were isolated from subfraction A₂ (655 mg) using radial planar chromatography with the same eluent. Fraction B (800 g) was

separated by CC chromatography on polyamide, and eluted *n*-hexane-EtOAc (from 4:1 to 1:1) gave subfractions B₁-B₂. Compounds **1** (5 mg) and **2** (7 mg) was obtained from subfraction B₁ (265 mg) by radial planar chromatography, eluted with *n*-hexane-diisopropylether (from 3:9 to 7:3), and diisopropylether as a solvent system. By the same methods, compound **5** (16 mg) isolated from subfraction B₂ (148 mg).

3.3. Spectral data

Meliquercifolin A (**1**): yellow solid, UV/Vis (MeOH) λ_{\max} (nm) (log ϵ): 217 (3.03), 269 (3.08), 289 (3.20) and 325 (2.81) nm. IR (KBr) ν_{\max} (cm⁻¹): 3411, 2972, 2925, 2852, 1649, 1604, 1579 and 1159. ¹H-NMR (400 MHz, CDCl₃) δ_{H} ppm: 6.01 (1H, *d*, *J* = 9.8 Hz, H-3), 7.81 (1H, *d*, *J* = 9.8 Hz, H-4), 4.04 (3H, *s*, 5-OCH₃), 4.71 (1H, *t*, *J* = 6.9 Hz, H-6), 5.42 (1H, *ddd*, *J* = 8.2; 6.2; 2.2 Hz, H-7), 6.21 (1H, *s*, H-9), 3.05 (1H, *dd*, *J* = 9.8; 5.8 Hz, H-3), 3.96 (1H, *t*, *J* = 2.2 Hz, H-4), 3.67 (3H, *s*, 5'-OCH₃), 3.59 (3H, *s*, 7'-OCH₃), 5.98 (1H, *s*, H-8), 2.17 (3H, *s*, H-10), 1.19 (3H, *s*, H-11) and 1.61 (3H, *s*, H-12). ¹³C-NMR (100 MHz, CDCl₃) δ_{C} ppm: 161.1 (C-2), 109.7 (C-3), 139.5 (C-4), 108.1 (C-4a), 152.7 (C-5), 104.7 (C-5a), 44.1 (C-6), 84.9 (C-7), 168.5 (C-8a), 91.7 (C-9), 156.7 (C-9a), 74.0 (C-2), 45.5 (C-3), 34.7 (C-4), 106.6 (C-4a), 155.7 (C-5), 62.6 (5'-OCH₃), 117.8 (C-6), 156.8 (C-7), 55.8 (7'-OCH₃), 96.8 (C-8), 156.4 (C-8a), 202.5 (C-9), 32.3 (C-10), 25.3 (C-11) and 25.7 (C-12). HRESIMS: *m/z* [M + H]⁺ calcd. for C₂₇H₂₇O₈ 479.1706, found 479.1729.

Meliquercifolin B (**2**): yellow solid, UV/Vis (MeOH) λ_{\max} (nm) (log ϵ): 217 (3.33), 234 (3.12), 255 (3.02), 268 (2.98) and 299 (2.69) nm. IR (KBr) ν (cm⁻¹): 3448, 2966, 2923, 2854, 1651, 1627, 1460 and 1186. ¹H-NMR (400 MHz, CDCl₃) δ_{H} ppm: 6.02 (1H, *d*, *J* = 9.6 Hz, H-3), 7.79 (1H, *d*, *J* = 9.6 Hz, H-4), 3.68 (3H, *s*, 5-OCH₃), 4.75 (1H, *t*, *J* = 6.8 Hz, H-6), 5.50 (1H, *dd*, *J* = 6.8; 3.9 Hz, H-7), 3.98 (3H, *s*, 9-OCH₃), 3.09 (1H, *dd*, *J* = 9.9; 5.8 Hz, H-3), 3.99 (1H, *dd*, *J* = 5.8; 1.8 Hz, H-4), 3.68 (3H, *s*, 5'-OCH₃), 3.57 (3H, *s*, 7'-OCH₃), 5.92 (1H, *s*, H-8), 2.21 (3H, *s*, H-10), 1.12 (3H, *s*, H-11), and 1.66 (3H, *s*, H-12). ¹³C-NMR (100 MHz, CDCl₃) δ_{C} ppm: 160.9 (C-2), 109.9 (C-3), 139.6 (C-4), 104.4 (C-4a), 126.0 (C-5), 106.9 (C-5a), 44.4 (C-6), 85.3 (C-7), 159.8 (C-8a), 148.1 (C-9), 149.1 (C-9a), 74.1 (C-2), 45.7 (C-3), 34.5 (C-4), 106.2 (C-4a), 156.9 (C-5), 62.7 (5'-OCH₃), 117.3 (C-6), 156.8 (C-7), 55.8 (7'-OCH₃), 96.8 (C-8), 163.8 (C-8a), 209.2 (C-9), 32.3 (C-10), 25.4 (C-11), and 25.7 (C-12). HRESIMS: *m/z* [M + H]⁺ calcd. for C₂₈H₂₉O₉ 509.1812, found 509.1802.

3.4. Cytotoxic assay

Effect cytotoxic of compounds (**1-5**) against HeLa (human cervical cancer cells), MCF-7 (human breast cancer cells), P-388 (human murine leukemia cells) were evaluated using the MTT colorimetric methods (Mah et al. 2015; Tanjung et al. 2017, 2018; Segun et al. 2019). Doxorubicin used as a control positive for HeLa, and MCF-7 as well as artonin for P-388 cells.

4. Conclusions

Meliquercifolins A (**1**) and B (**2**), two new coumarins bearing an acetophenone derivative were isolated from the leaves of *M. quercifolia* together with three known

compounds, melicodenines E (3), F (4) and I (5). Compounds (1 and 2) are the first time coumarin bearing an acetophenone derivative found on *Melicope*.

Acknowledgments

Universitas Airlangga, Surabaya, Indonesia supported this work, through Penelitian Disertasi Doktor grant 2019.

21

Disclosure statement

No potential conflict of interest reported by the authors.

References

- Appelhans MS, Wen J, Duretto M, Crayn D, Wagner WL. 2018. Historical biogeography of *Melicope* and its close relatives with a special emphasis on Pacific dispersals. *J Syst Ecol.* 9999: 1–24. [CrossRef]
- George S, Venkataraman R, Baby S. 2017. Melicodenine I, a new quinolinone alkaloid from *Melicope denhamii* leaves. *Nat Prod Res.* 31(8):890–895.
- Mah SH, Ee GCL, Teh SS, Sukari MA. 2015. *Calophyllum inophyllum* and *Calophyllum soulattri* source of anti-proliferative xanthenes and their structure–activity relationships. *Nat Prod Res.* 29(1):98–101.
- Nakashima K, Oyama M, Ito T, Akao Y, Witono JR, Darnaedi D, Tanaka T, Murata J, Iinuma M. 2012. Novel quinolinone alkaloids bearing a lignoid moiety and related constituents in the leaves of *Melicope denhamii*. *Tetrahedron.* 68(10):2421–2428.
- Nguyen NH, Ha TKQ, Choi S, Eum S, Lee CH, Bach TT, Chinh VT, Oh WK. 2016. Chemical constituents from *Melicope pteleifolia* leaves. *Phytochem.* 130:291–300.
- Saputri RD, Tjahjandarie TS, Tanjung M. 2018. Meliglabin, a new flavonol derivative from the leaves of *Melicope glabra* (Blume) T.G. Hartley. *Nat Prod Sci.* 24(3):155–158.
- Segun PA, Ogbole OO, Ismail FMD, Nahar L, Evans AR, Ajaiyeoba EO, Sarker SD. 2019. Bioassay-guided isolation and structure elucidation of cytotoxic stilbenes and flavonols from the leaves of *Macaranga barteri*. *Fitoterapia.* 134:151–157.
- Tanjung M, Hakim EH, Syah YM. 2017. Prenylated dihydrostilbenes from *Macaranga rubiginosa*. *Chem Nat Compd.* 53(2):215–218.
- Tanjung M, Rachmadiarti F, Saputri RD, Tjahjandarie TS. 2018. Mesucalophylloidin, a new isoprenylated 4-phenylcoumarin from *Mesua calophylloides* (Ridl.) Kosterm. *Nat Prod Res.* 32(9): 1062–1067.
- Xu J, Sun X, Liu X, Peng M, Li S, Jin D-Q, Lee D, Bartlam B, Guo Y. 2016. Phytochemical constituents from *Melicope pteleifolia* that promote neurite outgrowth in PC12 cells. *J. Funct Foods.* 23:565–572.

Two novel coumarins bearing an acetophenone derivative from the leaves of Melicope Quercifolia

ORIGINALITY REPORT

25%

SIMILARITY INDEX

13%

INTERNET SOURCES

24%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1

Carina Weber, Till Opatz.
"Bisbenzylisoquinoline Alkaloids", Elsevier BV,
2019

Publication

2%

2

Mulyadi Tanjung, Intan Nurmalasari, Aisyah
Kanti Wilujeng, Ratih Dewi Saputri, Fida
Rachmadiarti, Tjitjik Srie Tjahjandarie. "
Acronyculatin P, A New Isoprenylated
Acetophenone from the Stem Bark of ",
Natural Product Sciences, 2018

Publication

1%

3

Ratih Dewi Saputri, Mulyadi Tanjung, Tjitjik
Srie Tjahjandarie. " Cytotoxic activity of
quinolinone Alkaloids and acylphloroglucinol
from the leaves of ", Journal of Physics:
Conference Series, 2018

Publication

1%

4

Tjitjik Srie Tjahjandarie, Ratih Dewi Saputri,
Ryan Ayub Wahjoedi, Mulyadi Tanjung. "
Melimoluccanin, A new isoprenylated
quinolone alkaloid from the leaves of T.G.
Hartley ", Journal of Physics: Conference

1%

Series, 2018

Publication

-
- | | | |
|----|---|----|
| 5 | Mulyadi Tanjung, Tjitjik Srie Tjahjandarie, Ratih Dewi Saputri, Muhammad Fajar Aldin, Hery Purnobasuki. " Two new pyranoxanthenes from the stem bark of P.F. Stevens ", Natural Product Research, 2020 Publication | 1% |
| 6 | Tjitjik Srie Tjahjandarie, Mulyadi Tanjung, Ratih Dewi Saputri, Devina Oktari Rahayu et al. " Two new 2-arylbenzofurans from L. and their cytotoxicity towards cancer cell ", Natural Product Research, 2020 Publication | 1% |
| 7 | qmro.qmul.ac.uk Internet Source | 1% |
| 8 | www.mdpi.com Internet Source | 1% |
| 9 | Clynton W. Halstead, Paul I. Forster, Peter G. Waterman. " Alkaloids from the stem bark of an Australian population of ", Natural Product Research, 2006 Publication | 1% |
| 10 | Harkat, H.. "An unusual lignan sulfate and aromatic compounds from Frankenia thymifolia Desf.", Biochemical Systematics and Ecology, 200703 Publication | 1% |
-

| | | |
|----|--|----|
| 11 | www.wfsbp.org Internet Source | 1% |
| 12 | Marina Dellagreca. "A new xyloside from <i>Chenopodium album</i> ", <i>Natural Product Research</i> , 1/1/2005 Publication | 1% |
| 13 | Sony George, Ramaswamy Venkataraman, Sabulal Baby. " Melicodenine I, a new quinolinone alkaloid from leaves ", <i>Natural Product Research</i> , 2016 Publication | 1% |
| 14 | repository.hkbu.edu.hk Internet Source | 1% |
| 15 | Park, J.W.. "Unidirectional @ α -cyclodextrin-based [2]rotaxanes bearing viologen unit on axle", <i>Tetrahedron Letters</i> , 20060605 Publication | 1% |
| 16 | Hui-Min Wang, Wen-Li Lo, Lee-Yu Huang, Yau-Der Wang, Chung-Yi Chen. " Chemical constituents from the leaves of ", <i>Natural Product Research</i> , 2010 Publication | 1% |
| 17 | Barry V L Potter, Marco Mottinelli, Masa Sinreh, Tea Lanisnik-Rizner, Mathew Leese. "N-Phenyl-1,2,3,4-tetrahydroisoquinoline:an alternative scaffold for design of 17 β -hydroxysteroid dehydrogenase 1 inhibitors", <i>ChemMedChem</i> , 2020 | 1% |

-
- 18 www.freepatentsonline.com <1 %
Internet Source
-
- 19 Mulyadi Tanjung, Ratih Saputri, Ryan Wahjoedi, Tjitjik Tjahjandarie. "4-Methoxy-3-(3-methylbut-2-en-1-yl)-7-[(3-methylbut-2-en-1-yl)oxy]quinolin-2(1H)-one from Melicope Moluccana T.G. Hartley", Molbank, 2017 <1 %
Publication
-
- 20 Stevenson, P.C.. "Insect antifeedant furanocoumarins from *Tetradium daniellii*", *Phytochemistry*, 200305 <1 %
Publication
-
- 21 www.pse-journal.hr <1 %
Internet Source
-
- 22 Hitotsuyanagi, Y.. "A structure-activity relationship study of brusatol, an antitumor quassinoid", *Tetrahedron*, 20060424 <1 %
Publication
-
- 23 patents.justia.com <1 %
Internet Source
-
- 24 Nisha Singh, Nawed Anjum, Ramesh Chandra. "Combating influenza: natural products as neuraminidase inhibitors", *Phytochemistry Reviews*, 2018 <1 %
Publication
-
- 25 Ahmed I. Khodair, Najim A. Al-Masoudi, Jean-Pierre Gesson. "A New Approach to the <1 %

Synthesis of Benzothiazole, Benzoxazole, and Pyridine Nucleosides as Potential Antitumor Agents", Nucleosides, Nucleotides and Nucleic Acids, 2003

Publication

26

Xuechen Li. "Mapping the Active Site of the Bacterial Enzyme LpxC Using Novel Carbohydrate-Based Hydroxamic Acid Inhibitors*", Journal of Carbohydrate Chemistry, 8/1/2005

Publication

<1%

27

Wendong Wang, Pornpun Rattananakin, Peter G. Goekjian. " Synthesis of -Glycoside Analogs via Thionolactones ", Journal of Carbohydrate Chemistry, 2007

Publication

<1%

28

www.fedoa.unina.it

Internet Source

<1%

29

Aulia Ilmiawati, Euis H. Hakim, Yana M. Syah. "Prenylated 9,10-dihydrophenanthrenes from Macaranga javanica", Zeitschrift für Naturforschung B, 2015

Publication

<1%

30

U Hasanah, T S Tjahjandarie, M Tanjung. " Chromanone Acid Derivatives from the Stem Bark of ", IOP Conference Series: Earth and Environmental Science, 2019

Publication

<1%

Canjun Li, Yiming Li, Hao H. Sun. " New

31 ganoderic acids, bioactive triterpenoid metabolites from the mushroom ", Natural Product Research, 2006 <1 %
Publication

32 Alka Elizabeth Varghese, Balaji Govindan, Jyothilakshmi Madhavankutty, Antony Thomas Valiyaveetil et al. " A new antifungal benzoic acid ester from ", Natural Product Research, 2017 <1 %
Publication

33 people.uea.ac.uk <1 %
Internet Source

34 Shih-Chuan Chan, Yuan-Shiun Chang, Sheng-Chu Kuo. "Neoflavonoids from *Dalbergia odorifera*", Phytochemistry, 1997 <1 %
Publication

35 Alain Guntern, Jean-Robert Ioset, Emerson F Queiroz, Chris M Foggin, Kurt Hostettmann. "Quinones from *Heliotropium ovalifolium*", Phytochemistry, 2001 <1 %
Publication

36 Yan-Hui Li, Fan Luo, Shu-Lin Peng, Jian Liang, Li-Sheng Ding. " A new dihydroisocoumarin from the rhizomes of ", Natural Product Research, 2006 <1 %
Publication

37 Ry[icaron]hi Masuda, Masaru Hojo, Tadaaki Ichi, Fujio Adachi, Kenji Yoshinaga. "SYNTHESIS AND REACTIONS OF HALO <1 %

DERIVATIVES OF 4-ISOPROPYL-2H-1,4-
THIAZIN-3-ONE", Phosphorus and Sulfur and
the Related Elements, 2006

Publication

38

www.tandfonline.com

Internet Source

<1 %

39

Jawhar Abd'elmonem Fakhfakh, Mohamed Damak. " Sesquioneolignans from the flowers of Coss. et Dur. (Asteraceae) ", Natural Product Research, 2007

Publication

<1 %

40

May Ying Shen, José Angelo Zuanazzi, Christiane Kan, Jean-Charles Quirion, Henri-Philippe Husson, I. R. C. Bick. " Alkaloids from ", Natural Product Letters, 1995

Publication

<1 %

41

A. Sajeli Begum, S. Santhosh Kumar. "Advances in the chemistry and pharmacological potential of coumarinolignans", Topics in Current Chemistry, 2018

Publication

<1 %

42

Chanmahasathien, W.. "Prenylated xanthenes with NGF-potentiating activity from *Garcinia xanthochymus*", Phytochemistry, 200311

Publication

<1 %

43

Kiyoaki Yonesu. "Ascotricins A and B, novel antagonists of sphingosine-1-phosphate receptor 1 from *Ascotricha chartarum* Berk.

<1 %

SANK 14186", The Journal of Antibiotics,
05/22/2009

Publication

44

Jiang-Miao Hu. "Two novel bibenzyls from *Dendrobium trigonopus*", *Journal of Asian Natural Products Research*, 07/2008

Publication

45

B. D'abrosca, M. Dellagreca, A. Fiorentino, A. Golino, P. Monaco, A. Zarrelli. " Isolation and characterization of new lignans from the leaves of ", *Natural Product Research*, 2007

Publication

46

Jing Xu, Xiaocong Sun, Xingyu Liu, Maoqin Peng, Shen Li, Da-Qing Jin, Dongho Lee, Mark Bartlam, Yuanqiang Guo.

"Phytochemical constituents from *Melicope pteleifolia* that promote neurite outgrowth in PC12 cells", *Journal of Functional Foods*, 2016

Publication

47

Tripetch Kanchanapoom, Ryoji Kasai, Phannipha Chumsri, Yoshikazu Hiraga, Kazuo Yamasaki. "Megastigmane and iridoid glucosides from *Clerodendrum inerme*", *Phytochemistry*, 2001

Publication

48

Ping-Jyun Sung, Wan-Ping Hu, Lee-Shing Fang, Tung-Yung Fan, Jeh-Jeng Wang. " Briarenol A, a new diterpenoid from a

<1 %

<1 %

<1 %

<1 %

<1 %

gorgonian ' sp. (Briareidae) ", Natural Product Research, 2005

Publication

49

Ding-Yong Wang, En-Gui Liu. " A new diaryheptanoid from the bark of ", Natural Product Research, 2008

Publication

<1 %

50

www.icmag.com

Internet Source

<1 %

51

Peter A. Segun, Omonike O. Ogbole, Toluwanimi E. Akinleye, Temitope O. C. Faleye, Adekunle J. Adeniji. " anti-enteroviral activity of stilbenoids isolated from the leaves of ", Natural Product Research, 2019

Publication

<1 %

52

Lei Wang. "Five new compounds from Dendrobium crystallinum", Journal of Asian Natural Products Research, 11/2009

Publication

<1 %

53

theses.gla.ac.uk

Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Two novel coumarins bearing an acetophenone derivative from the leaves of Melicope Quercifolia

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
