

# A New Cinnamyl Acid Derivative from the Roots of *Willughbeia coriacea* Wall.

*by* Mulyadi Tanjung

---

**Submission date:** 16-Dec-2020 02:24PM (UTC+0800)

**Submission ID:** 1476556711

**File name:** 4.\_Willughbeia\_coriceae.pdf (307.29K)

**Word count:** 1986

**Character count:** 9553



## A New Cinnamyl Acid Derivative from the Roots of *Willughbeia coriacea* Wall.

Mulyadi Tanjung\*, Tjitjik Sri Tjahjandarie, Ratih Dewi Saputri, Andre Harsono, and Muhammad Fajar Aldin

Natural Products Chemistry Research Group, Organic Chemistry Division, Department of Chemistry,  
Faculty of Science and Technology, Universitas Airlangga, Surabaya, Indonesia

**Abstract** – A new cinnamyl acid derivative, willughbein A (**1**) along with pinoresinol (**2**), alyterinate A (**3**), and scopoletin (**4**), were isolated from the roots of *Willughbeia coriacea* Wall. The structure of **1** has been determined based on HRESIMS, 1D, and 2D NMR spectral data. All of the isolates were evaluated for their cytotoxicity against three human cancer cells (HeLa, T47D, MCF-7, and P-388). Compound **3** showed moderate activity against P-388 cells with an IC<sub>50</sub> value of 3.04 µg/mL.

**Keywords** – Willughbein A, Cinnamyl acid, *Willughbeia coriacea*, Cytotoxic

### Introduction

*Willughbeia coriacea* Wall (Apocynaceae) is one of Kalimantan island endemic plants, and local people consume its fruit. The public knows *W. coriacea* as Dangu, and its stew used for diarrhea. Information on secondary metabolite compounds from *W. coriacea* has no scientific report.<sup>1</sup>

*W. cochinchinensis* from Vietnam is the only one reported about the content of secondary metabolites. *W. cochinchinensis* produces phenolic compounds, including cinnamic acids, lignans, coumarins, and diarylheptanoids. The phenolic compound of *W. cochinchinensis* shows activity as an inhibitor of acetylcholinesterase (AChE) and butylocholinesterase (BChE), which causes Alzheimer's disease.<sup>2</sup>

In summary, we reported the isolation of a new cinnamyl acid derivative, willughbein A (**1**), together with three known compounds from the roots of *W. coriacea*. The cytotoxic activity of compounds **1-4** against four human cancer cells (HeLa, T47D, MCF-7, and P-388) also reported.

### Experimental

**General experimental procedures** – UV spectra measured with a Shimadzu UV-1800 recording spectrophoto-

meter. IR spectra measured with a Shimadzu IR Tracer-100 spectrophotometer, respectively. 1D and 2D NMR run on a JEOL ECA 400 spectrometer in CDCl<sub>3</sub>. HRESIMS measured on a Waters LCT Premier XE ESI-TOF mass spectrometer. Gravity column chromatography and planar radial chromatography were carried out using silica gel 60, Sephadex LH-20, and silica gel 60 F<sub>254</sub>. The Spot of compounds on TLC visualized under a UV lamp and anisaldehyde reagent.

**Plant materials** – The roots of *W. coriacea* collected from Hajak Village, Muara Teweh, North Barito, Central Kalimantan, Indonesia, in September 2018. Mr. Ismail Rachman, a senior botanist from Herbarium Bogoriense, identified plant material. Voucher specimens (WC 20183) stored at the Bogoriense Herbarium, Center for Research and Development of Biology, National Science Institute, Bogor, Indonesia.

**Extraction and isolation** – The powdered and dried roots of *W. coriacea* (3.48 kg) were extracted with methanol for 24 hours at room temperature using the maceration method. The extraction of the material was carried out twice. The maceration results were filtered, and the solvent was evaporated with a rotary vacuum evaporator so that the methanol extract (400 g) was obtained. The MeOH extract redissolved in MeOH-water (9:1) and was partitioned with *n*-hexane (135 g) and EtOAc (12.4 g). The EtOAc extract (12 g) was further fractionated by gravity column chromatography on silica gel (150 g) eluted with *n*-hexane-EtOAc by increasing polarity (9:1, 4:1, and 7:3) to give three significant fractions A-C. Fraction B (2.15 g) separated by Sephadex LH-20 eluted with methanol to provide subfractions B<sub>1</sub>-

\*Author for correspondence

Mulyadi Tanjung, Natural Products Chemistry Research Group, Organic Chemistry Division, Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Surabaya, Indonesia  
Tel: +62-31-5936501; E-mail: mulyadi-t@fst.unair.ac.id




Fig. 1. The phenolic compounds 1 - 4 from the roots of *W. coriacea*.

B<sub>2</sub>. The purification of subfraction B<sub>2</sub> by planar radial chromatography using *n*-hexane-diisopropyl ether (from 4:1 to 3:7) to yield compounds 2 (14 mg), and 3 (17 mg). Fraction C (13 g) was fractionated using Sephadex LH-20 and eluted with methanol to give subfractions C<sub>1</sub>-C<sub>2</sub>. Subfraction C<sub>2</sub> was purified by planar radial chromatography using *n*-hexane-acetone (from 9:1 to 1:1) to yield compounds 1 (17 mg), and 4 (32 mg).

**Willughbein A (1)** – Colorless oil, UV (MeOH)  $\lambda_{\max}$  nm (log  $\epsilon$ ): 220 (4.14), 244 (3.56), and 268 (3.79). IR (KBr)  $\nu_{\max}$  cm<sup>-1</sup>: 3525, 1585, and 1126. <sup>1</sup>H and <sup>13</sup>C NMR see Table 1. HRESIMS:  $m/z$  [M+H]<sup>+</sup> calcd. for C<sub>12</sub>H<sub>17</sub>O<sub>4</sub> 225.1121, found 225.1127.

**Pinoresinol (2)** – Yellow solid, mp. 125 - 126 °C. The <sup>1</sup>H and <sup>13</sup>C NMR spectral data in CDCl<sub>3</sub> of 2 very similar to published data.<sup>3</sup>

**Alyterinate A (3)** – White solid, mp. 114 - 115 °C. The <sup>1</sup>H and <sup>13</sup>C NMR spectral data in CDCl<sub>3</sub> of 3 very similar to published data.<sup>4</sup>

**Scopoletin (4)** – White solid, mp. 203 - 204 °C. The <sup>1</sup>H and <sup>13</sup>C NMR spectral data in CDCl<sub>3</sub> of 4 very similar to published data.<sup>3</sup>

**Cytotoxic activity** – All of the compounds (1 - 4) were assayed for their cytotoxicity against HeLa (human cervical cells), P-388 (murine leukemia cells), MCF-7, and T47D (human breast cells) according to the MTT method. Each cell-cultured RPMI-1640 medium was supplemented by 10% fetal bovine serum at 37 °C for 48 h in a 5% CO<sub>2</sub> incubator. Briefly, before the active


compounds were added, approximately  $4 \times 10^4$  cells/well were seeded in 96-well and incubated at 37 °C for 24 h in a CO<sub>2</sub> incubator. The death cells by each the active compounds 1 - 4 were measured using a microplate reader spectrometer at  $\lambda$  540 nm. The IC<sub>50</sub> values can be calculated through extrapolation 50% percentage cells vs. various concentrations of the active compounds using regression analysis. Doxorubicin was used as the positive control for HeLa, MCF-7, T47D, and artonin E for P-388 cells.<sup>5-8</sup>

## Result and Discussion

Compound 1 or willughbein A isolated as a colorless oil, and showed a positive molecular ion peak [M+H]<sup>+</sup> at  $m/z$  225.1127 (calculated 225.1121), indicating a chemical formula of C<sub>12</sub>H<sub>16</sub>O<sub>4</sub> by HRESIMS spectra. The UV spectra showed maximum absorption at  $\lambda_{\max}$  (log  $\epsilon$ ): 220 (4.14), 244 (3.56), and 268 nm (3.79) characteristic for propenyl phenol derivative.<sup>9</sup> Compound (1) showed absorptions for hydroxyl (3456 cm<sup>-1</sup>), aromatic (1601 - 1496 cm<sup>-1</sup>), and ether (1178 cm<sup>-1</sup>) groups, respectively by IR spectra.<sup>10</sup> The <sup>1</sup>H NMR spectra (Table 1) of 1 showed the presence of a *trans* 3-methoxy-1-propenyl signal at  $\delta_{\text{H}}$  6.53 (1H, d,  $J$  = 15.9 Hz, H-3),  $\delta_{\text{H}}$  6.20 (1H, dt,  $J$  = 15.9; 6.1 Hz, H-2),  $\delta_{\text{H}}$  4.09 (2H, dd,  $J$  = 6.1; 1.4 Hz, H-1), and  $\delta_{\text{H}}$  3.39 (3H, s, 1-OCH<sub>3</sub>).<sup>11</sup> The <sup>1</sup>H NMR spectra of 1 also showed that the presence of a 1,3,4,5-tetrasubstituted benzene at  $\delta_{\text{H}}$  6.62 (2H, s, H-2'/H-6'), a hydroxyl signal at  $\delta_{\text{H}}$  5.95 (1H, s, 5'-OH), two methoxyl signals at  $\delta_{\text{H}}$  3.84

**Table 1.** NMR data (400 MHz, CDCl<sub>3</sub>) of willughbein A (**1**)

No.C	$\delta_H$ (mult, <i>J</i> in Hz)	$\delta_C$	HMBC
1	4.09 ( <i>dd</i> , 6.1; 1.4)	73.1	C-2; C-3; 1-OCH <sub>3</sub>
2	6.20 ( <i>dt</i> , 15.9; 6.1)	125.6	C-1; C-1'
3	6.53 ( <i>d</i> , 15.9)	132.4	C-1; C-2'/6'
1'	-	132.5	-
2'/6'	6.62 ( <i>s</i> )	103.5	C-1'; C-3'/5'; C-2'/6'; C-4'
3'	-	153.4	-
4'	-	149.2	-
5'	-	137.9	-
5'-OH	5.95 ( <i>s</i> )	-	C-5'
1-OCH <sub>3</sub>	3.39 ( <i>s</i> )	58.1	C-1
3'-OCH <sub>3</sub>	3.87 ( <i>s</i> )	56.1	C-3'
4'-OCH <sub>3</sub>	3.84 ( <i>s</i> )	61.0	C-4'

**Fig. 2.** Selected HMBC correlations of willughbein A (**1**).

(3H, *s*, 4'-OCH<sub>3</sub>), and  $\delta_H$  3.87 (3H, *s*, 3'-OCH<sub>3</sub>). The <sup>13</sup>C NMR spectra of compound **1** exhibited 12 carbon signals, consisting of three methoxyl carbons, one methylene carbon, four methine carbons, and four quaternary carbons. The 2D NMR (HMQC and HMBC) explained the position of hydroxyl and methoxyl groups (Fig. 2). The *trans* ethene signal at  $\delta_H$  6.20 (H-2) correlated with a quaternary carbon at  $\delta_C$  132.5 (C-1'), and a methylene carbon at  $\delta_C$  73.1 (C-1). The *trans* ethene signal at  $\delta_H$  6.53 (H-3) correlated with a methine carbons of aromatic at  $\delta_C$  103.5 (C-2'/C-6') and a methylene carbon at  $\delta_C$  73.1 (C-1). The methylene signal at  $\delta_H$  4.09 (H-1) correlated to  $\delta_C$  125.6 (C-2),  $\delta_C$  132.4 (C-3), and  $\delta_C$  58.1 (1-OCH<sub>3</sub>). The

methoxyl group at  $\delta_H$  3.39 (1-OCH<sub>3</sub>) showed correlations with a methylene carbon at  $\delta_C$  73.1. The signal of aromatic at  $\delta_H$  6.62 (H-2'/H-6') showed correlations with three oxyaryl carbons at  $\delta_C$  153.4 (C-3'),  $\delta_C$  149.2 (C-4'),  $\delta_C$  137.9 (C-5'), a methine carbon at 103.5 (C-2'/C-6'), and a quaternary carbon at  $\delta_C$  132.5 (C-1'). Furthermore, the signal of the hydroxyl group at  $\delta_H$  5.95 (5'-OH) correlated to  $\delta_C$  137.9 (C-5'). The signal of the methoxyl at  $\delta_H$  3.87 (3'-OCH<sub>3</sub>) correlated to  $\delta_C$  153.4 (C-3'), and  $\delta_H$  3.84 (4'-OCH<sub>3</sub>) showed correlations with an oxyaryl carbon at  $\delta_C$  149.2 (C-4').

From the HRESIMS, and NMR spectrum, the structure of **1** was assigned as willughbein A. Other HMBC correlations were supporting the structure of **1**, shown in Table 1 and Fig. 2. All of the compounds (**1** - **4**) were analyzed for their cytotoxicity against HeLa (human cervical cells), P-388 (murine leukemia cells), MCF-7, and T47D (human breast cells) by MTT method. Artonin E and doxorubicin were used as a positive control.<sup>12</sup> All of the compounds were inactive against three human cells, except compound **3** showed moderate activity against P-388 cells.

**Table 2.** Cytotoxicity data of compounds **1** - **4**

Compounds	IC <sub>50</sub> (μg/mL)			
	HeLa	T47D	MCF-7	P-388
Willughbein A ( <b>1</b> )	> 100	13.33 ± 1.42	> 100	11.95 ± 0.89
Pinoresinol ( <b>2</b> )	90.21 ± 1.35	5.09 ± 0.37	> 100	17.38 ± 0.95
Alyterinate A ( <b>3</b> )	83.04 ± 1.82	34.41 ± 1.42	> 100	3.04 ± 0.27
Scopoletin ( <b>4</b> )	> 100	> 100	> 100	> 100
Doxorubicin	46.11 ± 0.45	23.18 ± 0.45	57.70 ± 0.51	-
Artonin E	-	-	-	1.33 ± 0.02

### Acknowledgments

The research funded by the Ministry of Research, Technology, and Higher Education, the Republic of Indonesia through Penelitian Dasar, Universitas Airlangga, 2019.

### References

- (1) Middleton, D. J. *Tree Flora of Sabah and Sarawak*. **2004**, *5*, 1-61.
- (2) Can, M. V.; Tran, A. H.; Pham, D. M.; Dinh, B. Q.; Le, Q. V.; Nguyen, B. V.; Nguyen, M. T. T.; Nguyen, H. X.; Nguyen, N. T.; Nishijo, H. *J. Ethnopharmacol.* **2018**, *214*, 99-105.
- (3) Sribuham, T.; Sriphana, U.; Thongsri, Y.; Yenjai, C. *Phytochem. Lett.* **2015**, *11*, 80-84.
- (4) Sriphana, U.; Thongsri, Y.; Ardwichai, P.; Poopasit, K.; Prariyachatigul, C.; Simasathiansophon, S.; Yenjai, C. *Fitoterapia*. **2013**, *91*, 39-43.
- (5) Saputri, R. D.; Tjahjandarie, T. S.; Tanjung, M. *Nat. Prod. Sci.* **2018**, *24*, 155-158.
- (6) Tjahjandarie, T. S.; Tanjung, M.; Saputri, R. D.; Nadar, P. B.; Aldin, M. F.; Marlina, E.; Permadi, A.; *Nat. Prod. Sci.* **2019**, *25*, 244-247.
- (7) Tanjung, M.; Rachmadiarti, F.; Saputri, R. D.; Tjahjandarie, T. S. *Nat. Prod. Res.* **2018**, *32*, 1062-1067.
- (8) Tanjung, M.; Rachmadiarti, F.; Prameswari, A.; Agyani, V. U. W.; Saputri, R. D.; Tjahjandarie, T. S.; Syah, Y. M. *Nat. Prod. Res.* **2018**, *32*, 1493-1498.
- (9) Zhao, L.; Chen, L-Y.; Liang, J. Y. *Chin. J. Nat. Med.* **2012**, *10*, 370-373.
- (10) Tanjung, M.; Mujahidin, D.; Hakim, E. H.; Damawan, A.; Syah, Y. M. *Nat. Prod. Commun.* **2010**, *5*, 1209-1211.
- (11) Huang, H. Y.; Ishikawa, T.; Peng, C. F.; Tsai, I. L.; Chen, I. S. *J. Nat. Prod.* **2008**, *71*, 1146-1151.
- (12) Tanjung, M.; Numalasari, I.; Wilujeng, A. K.; Saputri, R. D.; Rachmadiarti, F.; Tjahjandarie, T. S. *Nat. Prod. Sci.* **2018**, *24*, 284-287.

Received November 25, 2019

Revised February 15, 2020

Accepted February 23, 2020

# A New Cinnamyl Acid Derivative from the Roots of *Willughbeia coriacea* Wall.

## ORIGINALITY REPORT

23%

SIMILARITY INDEX

13%

INTERNET SOURCES

21%

PUBLICATIONS

4%

STUDENT PAPERS

## PRIMARY SOURCES

- 1** [doczz.net](http://doczz.net) 4%  
Internet Source
- 2** Ya-Ching Shen, Kuang-Liang Lo, Yao-Haur Kuo, Yuh-Chi Kuo, Chung-Hsiung Chen, Ashraf T. Khalil. " Cespiphytins Q–V, Verticillene Diterpenoids from ", *Journal of Natural Products*, 2008 2%  
Publication
- 3** E Marliana, R Ruga, R Hairani, T S Tjahjandarie, M Tanjung. " Antioxidant activity of flavonoid constituents from the leaves of ", *Journal of Physics: Conference Series*, 2019 2%  
Publication
- 4** [www.tandfonline.com](http://www.tandfonline.com) 2%  
Internet Source
- 5** Mulyadi Tanjung, Euis H. Hakim, Yana M. Syah. "Prenylated Dihydrostilbenes from *Macaranga rubiginosa*", *Chemistry of Natural Compounds*, 2017 1%

6

[repositorio.ufsc.br](https://repositorio.ufsc.br)

Internet Source

1%

---

7

Koh Hashida, Seiji Ohara, Rei Makino. "Base-Catalyzed Reactions of (-)-Epicatechin: Formation of Enantiomers of Base-Catalyzed Reaction Products from (+)-Catechin", *Journal of Wood Chemistry and Technology*, 2003

Publication

1%

---

8

N. S. Aminah, A. Yulvia, M. Tanjung. "Methyl-3,4-dihydroxybenzoate and 9-10-dihydrophenanthrene-2,4,7-triol two phenolic compounds from *Dioscorea alata* L. and their antioxidant activity", AIP Publishing, 2017

Publication

1%

---

9

Mulyadi Tanjung, Fida Rachmadiarti, Anindita Prameswari, Villa Ultha Wustha Agyani et al. "Airlanggins A-B, two new isoprenylated benzofuran-3-ones from the stem bark of ", *Natural Product Research*, 2017

Publication

1%

---

10

[www.beilstein-journals.org](http://www.beilstein-journals.org)

Internet Source

1%

---

11

Le Thi Hong Nhung, Pham Thi Mai Huong, Nguyen Tuan Anh, Bui Huu Tai et al. " Two new norlignans from the aerial parts of (*Blume*) Merr

1%



12

Mulyadi Tanjung, Tjitjik Srie Tjahjandarie, Mulya Hadi Sentosa. "Antioxidant and cytotoxic agent from the rhizomes of *Kaempferia pandurata*", *Asian Pacific Journal of Tropical Disease*, 2013

Publication

---

1%

13

Desi Harneti, Asep Supriadin, Mariam Ulfah, Agus Safari, Unang Supratman, Khalijah Awang, Hideo Hayashi. "Cytotoxic constituents from the bark of *Aglaia eximia* (Meliaceae)", *Phytochemistry Letters*, 2014

Publication

---

1%

14

Laxinamujila Bai, Hai Ying Bao, Tolgor Bau. " Isolation and identification of a new benzofuranone derivative from ", *Natural Product Research*, 2014

Publication

---

1%

15

Fang-Pin Chang, Shyh-Shyun Huang, Tzong-Huei Lee, Chi-I Chang, Tzong-Fu Kuo, Guan-Jhong Huang, Yueh-Hsiung Kuo. "Four New Iridoid Metabolites Have Been Isolated from the Stems of *Neonauclea reticulata* (Havil.) Merr. with Anti-Inflammatory Activities on LPS-Induced RAW264.7 Cells", *Molecules*, 2019

Publication

---

1%

16

[mro.massey.ac.nz](http://mro.massey.ac.nz)



17

[www.mdpi.com](http://www.mdpi.com)

Internet Source

1%

18

[docplayer.net](http://docplayer.net)

Internet Source

<1%

19

Desi Harneti, Roekmiati Tjokronegoro, Agus Safari, Unang Supratman et al. "Cytotoxic triterpenoids from the bark of *Aglaia smithii* (Meliaceae)", *Phytochemistry Letters*, 2012

Publication

<1%

20

Yi Liu, Jun Zhang, Ran Wen, Guang-Zhong Tu, Hu-Biao Chen, Hong Liang, Yu-Ying Zhao. "Anti-inflammatory and antiproliferative prenylated chalcones from ", *Journal of Asian Natural Products Research*, 2018

Publication

<1%

21

[osdir.com](http://osdir.com)

Internet Source

<1%

22

Feng-Xian Yang, Jian-Ping Huang, Zhixiang Liu, Zhiyan Wang et al. " Benwamycins A–G, Trialkyl-Substituted Benzene Derivatives from a Soil-Derived ", *Journal of Natural Products*, 2020

Publication

<1%

Pan Deng, Tiangeng You, Xiaoyan Chen, Tao

23 Yuan, Haihua Huang, Dafang Zhong. "Identification of Amiodarone Metabolites in Human Bile by Ultraperformance Liquid Chromatography/Quadrupole Time-of-Flight Mass Spectrometry", Drug Metabolism and Disposition, 2011  
Publication <1%

---

24 repository.unair.ac.id  
Internet Source <1%

---

25 Jiayuan Li, Ning Li, Xuezheng Li, Gang Chen, Cungang Wang, Bin Lin, Yue Hou. "Characteristic  $\alpha$ -Acid Derivatives from with Antineuroinflammatory Activities ", Journal of Natural Products, 2017  
Publication <1%

---

26 Vitaly A. Osyanin, Dmitry V. Osipov, Yuri N. Klimochkin. " Reactions of -Quinone Methides with Pyridinium Methylides: A Diastereoselective Synthesis of 1,2-Dihydronaphtho[2,1- ]furans and 2,3-Dihydrobenzofurans ", The Journal of Organic Chemistry, 2013  
Publication <1%

---

Exclude quotes Off  
Exclude bibliography On

Exclude matches Off

# A New Cinnamyl Acid Derivative from the Roots of *Willughbeia coriacea* Wall.

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---