



MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>

Your final paper in AOB

6 pesan

Agoes Soegianto <agoes_soegianto@fst.unair.ac.id>
Kepada: MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>

23 Mei 2020 12.29

Your final paper in AOB

Prof. Agoes Soegianto

Dept. Biology, Airlangga University

Kampus C, Jl. Mulyorejo, Surabaya 60115

INDONESIA

2 lampiran

36(2) Contents.pdf
109K **220-226.pdf**
201K

MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>
Kepada: priyawan rachmadi <priyawan-r@fkg.unair.ac.id>

26 Mei 2020 17.34

----- Forwarded message -----

Dari: **Agoes Soegianto** <agoes_soegianto@fst.unair.ac.id>

Date: Sab, 23 Mei 2020 12:29

Subject: Your final paper in AOB

To: MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>

Your final paper in AOB

Prof. Agoes Soegianto

Dept. Biology, Airlangga University

Kampus C, Jl. Mulyorejo, Surabaya 60115

INDONESIA

2 lampiran

36(2) Contents.pdf
109K **220-226.pdf**
201K

MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>
Kepada: priyawan rachmadi <priyawan-r@fkg.unair.ac.id>

26 Mei 2020 17.37

Matur nuwon Prof. Agoes 🙏

Pada tanggal Sel, 26 Mei 2020 17:34, MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id> menulis:

----- Forwarded message -----

Dari: **Agoes Soegianto** <agoes_soegianto@fst.unair.ac.id>
Date: Sab, 23 Mei 2020 12:29
Subject: Your final paper in AOB
To: MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>

Your final paper in AOB

Prof. Agoes Soegianto
Dept. Biology, Airlangga University
Kampus C, Jl. Mulyorejo, Surabaya 60115
INDONESIA

MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>
Kepada: Aqsa Sjuhada <aqsa@fkg.unair.ac.id>

26 Mei 2020 18.20

----- Forwarded message -----


Dari: **Agoes Soegianto** <agoes_soegianto@fst.unair.ac.id>
Date: Sab, 23 Mei 2020 12:29
Subject: Your final paper in AOB
To: MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>

Your final paper in AOB

Prof. Agoes Soegianto
Dept. Biology, Airlangga University
Kampus C, Jl. Mulyorejo, Surabaya 60115
INDONESIA

2 lampiran

 **36(2) Contents.pdf**
109K

 **220-226.pdf**
201K

priyawan rachmadi <priyawan-r@fkg.unair.ac.id>
Kepada: MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>

27 Mei 2020 10.29

Assalamualaikum pa Lutfi itu jurnal yg baru lagi ?

On Tue, May 26, 2020 at 5:34 PM MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id> wrote:

Matur nuwon Prof. Agoes 🙏

Pada tanggal Sel, 26 Mei 2020 17:34, MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id> menulis:

----- Forwarded message -----

Dari: **Agoes Soegianto** <agoes_soegianto@fst.unair.ac.id>
Date: Sab, 23 Mei 2020 12:29
Subject: Your final paper in AOB
To: MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>

Your final paper in AOB

Prof. Agoes Soegianto
Dept. Biology, Airlangga University

**Kampus C, Jl. Mulyorejo, Surabaya 60115
INDONESIA**

MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>
Kepada: priyawan rachmadi <priyawan-r@fkg.unair.ac.id>

27 Mei 2020 14.38

Yang dulu, legalisasinya

Pada tanggal Rab, 27 Mei 2020 10:29, priyawan rachmadi <priyawan-r@fkg.unair.ac.id> menulis:
Assalamualaikum pa Lutfi itu jurnal yg baru lagi ?

On Tue, May 26, 2020 at 5:34 PM MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id> wrote:

Matur nuwon Prof. Agoes 🙏

Pada tanggal Sel, 26 Mei 2020 17:34, MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id> menulis:

----- Forwarded message -----

Dari: **Agoes Soegianto** <agoes_soegianto@fst.unair.ac.id>

Date: Sab, 23 Mei 2020 12:29

Subject: Your final paper in AOB

To: MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>

Your final paper in AOB

Prof. Agoes Soegianto

Dept. Biology, Airlangga University

Kampus C, Jl. Mulyorejo, Surabaya 60115

INDONESIA



MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>

Your final paper in AOB

6 pesan


Agoes Soegianto <agoes_soegianto@fst.unair.ac.id>
Kepada: MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>

23 Mei 2020 12.29

Your final paper in AOB

Prof. Agoes Soegianto**Dept. Biology, Airlangga University
Kampus C, Jl. Mulyorejo, Surabaya 60115
INDONESIA**

2 lampiran


 **36(2) Contents.pdf**
109K **220-226.pdf**
201K

MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>
Kepada: priyawan rachmadi <priyawan-r@fkg.unair.ac.id>

26 Mei 2020 17.34

[Kutipan teks disembunyikan]

2 lampiran

 **36(2) Contents.pdf**
109K **220-226.pdf**
201K

MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>
Kepada: priyawan rachmadi <priyawan-r@fkg.unair.ac.id>

26 Mei 2020 17.37

Matur nuwon Prof. Agoes 🙏

[Kutipan teks disembunyikan]

MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>
Kepada: Aqsa Sjuhada <aqsa@fkg.unair.ac.id>

26 Mei 2020 18.20

----- Forwarded message -----

Dari: **Agoes Soegianto** <agoes_soegianto@fst.unair.ac.id>


Date: Sab, 23 Mei 2020 12:29

[Kutipan teks disembunyikan]

[Kutipan teks disembunyikan]

2 lampiran

 **36(2) Contents.pdf**
109K

 **220-226.pdf**
201K

priyawan rachmadi <priyawan-r@fkg.unair.ac.id>
Kepada: MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>

27 Mei 2020 10.29

Assalamualaikum pa Lutfi itu jurnal yg baru lagi ?

[Kutipan teks disembunyikan]

MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>
Kepada: priyawan rachmadi <priyawan-r@fkg.unair.ac.id>

27 Mei 2020 14.38

Yang dulu, legalisasinya

[Kutipan teks disembunyikan]



MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>

Revision of your papers

4 pesan

Agoes Soegianto <agoes_soegianto@fst.unair.ac.id>

21 Februari 2020 14.46

Kepada: m.luthfi@fkg.unair.ac.id

Dear Author,
Your papers have been selected to be published in Annals of Biology, However
Please revise your article according to the standard below:

As per NAAS, agency scoring journals for Impact factor, references older than 10 years have negative impact on quality of the journal. **Hence check and delete all old references up to 2011 if possible.**

Ensure that there are at least 4 to 5 latest references of 2017, 2018 and 2019 both in text and references. Please delete older ones when latest are available if possible.

Mention name, email and mobile of corresponding author.

Be sure that each reference is in text and vice versa

Article is written only in **past tense**, as it is written after conducting the experiment. **Never write in present tense.**

Instruction to authors enclosed for ready reference especially for references.
In each reference, name of authors, journal, volume and pages must be mentioned.
Please send it back to me in 48 hours.

Best Regards

Prof. Agoes Soegianto

Dept. Biology, Airlangga University
Kampus C, Jl. Mulyorejo, Surabaya 60115
INDONESIA

2 lampiran



AOB 16 - 1176-Expression of lymphocyte t (CD8+) cells in severe early childhood caries and free caries 2020.docx

855K



AOB 15-0001 Effects Centella Asiatica Extract.docx

5534K

MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>

21 Februari 2020 21.45

Kepada: Agoes Soegianto <agoes_soegianto@fst.unair.ac.id>

Nggeh Prof Agus. Matur nuwon 🙏🙏🙏

Pada tanggal Jum, 21 Feb 2020 10:47 AM, Agoes Soegianto <agoes_soegianto@fst.unair.ac.id> menulis:

Dear Author,

Your papers have been selected to be published in Annals of Biology, However
Please revise your article according to the standard below:

As per NAAS, agency scoring journals for Impact factor, references older than 10 years have negative impact on quality of the journal. **Hence check and delete all old references up to 2011 if possible.**

Ensure that there are at least 4 to 5 latest references of 2017, 2018 and 2019 both in text and references. Please delete older ones when latest are available if possible.

Mention name, email and mobile of corresponding author.

Be sure that each reference is in text and vice versa

Article is written only in **past tense**, as it is written after conducting the experiment. **Never write in present tense.**

Instruction to authors enclosed for ready reference especially for references.
In each reference, name of authors, journal, volume and pages must be mentioned.
Please send it back to me in 48 hours.

Best Regards

Prof. Agoes Soegianto

Dept. Biology, Airlangga University
Kampus C, Jl. Mulyorejo, Surabaya 60115
INDONESIA

Agoes Soegianto <agoes_soegianto@fst.unair.ac.id>
Kepada: MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>

25 Februari 2020 10.18

Dear Author,

Please add your correction in this attached files (C15, C16.docx, this files have been corrected by Publisher).

And send back it to me in 24-48 h.

Thank you in advance for your cooperation.

Best regard,

Prof. Agoes Soegianto

Dept. Biology, Airlangga University
Kampus C, Jl. Mulyorejo, Surabaya 60115
INDONESIA

On Fri, Feb 21, 2020 at 9:46 PM MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id> wrote:

Nggeh Prof Agus. Matur nuwon 🙏🙏🙏

Pada tanggal Jum, 21 Feb 2020 10:47 AM, Agoes Soegianto <agoes_soegianto@fst.unair.ac.id> menulis:

Dear Author,

Your papers have been selected to be published in Annals of Biology, However
Please revise your article according to the standard below:

As per NAAS, agency scoring journals for Impact factor, references older than 10 years have negative impact on quality of the journal. **Hence check and delete all old references up to 2011 if possible.**

Ensure that there are at least 4 to 5 latest references of 2017, 2018 and 2019 both in text and

references. Please delete older ones when latest are available if possible.

Mention name, email and mobile of corresponding author.

Be sure that each reference is in text and vice versa

Article is written only in **past tense**, as it is written after conducting the experiment. **Never write in present tense.**


Instruction to authors enclosed for ready reference especially for references.
In each reference, name of authors, journal, volume and pages must be mentioned.
Please send it back to me in 48 hours.


Best Regards

Prof. Agoes Soegianto

Dept. Biology, Airlangga University
Kampus C, Jl. Mulyorejo, Surabaya 60115
INDONESIA

2 lampiran

 **C 16.docx**
856K

 **C 15.docx**
5533K

MUHAMMAD LUTHFI <m.luthfi@fkg.unair.ac.id>
Kepada: Agoes Soegianto <agoes_soegianto@fst.unair.ac.id>

25 Februari 2020 22.08

Assalamualaikum warakhmatulohi wabarokatuh
berikut Prof kami kirim naskah yang telah kami revsi setelah pembenahan refferensinya sesuai permintaan editor.
t kasih

wassalamualaikum...
luthfi

Pada tanggal Jum, 21 Feb 2020 pukul 14.47 Agoes Soegianto <agoes_soegianto@fst.unair.ac.id> menulis:

Dear Author,
Your papers have been selected to be published in Annals of Biology, However
Please revise your article according to the standard below:

As per NAAS, agency scoring journals for Impact factor, references older than 10 years have negative impact on quality of the journal. **Hence check and delete all old references up to 2011 if possible.**

Ensure that there are at least 4 to 5 latest references of 2017, 2018 and 2019 both in text and references. Please delete older ones when latest are available if possible.

Mention name, email and mobile of corresponding author.

Be sure that each reference is in text and vice versa

Article is written only in **past tense**, as it is written after conducting the experiment. **Never write in**

present tense.


Instruction to authors enclosed for ready reference especially for references.
In each reference, name of authors, journal, volume and pages must be mentioned.
Please send it back to me in 48 hours.


Best Regards

Prof. Agoes Soegianto

Dept. Biology, Airlangga University
Kampus C, Jl. Mulyorejo, Surabaya 60115
INDONESIA

2 lampiran

 **C 16.docx**
858K

 **C 15.docx**
5535K

Expression Analysis of T Lymphocyte (CD8⁺) in Severe Early Childhood Caries

MUHAMMAD LUTHFI^{1*}, PRIYAWAN RACHMADI², AQSA SJUHADA OKI³
and AGUNG SOSIAWAN⁴

^{1,3}Department of Oral Biology, Faculty of Dental Medicine, Universitas Airlangga,
Surabaya, Indonesia

²Department of Dental Material, Faculty of Dental Medicine, Universitas Airlangga,
Surabaya, Indonesia

⁴Department of Public Health, Faculty of Dental Medicine, Universitas Airlangga,
Surabaya, Indonesia

*(e-Email : m.luthfi@fkg.unair.ac.id, Phone: +6281357898957)

ABSTRACT

In children younger than 3 years old, is considered as Severe Early Childhood Caries (S-ECC). This analytic observational study compared the level of CD8⁺ expression in saliva of children with S-ECC and free-caries children. Lymphocyte cells contained in saliva acquired by instructing the experimental subjects to gargle 10ml 1.5% NaCl solution. The expression of T lymphocyte (CD8⁺) were analyzed by means of flow cytometry. The expression of T lymphocyte (CD8⁺) in children with S-ECC (0.3400±0.14726) was higher than free-caries group (0.3250±0.11301) with P value of 0.655.

Key words : T lymphocyte (CD8⁺), S-ECC, adaptive immunity

INTRODUCTION

Early childhood caries (ECC) is a condition found ≥ 1 decay, loss (due to caries), or the presence of fillings in primary teeth in children aged 71 months or younger, whereas severe ECC (S-ECC) occurs in children children < 3 years with ≥ 1 rot, missing (due to caries), or filled tooth surfaces and in children aged 4-6 years with high caries scores (Colak et al., 2013). Severe dental caries that affect children younger than 3 years old is considered as Severe Early Childhood Caries (S-ECC) (Sukuraman and Pradeep, 2017). The effect of ECC in children is not limited in oral health only, but

also the general health (Naidu *et al.*, 2016). In terms of dental and oral health, ECC may give rise to pain, orthodontic problem, enamel damage, and also disturbance in oral function, such as mastication and speech. Besides, the development of permanent dentition in children with S-ECC may also be affected (Abanto *et al.*, 2016).

Cariogenic bacteria are one of the risk factor of ECC. The pathophysiology of ECC is correlated to the early colonization and the high level of cariogenic bacteria, *Streptococcus mutans*, and *Streptococcus sobrinus*, aside from the high level of sugar contained in dental plaque. Salivary protein, glucan, will breakdown the sugar, creating acidic environment in the oral cavity, thus, give rise to enamel and dentin demineralization (Jayabal and Mahesh, 2014).

The immune response is the body's ability to stay healthy by providing protection against harmful agents, most of which are microbes and is a specific and very specific response to pathogens. The innate response is the first line of defense in defending the body against pathogens in the same way at all times. These natural mechanisms include skin, saliva, tears, various cytokines, complement proteins, lysozyme, bacterial flora, and many cells including neutrophils, basophils, eosinophils, monocytes, macrophages, reticuloendothelial system, natural killer cells (NK cells), epithelial cells, endothelial cells, red blood cells, and platelets (Arce-Sillas *et al.*, 2016). Adaptive immune responses that are obtained adaptively will utilize the abilities of specific lymphocytes and their products (immunoglobulin, and cytokines) to produce responses to microbes that attack (Lawrence *et al.*, 2016).

During infection, the immune response should be able to eliminate pathogenic microbial invasion with minimal tissue damage. Both innate and adaptive immune response, including T cells play important role in eliminating pathogenic microbes by releasing pro-inflammatory cytokine and activating cytotoxic T lymphocyte (CTL). Besides, T helper (TH) and regulatory T cells also necessary in secreting antibody by plasma cells, along with the immunomodulatory cytokine such as interleukin. Recent studies found the new important function of TH, including T Follicle (Th17, Th22), in immunity toward infection, also in disease progression and prognosis. Based on the aforementioned background, this study was conducted to observe and compare the expression of T lymphocyte (CD8⁺) in children with S-ECC and free-caries children.

MATERIALS AND METHODS

This study was an analytic observational study, with cross-sectional analysis on two groups of sample; children with S-ECC and free caries children. All the procedures in this study had been reviewed and approved by the Health Research Ethical Clearance Commission of Universitas Airlangga, Faculty of Dental Medicine, with certificate no 209/HRECC. FODM/IX/2017. The samples, in this case, the parents had given their consent to participate in after had been given a complete information about the study. Saliva were collected from the subject aging from 4 to 6 years. The subjects were instructed to gargle 10 ml of 1.5% sterile NaCl solution for 30 seconds, and expectorate it into a sterile glass. This procedure was repeated four times. The saliva was then stored in cooler at 4 °C before freezing at or below -20 °C. Saliva was collected during 9 to 11 am.

The collected saliva was centrifuged at 450 g for 15 minutes at 40 °C. The pellet was then mixed into 2ml of Roswell Park Memorial Institute (RPMI) medium (TranGen biotech, Beijing, China) and vortexed (Luthfi *et al.*, 2019). The cell suspension was subsequently taken and counted using hemocytometer (JSQA hemocytometer, Hunan, China).

An equal number of volume from the cell suspension and 0.2% of trypan blue solution (PubChem, Bathesda, USA) staining were mixed in Eppendorf and vortexed. An aliquot of the same suspension (20 µl) were added into both chamber of the hemocytometer and observed under a light microscope (digital microscope XSZ-107BN, Hunan, China) with 10 x magnifications. The mixtures were subsequently transferred under the cover slip by capillary action to cover the area of the grid. The cells were counted in the area of 16 squares, which was equal to cell count $\times 10^4$ /ml. The viable cells in the squares were counted, and the cells count per ml were calculated using the following formula:

Cell/ml = Means of cell count per primary square $\times 10^4 \times$ diluting factors.

Lymphocyte cells (3×10^5 cells/ml) were cultured in the tissue culture flask (Greiner) 75 cm^2 with complete culture medium (RPMI-1640, 10% fetal calf serum (FCS), and 1% penicillin/streptomycin) in 5% CO_2 and atmosphere humidity 95% at 37 °C for 24 hours. The cultures were checked daily to observe the changes in color, turbidity, density, and growth pattern using inverted light microscope (Nikon).

The expressions of CD8⁺ were observed by means of flow cytometry method adapted from Luthfi *et al.*, 2019. Fluorescein isothiocyanate (FITC), phycoerythrin (PE), allophycocyanin (APC), Peridinin chlorophyll protein (PerCP), PerCP-Cy5.5-conjugated monoclonal antibodies (mAbs) from Becton Dickinson (San Jose, CA, USA). The optimal concentration of mAbs was determined for each mAb with titration. Flow cytometry simultaneously measured and analyzed for the physical properties of particles such as cells because it flew through the flow of fluid through a beam of light. The nature of scattering cell light was used to analyze changes in size, granularity, internal complexity and relative fluorescence intensity.

The stained lymphocytes were analyzed using flow cytometer (LSR 11 Sorvall RT7 Plus, Becton Dickinson, USA) with cell quest software (Becton Dickinson, USA). The results were analyzed using flow Jo 7.0 (USA) software. The expressions of CD8⁺ were analyzed using standard FACScan procedure with mAbs according to the producer protocol. The results were calculated and presented in mean. The acquired data was analyzed the normality and homogeneity, then followed by T-test to find the difference between two groups, with the level of significance at 0.05.

RESULTS AND DISCUSSION

Kolmogorov-Smirnov test showed p value of 0.200, and the Shapiro-Wilk test showed p value of 0.452, which meant that the data was normally distributed with homogeneity (Table 1). Therefore, the data was subsequently analyzed using independent t-test to find any difference between groups.

Table 1. Kolmogorov-smirnov and Shapiro-Wilk test result of T lymphocyte (CD8⁺) expression after 24 hours incubation and analyzed using flow cytometry

Variable	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
CD8 ⁺	,173	16	0.200	0.948	16	0.452

Table 2 showed that the expression of T lymphocyte (CD8⁺) of children with S-ECC was higher than the free-carries children. However, the independent t-test showed no significant difference of T lymphocyte (CD8⁺) between children with S-ECC and free caries children.

Table 2. Mean and standard deviation of T lymphocyte (CD8⁺) expression after 24 hours incubation and analyzed using flow cytometry

No	Group	N	CD8 ⁺ Expression	P-Value
			Mean ± SD	
1	S-ECC	8	0.3400 ± 0.14726	0.655
2	Free Caries	8	0.3250 ± 0.11301	




Figure 1. The expression of T lymphocyte (CD8⁺) (0.31%) in the saliva of children with S-ECC




Figure 2. The ekspresion of T lymphocyte (CD8⁺) (0.35%) in the saliva of children with S-ECC




Figure 3. The ekspresion of T lymphocyte (CD8⁺) (0.34%) in the saliva of free-carries children




Figure 4. The ekspresion of T lymphocyte (CD8⁺) (0.31%) in the saliva of free-carries children

CD8 + T cells are killer T cells (or cytotoxic T cells), are effector cells that function as cell-mediated immunity. CD8 + T cells are naive and must be activated in order to function effector cells (ie immune function). This activation occurs through interactions with pro-APC ("professional" antigen presenting cells), especially

dendritic cells in lymph nodes / follicles, and leads to intracellular pathways that regulate more TCR specific antigens on T cells and lead to effector functions. T cells can only recognize protein-based antigens (Lyu et. al., 2019).

S. mutans was the main bacteria that strongly correlated to ECC, while the other oral bacteria in dental biofilm were involved in caries initiation and development (Hajishengallis et al., 2017). Those bacteria were *Lactobacillus* sp. which play important role in the caries development (Li and Tanner, 2015). *Actinomyces* sp., specifically *Actinomyces gerenseriae* were also found in initial caries. *Bifido bacterium* are found in deep carious lesion, while several other *Streptococcus* non-mutans were acidogenic and aciduric bacteria, also playing role in caries. Epidemiology data showed that during the pathogenesis of dental caries, *Candida albicans* was also involved (Sukuraman and Pradeep, 2014). The previous study showed that the presence of *C. albicans* in children with ECC was significantly higher than those without caries. Besides, children with more population of *C. albicans* in oral cavity had a higher risk of ECC compared to those with low level of *C. albicans* in the oral cavity (Xiao et al., 2017).

Based on the result of this study, the expression of T lymphocyte (CD8⁺) was higher in children with S-ECC than in those without caries, however, there was no significant difference. This showed the importance of host defense against microbes, both in ECC and S-ECC. T cell (CD8⁺) denoted one of the cells which was able to identify and eliminate the infected cells, therefore, it was an important host defense component toward pathogenic agent. Recent studies found that in infectious diseases, it was not the main pathological agent that directly correlated to the aggressor, but to the abnormal immune response. Thus, initiate hypersensitive reaction, an excessive and uncontrolled immune response, which caused tissue damage. Dysregulation of the innate immune response causes excess production of proinflammatory cytokines such as IL-1 β and TNF α , or excessive response to the level of stimulation of proinflammatory cytokines that can trigger the release of endogenous stimuli, including damage associated molecular patterns (DAMP) which have an impact on the immune (de Jesus et al., 2015).

This explained the higher expression of T lymphocyte (CD8⁺) in children with S-ECC, compared to the free-caries group. Increase in lymphocyte proliferation and IFN- γ expression up to 6 hours incubation was used as the indicator of early detection marker of severe early childhood caries (luthfi et al., 2019)

The killing function of CD8 + T cells is mediated by two mechanisms. The first mechanism involves the use of Fas / Fas Ligand (FasL). Activated CD8 + T cells express FasL bound to Fas (CD95) which is a receptor found in many cell types, which leads to activation of caspases and apoptosis of target cells. The second mechanism is activating CD8 + T cells which can be used to kill antigens by releasing granzymes and perforins which are two compounds that have the ability to cut the cell wall and caspases that are active. CD8 + T cells that are activated also release IFN- γ which is a cytokine used in the macrophage activation process (Wu and Lyu, 2019; Nakiboneka et. al., 2019).

Among various cytokine that involved in defense toward bacteria, the pro-inflammatory cytokine, such as TNF- α , IL- β , and IL-6 were produced in the initial stage of infection, causing fever that inhibited bacterial multiplication. Cytokine also increased the expression of adhesion molecule (P-selectine and ICAM), that eased the migration of cells from the vessels to the infected area, and induced neutrophils and macrophages to secrete NO and destroyed bacteria. Other cytokine that was produced in the initial phase of infection disturbed the adaptive immune response, were the cytokine produced by macrophage, IL-12, that played role in differentiation of Th0 into Th1. While IL-4, produced by basophil, mastocyte, and macrophage, induced the differentiation of Th0 into Th2, that collaborated with B lymphocyte to produce antibody (Bacharier and Geha, 2000).

CONCLUSION

The level of T lymphocyte (CD8⁺) in children with S-ECC was higher than the free-carries children.

ACKNOWLEDGMENT

The authors are highly thankful to Prof. Muhaimin Rifa'i, PhD. Med. Sc for his help in conducting this research. The authors would like to thank Directorate of Research and Community Services of Directorate General of Research and Development Strengthening from Ministry of Research, Technology and Higher Education of the Republic of Indonesia for granting funds provided for this research.

REFERENCES

- Abanto, J. Carvalho, T. S., Mendes, F.M., Wanderley, M. T., Bonecker, M. and Raggio, D. P. (2011). Impact of oral diseases and disorders on oral health-related quality of life of preschool children. *Community Dentistry and Oral Epidemiology* 39 : 105–114
- Arce-Sillas A, Álvarez-Luquín DD, Tamaya-Domínguez B, Gomez-Fuentes S, Trejo-García A, Melo-Salas M, Cárdenas G, Rodríguez-Ramírez J, Adalid-Peralta L. (2016). Regulatory T Cells: Molecular Actions on Effector Cells in Immune Regulation. *J Immunol Res.* 2016:1720827.
- Colak H, Dülgergil CT, Dalli M, Hamidi MM. (2013). Early childhood caries update: A review of causes, diagnoses, and treatments. *J Nat Sci Biol Med*;4(1):29–38.
- de Jesus, A. A., Canna, S. W., Liu, Y. & Goldbach-Mansky, R. Molecular mechanisms in genetically defined autoinflammatory diseases: disorders of amplified danger signaling. *Annu. Rev. Immunol.* 33, 823–874 (2015)
- Hajishengallis, E., Parsaei, Y., Klein, M. I. and Koo, H. (2017). Advances in the microbial etiology and pathogenesis of early childhood caries. *Molecular Oral Microbiology* 32 : 24-34
- Jayabal, J. and Mahesh, R. (2014). Current State of Topical Antimicrobial Therapy in Management of Early Childhood Caries. *ISRN Dentistry* 762458. <https://doi.org/10.1155/2014/762458>
- Lawrence H, Mawdesley AE, Holland JP, Kirby JA, Deehan DJ, Tyson-Capper AJ. (2016). Targeting Toll-like receptor 4 prevents cobalt-mediated inflammation. *Oncotarget.* 7(7):7578-85
- Lyu F, Ozawa T, Hamana H, Kobayashi E, Muraguchi A, Kishi H. (. 2019). A novel and simple method to produce large amounts of recombinant soluble peptide/major histocompatibility complex monomers for analysis of antigen-specific human T cell receptors. *N Biotechnol.* 49:169-177

- Luthfi, M., Setijanto, D., Rahardjo, M. B., Indrawati, R., Rachmadi, P. and Ruth, M. S. (2019). Correlation between human neutrophil peptide 1–3 secretion and azurophilic granule (CD63) expression in early childhood caries. *Dent. Res. J.* 16 : 81-6
- Luthfi, M., Oki, A. S., Indrawati, R. and Latuamury, N. S. (2019). Analysis of Lymphocyte Cell Proliferation and IFN- γ Expression In Saliva of Severe Early Childhood Caries and Caries-Free in Surabaya. *Malaysian Journal of Medicine and Health Sciences* 15 (Supp 3) : 8-10
- Li, Y. and Tanner, A. E. (2015). Effect of antimicrobial intervention on oral microbiota associated with early childhood caries. *Pediatric Dentistry* 37 : 226-244
- Naidu, R., Nunn, J. and Donnelly, E. (2016). Oralhealth-related quality of life and early childhood caries among preschool children in Trinidad. *BMC Oral Health*, 16 : 128, DOI: 10.1186/s12903-016-0324-7
- Nakiboneka R, Mugaba S, Auma BO, Kintu C, Lindan C, Nanteza MB, Kaleebu P, Serwanga J. (2019). Interferon gamma (IFN- γ) negative CD4+ and CD8+ T-cells
- Sukuraman, A. and Pradeep, S. A. (2017). Early childhood caries: Prevalence, risk factors, and prevention. *Frontiers in Pediatrics* 5 : 157 doi: 10.3389/fped.2017.00157.
- Yuejin Liang, Xiaojun Chen, Jinling Chen and Fengliang Liu, (2018). Differentiation and Function of T Cell Subsets in Infectious Diseases. *Journal of Immunology Research* 2018 : 1-2
- Xiao, J., Huang, X., Alkhers, N., Alzamil, H., Alzoubi, S. and Wu, T. T. (2017). *Candida albicans* and early childhood caries: A systematic review and meta-analysis. *Caries Research* 52 : 102-112
- Wu B, Lyu FL. (2019). Progress of CD8+ T cell-mediated immune response to *Toxoplasma gondii* infection]. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi.* 32(2):143-7.
- Nakiboneka R, Mugaba S, Auma BO, Kintu C, Lindan C, Nanteza MB, Kaleebu P, Serwanga J. (2019). Interferon gamma (IFN- γ) negative CD4+ and CD8+ T-cells can produce immune mediators in response to viral antigens. *Vaccine.* Jan 03;37(1):113-122.