

The influence of lead (Pb), Zinc (Zn), Ratio Lead (Pb) to Zinc (Zn) in Attention deficit hyperactivity disorder (ADHD)

by Yunias Setiawati

Submission date: 01-Dec-2020 02:47PM (UTC+0800)

Submission ID: 1461175628

File name: to_Zinc_Zn_in_Attention_deficit_hyperactivity_disorder_ADHD.pdf (6.71M)

Word count: 3819

Character count: 20299

ISSN-0976-0245 (Print) • ISSN-0976-5506 (Electronic)

Volume 10 Number 8 August 2019



Indian Journal of Public Health Research & Development

An International Journal

SCOPUS IJPHRD CITATION SCORE

Indian Journal of Public Health Research and Development
Scopus coverage years: from 2010 to 2017 Publisher:
R.K. Sharma, Institute of Medico-Legal Publications
ISSN:0976-0245E-ISSN: 0976-5506 Subject area: Medicine:
Public Health, Environmental and Occupational Health
CiteScore 2015-0.02
SJR 2015-0.105
SNIP 2015-0.034



Website:

www.ijphrd.com

Indian Journal of Public Health Research & Development

EXECUTIVE EDITOR

Prof Vidya Surwade

Associate Professor, Dr Baba Saheb Ambedkar, Medical College & Hospital, Rohinee, Delhi

INTERNATIONAL EDITORIAL ADVISORY BOARD

- Dr. Abdul Rashid Khan B. Md Jagar Din**, (*Associate Professor*)
Department of Public Health Medicine, Penang Medical College, Penang, Malaysia
- Dr. V Kumar** (*Consulting Physician*)
Mount View Hospital, Las Vegas, USA
- Basheer A. Al-Sum**,
Botany and Microbiology Deptt., College of Science, King Saud University,
Riyadh, Saudi Arabia
- Dr. Ch Vijay Kumar** (*Associate Professor*)
Public Health and Community Medicine, University of Burami, Oman
- Dr. VMC Ramaswamy** (*Senior Lecturer*)
Department of Pathology, International Medical University, Bukit Jalil, Kuala Lumpur
- Kartavya J. Vyas** (*Clinical Researcher*)
Department of Deployment Health Research,
Naval Health Research Center, San Diego, CA (USA)
- Prof. PK Pokharel** (*Community Medicine*)
BP Koirala Institute of Health Sciences, Nepal

NATIONAL SCIENTIFIC COMMITTEE

- Dr. Anju Ade** (*Associate Professor*)
Navodaya Medical College, Raichur, Karnataka
- Dr. E. Venkata Rao** (*Associate Professor*) Community Medicine,
Institute of Medical Sciences & SUM Hospital, Bhubaneswar, Orissa.
- Dr. Amit K. Singh** (*Associate Professor*) Community Medicine,
VCSG Govt. Medical College, Srinagar – Garhwal, Uttarakhand
- Dr. R G Viveki** (*Professor & Head*) Community Medicine,
Belgaum Institute of Medical Sciences, Belgaum, Karnataka
- Dr. Santosh Kumar Mulage** (*Assistant Professor*)
Anatomy, Raichur Institute of Medical Sciences Raichur (RIMS), Karnataka
- Dr. Gourl Ku. Padhy** (*Associate Professor*) Community and Family
Medicine, All India Institute of Medical Sciences, Raipur
- Dr. Ritu Goyal** (*Associate Professor*)
Anaesthesia, Saraswati Institute of Medical Sciences, Panchsheel Nagar
- Dr. Anand Kalaskar** (*Associate Professor*)
Microbiology, Prathima Institute of Medical Sciences, AP
- Dr. Md. Amirul Hassan** (*Associate Professor*)
Community Medicine, Government Medical College, Ambedkar Nagar, UP
- Dr. N. Girish** (*Associate Professor*) Microbiology, VIMS&RC, Bangalore
- Dr. BR Hungund** (*Associate Professor*) Pathology, JNMC, Belgaum.
- Dr. Sartaj Ahmad** (*Assistant Professor*).
Medical Sociology, Department of Community Medicine, Swami Vivekananda Subharti
University, Meerut, Uttar Pradesh, India
- Dr Sumeeta Soni** (*Associate Professor*)
Microbiology Department, B. J. Medical College, Ahmedabad, Gujarat, India

NATIONAL EDITORIAL ADVISORY BOARD

- Prof. Sushanta Kumar Mishra** (Community Medicine)
GSL Medical College – Rajahmundry, Karnataka
- Prof. D.K. Srivastava** (*Medical Biochemistry*)
Jamia Hamdard Medical College, New Delhi
- Prof. M Sriharibabu** (*General Medicine*) GSL Medical College, Rajahmundry,
Andhra Pradesh
- Prof. Pankaj Datta** (*Principal & Prosthodontist*)
Indraprastha Dental College, Ghaziabad

NATIONAL EDITORIAL ADVISORY BOARD

- Prof. Samarendra Mahapatro** (*Pediatrician*)
HI-Tech Medical College, Bhubaneswar, Orissa
- Dr. Abhiruchi Galhotra** (*Additional Professor*) Community and Family
Medicine, All India Institute of Medical Sciences, Raipur
- Prof. Deepti Pruthvi** (*Pathologist*) SS Institute of Medical Sciences &
Research Center, Davangere, Karnataka
- Prof. G S Meena** (*Director Professor*)
Maulana Azad Medical College, New Delhi
- Prof. Pradeep Khanna** (*Community Medicine*)
Post Graduate Institute of Medical Sciences, Rohtak, Haryana
- Dr. Sunil Mehra** (*Paediatrician & Executive Director*)
MAMTA Health Institute of Mother & Child, New Delhi
- Dr Shailendra Handu**, *Associate Professor*, Phrma, DM (Pharma, PGI
Chandigarh)
- Dr. A.C. Dhariwal**: *Directorate of National Vector Borne Disease*
Control Programme, Dte. DGHS, Ministry of Health Services, Govt. of
India, Delhi

Print-ISSN: 0976-0245-Electronic-ISSN: 0976-5506, Frequency: Quarterly
(Four issues per volume)

Indian Journal of Public Health Research & Development is a double blind peer reviewed international journal. It deals with all aspects of Public Health including Community Medicine, Public Health, Epidemiology, Occupational Health, Environmental Hazards, Clinical Research, and Public Health Laws and covers all medical specialties concerned with research and development for the masses. The journal strongly encourages reports of research carried out within Indian continent and South East Asia.

The journal has been assigned International Standards Serial Number (ISSN) and is indexed with Index Copernicus (Poland). It is also brought to notice that the journal is being covered by many international databases. The journal is covered by EBSCO (USA), Embase, EMCare & Scopus database. The journal is now part of DST, CSIR, and UGC consortia.

Website : www.ijphrd.com

©All right reserved. The views and opinions expressed are of the authors and not of the Indian Journal of Public Health Research & Development. The journal does not guarantee directly or indirectly the quality or efficacy of any product or service featured in the advertisement in the journal, which are purely commercial.

Editor

Dr. R.K. Sharma
Institute of Medico-legal Publications
501, Manisha Building, 75-76, Nehru Place,
New Delhi-110019

Printed, published and owned by

Dr. R.K. Sharma
Institute of Medico-legal Publications
501, Manisha Building, 75-76, Nehru Place,
New Delhi-110019

Published at

Institute of Medico-legal Publications
501, Manisha Building, 75-76, Nehru Place,
New Delhi-110019

The Influence of Lead (Pb), Zinc (Zn), Ratio Lead (Pb) to Zinc (Zn) in Attention Deficit Hyperactivity Disorder (ADHD)

Yunias Setiawati¹, H J Mukono², Joni Wahyuhadi³, Endang Warsiki⁴

¹Doctoral Program of Health Science, Faculty of Public Health, Airlangga University; Indonesia;

²Department of Environmental Health, Faculty of Public Health, Airlangga University; Indonesia, ³Neurosurgery Department, Faculty of Medicine, Airlangga University; Indonesia, ⁴Psychiatric Department, Faculty of Medicine, Airlangga University; Indonesia

ABSTRACT

Background: Attention Deficit Hyperactivity Disorder (ADHD) is a common childhood psychiatric disorder often encountered in clinical practice with major symptoms of inattention, hyperactivity, and impulsivity. Pollution of lead (Pb) is thought to be cause of ADHD that affects the cognitive deficit of the brain. Zinc supplementation (Zn) can improve the symptoms of ADHD by increasing dopamine transporters binding. The aims of this study is to compare the level of Pb and Pb to Zn ratio on the subject of ADHD and normal children.

Method: This is an observational analytic study with case control design on 44 respondents. Statistical analysis using non-parametric test Man Whitney.

Result: There is no significant difference of Pb level in children with ADHD and normal children with $p = 0,431$ and there is a significant difference of Zn level in ADHD and normal children with $p = 0,011$ and significant difference in Pb to Zn ratio with $p = 0,015$.

Keyword: Lead (Pb), Zinc (Zn), Ratio Pb to Zinc (Zn), ADHD

BACKGROUND

Exposure to lead (Pb) in the environment has a neurotoxic effect resulting in behavioral disorders and cognitive deficits⁽¹⁾. For decades there has been considerable effort to reduce lead levels in the environment⁽²⁾. Lead may adversely affect cognitive disorders, decreased IQ scores, learning disabilities in mathematics, reading and verbal memory⁽³⁾.

Metaanalysis study by Goodlad et al concluded that the diagnosis of ADHD is increasing, thus attracting researchers to analyze the factors of heavy metals as factors causing ADHD⁽⁴⁾. Lead is the most responsible for mental disorders and thus requires clinical attention⁽⁵⁾. In humans, small amounts of metal can

adversely affect health and psychological disorders such as ADHD⁽⁶⁾. Pb adversely affects the child in the form of mental retardation disorder and impulsivity^(5,7-9). Pb exposure is more common in hair than blood. Pb levels in hair tend to be more sedentary and reflect Pb levels in the subjects compared with temporary blood⁽¹⁰⁻¹⁵⁾. While the trace elements Zinc (Zn), is an essential nutrient in animals and humans^(16,17). In ADHD children it is suspected that there is a decrease in Zinc levels that cause concentration and attention disorder^(11,18). The impact of high lead levels and low levels of Zinc can lead to impaired cognitive function and learning disabilities in children⁽¹⁹⁾. Measurement of Pb and Zinc levels in hair and Zinc to Pb ratio is expected to help reduce the risk of ADHD severity⁽¹¹⁾. Zinc plays an important role in the metabolism of neurotransmitters and prostaglandins, by maintaining the function and structure of the brain. Dopamine plays an important role in the pathophysiology of ADHD and the melatonin hormone plays an important role in the regulation of

Corresponding author:

Yunias

yunias.setiawati@gmail.com

dopamine, as of the administration of Zinc is expected to improve ADHD symptoms. Zinc levels of children with ADHD are thought to be lower than normal children. With the administration of Zinc supplementation, it will be able to correct the existing disorder^(17,20,21).

METHOD

This is a a case control study in grade 1 to grade 6 students of Bina Karya Elementary School Surabaya. With the use of random sampling formula, we get the number of samples of 23 ADHD subjects and 21 non ADHD subjects who meet the inclusion criteria. The inclusion criteria of the study were ADHD patients aged 6-12 years old, mothers of ADHD patients with minimum

junior high school education, able to communicate using Indonesian, cooperative, and did not experience severe psychosocial stressors.

Examination of lead and zinc levels were used hair samples by Atomic Absorption Spectrophotometry (AAS) method. Abbreviated Conners Teacher Rating Scale is used to assess ADHD scores. This scale was developed by C. Keith Conners, Ph.D. While the validity and reliability in the Indonesian has been done by Sasanti Juniar in 1999.

To assess the psychosocial stressors, it used Social Readjustment Rating Scale by Holmes and Rahe (1967).

RESULTS

Table 1 Mother and respondent demographic data

	ADHD Category				
	Normal		ADHD		
	Frequency	%	Frequency	%	
Gender	Girl	6	28,57%	7	30,43%
	Boy	15	71,43%	16	69,57%
Occupation	Not occupied	16	76,19%	15	65,22%
	Occupied	5	23,81%	8	34,78%
Income	<1.000.000 / month	11	52,38%	19	82,61%
	1.000.000-3.000.000 / month	6	28,57%	3	13,04%
	>3.000.000 / month	4	19,05%	1	4,35%
Husband and wife relationship	Poor	0	0,00%	3	13,04%
	Good	21	100,00%	20	86,96%
Relationship among family member	Poor	0	0,00%	5	21,74%
	Good	21	100,00%	18	78,26%
Maternal age	25-31 y.o	1	4,76%	2	8,70%
	32-38 y.o	9	42,86%	5	21,74%
	39-45 y.o	8	38,10%	11	47,83%
	46-52 y.o	2	9,52%	4	17,39%
	53-59 y.o	1	4,76%	1	4,35%
Children age	≤12 tahun	19	90,48%	22	95,65%
	> 12 tahun	2	9,52%	1	4,35%

The results in table 1 show 44 respondents consisting of 23 respondents ADHD and 21 respondents non ADHD. From the age of children found the most age is less than 12 years with a percentage of 46.3% in non-ADHD respondents and 53.7% of ADHD respondents. The order of the most children is the first child of 56.52%. The highest maternal education in ADHD and non ADHD respondents was others each was 11 (50%). Maternal age in respondents ADHD mostly in the range 39-45 years of 11 people (57.9%), while the mother of respondents non ADHD as many as 9 people (64.3%). Most mothers of ADHD and non-ADHD children do not work and with the most income is less than Rp 1000,000 per month. Husband and wife relationship is poor on the mother of ADHD respondents of 3 people (100%) and the relationship between family members is poor as many as 5 people.

Table 2. Statistical Analysis Results of Pb level, Zn level, and Pb to Zn ratio

	Median (Minimum – maximum)	p value
Pb on ADHD (n = 23)	10,74 (4,76-18,05)	0,431
Pb on non ADHD (n=21)	11,39 (4,78-24,08)	
Zn on ADHD (n = 23)	307,84 (82,75-511,96)	0,011
Zn on non ADHD (n=21)	177,55 (17,56-448,39)	
Pb/Zn on ADHD (n = 23)	0,03 (0,02-0,15)	0,015
Pb/Zn on non ADHD (n=21)	0,06 (0,02-0,22)	

significant $p < 0,05$

DISCUSSION

From the demographic data of the respondents, the result of the sample of ADHD, boys patients were 16 children and girls were 7 children. This result is consistent with previous findings that the prevalence of ADHD is greater in males in comparison between 3-4 boys versus 1 girl⁽²²⁻²⁴⁾. From the demographic characteristics of

mothers that consist of age, education, occupation, and income, it is found that the age of mothers in ADHD children is majority older than mothers of non-ADHD children. This result is inconsistent with previous studies which concluded that the maternal age of ADHD children was younger than that of non-ADHD children⁽²⁵⁾. This result can be obtained because the parents' knowledge of the problems is still low so they do not understand how to handle their children disorder.

The results of the research was the majority of mothers are other education, namely elementary and junior high school, this result obtained based on interviews that many mothers was married at a young age and can not continue to higher education because there is no cost and some others mothers experience pregnancy.

From the data about the occupation of the mother found that most mothers was unoccupied. In parent interviews it is found that most mothers should take care of their children at home because many father work outside as a driver, private employee and work as a handyman. This result is also in accordance with previous research that the mother of many ADHD children who have lost their job or can not work because they have to take care of their own ADHD child⁽²⁵⁾. The results of maternal earnings working on mothers of ADHD and non-ADHD children were the largest in less than a million rupiah but the majority stated the results were sufficient. This result is because most of the respondents are Javanese who put togetherness, *gotong royong* and *nrimo* culture and emphasize the existence of life as flowing water surrender to God and shame to express the shortcomings^(26,27).

Statistical analysis used Mann Whitney non parametric comparative test. The data were not normally distributed and there was no significant difference between Pb levels in ADHD and non ADHD children. This result is consistent with the theory that causes of ADHD this day is unclear. High levels of Pb may cause cognitive, emotional and behavioral disorders but Pb is not the determining factor that causes ADHD. This theory also corresponds to results in the field showing high Pb levels in non-ADHD subjects⁽²⁸⁾. The results showed that Pb levels in ADHD respondents were lower than those for non-ADHD respondents but this condition still requires clinical attention because Pb is toxic and a global health problem in the environment and may

interfere with biochemical processes in the brain⁽²⁹⁾. Lead exposure still requires more research because it is an important issue in the global health of the world, especially in developing countries^(3,30).

There was a significant difference in Zinc levels in ADHD children and non ADHD. From the above results it can be considered to administer zinc on ADHD subjects. This result is consistent with the literature which states that in ADHD subjects there is deficiency or excess of Zn level⁽³¹⁾. Zinc plays an important role in protein and DNA synthesis, wound healing, bone structure and improvement of the immune system. Zinc deficiency may involve impaired growth, loss of hair, diarrhea, immune system decline and dermatitis⁽¹¹⁾. Zinc deficiency also causes late growth process, late puberty, erectile dysfunction and hypogonadism in men. It is said that 5% of the world's population suffers from zinc deficiency⁽³¹⁾. Zinc is a trace element because the concentration in plasma is only about 12-16µM, in the body there are 2-4 grams of zinc with a majority in skeletal bone and muscle. The body does not store zinc and zinc intake is obtained from food. The recommended dosage for girls and boys ages 4 to 8 months is 5mg, 14 to 18 years of age in women by 9 mg and men by 11 mg⁽³¹⁾. The concentration of zinc persists constantly in hair, skin, heart, muscle, as in plasma, liver, bone and testes. Chronic zinc deficiency is more sedentary in hair than in plasma⁽³¹⁾. Zinc is an essential micronutrient of physiological metabolism. Zinc plays an important role in catalyzing enzyme activity, contributing to protein structure and gene expression regulation⁽²⁰⁾. Other studies suggest excess zinc levels may also cause ADHD.

There was a significant difference between the ratio of Pb to Zn in ADHD and non ADHD children. From the examination there are differences in the ratio of Pb and Zn where the high Pb level come with lower Zn level. Chronic lead exposure adversely affects physical growth, development of the nervous system, memory impairment, learning disorders, cognitive deficits, psychological and behavioral disorders⁽⁹⁾. Infants and children are at high risk of lead pollution, with levels below 10 µg/100 ml may cause growth disturbance⁽²⁹⁾. Much effort has been made to reduce lead exposure in infants and children. The chelation program is used as the ultimate management but the therapeutic process is risky, with 50% suffering the side effects of losing the minerals needs⁽¹³⁾. The chelation is also not useful in improving cognitive, behavioral and neurocognitive

disorders⁽³²⁾. Therapy on lead can increase the cytotoxicity and astrocyte activity. Lead exposure causes astrocyte damage. Lead can increase lysosomal density and nucleus inclusions, increase the area of reticulum of endoplasma, increase lactic dehydrogenase (LDH) damage, and decrease the levels of the sulfhydryl group that can be overcome by antioxidant administration. The lead deposit in astrocyte leads to the accumulation of protein in the endoplasmic reticulum, increased protein response, and breakage of cell cycles⁽²⁹⁾.

CONCLUSION

There was a significant difference between Zinc levels in ADHD and non ADHD subjects and Pb/Zn ratios in ADHD and non ADHD subjects.

Ethical Clearance: Taken from Health Research Ethics Committee Faculty of Public Health Airlangga University No. 657KEPK

Source of Funding: Self

Conflict of Interest: Nil

REFERENCES

1. Nicolescu R, Petcu C, Cordeanu A, Fabritius K, Schlumpf M, Winneke G, et al. Environmental exposure to lead, but not other neurotoxic metals, relates to core elements of ADHD in Romanian children: Performance and questionnaire data. *PLoS One*. 2010;11(4):476–83.
2. Senut M, Cingolani P, Sen A, Kruger A, Shaik A, Hirsch H, et al. Epigenetics of early-life lead exposure and effects on brain development. *PLoS One*. 2012;4:665–74.
3. Yousef S, Adem A, Zoubeidi T, Kosanovic M, Mabrouk AA, Eapen V. Attention Deficit Hyperactivity Disorder and Environmental Toxic Metal Exposure in the United Arab Emirates. *PLoS One*. 2011;57(6):457–60.
4. Children S. Research | Children's Health Environmental Lead Exposure and Attention Deficit / Hyperactivity Disorder Symptom Domains in a Community Sample of South Korean. *PLoS One*. 2015;123(3):271–6.
5. Goodlad JK, Marcus DK, Fulton JJ. Clinical Psychology Review Lead and Attention-Deficit / Hyperactivity Disorder (ADHD) symptoms:

- A meta-analysis. *Clin Psychol Rev* [Internet]. Elsevier Ltd; 2013;33(3):417–25. Available from: <http://dx.doi.org/10.1016/j.cpr.2013.01.009>
6. Landis Wayne. *Environmental Toxicology*. second edi. Maggie M, editor. Florida: CRC Press LLC; 1999. 177-188 p.
 7. Macedoni-luk M, Gosar D, Bjørklund G, Ora J, Kodri J. Levels of Metals in the Blood and Specific Porphyrins in the Urine in Children with Autism Spectrum Disorders. 2014;
 8. Viktorinova A, Ursinyova M, Trebaticka J, Uhnakova I. Changed Plasma Levels of Zinc and Copper to Zinc Ratio and Their Possible Associations with Parent- and Teacher-Rated Symptoms in Children with Attention-Deficit Hyperactivity Disorder. 2016;1–7.
 9. Marques RC, Bernardi JVE, Dórea JG, Fatima M De, Moreira R. Perinatal multiple exposure to neurotoxic (lead , methylmercury , ethylmercury , and aluminum) substances and neurodevelopment at six and 24 months of age. *Environ Pollut* [Internet]. Elsevier Ltd; 2014;187:130–5. Available from: <http://dx.doi.org/10.1016/j.envpol.2014.01.004>
 10. Lead I, Times A, Advisory L, News S. Lead in Hair - Concern Over its Use as an Indicator of Non-Occupational Exposure LEAD Action News Vol 3 no 3 Winter 1995 ISSN 1324-6011 Lead in Hair - Concern Over its Use as an Indicator of Non-Occupational Exposure. 2018;3(3):13–4.
 11. Shin D, Ph D, Kim E, Oh K, Ph D. The Relationship between Hair Zinc and Lead Levels and Clinical Features of Attention-Deficit Hyperactivity Disorder. 2014;25(1):28–36.
 12. Marques RC, Bernardi JVE, Dórea JG, De Fatima R Moreira M, Malm O. Perinatal multiple exposure to neurotoxic (lead, methylmercury, ethylmercury, and aluminum) substances and neurodevelopment at six and 24 months of age. *Environ Pollut*. Elsevier Ltd; 2014;187:130–5.
 13. El F, Mohamed B, Zaky EA, El-sayed AB, Elhossieny RM, Zahra SS, et al. Assessment of Hair Aluminum , Lead , and Mercury in a Sample of Autistic Egyptian Children : Environmental Risk Factors of Heavy Metals in Autism. 2015;2015.
 14. Heavy Metals and Trace Elements in Hair and Urine of a Sample of Arab Children with Autistic Spectrum Disorder. 2011;6(4):247–57.
 15. Chan AYW, Mak TWL. Use of hair analysis in the diagnosis of heavy metal poisoning : report of three. 2004;
 16. Froehlich TE, Loe IM, Gilman RC, Anixt JS, Chirdkiatgumchai V, Kuan L. Update on Environmental Risk Factors for Attention-Deficit/Hyperactivity Disorder. *Curr Psychiatr*. 2012;13(5):333–44.
 17. Villagomez A, Ramtekkar U. Iron, Magnesium, Vitamin D, and Zinc Deficiencies in Children Presenting with Symptoms of Attention-Deficit/Hyperactivity Disorder. *Children* [Internet]. 2014;1(3):261–79. Available from: <http://www.mdpi.com/2227-9067/1/3/261/>
 18. Shin D-W, Kim E-J, Oh K-S, Shin Y-C, Lim S-W. The Relationship between Hair Zinc and Lead Levels and Clinical Features of Attention-Deficit Hyperactivity Disorder. *소아청소년정신의학*. 2014;25(1):28–36.
 19. Tippairote T, Temviriyankul P, Benjapong W. Hair Zinc and Severity of Symptoms Are Increased in Children with Attention Deficit and Hyperactivity Disorder : a Hair Multi-Element Profile Study. 2017;
 20. Croatica AM. [The role of zinc in the treatment of hyperactivity disorder in children]. 2018;63(4):1–2.
 21. Elbaz F, Zahra S, Hanafy H. Magnesium , zinc and copper estimation in children with attention deficit hyperactivity disorder. *Egypt J Med Hum Genet* [Internet]. Ain Shams University; 2017;18(2):153–63. Available from: <http://dx.doi.org/10.1016/j.ejmhg.2016.04.009>
 22. Muñoz-Silva A, Lago-Urbano R, Sanchez-Garcia M, Carmona-Márquez J. Child/adolescent's ADHD and parenting stress: The mediating role of family impact and conduct problems. *Front Psychol*. 2017;8(DEC):1–12.
 23. Knouse LE, Knouse LE. *Cognitive-Behavioral Therapies for ADHD Cognitive-Behavioral Therapies for ADHD*. 2014;
 24. Andrade EM, Geha LM, Duran P, Suwvan R, Machado F, do Rosário MC. Quality of life in caregivers of ADHD children and diabetes patients. *Front Psychiatry*. 2016;7(JUL):7–12.

The influence of lead (Pb), Zinc (Zn), Ratio Lead (Pb) to Zinc (Zn) in Attention deficit hyperactivity disorder (ADHD)

ORIGINALITY REPORT

16%

SIMILARITY INDEX

14%

INTERNET SOURCES

10%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

1	www.e-sciencecentral.org Internet Source	2%
2	repository.unair.ac.id Internet Source	2%
3	bibliosjd.files.wordpress.com Internet Source	1%
4	worldwidescience.org Internet Source	1%
5	www.indianjournals.com Internet Source	1%
6	Balmer, Nina V., and Marcel Leist. "Epigenetics and Transcriptomics to Detect Adverse Drug Effects in Model Systems of Human Development", Basic & Clinical Pharmacology & Toxicology, 2014. Publication	1%
7	ehp.niehs.nih.gov Internet Source	1%

8	stacks.cdc.gov Internet Source	1%
9	zombiedoc.com Internet Source	1%
10	sinta3.ristekdikti.go.id Internet Source	1%
11	eprints.ners.unair.ac.id Internet Source	<1%
12	www.medicopublication.com Internet Source	<1%
13	dergipark.org.tr Internet Source	<1%
14	"Networking of Mutagens in Environmental Toxicology", Springer Science and Business Media LLC, 2019 Publication	<1%
15	my.psychologytoday.com Internet Source	<1%
16	Joanna Fiłon, Jolanta Ustymowicz-Farbiszewska, Elżbieta Krajewska-Kułak. "Analysis of lead, arsenic and calcium content in the hair of children with autism spectrum disorder", BMC Public Health, 2020 Publication	<1%

journals.plos.org

17

Internet Source

<1%

18

eprints.undip.ac.id

Internet Source

<1%

19

Tyler C. Smith, Isabel G. Jacobson, Besa Smith, Tomoko I. Hooper, Margaret A. K. Ryan, For The Millennium Cohort Study Tea. "The occupational role of women in military service: Validation of occupation and prevalence of exposures in the Millennium Cohort Study", International Journal of Environmental Health Research, 2007

Publication

<1%

20

www.omicsonline.org

Internet Source

<1%

21

Cabaña-Muñoz, María Eugenia, José María Parmigiani-Izquierdo, Luis Alberto Bravo-González, Hee-Moon Kyung, and José Joaquín Merino. "Increased Zn/Glutathione Levels and Higher Superoxide Dismutase-1 Activity as Biomarkers of Oxidative Stress in Women with Long-Term Dental Amalgam Fillings: Correlation between Mercury/Aluminium Levels (in Hair) and Antioxidant Systems in Plasma", PLoS ONE, 2015.

Publication

<1%

22

"IUNS. 21st International Congress of Nutrition. Buenos Aires, Argentina, October 15-20, 2017: Abstracts", *Annals of Nutrition and Metabolism*, 2017

Publication

<1%

23

Amelia Villagomez, Ujjwal Ramtekkar. "Iron, Magnesium, Vitamin D, and Zinc Deficiencies in Children Presenting with Symptoms of Attention-Deficit/Hyperactivity Disorder", *Children*, 2014

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

The influence of lead (Pb), Zinc (Zn), Ratio Lead (Pb) to Zinc (Zn) in Attention deficit hyperactivity disorder (ADHD)

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
